

# Knockout of phospholipase C $\epsilon$ attenuates *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine-induced bladder tumorigenesis

TAIMAO JIANG<sup>1,2\*</sup>, TAO LIU<sup>1\*</sup>, LIN LI<sup>3</sup>, ZHIJUN YANG<sup>4</sup>, YUNFENG BAI<sup>5</sup>, DONGYE LIU<sup>2</sup> and CHUIZE KONG<sup>1</sup>

<sup>1</sup>Department of Urology, The First Affiliated Hospital of China Medical University, Shenyang, Liaoning 110001;

<sup>2</sup>Department of Urology, The 463 Hospital of Chinese People's Liberation Army, Shenyang, Liaoning 110042;

<sup>3</sup>Department of Rehabilitation Medicine, Shengjing Hospital of China Medical University, Shenyang, Liaoning 110036;

<sup>4</sup>Department of Neurosurgery, Hospital of Beijing Military District of People's Liberation Army, Beijing 100070;

<sup>5</sup>Infection Section 15, The 302 Hospital of Chinese People's Liberation Army, Beijing 100039, P.R. China

Received December 9, 2014; Accepted September 1, 2015

DOI: 10.3892/mmr.2016.4762

**Abstract.** Bladder cancer frequently shows mutational activation of the oncogene Ras, which is associated with bladder carcinogenesis. However, the signaling pathway downstream of Ras remains to be fully elucidated. *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine (BBN) is able to induce bladder cancer by driving the clonal expansion of initiated cells carrying the activated form of Ras. Phospholipase C $\epsilon$  (PLC $\epsilon$ ) is the main target of BBN, while the tumor promoting role of PLC $\epsilon$  remains controversial. The present study examined the role of PLC $\epsilon$  in BBN-induced bladder carcinogenesis of mice with genetically inactivated PLC $\epsilon$ . Using light and electron microscopy, the present study demonstrated that PLC $\epsilon^{-/-}$  mice were resistant to BBN-induced bladder carcinogenesis. Furthermore, it was demonstrated that cyclooxygenase 2 and vascular endothelial growth factor-A were affected by the PLC $\epsilon$  background of the mice, suggesting that the role of PLC $\epsilon$  in tumor promotion may be ascribed to augmentation of inflammatory responses and angiogenesis. These results indicated that PLC $\epsilon$  is crucial for BBN-induced bladder carcinogenesis as well as signaling downstream of Ras, and that PLC $\epsilon$  is a candidate molecular target for the development of anti-cancer drugs.

## Introduction

Bladder cancer is one of the most common tumor types of the urinary system. However, its etiopathogenesis has remained to be fully elucidated. Study of the multifocal and polyclonal origins of bladder cancer as well as various experiments and epidemiological studies have shown that the persistence of carcinogenic substances in urine is an important cause of bladder cancer and also a major reason for bladder tumor recurrence after treatment (1-3).

The activation of numerous small guanine triphosphatases (GTPases) of the Ras superfamily is a crucial step in the regulation of a variety of cellular processes via complex cellular signaling networks. Phosphatidylinositol-specific phospholipase C (PLC) has emerged as one of the important signaling nodes in these complex networks, acting as a target as well as a regulator of small GTPases. Six major families of PLC (PLC $\beta$ , PLC $\gamma$ , PLC $\delta$ , PLC $\epsilon$ , PLC $\zeta$  and PLC $\eta$ ) are known (4), which are characterized by regulatory regions unique to each family. PLC $\epsilon$  is a recent addition to the growing list of Ras-associated effectors (5). Most suggested Ras effectors show either kinase or guanine nucleotide exchange factor activity. However, Ras has been implicated in regulating another class of enzyme, namely phosphoinositide-specific PLC, which is involved in the generation of well-characterized secondary messengers.

Studies have demonstrated that PLC $\epsilon$  is an effector of the Ras family of small GTPases, which bind directly to the Ras-associated domain of PLC $\epsilon$  (6-9). Further studies have revealed that PLC $\epsilon$  is also activated by small GTPase Ras homolog gene family, member A as well as heterotrimeric G proteins G $\alpha$ 12 and G $\beta$ 1 $\gamma$ 2 (9,10). Via these multiple regulatory mechanisms, PLC $\epsilon$  mediates signals originating from a large variety of cell surface receptors (11,12). In addition, PLC $\epsilon$  exerts a function as a guanine nucleotide exchange factor for Ras-related protein 1 (Rap1) via its CDC25 homology domain (13). Mice homozygous regarding the functionally inactivated PLC $\epsilon$  allele (PLC $\epsilon^{-/-}$  mice) exhibit semilunar valvulogenetic defects, which lead to cardiac dilation (14-16). Bai *et al* (17) demonstrated that PLC $\epsilon^{-/-}$  mice were resistant

---

*Correspondence to:* Dr Chuize Kong, Department of Urology, The First Affiliated Hospital of China Medical University, 155 North Nanjing Street, Shenyang, Liaoning 110001, P.R. China  
E-mail: chuizekong163@163.com

\*Contributed equally

**Key words:** phospholipase C $\epsilon$ , *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine, cyclooxygenase-2, vascular endothelial growth factor-A

Table I. Statistics of control mice and mouse models of bladder cancer.

Group	Type	Treatment	Normal n (%)	Atypical hyperplasia, n (%)	Transitional cell carcinoma, n (%)	Dead animals (n)	Surviving animals (n)
A	PLC $\epsilon^{+/+}$	Control, none	24 (100)	0 (0)	0 (0)	1	23
B	PLC $\epsilon^{+/+}$	BBN (0.1 %) <sup>a</sup>	8 (18.18)	8 (18.18)	28 (63.64)	4	44
C	PLC $\epsilon^{-/-}$	Control, none	24 (100)	0 (0)	0 (0)	0	24
D	PLC $\epsilon^{-/-}$	BBN (0.1 %) <sup>a</sup>	12 (26.09)	14 (30.43)	20 (43.48)	2	46

<sup>a</sup>Administered via drinking water supplementation. BBN, *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine; PLC, phospholipase C.

to 7,12-dimethylbenz(a)anthracene-induced skin tumor formation with 12-*O*-tetradecanoylphorbol-13-acetate (TPA) as a tumor promoter. In addition, PLC $\epsilon^{-/-}$  mice were void of basal layer cell proliferation and epidermal hyperplasia, suggesting the role of PLC $\epsilon$  in tumor promotion.

The Ras gene is one of the most common genes associated with bladder cancer. It has been confirmed that PLC $\epsilon$  is an effector of Ras and is in turn regulated by Ras in a GTP-dependent manner. The association between bladder transitional cell carcinoma and PLC $\epsilon$  has not been fully elucidated. The present study hypothesized that PLC $\epsilon$  has a significant role in the development of bladder transitional cell carcinoma. In order to verify this hypothesis, the effects of PLC $\epsilon$  knockdown on *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine (BBN)-mediated induction of bladder cancer were investigated in mice. BBN is able to induce bladder cancer by targeting PLC $\epsilon$  and activating Ras to drive the clonal expansion of initiated cells (18). The present study suggested that PLC $\epsilon$  has a crucial role in the development of cancer downstream of Ras signaling.

## Materials and methods

**Animals.** PLC $\epsilon^{-/-}$  mice were obtained from the Laboratory of Experimental Molecular Biology at Kobe University (Kobe, Japan) and maintained at the 302 Hospital of the Chinese People's Liberation Army (Beijing, China). PLC $\epsilon^{-/-}$  homozygous mice and wild-type (PLC $\epsilon^{+/+}$ ) littermates were obtained by crossing C57BL/6J PLC $\epsilon^{+/+}$  and 129S4 PLC $\epsilon^{-/-}$  mice. Mice carrying inactivated PLC allele (PLC $\epsilon^{-}$ ) were generated using in-frame deletion of an exon encoding the catalytic X domain, as described previously (14). All animals were housed under standard conditions (22±1°C; 12-h light/dark cycle; 50-55% humidity) with free access to food pellets and tap water. Experiments were performed on 6-8 week-old male mice, and a total of 72 PLC $\epsilon^{+/+}$  and 72 PLC $\epsilon^{-/-}$  mice were used in the present study. The mean weight of the animals was 20.3±0.1 g for the PLC $\epsilon^{+/+}$  mice and 20.3±0.15 g for the PLC $\epsilon^{-/-}$  mice. Disparate groups of mice were used in each experiment. Animal experiments were approved by the Animal Care and Use Committee of The 463 hospital of Chinese People's Liberation Army (Shenyang, China).

**Materials.** Antibodies used included anti-mouse PLC $\epsilon$  antibody (Dako Cytomation, Copenhagen, Denmark), rabbit anti-mouse polyclonal anti-vascular endothelial

growth factor-A (VEGF-A) antibody (cat. no. ab51745; Abcam, Cambridge, MA, USA), rabbit anti-mouse polyclonal anti-GAPDH antibody (cat. no. sc-25778; Santa Cruz Biotechnologies, Dallas, TX, USA), rabbit anti-mouse anti-cyclooxygenase-2 (COX-2) antibody (cat. no. 160126; Cayman, Ann Arbor, MI, USA) and goat anti-rabbit immunoglobulin G-horse radish peroxidase conjugated secondary antibodies (Santa Cruz Biotechnologies).

**BBN-induced mouse model of bladder cancer.** A BBN-induced mouse model of bladder cancer was prepared according to the method described by Vecchione *et al* (19) with certain modifications. Briefly, six-week-old male PLC $\epsilon^{-/-}$  mice (n=72) and PLC $\epsilon^{+/+}$  mice (n=72) were sub-divided into BBN treatment groups (n=48) and control groups (n=24) without treatment of BBN (Table I). BBN-treated mice were given tap water containing 0.1% BBN for 12 weeks. Thereafter, they had access to tap water without BBN. Control mice were given water without BBN throughout the experiment. Mice were sacrificed at 8, 12 and 18 weeks after the cessation of BBN treatment. Bladder specimens were harvested and analyzed for pathology [hematoxylin-eosin (HE; Beyotime Institute of Biotechnology, Shanghai, China) staining and ultrastructural assessment] and protein (western blot and immunofluorescence).

**Pathological analysis.** The mice were anesthetized with sodium pentobarbitone (40 mg/kg, i.p.; Sigma-Aldrich, St. Louis, MO, USA) and then transcardially perfused with 10 ml 0.9% saline, followed by 30 ml 0.1 M phosphate buffer (PB; pH 7.4) containing 4% paraformaldehyde for 5 min. At necropsy, urinary bladders were removed and placed in 4% paraformaldehyde for HE staining immunofluorescence or in 2.5% glutaraldehyde (Sigma-Aldrich) for ultrastructural study. For HE staining, each bladder was dissected, processed for routine paraffin embedding, cut into 5  $\mu$ m sections and mounted onto polylysine-coated slides. Sections were de-waxed in xylene (Sinopharm, Shanghai, China), re-hydrated in a descending series of ethanols and processed for routine HE. For ultrastructural study, sections were washed with 0.01 M phosphate-buffered saline (PBS), post-fixed with 10 g/l OsO<sub>4</sub> in PB for 45 min, de-hydrated through a graded ethanol series and propylene oxide, and flat-embedded with Epon-812 (Nanjing Tansi Technology Co., Ltd., Nanjing, China). The sections were examined under a light microscope (AE31; Motic, Xiamen, China) and regions containing all layers of the bladder were investigated under an electron




Figure 1. Bladder anatomy of mice in the experimental groups. Representative images of (A) mice in Group B at week eight, (B) mice in Group D at week eight, (C) mice in Group B at week 12 (the arrow indicates hematuria) and (D) mice in Group D at week 12. (E) Bladder tumors of Group A (left) and B (right) at week 12. (F) Bladder tumors of Group C (left) and D (right) at week 12. Groups: A, untreated PLC $\epsilon^{+/+}$  control mice; B, PLC $\epsilon^{+/+}$  mice induced with 0.1% BBN; C, untreated PLC $\epsilon^{-/-}$  mice; D, PLC $\epsilon^{-/-}$  mice induced with 0.1% BBN. BBN, *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine; PLC, phospholipase C.

microscope. Tissue samples from the selected regions were cut into sections on an ultramicrotome (EM UC7; Leica, Wetzlar, Germany) and prepared for electron microscopic analysis (H-7100; Hitachi, Tokyo, Japan).

**Western blot analysis.** Protein was extracted from each sample using radioimmunoprecipitation assay lysis buffer (Beyotime Institute of Biotechnology). The concentrations of proteins were detected using the bicinchoninic acid assay (Beyotime Institute of Biotechnology). Protein samples were separated by 10% SDS-PAGE and then transferred onto nitrocellulose membranes. Then membranes were blocked with 5% non-fat milk followed by incubation with antibodies against COX2 (1:500), VEGF-A (1:5,000) and GAPDH (1:1,000), respectively, at 37°C for another 1.5 h. Membranes were subsequently washed with Tris-buffered saline containing Tween 20 (TBST) and incubated with the corresponding secondary antibodies conjugated with horseradish peroxidase (1:5,000) for 45 min at 37°C. Following washing with TBST, a RapidStep™ ECL reagent (Millipore, Bedford, MA, USA) and x-ray film were used to capture images of the blots.

**Immunofluorescence.** Detection of PLC $\epsilon$ , COX2 and VEGF-A was performed on consecutive sections obtained from

paraffin-embedded tissues using an immunofluorescence double labeling method. Sections were dried at room temperature, de-paraffinized, re-hydrated and then treated with 2% hydrogen peroxide at 37°C for 5 min. Antigen retrieval was performed by pepsin (Biosharp, Hefei, China) treatment at 40°C for 10 min. After blocking with 5% bovine serum albumin, the sections were incubated with anti-PLC $\epsilon$  and COX2 (1:200) or VEGF-A antibody (1:500) overnight at 4°C. Sections were washed with 0.01 M PBS and incubated with fluorescein isothiocyanate-labeled anti-goat immunoglobulin (Ig)G antibody (1:500; Sigma-Aldrich) for PLC $\epsilon$  and Texas Red-labeled anti-rabbit IgG antibody (1:1,000; Molecular Probes, Breda, Netherlands) for COX2 and VEGF-A at 37°C for 2 h. The sections were thoroughly rinsed in 0.01 M PBS between and after the incubation steps. The sections were examined using confocal laser-scanning microscopy (FV1000; Olympus, Tokyo, Japan). Digital images were captured using EZ-C1 3.50 software (Nikon).

**Statistical analysis.** All statistical analyses were performed using SPSS 10.0 software (SPSS, Inc., Chicago, IL, USA). The Kaplan-Meier test was used to analyze survival data. The incidence and multiplicity of urinary bladder cancer were analyzed by the Fisher's exact and Wilcoxon's rank-sum tests,




Figure 2. Representative histological HE staining images and electron micrographs of bladder mucosa of the mice in the experimental groups at 12 weeks. HE staining of normal bladder mucosa in (A) group A and (B) group C. (C) In group B, HE staining revealed significantly heterogeneous bladder mucosa. (D) HE staining of bladder mucosa in group D, revealing thickened, carcinomatous bladder mucosa (magnification,  $\times 200$ ). Ultrastructures of bladder mucosa of mice in (E) group A and (F) group C. (G) Ultrastructure of transitional epithelial membrane cells of mice of group B. The chromatin was distributed across the nuclear membrane, nucleolar hypertrophy was observed and fiber structure around the nuclei could not be distinguished. (H) Ultrastructure of bladder mucosa of mice in group D, revealing significantly heterogeneous transitional cells in the mucosa. Scale bar, 5000 nm. Groups: A, untreated PLC $\epsilon^{+/+}$  control mice; B, PLC $\epsilon^{+/+}$  mice induced with 0.1% BBN; C, untreated PLC $\epsilon^{-/-}$  mice; D, PLC $\epsilon^{-/-}$  mice induced with 0.1% BBN. BBN, *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine; PLC, phospholipase C; HE, hematoxylin and eosin.

respectively. Each experiment was repeated three times and representative results are presented.

## Results

**General observations.** In the present study, one hundred and forty-four mice were used, out of which seven died during the experiment (5%; Group A, 1; group B, 4; group D, 2). These mice showed no significant difference from others in weight and activity levels prior to their death. A higher susceptibility of PLC $\epsilon^{+/+}$  mice to BBN-induced tumor development was observed. The tumors of PLC $\epsilon^{-/-}$  mice were smaller than those of PLC $\epsilon^{+/+}$  mice at all time-points examined (Fig. 1A-D). In Group B, eight mice (18.18%) had normal urinary bladder mucous membranes, eight mice (18.18%) showed atypical hyperplasia and 63.6% (28/44) presented with neoplastic lesions (with *in situ* or invasive carcinoma), and in group D, these numbers were 26.09% (12/46), 30.43% (14/46) and 43.5% (20/46), respectively (Table I). Pre-neoplastic lesions in the urothelium adjacent to advanced tumors were frequently observed. By contrast, in the control groups A and C, atypical hyperplasia and neoplastic lesions were not observed. The size of the tumors in group B was significantly larger than that in group D (Fig. 1E and F). No significant differences were observed between groups A and C.

**Pathological changes.** In the present study, pathological changes in the development of BBN-induced bladder carcinoma in PLC $\epsilon^{-/-}$  and PLC $\epsilon^{+/+}$  mice were monitored. The following pathological changes were observed by light microscopy: Smooth mucosa of bladder walls were present in groups A and C without the presence of ulcers, congestion or neoplasms (Fig. 2A and B). The morphology of the mucosa of the bladder walls in groups B and D changed gradually with increasing time of BBN intake. In general, tumor formation




Figure 3. Western blot analysis for COX-2 and VEGF-A protein expression in bladder tumors from experimental mice at week 18 treated with *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine. GAPDH was used as a reference. A representative of three blots is shown. COX, cyclooxygenase; VEGF, vascular endothelial growth factor; PLC, phospholipase C.

in PLC $\epsilon^{-/-}$  mice was delayed and its incidence was reduced compared with that in PLC $\epsilon^{+/+}$  mice. Electron microscopy further confirmed the morphological findings in the experimental groups: While no differences in morphology of the bladder transitional epithelium were observed between groups A and C (Fig. 2E and F), the following pathological changes were visible in group B at week 12: Chromatin was distributed across the nuclear membrane, nucleolar hypertrophy was present and fiber structure around the nuclei was disordered (Fig. 2G). However, heterogeneous transitional cells were observed in the mucosa of mice in group D at week 12 (Fig. 2H). All of the above results demonstrated that knockout of PLC $\epsilon$  attenuated BBN-induced bladder tumorigenesis.

**Role of PLC $\epsilon$  in BBN-induced expression of inflammatory and angiogenesis-associated molecules.** In order to investigate the underlying mechanisms of the involvement of PLC $\epsilon$  in tumor development, the present study examined




Figure 4. Representative immunofluorescence images. (A) Expression of VEGF-A in the bladder mucosa of mice in Groups A, B, C and D at week 18. Green fluorescence, PLC $\epsilon$ ; red fluorescence, VEGF-A. (B) Expression of COX-2 in the bladder mucosa of mice in Groups A, B, C and D at week 18. Green fluorescence, PLC $\epsilon$ ; red fluorescence, COX-2. Scale bar, 100  $\mu$ m. Groups: A, untreated PLC $\epsilon^{+/+}$  control mice; B, PLC $\epsilon^{+/+}$  mice induced with 0.1% BBN; C, untreated PLC $\epsilon^{-/-}$  mice; D, PLC $\epsilon^{-/-}$  mice induced with 0.1% BBN. BBN, *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine; PLC, phospholipase; C COX, cyclooxygenase; VEGF, vascular endothelial growth factor.

the expression of representative inflammatory and angiogenesis-associated proteins in tumors from the bladders of mice in groups B and D. Previous studies have revealed that PLC $\epsilon$  is important in TPA-induced skin inflammation and tumor promotion (17,20). Inflammation is known to exert its highest effects on tumorigenesis in the later stage, in which tumors progress to high-grade adenomas (21-23). Therefore, the present study investigated the role of PLC $\epsilon$  in inflammation associated with late-stage bladder tumorigenesis. In grade-matched tumors from PLC $\epsilon^{+/+}$  and PLC $\epsilon^{-/-}$  mice at week 18, the expression of COX-2 and VEGF-A was assessed; these proteins were selected to be analyzed as representative signaling molecules due to their intrinsic functions in inflammation and angiogenesis, respectively, and because they have been implicated in bladder tumorigenesis (24). Western blot analysis revealed that the expression levels of COX-2 and VEGF-A were influenced by the PLC $\epsilon$  background of the mice, with upregulation in the BBN-treated PLC $\epsilon^{+/+}$  mice, but not in BBN-treated PLC $\epsilon^{-/-}$  mice (Fig. 3). In addition, the expression of COX-2 and VEGF-A was detected by immunofluorescence analysis. Groups A and B showed stronger green fluorescence (PLC $\epsilon$ ) than groups C and D, which was consistent with their genetic backgrounds. Consistent with the western blot results, marked increased in the VEGF-A signals in tumor epithelial cells of high-grade adenomas in PLC $\epsilon^{+/+}$  mice were observed, which were less apparent in PLC $\epsilon^{-/-}$  mice (Fig. 4A). The expression patterns of COX-2 were similar to those of VEGF-A, with increased expression in groups B and D (Fig. 4B). However, a locally elevated level of COX-2 was also identified. These results demonstrated that the protective role of PLC $\epsilon$  knockout against

bladder carcinogenesis may be associated with inflammatory responses and angiogenesis.

## Discussion

Numerous chemicals are known to evoke bladder carcinoma, among which nitroso compounds, including BBN and *N*-methyl-*N*-nitrosourea (MNU), and nitrofurans, such as *N*-[4-(5-nitro-2-furyl)-2-thiazolyl]-formamide (FANFT) are most potent (25,26). BBN and FANFT are indirect carcinogens through oral intake, while MNU is a carcinogen requiring direct bladder instillation. Due to its high potency to induce bladder cancer, BBN is the most suitable reagent to generate *in vivo* models of bladder cancer and to study bladder carcinogenesis. The carcinogenicity of BBN is limited to the bladders of rats, mice and dogs (27). No marked difference between the bladder cancer was observed between humans and mice, and rats and dogs. Therefore, BBN-induced bladder cancer is similar to transitional cell carcinoma in patients in both kinetic and histological features (28). PLC $\epsilon$  is important in the development and progression of human cancer types (29). The present study used BBN to induce bladder cancer, and knockout of PLC $\epsilon$  attenuated BBN-induced tumorigenesis of bladder cancer. This indicated that PLC $\epsilon$  is an oncogene and may be a therapeutic target for the treatment and prevention of bladder cancer. The downstream metabolite of BBN, *N*-butyl-*N*-(3-carboxy-propyl) nitrosamine (BCPN), is subjected to urinary excretion and comes in direct contact with the urinary tract, resulting in epithelial cell DNA damage and carcinogenesis (30). Compared with a high, single dose of BBN, smaller and multiple doses of BBN induce a higher rate

of bladder cancer with larger tumor size, poorer differentiation and a higher rate of infiltration.

Bladder carcinoma is the most common tumor type of the urinary system. Its occurrence and development involve numerous genes and processes, as well as congenital and acquired factors (31-33). Ras was the first oncogene identified in human bladder carcinoma and has been proved to be highly relevant to its development (34-36). PLC $\epsilon$ , as a downstream effector protein of Ras, may have an important role in the occurrence and development of bladder carcinoma.

PLC $\epsilon$  was first identified in *Caenorhabditis elegans* by Shibatohe *et al* (37) in 1998 as a novel sub-type of the PLC family. PLC $\epsilon$  has been reported to act as an effector protein for the products of the oncogene Ras and the tumor suppressor gene Rap (5,7,8). In recent years, the role of PLC $\epsilon$  in tumors has received increasing attention. PLC $\epsilon^{-/-}$  mice were successfully established in Kataoka's laboratory at Kobe University (Kobe, Japan) by Bai *et al* (17). In these PLC $\epsilon^{-/-}$  mice, the tumor incidence was significantly decreased and the progression of chemically induced skin tumors was inhibited (17), suggesting that PLC $\epsilon$  has an important role in tumor development. Consistent with the hypothesis of the present study, Bourguignon *et al* (38) found that PLC $\epsilon$  is involved in human head and neck squamous cell carcinoma (38); furthermore, Cheng *et al* (39) and Ling *et al* (40) reported that small hairpin RNA-mediated knockdown of PLC $\epsilon$  inhibited bladder cancer cell proliferation and cell cycle progression *in vitro*. PLC $\epsilon$  was also demonstrated to be associated with invasion and migration of bladder cancer (41).

The present study observed almost complete squamous commitment of the neoplastic lesions. After 18 weeks of BBN treatment, 43.48% of PLC $\epsilon$ -knockout mice developed invasive bladder cancer, which was significantly lower than the incidence observed in wild-type mice. This result is consistent with the notion that wild-type PLC $\epsilon$  mice develop bladder tumors more rapidly and frequently after chronic intake of BBN than PLC $\epsilon^{-/-}$  mice, further suggesting that the presence of PLC $\epsilon$  sensitizes tissues to carcinogenesis. The present study also observed that the incidence of atypical hyperplasia in PLC $\epsilon^{-/-}$  mice was significantly higher than that observed in wild-type mice. This indicates that wild-type mice are pre-disposed to developing bladder cancer. Differences in the incidence of bladder tumors between these groups of mice indicated that urothelial cells require two intact PLC $\epsilon$  alleles in order to respond efficiently to the damaging action of chemical carcinogens.

PLC $\epsilon$  has been demonstrated to be associated with inflammatory responses (42). Ikuta *et al* (20) have shown that two TPA targets, Ras guanyl-releasing protein 3 and protein kinase C, are involved in TPA-induced inflammation through the activation of PLC $\epsilon$ , leading to tumor promotion. Li *et al* (43) suggested that PLC $\epsilon$  has crucial roles in intestinal tumorigenesis through two distinct mechanisms - augmentation of angiogenesis and inflammation. In order to determine whether PLC $\epsilon$  has a role in bladder tumorigenesis through augmentation of angiogenesis and inflammation in BBN-induced bladder cancer, the present study also examined inflammatory and angiogenesis-associated factors COX-2 and VEGF-A, respectively. The results showed that VEGF-A was upregulated by BBN treatment in the tumors of PLC $\epsilon^{+/+}$  mice, while no marked changes in PLC $\epsilon^{-/-}$  mice were observed. COX-2

showed a similar pattern to that of VEGF-A, however, with local elevation. These results suggested that bladder cancer is induced by BBN through two distinct mechanisms, augmentation of angiogenesis and inflammation, and that PLC $\epsilon$  has a pivotal role in tumorigenesis of bladder cancer induced by BBN.

In conclusion, the present study revealed that PLC $\epsilon$  has a crucial role in BBN-induced carcinogenesis of bladder epithelial cells, as knockout of PLC $\epsilon$  attenuated bladder carcinogenesis induced by BBN. The present study also provided solid *in vivo* evidence for the importance of PLC $\epsilon$  signaling in carcinogenesis. These results indicated that specific inhibitors of PLC $\epsilon$  may be useful for the treatment and prevention of certain types of cancer.

### Acknowledgements

The present study was supported by grants from the National Natural Science Foundation of China (nos. 81101938 and 81271316) and the Science and Technology Development Project of Liaoning Province (no. 2012225020).

### References

1. Beland FA, Beranek DT, Dooley KL, Heflich RH and Kadlubar FF: Arylamine-DNA adducts in vitro and in vivo: Their role in bacterial mutagenesis and urinary bladder carcinogenesis. *Environ Health Perspect* 49: 125-134, 1983.
2. Akaza H, Murphy WM and Soloway MS: Bladder cancer induced by noncarcinogenic substances. *J Urol* 131: 152-155, 1984.
3. Cohen SM: Role of urinary physiology and chemistry in bladder carcinogenesis. *Food Chem Toxicol* 33: 715-730, 1995.
4. Katan M: New insights into the families of PLC enzymes: Looking back and going forward. *Biochem J* 391: e7-e9, 2005.
5. Kelley GG, Reks SE, Ondrako JM and Smrcka AV: Phospholipase C(epsilon): A novel Ras effector. *EMBO J* 20: 743-754, 2001.
6. Mitin N, Rossman KL and Der CJ: Signaling interplay in Ras superfamily function. *Curr Biol* 15: R563-R574, 2005.
7. Lopez I, Mak EC, Ding J, Hamm HE and Lomasney JW: A novel bifunctional phospholipase c that is regulated by Galpha 12 and stimulates the Ras/mitogen-activated protein kinase pathway. *J Biol Chem* 276: 2758-2765, 2001.
8. Song C, Hu CD, Masago M, Kariyai K, Yamawaki-Kataoka Y, Shibatohe M, Wu D, Satoh T and Kataoka T: Regulation of a novel human phospholipase C, PLCepsilon, through membrane targeting by Ras. *J Biol Chem* 276: 2752-2757, 2001.
9. Bunney TD and Katan M: Phospholipase C epsilon: Linking second messengers and small GTPases. *Trends Cell Biol* 16: 640-648, 2006.
10. Smrcka AV, Brown JH and Holz GG: Role of phospholipase C $\epsilon$  in physiological phosphoinositide signaling networks. *Cell Signal* 24: 1333-1343, 2012.
11. Edamatsu H, Satoh T and Kataoka T: Ras and Rap1 activation of PLCepsilon lipase activity. *Methods Enzymol* 407: 99-107, 2006.
12. Kelley GG, Kaproth-Joslin KA, Reks SE, Smrcka AV and Wojcikiewicz RJ: G-protein-coupled receptor agonists activate endogenous phospholipase Cepsilon and phospholipase Cbeta3 in a temporally distinct manner. *J Biol Chem* 281: 2639-2648, 2006.
13. Satoh T, Edamatsu H and Kataoka T: Phospholipase Cepsilon guanine nucleotide exchange factor activity and activation of Rap1. *Methods Enzymol* 407: 281-290, 2006.
14. Tadano M, Edamatsu H, Minamisawa S, Yokoyama U, Ishikawa Y, Suzuki N, Saito H, Wu D, Masago-Toda M, Yamawaki-Kataoka Y, *et al*: Congenital semilunar valvulogenesis defect in mice deficient in phospholipase C epsilon. *Mol Cell Biol* 25: 2191-2199, 2005.
15. Wang H, Oestreich EA, Maekawa N, Bullard TA, Vikstrom KL, Dirksen RT, Kelley GG, Blaxall BC and Smrcka AV: Phospholipase C epsilon modulates beta-adrenergic receptor-dependent cardiac contraction and inhibits cardiac hypertrophy. *Circ Res* 97: 1305-1313, 2005.

16. Kariya K, Bui YK, Gao X, Sternberg PW and Kataoka T: Phospholipase Cepsilon regulates ovulation in *Caenorhabditis elegans*. *Dev Biol* 274: 201-210, 2004.
17. Bai Y, Edamatsu H, Maeda S, Saito H, Suzuki N, Satoh T and Kataoka T: Crucial role of phospholipase Cepsilon in chemical carcinogen-induced skin tumor development. *Cancer Res* 64: 8808-8810, 2004.
18. Fujita J, Ohuchi N, Ito N, Reynolds SH, Yoshida O, Nakayama H and Kitamura Y: Activation of H-ras oncogene in rat bladder tumors induced by N-butyl-N-(4-hydroxybutyl)nitrosamine. *J Natl Cancer Inst* 80: 37-43, 1988.
19. Vecchione A, Sevignani C, Giarnieri E, Zanasi N, Ishii H, Cesari R, Fong LY, Gomella LG, Croce CM and Baffa R: Inactivation of the FHIT gene favors bladder cancer development. *Clin Cancer Res* 10: 7607-7612, 2004.
20. Ikuta S, Edamatsu H, Li M, Hu L and Kataoka T: Crucial role of phospholipase C epsilon in skin inflammation induced by tumor-promoting phorbol ester. *Cancer Res* 68: 64-72, 2008.
21. Mantovani A, Allavena P, Sica A and Balkwill F: Cancer-related inflammation. *Nature* 454: 436-444, 2008.
22. Greten FR, Eckmann L, Greten TF, Park JM, Li ZW, Egan LJ, Kagnoff MF and Karin M: IKKbeta links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* 118: 285-296, 2004.
23. Karin M and Greten FR: NF-kappaB: Linking inflammation and immunity to cancer development and progression. *Nat Rev Immunol* 5: 749-759, 2005.
24. Bartoletti R, Cai T, Nesi G, Sardi I and Rizzo M: Qualitative and quantitative analysis of angiogenic factors in transitional cell bladder carcinoma: Relationship with clinical course at 10 years follow-up. *Oncol Rep* 14: 251-255, 2005.
25. Cui L, Shi Y, Dai G, Pan H, Chen J, Song L, Wang S, Chang HC, Sheng H and Wang X: Modification of N-Methyl-N-Nitrosourea initiated bladder carcinogenesis in Wistar rats by terephthalic acid. *Toxicol Appl Pharmacol* 210: 24-31, 2006.
26. Ariel I, Ayesh S, Gofrit O, Ayesh B, Abdul-Ghani R, Pizov G, Smith Y, Sidi AA, Birman T, Schneider T, *et al*: Gene expression in the bladder carcinoma rat model. *Mol Carcinog* 41: 69-76, 2004.
27. Ding J, Xu D, Pan C, Ye M, Kang J, Bai Q and Qi J: Current animal models of bladder cancer: Awareness of translatability (Review). *Exp Ther Med* 8: 691-699, 2014.
28. Gofrit ON, Birman T, Dinaburg A, Ayesh S, Ohana P and Hochberg A: Chemically induced bladder cancer-a sonographic and morphologic description. *Urology* 68: 231-235, 2006.
29. Zhang RY, Du WQ, Zhang YC, Zheng JN and Pei DS: PLCε signaling in cancer. *J Cancer Res Clin Oncol*: June 25, 2015 (Epub ahead of print).
30. Gofrit ON, Birman T, Dinaburg A, Ayesh S, Ohana P and Hochberg A: Chemically induced bladder cancer-a sonographic and morphologic description. *Urology* 68: 231-235, 2006.
31. Chu H, Wang M and Zhang Z: Bladder cancer epidemiology and genetic susceptibility. *J Biomed Res* 27: 170-178, 2013.
32. Murta-Nascimento C, Schmitz-Drager BJ, Zeegers MP, Steineck G, Kogevinas M, Real FX and Malats N: Epidemiology of urinary bladder cancer: from tumor development to patient's death. *World J Uro* 25: 285-295, 2007.
33. Malats N and Real FX: Epidemiology of bladder cancer. *Hematol Oncol Clin North Am* 29: 177-189, vii, 2015.
34. Burchill SA, Neal DE and Lunec J: Frequency of H-ras mutations in human bladder cancer detected by direct sequencing. *Br J Urol* 73: 516-521, 1994.
35. Traczyk M, Borkowska E, Rozniecki M, Purpurowicz R, Jędrzejczyk A, Marks P, Pietrusiński M, Jabłonowski Z, Sosnowski M and Kałużewski B: Polymorphic variants of H-RAS protooncogene and their possible role in bladder cancer etiology. *Cent European J Urol* 65: 84-87, 2012.
36. Ahmad I, Patel R, Liu Y, Singh LB, Taketo MM, Wu XR, Leung HY and Sansom OJ: Ras mutation cooperates with β-catenin activation to drive bladder tumorigenesis. *Cell Death Dis* 2: e124, 2011.
37. Shibatohe M, Kariya K, Liao Y, Hu CD, Watari Y, Goshima M, Shima F and Kataoka T: Identification of PLC210, a *Caenorhabditis elegans* phospholipase C, as a putative effector of Ras. *J Biol Chem* 273: 6218-6222, 1998.
38. Bourguignon LY, Gilad E, Brightman A, Diedrich F and Singleton P: Hyaluronan-CD44 interaction with leukemia-associated RhoGEF and epidermal growth factor receptor promotes Rho/Ras co-activation, phospholipase C epsilon-Ca<sup>2+</sup> signaling and cytoskeleton modification in head and neck squamous cell carcinoma cells. *J Biol Chem* 281: 14026-14040, 2006.
39. Cheng H, Luo C, Wu X, Zhang Y, He Y, Wu Q, Xia Y and Zhang J: shRNA targeting PLCε inhibits bladder cancer cell growth in vitro and in vivo. *Urology* 78: 474.e477-411, 2011.
40. Ling Y, Chunli L, Xiaohou W and Qiaoling Z: Involvement of the PLCε/PKCα pathway in human BIU-87 bladder cancer cell proliferation. *Cell Biol Int* 35: 1031-1036, 2011.
41. Du HF, Ou LP, Yang X, Song XD, Fan YR, Tan B, Luo CL and Wu X: A new PKCα/β/TBX3/E-cadherin pathway is involved in PLCε-regulated invasion and migration in human bladder cancer cells. *Cell Signal* 26: 580-593, 2014.
42. Nagano T, Edamatsu H, Kobayashi K, Takenaka N, Yamamoto M, Sasaki N, Nishimura Y and Kataoka T: Phospholipase ce, an effector of ras and rap small GTPases, is required for airway inflammatory response in a mouse model of bronchial asthma. *PLoS One* 9: e108373, 2014.
43. Li M, Edamatsu H, Kitazawa R, Kitazawa S and Kataoka T: Phospholipase Cepsilon promotes intestinal tumorigenesis of Apc (Min/+) mice through augmentation of inflammation and angiogenesis. *Carcinogenesis* 30: 1424-1432, 2009.