

Pathogenic *Acinetobacter*: from the Cell Surface to Infinity and Beyond

Brent S. Weber,^{a,b} Christian M. Harding,^b Mario F. Feldman^b

Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada^a; Department of Molecular Microbiology, Washington University School of Medicine in St. Louis, St. Louis, Missouri, USA^b

The genus *Acinetobacter* encompasses multiple nosocomial opportunistic pathogens that are of increasing worldwide relevance because of their ability to survive exposure to various antimicrobial and sterilization agents. Among these, *Acinetobacter baumannii*, *Acinetobacter nosocomialis*, and *Acinetobacter pittii* are the most frequently isolated in hospitals around the world. Despite the growing incidence of multidrug-resistant *Acinetobacter* spp., little is known about the factors that contribute to pathogenesis. New strategies for treating and managing infections caused by multidrug-resistant *Acinetobacter* strains are urgently needed, and this requires a detailed understanding of the pathobiology of these organisms. In recent years, some virulence factors important for *Acinetobacter* colonization have started to emerge. In this review, we focus on several recently described virulence factors that act at the bacterial surface level, such as the capsule, O-linked protein glycosylation, and adhesins. Furthermore, we describe the current knowledge regarding the type II and type VI secretion systems present in these strains.

Infections caused by pathogenic members of the genus *Acinetobacter* are emerging as a significant threat to human health. These Gram-negative bacteria are responsible for increasing numbers of infections encountered in hospitals, particularly among immunocompromised patients (1, 2), and community-acquired infections are also increasing in prevalence (3). A recently recognized population at particular risk for *Acinetobacter* infections is military service members who have suffered combat-related injuries, who often acquire these infections in field hospitals (4–6). Although *Acinetobacter* spp. primarily cause pneumonia, they are also frequent causes of wound and burn infections, meningitis, urinary tract infections, and sepsis (7). There is a growing trend for these isolates to display high levels of antibiotic resistance, with some being resistant to all clinically available antibiotics (8). Collectively, these attributes have led to a pressing need to elucidate the mechanisms used by pathogenic *Acinetobacter* to cause disease.

The bacterial cell surface plays essential roles in sensing of the environment, interactions with the host, and maintenance of cellular homeostasis (9). The molecular structures present on the cell surface, and those that extend beyond the surface, are of central importance for understanding the pathogenesis of an organism; these are often the key determinants that mediate bacterial virulence and thus represent important targets for novel antimicrobials and vaccines (10). This review highlights the recent research on pathogenic *Acinetobacter* spp., which has led to several key findings on the strategies used by these bacteria to elaborate cell surface and secretion components that are vital for causing disease (Fig. 1).

CELL SURFACE

The cell surface of Gram-negative bacteria plays myriad roles in the physiology of these organisms, including transport of molecules into and out of the cell, interaction with and sensing of the extracellular environment, and protection from external stresses (9). While all bacterial cell surfaces are composed primarily of lipids, carbohydrates, and proteins, the diversity in the molecular composition and arrangement of these structures has vast implications for virulence in pathogenic bacteria. Recent experimental

investigation into the arrangement and composition of these structures on *Acinetobacter* cell surfaces has provided important insights into their role in the pathobiology of these important human pathogens.

Lipooligosaccharide. Lipopolysaccharide (LPS), the major component of the outer leaflet of the outer membrane (OM) of many Gram-negative bacteria, is an immunostimulatory molecule that plays an important role in bacterial resistance to external stresses (11). LPS is composed of the endotoxic lipid A, a core oligosaccharide, and a repeating sugar structure called the O antigen (11). During biosynthesis of LPS, the core oligosaccharide is built onto the lipid A moiety in the cytoplasm and flipped into the periplasmic space. The repeat subunit of the O antigen is synthesized separately onto an undecaprenyl phosphate (Und-P) carrier, which is then flipped to the periplasm and ligated to the lipid A core by the WaaL ligase enzyme (12); thus, the WaaL ligase is essential for the production of O antigen. Many pathogens, such as *Escherichia coli* and *Salmonella* spp., produce diverse O-antigen structures that form the basis for serotyping schemes (13–15). However, other important bacterial pathogens, such as *Neisseria* and *Campylobacter*, lack a WaaL ligase ortholog and do not produce O antigen, elaborating only lipooligosaccharide (LOS) (16).

Whether *Acinetobacter* spp. elaborate LPS or LOS on their cell surface has been a topic of considerable debate. Several reports have described the presence of LPS in *Acinetobacter*, many of which have used structural or antibody-based methods to detect O-antigen carbohydrate moieties (17–26). However, most *Acinetobacter* spp. do not show “typical” LPS laddering upon silver

Accepted manuscript posted online 28 December 2015

Citation Weber BS, Harding CM, Feldman MF. 2016. Pathogenic *Acinetobacter*: from the cell surface to infinity and beyond. *J Bacteriol* 198:880–887.
doi:10.1128/JB.00906-15.

Editor: W. Margolin

Address correspondence to Mario F. Feldman, mariofeldman@wustl.edu.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.

coded by the gene near the pilin gene was found to exclusively glycosylate the cognate pilin protein, similar to the PilO/TfpO protein from *Pseudomonas aeruginosa* (47), while the second gene encoded a general O-OTase that was responsible for the glycosylation of multiple proteins, similar to PglL from *Neisseria* (48, 49). Although the functional significance of having two glycosylation systems remains to be determined, this represents the first known case of multiple O-OTases present in a single bacterium.

Capsule. Like many other pathogens, *Acinetobacter* spp. produce an extracellular capsule that provides a layer of protection from external threats such as complement-mediated killing (50). Capsule production and protein glycosylation are exquisitely linked in *Acinetobacter*, as the carbohydrate repeat unit found in the capsule is the same as the single repeat unit attached to proteins (46). The sugar subunits for capsule and protein glycosylation are derived from the same pathway, in which an initiating glycosyltransferase, PglC/ItrA, attaches the first carbohydrate to Und-P, followed by the addition of other sugar monomers by glycosyltransferase enzymes to complete the repeat unit (46). This repeat unit is then flipped to the periplasm, and the capsule and protein glycosylation pathways diverge at this point. In the case of protein glycosylation, this single repeat unit is attached to the target protein by the O-OTase (29). For capsule production, individual sugar repeat units are instead polymerized and exported to the cell surface. This bifurcated pathway represents a novel mechanism that illustrates the evolutionary connections between capsule and protein glycosylation, which may allow *Acinetobacter* to adapt rapidly to changing environments. How *A. baumannii* partitions a given carbohydrate repeat unit to the protein glycosylation pathway or capsule production remains to be determined. Although the carbohydrate structures produced by different strains are highly variable, functional studies have shown that capsule production is essential for *Acinetobacter* survival during infection and growth in serum (32, 46, 50, 51). It was recently reported that capsule production could be increased by the presence of subinhibitory levels of antibiotics, which increased resistance to complement-mediated killing and led to a hypervirulent phenotype in a mouse model of systemic infection (52). This capsule hyperproduction phenotype was shown to be controlled by the two-component BfmRS system, which regulates several other important virulence factors in *Acinetobacter* (53). *Acinetobacter* spp. also produce a surface-associated poly- β -1-6-*N*-acetylglucosamine (PNAG) polysaccharide, which is important for virulence and biofilm formation (54).

Pili. Filamentous bacterial surface appendages, termed pili, mediate interactions between the producing organism and their environment. *Acinetobacter* pili have been studied since 1975, when Henrichsen and Blom observed that *Acinetobacter calcoaceticus* strains displaying surface fimbrial structures exhibited twitching motility (55, 56), a form of bacterial locomotion now known to be dependent on functioning type IV pili (57). Furthermore, it has been shown that the nonpathogenic model organism *Acinetobacter baylyi* ADP1 produces both thin and thick pili (58); however, the roles pili play in the biology and pathobiology of pathogenic *Acinetobacter* spp. have been only partially elucidated.

A system of chaperone/usher pili, designated Csu pili, has been identified in all sequenced pathogenic *Acinetobacter* spp.; however, the Csu pili have been primarily studied in *A. baumannii* (59). The *Acinetobacter* Csu pili are required for biofilm formation and maintenance in *A. baumannii* ATCC 19606 but were found

not to play a role in adherence to human epithelial cells (60). Another study found that the CsuA/B pilin subunit was the most abundant protein identified within the pellicle matrix of multiple *A. baumannii* strains (61), further strengthening the role of Csu pili in biofilm formation and maintenance. Previous reports also identified a single nucleotide insertion in the *csuB* gene of *A. baumannii* ATCC 17978, suggesting that this system may be nonfunctional in this strain (62, 63); however, recent resequencing of the *A. baumannii* ATCC 17978 genome (GenBank accession no. CP012004) did not find the same insertion event (64). Lastly, other chaperone/usher pilus-like systems have been bioinformatically identified in many *A. baumannii* strains, but none of those systems has been functionally characterized (61, 62). Mass spectrometric characterization of pellicle-associated proteins did find that non-Csu-pilin subunits were present, indicating that these systems may be functional.

Medically relevant *Acinetobacter* spp. have also been shown to produce type IV pili (Tfp), which are dynamic bacterial surface appendages known to mediate twitching motility, horizontal gene transfer, and biofilm formation (65). Although bioinformatic studies have identified genes predicted to encode proteins required for the biogenesis of Tfp in *A. baumannii*, only *Acinetobacter nosocomialis* strain M2 has been shown conclusively to produce functioning Tfp (66, 67), which is glycosylated by a TfpO-like oligosaccharyltransferase (30). Many *A. baumannii* isolates have been found to be naturally transformable and to exhibit twitching motility (63, 68–70), two classic Tfp-associated phenotypes, which strongly indicates their presence. Tfp-like structures were also identified on *A. baumannii* ATCC 17978 (71); furthermore, mutants in predicted Tfp biogenesis components of *A. baumannii* ATCC 17978 exhibited impaired biofilm formation (71) but the major pilin subunit, PilA, has not been shown to be surface exposed and/or associated with the pilin structures observed. Although Tfp, with roles in *Acinetobacter* motility and natural transformation, has emerged as a possible virulence factor, no studies have conclusively linked Tfp to the pathobiology of *Acinetobacter*, as is the case for *Pseudomonas* and *Neisseria*.

PROTEIN SECRETION

Extracellular export of proteins is a fundamental process of all forms of life. Protein secretion systems of Gram-negative bacteria are extremely diverse in function and composition and often are important mediators of virulence. Recent research has elucidated several of the mechanisms *Acinetobacter* uses to secrete proteins and the role they play in the biology of *Acinetobacter*. Here we describe the recent insights into macromolecular protein secretion systems present in *Acinetobacter*, with a focus on the systems that export proteins out of the cell and that have been characterized experimentally.

Type II secretion. The most recently described secretion system is a functional type II secretion system (T2SS) identified in both *A. nosocomialis* strain M2 (119) and *A. baumannii* ATCC 17978 (72); moreover, it was shown that clinical isolates of *Acinetobacter pittii*, *A. baumannii*, *A. calcoaceticus*, and *Acinetobacter junii* all were able to secrete type II substrates, indicating that functioning type II secretion systems seem to be the rule and not the exception. With regard to the T2SS of *A. nosocomialis* strain M2, a two-dimensional differential gel electrophoresis approach identified multiple putative type II substrates; the LipA and LipH lipases and the CpaA metallopeptidase were vali-

dated as *bona fide* type II secretion substrates. Interestingly, both LipA and CpaA required specific membrane-associated chaperones for secretion, which indicates that T2SS chaperones are more widespread than previously recognized. Importantly, it was shown that an *A. nosocomialis* strain M2 *gspD* mutant lacking the outer membrane secretin of the T2SS was severely attenuated in both the *Galleria mellonella* and murine pulmonary infection models. Specifically, mice intranasally infected with the *gspD* mutant strain had ~ 2 -log lower bacterial burdens in both the lungs and the spleen after 36 h, compared to both the parental strain and the complemented mutant. In *A. baumannii* ATCC 17978, Johnson et al. identified a lipase, LipA, secreted in a T2SS-dependent manner that was required for growth on medium containing lipids as a sole carbon source (72). Mutants with mutations in *lipA* or the T2SS structural gene *gspD* were less competitive than the wild-type strain in a mixed-infection murine model of bacteremia. Collectively, these findings indicate that the *Acinetobacter* T2SS is a previously unrecognized virulence factor mediating pathogenesis in a relevant mammalian model. Interestingly, a recent study by Wang et al. utilized an *A. baumannii* ATCC 17978 *gspN* mutant for validation of their insertion sequencing murine pulmonary infection studies, and they subsequently found that the *gspN* mutant did not display any virulence defect in survival or competition models, compared to the parent strain (51). Although these data are in contrast to the newly defined role of type II secretion in *Acinetobacter*, it was demonstrated previously that *gspN* homologs were not required for a functioning T2SS in *Klebsiella oxytoca* (73); furthermore, *gspN* homologs are absent from numerous known T2SSs in other Gram-negative bacteria (74), indicating the dispensable nature of GspN in functioning T2SSs.

Autotransporters. A type V autotransporter has been characterized in *A. baumannii*. The *Acinetobacter* trimeric autotransporter (Ata) was found to be crucial for the ability of certain *A. baumannii* strains to adhere to extracellular matrix components, including collagen I, III, IV, and V (75). Ata is also an important mediator of *A. baumannii* biofilm formation and maintenance, as an *A. baumannii* ATCC 17978 *ata* mutant had significantly diminished biofilm production and was less virulent in a murine intraperitoneal infection model, compared to the parental and complemented strains (75).

Type VI secretion. Bacteria interact with each other in a multitude of ways; these interactions are often competitive in nature and play important roles in niche establishment (76). The bacterial type VI secretion system was first formally described for *Vibrio cholerae* and *P. aeruginosa* and was suggested to play a role against eukaryotic hosts (77, 78). While several T6SSs have been determined to secrete antieukaryotic toxins, it has recently been appreciated that many bacteria use their T6SSs to secrete antibacterial toxins to kill competing bacteria (79, 80). The T6SS is composed of approximately 15 conserved structural proteins and a variable number of accessory factors, which work in concert to secrete proteins extracellularly (81). Important components include Hcp, which forms a polymerized tubular structure that is secreted out of the cell and is essential for protein secretion, and VgrGs, which are present at the tip of this structure and can have effector activity or facilitate effector secretion (82). The T6SS bears striking similarity to bacteriophage, both structurally and functionally (83).

The presence of a T6SS was initially predicted bioinformatically for *A. baylyi*, and Hcp was subsequently detected in superna-

tants of *A. baumannii* ATCC 19606 (84, 85). The genetic organization and sequences of T6SS genes are remarkably well conserved across *Acinetobacter* spp. Based on homology with T6SS genes in other bacteria, the single *Acinetobacter* T6SS locus includes most of the genes required for apparatus assembly and function (86, 87). Notably, the main T6SS cluster does not contain *vgrG* genes, which are instead scattered throughout the genome. The VgrG proteins of *Acinetobacter*, which differ in number from strain to strain, do not seem to include effector domains (86, 88). Instead, bioinformatic analyses suggest that the proteins are most likely to mediate the secretion of adjacently encoded toxic effectors, with cognate immunity proteins being encoded nearby (88). However, no *bona fide* T6SS-dependent effectors have been experimentally characterized in *Acinetobacter*. In *A. baylyi*, mutation of three PAAR proteins, which interact with VgrG proteins, results in loss of Hcp secretion, and one of those PAAR proteins has been experimentally determined to be secreted, although it is not clear whether the proteins have any effector functions themselves (89). The primary function of the T6SS in *Acinetobacter* seems to be to kill competing bacteria, and *Acinetobacter* spp. with active T6SSs are able to kill a wide variety of other bacteria, including other strains of the same species (64, 87–89).

T6SS expression often is tightly controlled and is activated only under certain conditions; the molecular mechanisms used to achieve this regulation are extremely diverse and complex and differ from organism to organism and even between strains of a given species (82). Although little is known about T6SS regulation in *Acinetobacter*, recent studies have provided insight into some of the regulatory mechanisms used by these organisms. T6SS activities vary widely in different strains and species of *Acinetobacter*, with some strains showing robust T6SSs and bacterial killing and others seeming to have inactive systems under laboratory conditions (64, 86–88). However, the available data suggest that strains with T6SSs invariably express the main protein Hcp to at least some level, with variations in whether the protein is secreted (thus determining whether the system is active) (64, 86, 88). In *A. baumannii* ATCC 19606, Hcp is constitutively secreted in wild-type cells but is lost in mutants lacking lipid A, potentially due to membrane disruptions (85). In *A. baumannii* ATCC 17978, T6SS activity is controlled by a plasmid (see below), and a chromosomally encoded histone-like nucleoid-structuring (H-NS) protein may also regulate the T6SS (90). Several *A. baumannii* isolates harbor a resistance plasmid that encodes repressors of the T6SS (64). Upon spontaneous loss of this plasmid and subsequent loss of the repressors, the T6SS is activated and the resistance genes are lost. The functional significance of this remains to be elucidated but, considering the tremendous amount of energy required for T6SS activity (91) and the fitness defects often caused by harboring multiple antibiotic resistance genes (92), this may represent a mechanism to maintain both systems while avoiding potentially deleterious effects of having them be active at the same time. In the absence of antibiotic pressure, *Acinetobacter* strains do not require resistance genes but are more likely to encounter competitors, thus losing the resistance plasmid and activating the T6SS, which could provide a competitive advantage. Because the cells that lose the plasmid will lose resistance to antibiotics, this strategy could constitute an altruistic mechanism to ensure *Acinetobacter* population survival. It should be noted that several recent multidrug-resistant (MDR) *A. baumannii* strains seem to have permanently inactivated their T6SSs through chromosomal gene loss; it has

been suggested that this may be a result of the antibiotic pressure being great enough to make it evolutionarily advantageous to completely lose the T6SS, rather than maintaining it in an inactive state (93). Although there are limited data, there appears to be a link between antibiotic resistance and T6SS status in *Acinetobacter*; strains that are multidrug resistant express but do not secrete Hcp, while those that are not multidrug resistant are more likely to have an active T6SS (64, 86, 88).

Outer membrane vesicles. A special case of protein secretion is the production of outer membrane vesicles (OMVs), which are blebs of outer membrane (OM) released from the bacterial cell surface (94). There is significant debate regarding whether OMVs are produced by a directed process or simply represent cellular debris. Proteomic comparisons between the OM and OMVs of some bacteria have shown that the protein profiles differ between these two fractions, indicating that some OM proteins are excluded from OMV recruitment and suggesting that OMV formation is a directed process (94). However, many studies also detected cytoplasmic proteins in OMV preparations, indicating that cell lysis could also be a major contributor to OMV formation (94). OMVs have been implicated in numerous biological functions, with particular attention being devoted to their role in virulence (95). Several studies on OMVs in *Acinetobacter* have suggested that they have many functions, including roles in horizontal gene transfer, antibiotic resistance, and virulence. A wide variety of cargo types have been identified in OMVs from different *Acinetobacter* strains, including virulence proteins, antibiotic resistance determinants, and DNA (96–102). An important virulence factor of *A. baumannii*, OmpA, has also been found to be associated with OMVs, and OMVs have been suggested to act as a mechanism for delivery for this protein to host cells (103). Furthermore, OmpA has been suggested to be involved in the biogenesis of OMVs (104). OMVs from *Acinetobacter* may have an important role in the development of novel therapeutics, as they can stimulate a strong immune response and are protective when administered as a vaccine (105–107).

FUTURE DIRECTIONS

The surge of infections caused by *Acinetobacter* spp. has led to increased interest from the scientific and medical communities and attempts to understand the disease-causing mechanisms of these organisms. Many recent studies have greatly increased our understanding of *Acinetobacter* infection mechanisms but have also illuminated what a formidable pathogen the health care community is facing. There is intense interest in discovering novel strategies to fight this pathogen, which is quickly becoming untreatable with our current antibiotic repertoire. Although the vast majority of patients who become infected with *Acinetobacter* spp. are immunocompromised, which complicates intervention strategies, vaccines have been proposed as an alternative method to fight MDR *Acinetobacter* (108, 109), and several promising candidates have been described (107, 110–114). Individuals for whom immunocompromise could potentially be predicted *a priori*, such as patients undergoing cancer treatments or surgery and military personnel entering conflicts, may benefit from a prophylactic vaccination strategy (115). Carbohydrate structures present in *Acinetobacter*, such as the capsule and glycoproteins, represent attractive antigenic targets for vaccine development; since the carbohydrate moieties are the same in both, targeting these structures may provide broad protection (46). Indeed, capsule-based

vaccines have shown efficacy in soft tissue, pneumonia, and bacteremia rodent models (116, 117). A drawback to this approach, however, is that the strain-to-strain variations in carbohydrate structures are so great that a multivalent vaccine to target all pathogenic *Acinetobacter* strains is unrealistic. Extensive epidemiological data on the capsular/glycoprotein serotypes most prevalent in a given health care institution may allow for a more directed approach to vaccine design. Indeed, PCR-based schemes have been proposed to accomplish this, and advances in the speed and cost of genome sequencing may make this a feasible approach (118). Because the production of capsular polysaccharides and glycoproteins is essential for virulence, targeting common steps in the biosynthetic pathway of these structures may be more feasible. Given the genomic plasticity of *Acinetobacter*, it is likely that a “one size fits all” solution to the problem is not possible, and multiple strategies should be investigated in order to determine the most beneficial approach for a given health care setting.

FUNDING INFORMATION

Washington University School of Medicine provided funding to Brent S. Weber, Christian M. Harding, and Mario F. Feldman. Gouvernement du Canada | Natural Sciences and Engineering Research Council of Canada (NSERC) provided funding to Brent S. Weber.

M.F.F. is a Washington University School of Medicine Faculty Diversity Scholar. B.S.W. is a recipient of the NSERC Postgraduate Scholarship-Doctoral award.

REFERENCES

1. Peleg AY, Hooper DC. 2010. Hospital-acquired infections due to Gram-negative bacteria. *N Engl J Med* 362:1804–1813. <http://dx.doi.org/10.1056/NEJMra0904124>.
2. Gaynes R, Edwards JR. 2005. Overview of nosocomial infections caused by Gram-negative bacilli. *Clin Infect Dis* 41:848–854. <http://dx.doi.org/10.1086/432803>.
3. Dexter C, Murray GL, Paulsen IT, Peleg AY. 2015. Community-acquired *Acinetobacter baumannii*: clinical characteristics, epidemiology and pathogenesis. *Expert Rev Anti Infect Ther* 13:567–573. <http://dx.doi.org/10.1586/14787210.2015.1025055>.
4. Scott P, Deye G, Srinivasan A, Murray C, Moran K, Hulten E, Fishbain J, Craft D, Riddell S, Lindler L, Mancuso J, Milstrey E, Bautista CT, Patel J, Ewell A, Hamilton T, Gaddy C, Tenney M, Christopher G, Petersen K, Endy T, Petrucci B. 2007. An outbreak of multidrug-resistant *Acinetobacter baumannii*-calcoaceticus complex infection in the US military health care system associated with military operations in Iraq. *Clin Infect Dis* 44:1577–1584. <http://dx.doi.org/10.1086/518170>.
5. Turton JF, Kaufmann ME, Gill MJ, Pike R, Scott PT, Fishbain J, Craft D, Deye G, Riddell S, Lindler LE, Pitt TL. 2006. Comparison of *Acinetobacter baumannii* isolates from the United Kingdom and the United States that were associated with repatriated casualties of the Iraq conflict. *J Clin Microbiol* 44:2630–2634. <http://dx.doi.org/10.1128/JCM.00547-06>.
6. Sebeny PJ, Riddle MS, Petersen K. 2008. *Acinetobacter baumannii* skin and soft-tissue infection associated with war trauma. *Clin Infect Dis* 47:444–449. <http://dx.doi.org/10.1086/590568>.
7. Dijkshoorn L, Nemea A, Seifert H. 2007. An increasing threat in hospitals: multidrug-resistant *Acinetobacter baumannii*. *Nat Rev Microbiol* 5:939–951. <http://dx.doi.org/10.1038/nrmicro1789>.
8. Valencia R, Arroyo LA, Conde M, Aldana JM, Torres MJ, Fernandez-Cuenca F, Garnacho-Montero J, Cisneros JM, Ortiz C, Pachon J, Aznar J. 2009. Nosocomial outbreak of infection with pan-drug-resistant *Acinetobacter baumannii* in a tertiary care university hospital. *Infect Control Hosp Epidemiol* 30:257–263. <http://dx.doi.org/10.1086/595977>.
9. Silhavy TJ, Kahne D, Walker S. 2010. The bacterial cell envelope. *Cold Spring Harb Perspect Biol* 2:a000414.
10. Grandi G. 2010. Bacterial surface proteins and vaccines. *F1000 Biol Rep* 2:36.

11. Raetz CR, Whitfield C. 2002. Lipopolysaccharide endotoxins. *Annu Rev Biochem* 71:635–700. <http://dx.doi.org/10.1146/annurev.biochem.71.110601.135414>.
12. Whitfield C, Amor PA, Koplín R. 1997. Modulation of the surface architecture of Gram-negative bacteria by the action of surface polymer: lipid A-core ligase and by determinants of polymer chain length. *Mol Microbiol* 23:629–638. <http://dx.doi.org/10.1046/j.1365-2958.1997.2571614.x>.
13. Stenutz R, Weintraub A, Widmalm G. 2006. The structures of *Escherichia coli* O-polysaccharide antigens. *FEMS Microbiol Rev* 30:382–403. <http://dx.doi.org/10.1111/j.1574-6976.2006.00016.x>.
14. Liu B, Knirel YA, Feng L, Perepelov AV, Senchenkova SN, Reeves PR, Wang L. 2014. Structural diversity in *Salmonella* O antigens and its genetic basis. *FEMS Microbiol Rev* 38:56–89. <http://dx.doi.org/10.1111/1574-6976.12034>.
15. Orskov I, Orskov F, Jann B, Jann K. 1977. Serology, chemistry, and genetics of O and K antigens of *Escherichia coli*. *Bacteriol Rev* 41:667–710.
16. Preston A, Mandrell RE, Gibson BW, Apicella MA. 1996. The lipooligosaccharides of pathogenic Gram-negative bacteria. *Crit Rev Microbiol* 22:139–180. <http://dx.doi.org/10.3109/10408419609106458>.
17. Pantophlet R, Brade L, Brade H. 1999. Identification of *Acinetobacter baumannii* strains with monoclonal antibodies against the O antigens of their lipopolysaccharides. *Clin Diagn Lab Immunol* 6:323–329.
18. Haseley SR, Traub WH, Wilkinson SG. 1997. Structures of polymeric products isolated from the lipopolysaccharides of reference strains for *Acinetobacter baumannii* O23 and O12. *Eur J Biochem* 244:147–154. <http://dx.doi.org/10.1111/j.1432-1033.1997.00147.x>.
19. Traub WH. 1989. *Acinetobacter baumannii* serotyping for delineation of outbreaks of nosocomial cross-infection. *J Clin Microbiol* 27:2713–2716.
20. Haseley SR, Wilkinson SG. 1998. Structure of the O-7 antigen from *Acinetobacter baumannii*. *Carbohydr Res* 306:257–263. [http://dx.doi.org/10.1016/S0008-6215\(97\)10039-8](http://dx.doi.org/10.1016/S0008-6215(97)10039-8).
21. Haseley S, Wilkinson SG. 1997. Structure of the O18 antigen from *Acinetobacter baumannii*. *Carbohydr Res* 301:187–192. [http://dx.doi.org/10.1016/S0008-6215\(97\)00095-5](http://dx.doi.org/10.1016/S0008-6215(97)00095-5).
22. Haseley SR, Wilkinson SG. 1994. Structure of the putative O10 antigen from *Acinetobacter baumannii*. *Carbohydr Res* 264:73–81. [http://dx.doi.org/10.1016/0008-6215\(94\)00179-0](http://dx.doi.org/10.1016/0008-6215(94)00179-0).
23. MacLean LL, Perry MB, Chen W, Vinogradov E. 2009. The structure of the polysaccharide O-chain of the LPS from *Acinetobacter baumannii* strain ATCC 17961. *Carbohydr Res* 344:474–478. <http://dx.doi.org/10.1016/j.carres.2008.12.026>.
24. Vinogradov EV, Brade L, Brade H, Holst O. 2003. Structural and serological characterisation of the O-antigenic polysaccharide of the lipopolysaccharide from *Acinetobacter baumannii* strain 24. *Carbohydr Res* 338:2751–2756. <http://dx.doi.org/10.1016/j.carres.2003.05.001>.
25. Vinogradov EV, Duus JO, Brade H, Holst O. 2002. The structure of the carbohydrate backbone of the lipopolysaccharide from *Acinetobacter baumannii* strain ATCC 19606. *Eur J Biochem* 269:422–430. <http://dx.doi.org/10.1046/j.0014-2956.2001.02647.x>.
26. Vinogradov EV, Pantophlet R, Dijkshoorn L, Brade L, Holst O, Brade H. 1996. Structural and serological characterisation of two O-specific polysaccharides of *Acinetobacter*. *Eur J Biochem* 239:602–610. <http://dx.doi.org/10.1111/j.1432-1033.1996.06020.x>.
27. Pantophlet R. 2008. *Lipopolysaccharides of Acinetobacter*. Caister Academic Press, Norfolk, United Kingdom.
28. Fregolino E, Gargiulo V, Lanzetta R, Parrilli M, Holst O, Castro CD. 2011. Identification and structural determination of the capsular polysaccharides from two *Acinetobacter baumannii* clinical isolates, MG1 and SMAL. *Carbohydr Res* 346:973–977. <http://dx.doi.org/10.1016/j.carres.2011.03.024>.
29. Iwashkiw JA, Seper A, Weber BS, Scott NE, Vinogradov E, Stratilo C, Reiz B, Cordwell SJ, Whittall R, Schild S, Feldman MF. 2012. Identification of a general O-linked protein glycosylation system in *Acinetobacter baumannii* and its role in virulence and biofilm formation. *PLoS Pathog* 8:e1002758. <http://dx.doi.org/10.1371/journal.ppat.1002758>.
30. Harding CM, Nasr MA, Kinsella RL, Scott NE, Foster LJ, Weber BS, Fiester SE, Actis LA, Tracy EN, Munson RS, Jr, Feldman MF. 2015. *Acinetobacter* strains carry two functional oligosaccharyltransferases, one devoted exclusively to type IV pilin, and the other one dedicated to O-glycosylation of multiple proteins. *Mol Microbiol* 96:1023–1041. <http://dx.doi.org/10.1111/mmi.12986>.
31. Schulz BL, Jen FE, Power PM, Jones CE, Fox KL, Ku SC, Blanchfield JT, Jennings MP. 2013. Identification of bacterial protein O-oligosaccharyltransferases and their glycoprotein substrates. *PLoS One* 8:e62768. <http://dx.doi.org/10.1371/journal.pone.0062768>.
32. Kenyon JJ, Hall RM. 2013. Variation in the complex carbohydrate biosynthesis loci of *Acinetobacter baumannii* genomes. *PLoS One* 8:e62160. <http://dx.doi.org/10.1371/journal.pone.0062160>.
33. McQueary CN, Kirkup BC, Si Y, Barlow M, Actis LA, Craft DW, Zurawski DV. 2012. Extracellular stress and lipopolysaccharide modulate *Acinetobacter baumannii* surface-associated motility. *J Microbiol* 50:434–443. <http://dx.doi.org/10.1007/s12275-012-1555-1>.
34. Luke NR, Sauberan SL, Russo TA, Beanan JM, Olson R, Loehfelm TW, Cox AD, St Michael F, Vinogradov EV, Campagnari AA. 2010. Identification and characterization of a glycosyltransferase involved in *Acinetobacter baumannii* lipopolysaccharide core biosynthesis. *Infect Immun* 78:2017–2023. <http://dx.doi.org/10.1128/IAI.00016-10>.
35. Lin L, Tan B, Pantapalangkoor P, Ho T, Baquir B, Tomaras A, Montgomery JI, Reilly U, Barbacci EG, Hujer K, Bonomo RA, Fernandez L, Hancock RE, Adams MD, French SW, Buslon VS, Spellberg B. 2012. Inhibition of LpxC protects mice from resistant *Acinetobacter baumannii* by modulating inflammation and enhancing phagocytosis. *mBio* 3:e00312-12. <http://dx.doi.org/10.1128/mBio.00312-12>.
36. Kenyon JJ, Nigro SJ, Hall RM. 2014. Variation in the OC locus of *Acinetobacter baumannii* genomes predicts extensive structural diversity in the lipooligosaccharide. *PLoS One* 9:e107833. <http://dx.doi.org/10.1371/journal.pone.0107833>.
37. Boll JM, Tucker AT, Klein DR, Beltran AM, Brodbelt JS, Davies BW, Trent MS. 2015. Reinforcing lipid A acylation on the cell surface of *Acinetobacter baumannii* promotes cationic antimicrobial peptide resistance and desiccation survival. *mBio* 6(3):e00478-15. <http://dx.doi.org/10.1128/mBio.00478-15>.
38. Arroyo LA, Herrera CM, Fernandez L, Hankins JV, Trent MS, Hancock RE. 2011. The *pmrCAB* operon mediates polymyxin resistance in *Acinetobacter baumannii* ATCC 17978 and clinical isolates through phosphoethanolamine modification of lipid A. *Antimicrob Agents Chemother* 55:3743–3751. <http://dx.doi.org/10.1128/AAC.00256-11>.
39. Chin CY, Gregg KA, Napier BA, Ernst RK, Weiss DS. 2015. A *PmrB*-regulated deacetylase required for lipid A modification and polymyxin resistance in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 59:7911–7914. <http://dx.doi.org/10.1128/AAC.00515-15>.
40. Moffatt JH, Harper M, Harrison P, Hale JD, Vinogradov E, Seemann T, Henry R, Crane B, St Michael F, Cox AD, Adler B, Nation RL, Li J, Boyce JD. 2010. Colistin resistance in *Acinetobacter baumannii* is mediated by complete loss of lipopolysaccharide production. *Antimicrob Agents Chemother* 54:4971–4977. <http://dx.doi.org/10.1128/AAC.00834-10>.
41. Beceiro A, Llobet E, Aranda J, Bengoechea JA, Doumith M, Hornsey M, Dhanji H, Chart H, Bou G, Livermore DM, Woodford N. 2011. Phosphoethanolamine modification of lipid A in colistin-resistant variants of *Acinetobacter baumannii* mediated by the *pmrAB* two-component regulatory system. *Antimicrob Agents Chemother* 55:3370–3379. <http://dx.doi.org/10.1128/AAC.00079-11>.
42. Pelletier MR, Casella LG, Jones JW, Adams MD, Zurawski DV, Hazlett KR, Doi Y, Ernst RK. 2013. Unique structural modifications are present in the lipopolysaccharide from colistin-resistant strains of *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 57:4831–4840. <http://dx.doi.org/10.1128/AAC.00865-13>.
43. Hood MI, Becker KW, Roux CM, Dunman PM, Skaar EP. 2013. Genetic determinants of intrinsic colistin tolerance in *Acinetobacter baumannii*. *Infect Immun* 81:542–551. <http://dx.doi.org/10.1128/IAI.00704-12>.
44. Hug I, Feldman MF. 2011. Analogies and homologies in lipopolysaccharide and glycoprotein biosynthesis in bacteria. *Glycobiology* 21:138–151. <http://dx.doi.org/10.1093/glycob/cwq148>.
45. Scott NE, Kinsella RL, Edwards AV, Larsen MR, Dutta S, Saba J, Foster LJ, Feldman MF. 2014. Diversity within the O-linked protein glycosylation systems of *Acinetobacter* species. *Mol Cell Proteomics* 13:2354–2370. <http://dx.doi.org/10.1074/mcp.M114.038315>.
46. Lees-Miller RG, Iwashkiw JA, Scott NE, Seper A, Vinogradov E, Schild S, Feldman MF. 2013. A common pathway for O-linked protein-glycosylation and synthesis of capsule in *Acinetobacter baumannii*. *Mol Microbiol* 89:816–830. <http://dx.doi.org/10.1111/mmi.12300>.
47. Castric P. 1995. *pilO*, a gene required for glycosylation of *Pseudomonas aeruginosa* 1244 pilin. *Microbiology* 141:1247–1254. <http://dx.doi.org/10.1099/13500872-141-5-1247>.

48. Faridmoayer A, Fentabil MA, Haurat MF, Yi W, Woodward R, Wang PG, Feldman MF. 2008. Extreme substrate promiscuity of the *Neisseria* oligosaccharyl transferase involved in protein O-glycosylation. *J Biol Chem* 283:34596–34604. <http://dx.doi.org/10.1074/jbc.M807113200>.
49. Faridmoayer A, Fentabil MA, Mills DC, Klassen JS, Feldman MF. 2007. Functional characterization of bacterial oligosaccharyltransferases involved in O-linked protein glycosylation. *J Bacteriol* 189:8088–8098. <http://dx.doi.org/10.1128/JB.01318-07>.
50. Russo TA, Luke NR, Beanan JM, Olson R, Sauberan SL, MacDonald U, Schultz LW, Umland TC, Campagnari AA. 2010. The K1 capsular polysaccharide of *Acinetobacter baumannii* strain 307-0294 is a major virulence factor. *Infect Immun* 78:3993–4000. <http://dx.doi.org/10.1128/IAI.00366-10>.
51. Wang N, Ozer EA, Mandel MJ, Hauser AR. 2014. Genome-wide identification of *Acinetobacter baumannii* genes necessary for persistence in the lung. *mBio* 5(3):e01163-14. <http://dx.doi.org/10.1128/mBio.01163-14>.
52. Geisinger E, Isberg RR. 2015. Antibiotic modulation of capsular exopolysaccharide and virulence in *Acinetobacter baumannii*. *PLoS Pathog* 11:e1004691. <http://dx.doi.org/10.1371/journal.ppat.1004691>.
53. Tomaras AP, Flagler MJ, Dorsey CW, Gaddy JA, Actis LA. 2008. Characterization of a two-component regulatory system from *Acinetobacter baumannii* that controls biofilm formation and cellular morphology. *Microbiology* 154:3398–3409. <http://dx.doi.org/10.1099/mic.0.2008/019471-0>.
54. Choi AH, Slamti L, Avci FY, Pier GB, Maira-Litran T. 2009. The *pgaABCD* locus of *Acinetobacter baumannii* encodes the production of poly- β -1-6-*N*-acetylglucosamine, which is critical for biofilm formation. *J Bacteriol* 191:5953–5963. <http://dx.doi.org/10.1128/JB.00647-09>.
55. Henrichsen J. 1975. The occurrence of twitching motility among Gram-negative bacteria. *Acta Pathol Microbiol Scand B* 83:171–178.
56. Henrichsen J, Blom J. 1975. Correlation between twitching motility and possession of polar fimbriae in *Acinetobacter calcoaceticus*. *Acta Pathol Microbiol Scand B* 83:103–115.
57. Merz AJ, So M, Sheetz MP. 2000. Pilus retraction powers bacterial twitching motility. *Nature* 407:98–102. <http://dx.doi.org/10.1038/35024105>.
58. Gohl O, Friedrich A, Hoppert M, Averhoff B. 2006. The thin pili of *Acinetobacter* sp. strain BD413 mediate adhesion to biotic and abiotic surfaces. *Appl Environ Microbiol* 72:1394–1401. <http://dx.doi.org/10.1128/AEM.72.2.1394-1401.2006>.
59. Tomaras AP, Dorsey CW, Edelmann RE, Actis LA. 2003. Attachment to and biofilm formation on abiotic surfaces by *Acinetobacter baumannii*: involvement of a novel chaperone-usher pili assembly system. *Microbiology* 149:3473–3484. <http://dx.doi.org/10.1099/mic.0.26541-0>.
60. de Breij A, Gaddy J, van der Meer J, Koning R, Koster A, van den Broek P, Actis L, Nibbering P, Dijkshoorn L. 2009. CsuA/BABCDE-dependent pili are not involved in the adherence of *Acinetobacter baumannii* ATCC19606(T) to human airway epithelial cells and their inflammatory response. *Res Microbiol* 160:213–218. <http://dx.doi.org/10.1016/j.resmic.2009.01.002>.
61. Nait Chabane Y, Marti S, Rihouey C, Alexandre S, Hardouin J, Lesouhaitier O, Vila J, Kaplan JB, Jouenne T, De E. 2014. Characterisation of pellicles formed by *Acinetobacter baumannii* at the air-liquid interface. *PLoS One* 9:e111660. <http://dx.doi.org/10.1371/journal.pone.0111660>.
62. Eijkelkamp BA, Stroehner UH, Hassan KA, Paulsen IT, Brown MH. 2014. Comparative analysis of surface-exposed virulence factors of *Acinetobacter baumannii*. *BMC Genomics* 15:1020. <http://dx.doi.org/10.1186/1471-2164-15-1020>.
63. Eijkelkamp BA, Stroehner UH, Hassan KA, Papadimitriou MS, Paulsen IT, Brown MH. 2011. Adherence and motility characteristics of clinical *Acinetobacter baumannii* isolates. *FEMS Microbiol Lett* 323:44–51. <http://dx.doi.org/10.1111/j.1574-6968.2011.02362.x>.
64. Weber BS, Ly PM, Irwin JN, Pukatzki S, Feldman MF. 2015. A multidrug resistance plasmid contains the molecular switch for type VI secretion in *Acinetobacter baumannii*. *Proc Natl Acad Sci U S A* 112:9442–9447. <http://dx.doi.org/10.1073/pnas.1502966112>.
65. Burrows LL. 2012. *Pseudomonas aeruginosa* twitching motility: type IV pili in action. *Annu Rev Microbiol* 66:493–520. <http://dx.doi.org/10.1146/annurev-micro-092611-150055>.
66. Harding CM, Tracy EN, Carruthers MD, Rather PN, Actis LA, Munson RS, Jr. 2013. *Acinetobacter baumannii* strain M2 produces type IV pili which play a role in natural transformation and twitching motility but not surface-associated motility. *mBio* 4:e00360-13. <http://dx.doi.org/10.1128/mBio.00360-13>.
67. Carruthers MD, Harding CM, Baker BD, Bonomo RA, Hujer KM, Rather PN, Munson RS, Jr. 2013. Draft genome sequence of the clinical isolate *Acinetobacter nosocomialis* strain M2. *Genome Announc* 1:e00906-13. <http://dx.doi.org/10.1128/genomeA.00906-13>.
68. Ramirez MS, Don M, Merkier AK, Bistue AJ, Zorreguieta A, Centron D, Tolmasky ME. 2010. Naturally competent *Acinetobacter baumannii* clinical isolate as a convenient model for genetic studies. *J Clin Microbiol* 48:1488–1490. <http://dx.doi.org/10.1128/JCM.01264-09>.
69. Wilharm G, Piesker J, Laue M, Skiebe E. 2013. DNA uptake by the nosocomial pathogen *Acinetobacter baumannii* occurs during movement along wet surfaces. *J Bacteriol* 195:4146–4153. <http://dx.doi.org/10.1128/JB.00754-13>.
70. Yoon EJ, Chabane YN, Goussard S, Snesrud E, Courvalin P, De E, Grillot-Courvalin C. 2015. Contribution of resistance-nodulation-cell division efflux systems to antibiotic resistance and biofilm formation in *Acinetobacter baumannii*. *mBio* 6:e00309-15. <http://dx.doi.org/10.1128/mBio.00309-15>.
71. Tucker AT, Nowicki EM, Boll JM, Knauf GA, Burdis NC, Trent MS, Davies BW. 2014. Defining gene-phenotype relationships in *Acinetobacter baumannii* through one-step chromosomal gene inactivation. *mBio* 5:e01313-14. <http://dx.doi.org/10.1128/mBio.01313-14>.
72. Johnson TL, Waack U, Smith S, Mobley H, Sandkvist M. 14 December 2015. *Acinetobacter baumannii* is dependent on the type II secretion system and its substrate LipA for lipid utilization and in vivo fitness. *J Bacteriol* <http://dx.doi.org/10.1128/JB.00622-15>.
73. Possot OM, Vignon G, Bomchil N, Ebel F, Pugsley AP. 2000. Multiple interactions between pullulanase secretor components involved in stabilization and cytoplasmic membrane association of PulE. *J Bacteriol* 182:2142–2152. <http://dx.doi.org/10.1128/JB.182.8.2142-2152.2000>.
74. Campos M, Cisneros DA, Nivaskumar M, Francetic O. 2013. The type II secretion system: a dynamic fiber assembly nanomachine. *Res Microbiol* 164:545–555. <http://dx.doi.org/10.1016/j.resmic.2013.03.013>.
75. Bentancor LV, Camacho-Peiro A, Bozkurt-Guzel C, Pier GB, Maira-Litran T. 2012. Identification of Ata, a multifunctional trimeric auto-transporter of *Acinetobacter baumannii*. *J Bacteriol* 194:3950–3960. <http://dx.doi.org/10.1128/JB.06769-11>.
76. Hibbing ME, Fuqua C, Parsek MR, Peterson SB. 2010. Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* 8:15–25. <http://dx.doi.org/10.1038/nrmicro2259>.
77. Pukatzki S, Ma AT, Sturtevant D, Krastins B, Sarracino D, Nelson WC, Heidelberg JF, Mekalanos JJ. 2006. Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system. *Proc Natl Acad Sci U S A* 103:1528–1533. <http://dx.doi.org/10.1073/pnas.0510322103>.
78. Mougous JD, Cuff ME, Raunser S, Shen A, Zhou M, Gifford CA, Goodman AL, Joachimiak G, Ordenez CL, Lory S, Walz T, Joachimiak A, Mekalanos JJ. 2006. A virulence locus of *Pseudomonas aeruginosa* encodes a protein secretion apparatus. *Science* 312:1526–1530. <http://dx.doi.org/10.1126/science.1128393>.
79. Ho BT, Dong TG, Mekalanos JJ. 2014. A view to a kill: the bacterial type VI secretion system. *Cell Host Microbe* 15:9–21. <http://dx.doi.org/10.1016/j.chom.2013.11.008>.
80. Russell AB, Peterson SB, Mougous JD. 2014. Type VI secretion system effectors: poisons with a purpose. *Nat Rev Microbiol* 12:137–148. <http://dx.doi.org/10.1038/nrmicro3185>.
81. Pukatzki S, McAuley SB, Miyata ST. 2009. The type VI secretion system: translocation of effectors and effector-domains. *Curr Opin Microbiol* 12:11–17. <http://dx.doi.org/10.1016/j.mib.2008.11.010>.
82. Silverman JM, Brunet YR, Cascales E, Mougous JD. 2012. Structure and regulation of the type VI secretion system. *Annu Rev Microbiol* 66:453–472. <http://dx.doi.org/10.1146/annurev-micro-121809-151619>.
83. Brunet YR, Henin J, Celia H, Cascales E. 2014. Type VI secretion and bacteriophage tail tubes share a common assembly pathway. *EMBO Rep* 15:315–321. <http://dx.doi.org/10.1002/embr.201337936>.
84. de Berardinis V, Vallenet D, Castelli V, Besnard M, Pinet A, Craud C, Samair S, Lechaplais C, Gyapay G, Richez C, Durot M, Kreimeyer A, Le Fevre F, Schachter V, Pezo V, Doring V, Scarpelli C, Medigue C, Cohen GN, Marliere P, Salanoubat M, Weissenbach J. 2008. A complete collection of single-gene deletion mutants of *Acinetobacter baylyi* ADP1. *Mol Syst Biol* 4:174.

85. Henry R, Vithanage N, Harrison P, Seemann T, Coutts S, Moffatt JH, Nation RL, Li J, Harper M, Adler B, Boyce JD. 2012. Colistin-resistant, lipopolysaccharide-deficient *Acinetobacter baumannii* responds to lipopolysaccharide loss through increased expression of genes involved in the synthesis and transport of lipoproteins, phospholipids, and poly- β -1,6-*N*-acetylglucosamine. *Antimicrob Agents Chemother* 56:59–69. <http://dx.doi.org/10.1128/AAC.05191-11>.
86. Weber BS, Miyata ST, Iwashkiw JA, Mortensen BL, Skaar EP, Pukatzki S, Feldman MF. 2013. Genomic and functional analysis of the type VI secretion system in *Acinetobacter*. *PLoS One* 8:e55142. <http://dx.doi.org/10.1371/journal.pone.0055142>.
87. Carruthers MD, Nicholson PA, Tracy EN, Munson RS, Jr. 2013. *Acinetobacter baumannii* utilizes a type VI secretion system for bacterial competition. *PLoS One* 8:e59388. <http://dx.doi.org/10.1371/journal.pone.0059388>.
88. Repizo GD, Gagne S, Foucault-Grunenwald ML, Borges V, Charpentier X, Limansky AS, Gomes JP, Viale AM, Salcedo SP. 2015. Differential role of the T6SS in *Acinetobacter baumannii* virulence. *PLoS One* 10:e0138265. <http://dx.doi.org/10.1371/journal.pone.0138265>.
89. Shneider MM, Buth SA, Ho BT, Basler M, Mekalanos JJ, Leiman PG. 2013. PAAR-repeat proteins sharpen and diversify the type VI secretion system spike. *Nature* 500:350–353. <http://dx.doi.org/10.1038/nature12453>.
90. Eijkelkamp BA, Stroehrer UH, Hassan KA, Elbourne LD, Paulsen IT, Brown MH. 2013. H-NS plays a role in expression of *Acinetobacter baumannii* virulence features. *Infect Immun* 81:2574–2583. <http://dx.doi.org/10.1128/IAI.00065-13>.
91. Basler M. 2015. Type VI secretion system: secretion by a contractile nanomachine. *Philos Trans R Soc Lond B Biol Sci* 370:20150021. <http://dx.doi.org/10.1098/rstb.2015.0021>.
92. Andersson DI, Hughes D. 2010. Antibiotic resistance and its cost: is it possible to reverse resistance? *Nat Rev Microbiol* 8:260–271.
93. Wright MS, Haft DH, Harkins DM, Perez F, Hujer KM, Bajaksouzian S, Benard MF, Jacobs MR, Bonomo RA, Adams MD. 2014. New insights into dissemination and variation of the health care-associated pathogen *Acinetobacter baumannii* from genomic analysis. *mBio* 5:e00963-13. <http://dx.doi.org/10.1128/mBio.00963-13>.
94. Haurat MF, Elhenawy W, Feldman MF. 2015. Prokaryotic membrane vesicles: new insights on biogenesis and biological roles. *Biol Chem* 396:95–109.
95. Ellis TN, Kuehn MJ. 2010. Virulence and immunomodulatory roles of bacterial outer membrane vesicles. *Microbiol Mol Biol Rev* 74:81–94. <http://dx.doi.org/10.1128/MMBR.00031-09>.
96. Rumbo C, Fernandez-Moreira E, Merino M, Poza M, Mendez JA, Soares NC, Mosquera A, Chaves F, Bou G. 2011. Horizontal transfer of the OXA-24 carbapenemase gene via outer membrane vesicles: a new mechanism of dissemination of carbapenem resistance genes in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 55:3084–3090. <http://dx.doi.org/10.1128/AAC.00929-10>.
97. Liao YT, Kuo SC, Chiang MH, Lee YT, Sung WC, Chen YH, Chen TL, Fung CP. 2015. *Acinetobacter baumannii* extracellular OXA-58 is primarily and selectively released via outer membrane vesicles after Sec-dependent periplasmic translocation. *Antimicrob Agents Chemother* 59:7346–7354. <http://dx.doi.org/10.1128/AAC.01343-15>.
98. Li ZT, Zhang RL, Bi XG, Xu L, Fan M, Xie D, Xian Y, Wang Y, Li XJ, Wu ZD, Zhang KX. 2015. Outer membrane vesicles isolated from two clinical *Acinetobacter baumannii* strains exhibit different toxicity and proteome characteristics. *Microb Pathog* 81:46–52. <http://dx.doi.org/10.1016/j.micpath.2015.03.009>.
99. Fulsundar S, Kulkarni HM, Jagannadham MV, Nair R, Keerthi S, Sant P, Pardesi K, Bellare J, Chopade BA. 2015. Molecular characterization of outer membrane vesicles released from *Acinetobacter radioresistens* and their potential roles in pathogenesis. *Microb Pathog* 83–84:12–22.
100. Fulsundar S, Harms K, Flaten GE, Johnson PJ, Chopade BA, Nielsen KM. 2014. Gene transfer potential of outer membrane vesicles of *Acinetobacter baylyi* and effects of stress on vesiculation. *Appl Environ Microbiol* 80:3469–3483. <http://dx.doi.org/10.1128/AEM.04248-13>.
101. Dallo SF, Zhang B, Denno J, Hong S, Tsai A, Haskins W, Ye JY, Weitao T. 2012. Association of *Acinetobacter baumannii* EF-Tu with cell surface, outer membrane vesicles, and fibronectin. *ScientificWorldJournal* 2012:128705.
102. Kwon SO, Gho YS, Lee JC, Kim SI. 2009. Proteome analysis of outer membrane vesicles from a clinical *Acinetobacter baumannii* isolate. *FEMS Microbiol Lett* 297:150–156. <http://dx.doi.org/10.1111/j.1574-6968.2009.01669.x>.
103. Jin JS, Kwon SO, Moon DC, Gurung M, Lee JH, Kim SI, Lee JC. 2011. *Acinetobacter baumannii* secretes cytotoxic outer membrane protein A via outer membrane vesicles. *PLoS One* 6:e17027. <http://dx.doi.org/10.1371/journal.pone.0017027>.
104. Moon DC, Choi CH, Lee JH, Choi CW, Kim HY, Park JS, Kim SI, Lee JC. 2012. *Acinetobacter baumannii* outer membrane protein A modulates the biogenesis of outer membrane vesicles. *J Microbiol* 50:155–160. <http://dx.doi.org/10.1007/s12275-012-1589-4>.
105. Nho JS, Jun SH, Oh MH, Park TI, Choi CW, Kim SI, Choi CH, Lee JC. 2015. *Acinetobacter nosocomialis* secretes outer membrane vesicles that induce epithelial cell death and host inflammatory responses. *Microb Pathog* 81:39–45. <http://dx.doi.org/10.1016/j.micpath.2015.03.012>.
106. Jun SH, Lee JH, Kim BR, Kim SI, Park TI, Lee JC, Lee YC. 2013. *Acinetobacter baumannii* outer membrane vesicles elicit a potent innate immune response via membrane proteins. *PLoS One* 8:e71751. <http://dx.doi.org/10.1371/journal.pone.0071751>.
107. McConnell MJ, Rumbo C, Bou G, Pachon J. 2011. Outer membrane vesicles as an acellular vaccine against *Acinetobacter baumannii*. *Vaccine* 29:5705–5710. <http://dx.doi.org/10.1016/j.vaccine.2011.06.001>.
108. Garcia-Quintanilla M, Pulido MR, McConnell MJ. 2013. First steps towards a vaccine against *Acinetobacter baumannii*. *Curr Pharm Biotechnol* 14:897–902.
109. Pachon J, McConnell MJ. 2014. Considerations for the development of a prophylactic vaccine for *Acinetobacter baumannii*. *Vaccine* 32:2534–2536. <http://dx.doi.org/10.1016/j.vaccine.2013.10.064>.
110. Badmasti F, Ajdary S, Bouzari S, Fooladi AA, Shahcheraghi F, Siadat SD. 2015. Immunological evaluation of OMV(PagL)+Bap(1-487aa) and AbOmpA(8-346aa)+Bap(1-487aa) as vaccine candidates against *Acinetobacter baumannii* sepsis infection. *Mol Immunol* 67:552–558. <http://dx.doi.org/10.1016/j.molimm.2015.07.031>.
111. Lin L, Tan B, Pantapalangkoor P, Ho T, Hujer AM, Taracila MA, Bonomo RA, Spellberg B. 2013. *Acinetobacter baumannii* rOmpA vaccine dose alters immune polarization and immunodominant epitopes. *Vaccine* 31:313–318. <http://dx.doi.org/10.1016/j.vaccine.2012.11.008>.
112. Bentancor LV, Routray A, Bozkurt-Guzel C, Camacho-Peiro A, Pier GB, Maira-Litran T. 2012. Evaluation of the trimeric autotransporter Ata as a vaccine candidate against *Acinetobacter baumannii* infections. *Infect Immun* 80:3381–3388. <http://dx.doi.org/10.1128/IAI.06096-11>.
113. McConnell MJ, Pachon J. 2010. Active and passive immunization against *Acinetobacter baumannii* using an inactivated whole cell vaccine. *Vaccine* 29:1–5. <http://dx.doi.org/10.1016/j.vaccine.2010.10.052>.
114. Huang W, Wang S, Yao Y, Xia Y, Yang X, Long Q, Sun W, Liu C, Li Y, Ma Y. 2015. OmpW is a potential target for eliciting protective immunity against *Acinetobacter baumannii* infections. *Vaccine* 33:4479–4485. <http://dx.doi.org/10.1016/j.vaccine.2015.07.031>.
115. Rubin LG, Levin MJ, Ljungman P, Davies EG, Avery R, Tomblyn M, Bouvaros A, Dhanireddy S, Sung L, Keyserling H, Kang I, Infectious Diseases Society of America. 2014. 2013 IDSA clinical practice guideline for vaccination of the immunocompromised host. *Clin Infect Dis* 58:309–318. <http://dx.doi.org/10.1093/cid/cit816>.
116. Russo TA, Beanan JM, Olson R, MacDonald U, Cox AD, St Michael F, Vinogradov EV, Spellberg B, Luke-Marshall NR, Campagnari AA. 2013. The K1 capsular polysaccharide from *Acinetobacter baumannii* is a potential therapeutic target via passive immunization. *Infect Immun* 81:915–922. <http://dx.doi.org/10.1128/IAI.01184-12>.
117. Bentancor LV, O'Malley JM, Bozkurt-Guzel C, Pier GB, Maira-Litran T. 2012. Poly-*N*-acetyl- β -(1-6)-glucosamine is a target for protective immunity against *Acinetobacter baumannii* infections. *Infect Immun* 80:651–656. <http://dx.doi.org/10.1128/IAI.05653-11>.
118. Hu D, Liu B, Dijkshoorn L, Wang L, Reeves PR. 2013. Diversity in the major polysaccharide antigen of *Acinetobacter baumannii* assessed by DNA sequencing, and development of a molecular serotyping scheme. *PLoS One* 8:e70329. <http://dx.doi.org/10.1371/journal.pone.0070329>.
119. Harding CM, Kinsella RL, Palmer LD, Skaar EP, Feldman MF. 2016. Medically relevant *Acinetobacter* species require a type II secretion system and specific membrane-associated chaperones for the export of multiple substrates and full virulence. *PLoS Pathog* 12:e1005391. <http://dx.doi.org/10.1371/journal.ppat.1005391>.