

Estradiol, TGF- β 1 and hypoxia promote breast cancer stemness and EMT-mediated breast cancer migration

SEONG-JOON PARK^{1,2*}, JOONG-GOOK KIM^{1*}, NAM DEUK KIM²,
KWANGMO YANG^{1,3,4}, JAE WOONG SHIM¹ and KYU HEO¹

¹Research Center, Dongnam Institute of Radiological and Medical Sciences, Busan 46033;
²Department of Pharmacy, Pusan National University, Busan 46241; ³Department of Radiation Oncology,
Dongnam Institute of Radiological & Medical Sciences, Busan 46033; ⁴Department of Radiation Oncology,
Korea Institute of Radiological & Medical Sciences, Seoul 01812, Republic of Korea

Received February 2, 2015; Accepted December 11, 2015

DOI: 10.3892/ol.2016.4115

Abstract. Breast cancer is one of the most common cancer types among women, acting as a distinct cause of mortality, and has a high incidence of recurrence. External stimuli, including 17 β -estradiol (E2), transforming growth factor (TGF)- β 1 and hypoxia, may be important in breast cancer growth and metastasis. However, the effects of these stimuli on breast cancer stem cell (CSC) regulation have not been fully investigated. In the present study, the proportion of cluster of differentiation (CD)44⁺/CD24^{-low} cells increased following treatment with E2, TGF- β 1 and hypoxia in MCF-7 cells. The expression of CSC markers, including SOX2, KLF4 and ABCG2, was upregulated continually by E2, TGF- β 1 and hypoxia. In addition, the expression levels of epithelial-mesenchymal transition-associated factors increased following treatment with E2, TGF- β 1 and hypoxia. Therefore, the migration ability of E2-, TGF- β 1- and hypoxia-treated MCF-7 cells was enhanced compared with control cells. In addition, the enhancement of apoptosis by 5-fluorouracil or radiation was abolished following treatment with E2, TGF- β 1 and hypoxia. These results indicate that E2, TGF- β 1 and hypoxia are important for regulating breast CSCs, and that the modulation of the microenvironment in tumors may improve the efficiency of breast cancer therapy.

Introduction

Breast cancer is one of the most common cancer types among women, acting as a distinct cause of mortality, and has a high

incidence of recurrence (1). Consequently, numerous studies have focused on the development of breast cancer therapies, including surgery, chemotherapy and radiotherapy (2-4). The cancer stem cell (CSC) hypothesis proposes that the preferential targets of oncogenic transformation are tissue stem cells or progenitor cells that are able to self-renew (5-8). Therefore, numerous studies have developed CSC phenotypic assays in order to identify CSCs (9-11). Breast CSCs have been characterized as cluster of differentiation (CD)44⁺/CD24^{-low} cells that initiate carcinogenesis in NOD/SCID mice (12).

Hypoxia is involved in various tumors and often occurs when tumor growth surpasses blood supply (13). In particular, breast cancers are sensitive to hypoxia, as they outgrow nutrients or oxygen vascular supplies (14). In addition, hypoxia often causes chemo- and radiotherapy resistance, and contributes to tumor metastasis (15). Although the effect of hypoxia in patients with breast cancer has been widely reported, the regulatory mechanism of hypoxia in breast CSC and therapeutic resistance remain unknown (16-18).

17 β -estradiol (E2) is the most effective female estrogen hormone and is pivotal in male and female physiology (19). Breast cancer is susceptible to estrogen hormones. Estrogen hormones promote the development and progression of breast cancer, and induce the invasion and metastasis of breast cancer cells that express estrogen receptors (ERs) to distant organs or lymph nodes (20). E2 also enhances the movement and invasion of breast cancer cells (21). However, due to the complexity of ER-triggered estrogen signaling, the effects of estrogen hormones on cancer are occasionally divergent.

The anti-inflammatory cytokine transforming growth factor (TGF)- β 1 is associated with embryonic development and homeostasis in adult organisms (22,23). TGF- β 1 is critical for angiogenesis, immunoregulation and cancer progression (24). In addition, TGF- β 1 acts as a tumor suppressor in the early stages of breast carcinoma, and is involved in the progression of tumors by resisting inhibited cell growth during the later stages of disease (25). TGF- β 1 also enhances breast cancer metastasis by inducing Smad family member 2 (Smad2) (26). However, the regulatory mechanisms of breast CSCs following treatment with E2 or TGF- β 1 have not been investigated.

Correspondence to: Dr Kyu Heo or Mr. Jae Woong Shim, Research Center, Dongnam Institute of Radiological and Medical Sciences, 40 Jwadong, Gijang, Busan 46033, Republic of Korea
E-mail: khjhk33@gmail.com
E-mail: shimjaewoong5187@gmail.com

*Contributed equally.

Key words: breast cancer stem cell, estrogen, transforming growth factor- β 1, hypoxia

In the present study, treatment with E2, TGF- β 1 and hypoxia led to breast CSC (CD44⁺/CD24^{-low}) expansion. CSC markers and epithelial-mesenchymal transition (EMT)-associated factors were expressed in order to investigate the underlying mechanisms. Additionally, the effects of E2, TGF- β 1 and hypoxia on cell migration and drug and radiation resistance were determined. The results indicate that E2, TGF- β 1 and hypoxia are important for the regulation of breast CSCs, and that the modulation of the tumor microenvironment may improve the efficiency of breast cancer therapy.

Materials and methods

Cell culture. The human breast cancer MCF-7 cell line was obtained from the American Type Culture Collection (Rockville, MD, USA) and cultured at 37°C in 20% O₂ and 5% CO₂ in Dulbecco's modified Eagle's medium (DMEM; Welgene, Daegu, South Korea) that contained 10% HyClone fetal bovine serum (FBS; GE Healthcare Life Sciences, Logan, UT, USA) and 1% Gibco antibiotic-antimycotic (Thermo Fisher Scientific, Inc., Waltham, MA, USA). The cells were incubated in a chamber containing a 5% CO₂ and 1% O₂ atmosphere for 48, 72 and 96 h and 1 week to create hypoxic conditions. The cells were treated with 10 nM E2 or 1 ng/ml TGF- β 1 (R&D Systems Inc., Minneapolis, MN, USA).

Immunoblot analysis. The MCF-7 cells (2x10⁶) were collected using 5 ml cold phosphate buffer solution (PBS; Welgene) and lysed in 100 μ l lysis buffer containing 20 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1 mM Na₂ ethylenediaminetetraacetic acid, 1 mM ethylene glycol tetraacetic acid, 1% Triton, 2.5 mM sodium pyrophosphate, 1 mM β -glycerophosphate, 1 mM Na₃VO₄, 1 μ g/ml leupeptin and 1 mM phenylmethanesulfonyl-fluoride (all Sigma-Aldrich, St. Louis, MO, USA). Following 1 hour of incubation on ice, samples were centrifuged at 13,000 x g for 20 min at 4°C. Subsequently, the supernatant was removed and quantified using the Bio-Rad Protein Assay kit II (Bio-Rad Laboratories, Inc., Hercules, CA, USA) according to the manufacturers's protocol. A total of 50 μ g of total protein was loaded and electrophoresed on sodium dodecyl sulfate-polyacrylamide gels (Bio-Rad Laboratories, Inc.) and transferred to polyvinylidene difluoride membranes (GE Healthcare Life Sciences, Chalfont, UK). The transferred membranes were blocked using 5% milk (BD Biosciences, San Jose, CA, USA) dissolved in Tris-buffered saline (20 mM Tris; 137 mM NaCl; pH 7.6; Sigma-Aldrich) containing 0.02% Tween 20 (Sigma-Aldrich), and incubated overnight at 4°C with specific primary antibodies. The membranes were subsequently incubated with specific horseradish peroxidase-conjugated secondary antibodies (detailed below). The blots were developed with the Super Signal Chemiluminescence reagent (Pierce Biotechnology, Inc., Rockford, IL, USA) by enhanced chemiluminescence (Thermo Fisher Scientific, Inc.).

Immunoblot antibodies. The following mouse monoclonal anti-human primary antibodies (1 μ g; dilution, 1:1,000) were utilized for immunoblot analysis: Anti-CD44 (catalog no., ab78960), anti-CD24 (catalog no., ab76514), anti-ATP-binding cassette sub-family G member 2 (ABCG2;

catalog no., ab130244), anti-JMJD1A (catalog no., ab107234) and anti-Kruppel-like factor 4 (KLF4; catalog no., ab75486) (all purchased from Abcam, Cambridge, MA, USA); epithelial cell adhesion molecule (EpCAM; catalog no., 2929) antibody was from Cell Signaling Technology, Inc. (Danvers, MA, USA); the cytokeratin 5/8 [mouse anti-human immunoglobulin G (IgG)]; catalog no., 550505), SOX2 (mouse anti-human IgG; catalog no., 561469) and β -catenin (mouse anti-human IgG; catalog no., 610154) antibodies were from BD Biosciences; the c-Myc (catalog no., sc-40) and E- and N-cadherin (catalog nos., sc-71008 and sc-271386, respectively) antibodies were from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA); the hypoxia inducing factor (HIF)-1 α (catalog no., NB100-105) antibody was from Novus Biologicals, LLC (Littleton, CO, USA); and the β -actin (catalog no., A-5441) antibody was from Sigma-Aldrich. The 3 μ g of goat anti-mouse polyclonal secondary antibody (catalog no. 62-6520) was from Thermo Fisher Scientific, Inc. and was used at a dilution of 1:4,000.

Flow cytometry. MCF-7 cells (2x10⁶) were washed twice using PBS and incubated with 2 μ g/ml monoclonal mouse anti-human CD24-fluorescein isothiocyanate (FITC; catalog no., 555427) and monoclonal mouse anti-human CD44-allophycocyanin (APC; catalog no., 559250) antibodies (BD Biosciences; dilution, 1:100) in the dark on ice for 1 h, and washed twice using cold PBS. The labeled cells were analyzed using the fluorescence-activated cell sorting FACS Aria cell sorter (BD Biosciences).

Migration assay. An *in vitro* wound-healing assay was used to assess two-dimensional cell motility. The MCF-7 cells (2x10⁶) were treated for 96 h with E2 or TGF- β 1 and hypoxia in 6-well plates. Then, a scratch was made on the cell layer with a micropipette tip, and the cultures were washed twice with serum-free medium to remove the floating cells. The cells were incubated in a chamber containing an atmosphere of 20% O₂ or in hypoxic conditions (5% CO₂ and 1% O₂) in DMEM medium at 37°C. Wound healing was visualized by comparing photographs 48 h later using the Qimaging QI Click Camera system mounted on a phase-contrast Nikon microscope (TS100; Nikon, Inc., Tokyo, Japan). For the Transwell assay, MCF-7 cells (2x10⁴) were seeded onto 8- μ m Transwell-inserts (Costar brand; Corning Life Sciences, Tewksbury, MA, USA). The lower chambers were filled with DMEM containing 10% FBS. Migrated cells were stained with crystal violet and counted using the Image-Pro Plus 7.0 software (Media Cybernetics, Rockville, MD, USA) 24 h later.

Apoptosis assay. Cell apoptosis was assessed using the Annexin V/phycoerythrin (PE) Apoptosis Detection kit (BD Biosciences; catalog no. 559763). Briefly, the MCF-7 cells (1x10⁶) were seeded in 100-mm dishes and incubated overnight. Then, the cells were treated with 5-fluorouracil (100 μ g/ml) or radiation (10 Gy) for 48 or 72 h, respectively. Subsequently, the cells were washed twice with 5 ml PBS and stained with 2.5 μ g/ml Annexin V/PE-conjugate and 5 μ l 7-aminoactinomycin for 15 min on ice in the dark. Subsequent to staining, the cells were analyzed using the FACS Aria cell sorter.




Figure 1. E2, TGF-β1 and hypoxia treatment increased the CD44⁺/CD24^{-low} population of MCF-7 cells. (A) Schematic representation of the experimental procedure. (B-F) Representative FACS stained images. The FACS analysis was performed using specific cell-surface markers for basal (CD44-allophycocyanin) and luminal (CD24-phycoerythrin) epithelial cells, and the percentage of CD44⁺/CD24^{-low} cells in each condition was assessed. The ratio of CD44⁺/CD24^{-low} cells in E2 (10 nM), TGF-β1 (1 ng/ml) and hypoxia (1% O₂) treated MCF-7 cells is indicated. E2, estradiol; TGF-β1, transforming growth factor-β1; CD, cluster of differentiation; FACS, fluorescence-activated cell sorting.

Statistical analysis. Statistical analyses were performed using Excel (Microsoft Corporation, Redmond, WA, USA). A Student's *t*-test was used to make statistical comparisons and P<0.05 was considered to indicate a significant difference.

Results

Effect of E2, TGF-β1 and hypoxia on breast CSC expansion. The percentage of CD44⁺/CD24^{-low} cells is considered to be




Figure 2. Patterns of cancer stem cell and epithelial-mesenchymal transition-associated marker expression in E2, TGF- β 1 and hypoxia-treated MCF-7 cells. (A) The expression of CD44, CD24, Sox2, ABCG2 and KLF4 was determined in E2, TGF- β 1, combined E2/TGF- β 1 and hypoxia-treated MCF-7 cells by western blotting. β -actin was used as a loading control. Western blotting was performed at each time point (48, 72 and 96 h and 1 week). (B) EpCAM, cytokeratin 5/8, E-cadherin, N-cadherin, β -catenin, c-Myc and HIF-1 α expression was determined by western blotting using same conditions that are indicated in Fig. 2A. E2, estradiol; TGF- β 1, transforming growth factor- β 1; CD, cluster of differentiation; Sox2, sex determining region Y-box 2; ABCG2, ATP-binding cassette sub-family G member 2; KLF4, Kruppel-like factor 4; EpCAM, epithelial cell adhesion molecule; HIF, hypoxia-inducible factor.

the breast CSC subpopulation. This percentage was assessed in MCF-7 cells, in the presence or absence of E2 and TGF- β 1, in order to investigate whether E2 or TGF- β 1 treatment affects the size of the CSC population. ER⁺ MCF-7 cells were treated with 10 nM E2, 1 ng/ml TGF- β 1 or a combination of the two for 48, 72 and 96 h, and 1 week. Following treatment, the proportion of stem-like cells was evaluated using flow cytometry (Fig. 1A). The results indicated that the proportion of CD44⁺/CD24^{-/low} cells was significantly expanded in E2-treated MCF-7 cells (48 h, 2.2%; 72 h, 4.1%; 96 h, 13%; 1 week, 1.1%) compared with untreated control cells (0.2%; Fig. 1B and C). TGF- β 1-treated MCF-7 cells also demonstrated a notable increase in the percentage of CD44⁺/CD24^{-/low} cells at 48 (3.9%), 72 (6.5%) and 96 h (13.0%), and 1 week (8.0%) subsequent to treatment with TGF- β 1 (Fig. 1D). However, no synergistic induction of the CD44⁺/CD24^{-/low} population by the TGF- β 1/E2 combined treatment was observed (Fig. 1E). The effect of hypoxia on breast CSC expansion was also examined. The results show that the proportion of CD44⁺/CD24^{-/low} cells increased in hypoxia-treated MCF-7 cells (48 h, 2.5%; 72 h, 3.1%; 96 h, 7.8%; 1 week, 4.1%) compared with the proportion in normoxic conditions (Fig. 1B and F). These results indicate that treatment with E2, TGF- β 1 and hypoxia in isolation expanded the CSC population of MCF-7 cells, but that the combined treatment had no synergistic effect.

Effect of E2, TGF- β 1 and hypoxia on the expression of breast CSC markers and EMT-associated factors. CSC marker

expression in E2-, TGF- β 1- or hypoxia-treated MCF-7 cells was examined, as the E2, TGF- β 1 and hypoxia treatment increased the CSC population in MCF-7 cells. MCF-7 cells were treated and incubated under the same conditions exhibited in Fig. 1A, and the CD44, CD24, SOX2, ABCG2 and KLF4 protein levels were measured using western blot analysis (Fig. 2A). Consistent with the flow cytometry results, CD44 expression increased in E2-, TGF- β 1- or hypoxia-treated MCF-7 cells. However, CD24 expression decreased following E2, TGF- β 1 and hypoxia treatment. Notably, the expression of the pluripotency-associated proteins SOX2 and KLF4 and the putative CSC marker ABCG2 increased when the cells were treated with E2, TGF- β 1 and hypoxia. These results indicate that SOX2, KLF4 and ABCG2, which are associated with breast cancer stemness, were actively regulated by the E2, TGF- β 1, and hypoxia treatments.

Several studies have demonstrated that CSCs and EMT-phenotypic cells have a tumor aggressiveness phenotype (27-29). Therefore, the present study investigated whether treatment with E2, TGF- β 1 and hypoxia affects the expression of EMT-associated factors. The results demonstrate that EpCAM and E-cadherin expression was not affected by E2- or TGF- β 1 stimulation (Fig. 2B). However, expression of the EMT markers cytokeratin 5/8 and N-cadherin, which increase cell motility, clearly increased following treatment with E2 or TGF- β 1. In addition, the levels of β -catenin and the target gene c-Myc were upregulated in E2 and TGF- β 1-treated MCF-7 cells compared with untreated control cells. EpCAM




Figure 3. E2, TGF-β1 and hypoxia treatment enhances cell motility. (A) Representative images of the induction of MCF-7 cell migration by treatment with E2 (10 nM), TGF-β1 (1 ng/ml), combined E2 (10 nM)/TGF-β1 (1 ng/ml) and hypoxia (1% O₂). (B) Migration of MCF-7 cells treated with E2, TGF-β1, combined E2/TGF-β1 and hypoxia was evaluated using Transwell polystyrene membranes with 8-mm pores. The number of migrating cells was counted and scored in relative units. P-values were calculated using Student's *t*-test. *P<0.05, ***P<0.001. E2, estradiol; TGF-β1, transforming growth factor-β1.




Figure 4. E2, TGF-β1, and hypoxia treatment increases chemo- and radioresistance. Each set of MCF-7 cells was treated with (A) 100 μg/ml 5-FU or (B) 10 Gy radiation to induce apoptosis. Then, apoptosis was analyzed with an Annexin V kit. E2, estradiol; TGF-β1, transforming growth factor-β1; 5-FU, fluorouracil.

expression increased, whereas E-cadherin expression decreased, in hypoxia-treated MCF-7 cells (Fig. 2B). Cytokeratin 5/8, N-cadherin, β-catenin and c-Myc levels increased

under hypoxic conditions. These results indicate that the CSC expansion mediated by E2, TGF-β1 and hypoxia may promote EMT by regulating the expression of EMT-associated factors.

Enhancement of cell motility by E2, TGF- β 1 and hypoxia.

The increased number of CSCs and progression of the EMT are associated with cell motility and invasion. A wound-healing assay with the E2, TGF- β 1 and hypoxia treatments was performed to visualize the effect on MCF-7 cell motility (Fig. 3A). E2 and TGF- β 1-treated cells demonstrated increased cell migration compared with control cells. The hypoxia treatment also significantly enhanced cellular migration ability. Similarly, the Transwell assay results indicated that E2-, TGF- β 1- and hypoxia-treated MCF-7 cells migrated more compared with control MCF-7 cells (Fig. 3B). The data indicate that treatment with TGF- β 1, E2 and hypoxia induces migration ability in MCF-7 cells.

Effect of E2, TGF- β 1 and hypoxia on drug and radioresistance.

Fluorouracil (5-FU) is a chemotherapeutic agent used to treat various types of cancer, including breast cancer (30). In the present study, ABCG2 expression was induced by E2, TGF- β 1 and hypoxia treatments (Fig. 2A). The effect of E2, TGF- β 1 and hypoxia on 5-FU-induced-apoptosis was investigated as ABCG2 mediates multidrug resistance. The results showed that E2-, TGF- β - and hypoxia-treated MCF-7 cells were more resistant to 5-FU compared with untreated control MCF-7 cells. In particular, 5-FU-induced apoptosis was entirely abolished in hypoxia-treated cells (Fig. 4A). E2-, TGF- β 1- and hypoxia-treated cells demonstrated inhibited radiation-induced apoptosis (Fig. 4B). These results indicate that the TGF- β 1, E2 and hypoxia treatments enhance drug and radioresistance, which may be due to the increased expression of CSC-associated proteins, including ABCG2 and β -catenin.

Discussion

CSCs possess the stem cell properties of self-renewal and differentiation; therefore, tumors may be initiated, grow, invade and metastasize (31). Anticancer therapies at present emphasize killing CSCs or inhibiting the induction of the CSC population (32,33). However, CSC-targeted cancer therapies are problematic due to the characteristics of CSCs, including radio- and chemoresistance (34,35). The aim of the present study was to identify the regulatory mechanisms for breast CSC expansion. Treatment with E2 induces the proliferation of human ER⁺ breast cancer cells (36,37). In addition, a previous study indicated that breast CSCs are stimulated by estrogens, through paracrine fibroblast growth factor/Tbx3 signaling (38). Numerous studies have demonstrated that estrogen hormones act as a negative regulator of TGF- β 1-stimulated cellular responses (39,40). Therefore, the present study examined the effects of E2 and TGF- β 1 treatment on the regulation of breast CSCs. The results of the present study indicated that treatment with E2 or TGF- β 1 in isolation expanded the CD44⁺/CD24^{-/low} cell subpopulation, but that a combination of E2 and TGF- β 1 treatment did not exhibit a synergistic effect (Fig. 1). These results demonstrate that E2 and TGF- β 1 do not affect each other during regulation of the breast CSC population. Oxygen is a critical regulator of cellular metabolism and proliferation, and hypoxia regulates a variety of pro-angiogenic pathways and carcinogenesis (41,42). Although numerous studies have reported that hypoxia improves therapeutic

efficacy by eliminating the CSC population (43-45), other studies indicate that hypoxia induces CSC characteristics by upregulating stemness-associated factors and a more aggressive phenotype (45,46). Therefore, in the present study, the CD44⁺/CD24^{-/low} cell population was monitored at each time point (48, 72 and 96 h and 1 week) following hypoxia treatment. As expected, the CD44⁺/CD24^{-/low} cell population expanded following hypoxia treatment (Fig. 1F). Although the peak percentage value varied for each treatment, the induced ratio of CD44⁺/CD24^{-/low} cells by E2, TGF- β 1 and hypoxia decreased over time (Fig. 1). Therefore, breast CSCs with the CD44⁺/CD24^{-/low} phenotype appear to maintain a consistent CSC ratio from external stimuli.

EMT is an essential process in tumor metastasis and recurrence (47). Previous studies have demonstrated that the EMT is tightly linked to CSC biology (27-29). In addition, breast cancer is a distinct cause of mortality and has a high incidence of recurrence (48). Therefore, identifying the mechanisms or molecules used in the EMT is important for breast cancer therapy. In the present study, the E2, TGF- β 1 and hypoxia treatments induced the expression of EMT-associated factors (Fig. 2B). Similarly, the migration ability of MCF-7 cells increased following treatment (Fig. 3). These results indicate that E2, TGF- β 1 and hypoxia may be critical in EMT.

CSCs are hypothesized to demonstrate resistance to chemo- and radiotherapy (31), and ABCG2 is a key regulator of chemoresistance (49). The results of the present study indicated that ABCG2 expression increased significantly while screening target proteins that were induced by treatment with TGF- β 1, E2 and hypoxia (Fig. 2A). The Annexin V analysis demonstrated that the E2, TGF- β 1 and hypoxia treatments attenuated 5-FU-induced apoptosis (Fig. 4A). In addition, treatment with E2, TGF- β 1 or hypoxia effectively blocked ionizing radiation-induced apoptosis (Fig. 4B). These results indicate that CSCs, which have been expanded by TGF- β 1, E2 and hypoxia, may enhance chemo- and radioresistance. Components of the cell cycle machinery regulate asymmetric cell division, cell shape, and protein translation in stem cells (50). In addition, numerous studies have reported that the regulation of cell cycle-associated factors may contribute to inhibit the chemotherapeutic resistance of CSCs (50,51). The present study demonstrated that the expression of cell cycle-associated factors, including β -catenin and c-Myc, increased following treatment with E2, TGF- β 1 and hypoxia (Fig. 2A). The present data suggest that these factors are involved in the regulation of the breast CSC population.

In conclusion, the results of the present study indicate that E2, TGF- β 1 and hypoxia are important in breast CSC expansion and in the regulation of CSC-associated protein expression. In addition, the regulation of hormones, growth factors and the tumor microenvironment, which are potential breast cancer therapeutic targets, may improve the efficacy of breast cancer therapy by inhibiting chemo- and radioresistance.

Acknowledgements

The present study was supported by the National Research Foundation of Korea (DIRAMS) grant, funded by the Korea

government (grant no. 50590-2015), and the Basic Science Research Program of the National Research Foundation of Korea (grant no. NRF-2013 M2A2A 7043665), funded by the Ministry of Science, ICT & Future Planning.

References

- Marmot MG, Altman DG, Cameron DA, Dewar JA, Thompson SG and Wilcox M: The benefits and harms of breast cancer screening: An independent review. *Br J Cancer* 108: 2205-2240, 2013.
- Kaufmann M, von Minckwitz G, Mamounas EP, Cameron D, Carey LA, Cristofanilli M, Denkert C, Eiermann W, Gnant M, Harris JR, *et al*: Recommendations from an international consensus conference on the current status and future of neoadjuvant systemic therapy in primary breast cancer. *Ann Surg Oncol* 19: 1508-1516, 2012.
- Bear HD, Anderson S, Smith RE, Geyer CE Jr, Mamounas EP, Fisher B, Brown AM, Robidoux A, Margolese R, Kahlenberg MS, *et al*: Sequential preoperative or postoperative docetaxel added to preoperative doxorubicin plus cyclophosphamide for operable breast cancer: National Surgical Adjuvant Breast and Bowel Project Protocol B-27. *J Clin Oncol* 24: 2019-2027, 2006.
- EBCTCG (Early Breast Cancer Trialists' Collaborative Group), McGale P, Taylor C, Correa C, Cutter D, Duane F, Ewertz M, Gray R, Mannu G, Peto R, *et al*: Effect of radiotherapy after mastectomy and axillary surgery on 10-year recurrence and 20-year breast cancer mortality: Meta-analysis of individual patient data for 8135 women in 22 randomised trials. *Lancet* 383: 2127-2135, 2014.
- Charafe-Jauffret E, Ginestier C and Birnbaum D: Breast cancer stem cells: Tools and models to rely on. *BMC Cancer* 9: 202, 2009.
- Bonnet D and Dick JE: Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med* 3: 730-737, 1997.
- Glinksy GV: Stem cell origin of death-from-cancer phenotypes of human prostate and breast cancers. *Stem Cell Rev* 3: 79-93, 2007.
- Krivtsov AV, Twomey D, Feng Z, Stubbs MC, Wang Y, Faber J, Levine JE, Wang J, Hahn WC, Gilliland DG, *et al*: Transformation from committed progenitor to leukaemia stem cell initiated by MLL-AF9. *Nature* 442: 818-822, 2006.
- Zhao RC, Zhu YS and Shi Y: New hope for cancer treatment: Exploring the distinction between normal adult stem cells and cancer stem cells. *Pharmacol Ther* 119: 74-82, 2008.
- Stingl J, Eirew P, Ricketson I, Shackleton M, Vaillant F, Choi D, Li HI and Eaves CJ: Purification and unique properties of mammary epithelial stem cells. *Nature* 439: 993-997, 2006.
- O'Brien CA, Kreso A and Jamieson CH: Cancer stem cells and self-renewal. *Clin Cancer Res* 16: 3113-3120, 2010.
- Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ and Clarke MF: Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA* 100: 3983-3988, 2003.
- Chen H, Yan Y, Davidson TL, Shinkai Y and Costa M: Hypoxic stress induces dimethylated histone H3 lysine 9 through histone methyltransferase G9a in mammalian cells. *Cancer Res* 66: 9009-9016, 2006.
- Harris AL: Hypoxia - a key regulatory factor in tumour growth. *Nat Rev Cancer* 2: 38-47, 2002.
- Brown JM and Wilson WR: Exploiting tumour hypoxia in cancer treatment. *Nat Rev Cancer* 4: 437-447, 2004.
- Chaudary N and Hill RP: Hypoxia and metastasis in breast cancer. *Breast Dis* 26: 55-64, 2006.
- Nuyten DS, Hastie T, Chi JT, Chang HY and van de Vijver MJ: Combining biological gene expression signatures in predicting outcome in breast cancer: An alternative to supervised classification. *Eur J Cancer* 44: 2319-2329, 2008.
- Samanta D, Gilkes DM, Chaturvedi P, Xiang L and Semenza GL: Hypoxia-inducible factors are required for chemotherapy resistance of breast cancer stem cells. *Proc Natl Acad Sci USA* 111: E5429-E5438, 2014.
- Aconcia F and Marino M: The effects of 17beta-estradiol in cancer are mediated by estrogen receptor signaling at the plasma membrane. *Front Physiol* 2: 30, 2011.
- Yager JD and Davidson NE: Estrogen carcinogenesis in breast cancer. *N Engl J Med* 354: 270-282, 2006.
- Zheng S, Huang J, Zhou K, Zhang C, Xiang Q, Tan Z, Wang T and Fu X: 17 β -Estradiol enhances breast cancer cell motility and invasion via extra-nuclear activation of actin-binding protein ezrin. *PLoS One* 6: e22439, 2011.
- Li L, Ren C, Yang G, Goltsov AA, Tabata K and Thompson TC: Caveolin-1 promotes autoregulatory, Akt-mediated induction of cancer-promoting growth factors in prostate cancer cells. *Mol Cancer Res* 7: 1781-1791, 2009.
- Kasagi S and Chen W: TGF-beta1 on osteoimmunology and the bone component cells. *Cell Biosci* 3: 4, 2013.
- Prud'homme GJ: Pathobiology of transforming growth factor beta in cancer, fibrosis and immunologic disease, and therapeutic considerations. *Lab Invest* 87: 1077-1091, 2007.
- Akhurst RJ and Derynck R: TGF-beta signaling in cancer - a double-edged sword. *Trends Cell Biol* 11 (Suppl 1): S44-S51, 2001.
- Lv ZD, Kong B, Li JG, Qu HL, Wang XG, Cao WH, Liu XY, Wang Y, Yang ZC, Xu HM, *et al*: Transforming growth factor- β 1 enhances the invasiveness of breast cancer cells by inducing a Smad2-dependent epithelial-to-mesenchymal transition. *Oncol Rep* 29: 219-225, 2013.
- Kong D, Li Y, Wang Z and Sarkar FH: Cancer stem cells and epithelial-to-mesenchymal transition (EMT)-phenotypic cells: Are they cousins or twins? *Cancers (Basel)* 3: 716-729, 2011.
- Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, Brooks M, Reinhard F, Zhang CC, Shipitsin M, *et al*: The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* 133: 704-715, 2008.
- Santisteban B, Reiman JM, Asiedu MK, Behrens MD, Nassar A, Kalli KR, Haluska P, Ingle JN, Hartmann LC, Manjili MH, *et al*: Immune-induced epithelial to mesenchymal transition *in vivo* generates breast cancer stem cells. *Cancer Res* 69: 2887-2895, 2009.
- Longley DB, Harkin DP and Johnston PG: 5-fluorouracil: Mechanisms of action and clinical strategies. *Nat Rev Cancer* 3: 330-338, 2003.
- Soltanian S and Matin MM: Cancer stem cells and cancer therapy. *Tumour Biol* 32: 425-440, 2011.
- Tang C, Ang BT and Pervaiz S: Cancer stem cell: Target for anti-cancer therapy. *FASEB J* 21: 3777-3785, 2007.
- Zhang S, Cui B, Lai H, Liu G, Ghia EM, Widhopf GF II, Zhang Z, Wu CC, Chen L, Wu R, *et al*: Ovarian cancer stem cells express ROR1, which can be targeted for anti-cancer-stem-cell therapy. *Proc Natl Acad Sci USA* 111: 17266-17271, 2014.
- Sakariassen PØ, Immervoll H and Chekenya M: Cancer stem cells as mediators of treatment resistance in brain tumors: Status and controversies. *Neoplasia* 9: 882-892, 2007.
- Rycaj K and Tang DG: Cancer stem cells and radioresistance. *Int J Radiat Biol* 90: 615-621, 2014.
- Pattarozzi A, Gatti M, Barbieri F, Würth R, Porcile C, Lunardi G, Ratto A, Favoni R, Bajetto A, Ferrari A, *et al*: 17beta-estradiol promotes breast cancer cell proliferation-inducing stromal cell-derived factor-1-mediated epidermal growth factor receptor transactivation: Reversal by gefitinib pretreatment. *Mol Pharmacol* 73: 191-202, 2008.
- Liu YF, Wu Q, Xu XM, Ren Y, Yu LN, Quan CS and Li YL: Effects of 17 β -estradiol on proliferation and migration of MCF-7 cell by regulating expression of claudin-6. *Zhonghua Bing Li Xue Za Zhi* 39: 44-47, 2010 (In Chinese).
- Fillmore CM, Gupta PB, Rudnick JA, Caballero S, Keller PJ, Lander ES and Kuperwasser C: Estrogen expands breast cancer stem-like cells through paracrine FGF/Tbx3 signaling. *Proc Natl Acad Sci USA* 107: 21737-21742, 2010.
- Negulescu O, Bognar I, Lei J, Devarajan P, Silbiger S and Neugarten J: Estradiol reverses TGF-beta1-induced mesangial cell apoptosis by a casein kinase 2-dependent mechanism. *Kidney Int* 62: 1989-1998, 2002.
- Silbiger S, Lei J, Ziyadeh FN and Neugarten J: Estradiol reverses TGF-beta1-stimulated type IV collagen gene transcription in murine mesangial cells. *Am J Physiol* 274: F1113-F1118, 1998.
- Kimura H, Braun RD, Ong ET, Hsu R, Secomb TW, Papahadjopoulos D, Hong K and Dewhirst MW: Fluctuations in red cell flux in tumor microvessels can lead to transient hypoxia and reoxygenation in tumor parenchyma. *Cancer Res* 56: 5522-5528, 1996.
- Gillies RJ and Gatenby RA: Hypoxia and adaptive landscapes in the evolution of carcinogenesis. *Cancer Metastasis Rev* 26: 311-317, 2007.
- Li Z and Rich JN: Hypoxia and hypoxia inducible factors in cancer stem cell maintenance. *Curr Top Microbiol Immunol* 345: 21-30, 2010.

44. Liu J and Wang Z: Increased oxidative stress as a selective anti-cancer therapy. *Oxid Med Cell Longev* 2015: 294303, 2015.
45. Liang D, Ma Y, Liu J, Trope CG, Holm R, Nesland JM and Suo Z: The hypoxic microenvironment upgrades stem-like properties of ovarian cancer cells. *BMC Cancer* 12: 201, 2012.
46. Heddleston JM, Li Z, McLendon RE, Hjelmeland AB and Rich JN: The hypoxic microenvironment maintains glioblastoma stem cells and promotes reprogramming towards a cancer stem cell phenotype. *Cell Cycle* 8: 3274-3284, 2009.
47. Voulgari A and Pintzas A: Epithelial-mesenchymal transition in cancer metastasis: mechanisms, markers and strategies to overcome drug resistance in the clinic. *Biochim Biophys Acta* 1796: 75-90, 2009.
48. Cardoso F, Fallowfield L, Costa A, Castiglione M and Senkus E; ESMO Guidelines Working Group: Locally recurrent or metastatic breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 22 (Suppl 6): vi25-vi30, 2011.
49. An Y and Ongkeko WM: ABCG2: The key to chemoresistance in cancer stem cells? *Expert Opin Drug Metab Toxicol* 5: 1529-1542, 2009.
50. Velasco-Velázquez MA, Yu Z, Jiao X and Pestell RG: Cancer stem cells and the cell cycle: Targeting the drive behind breast cancer. *Expert Rev Anticancer Ther* 9: 275-279, 2009.
51. Ravandi F and Estrov Z: Eradication of leukemia stem cells as a new goal of therapy in leukemia. *Clin Cancer Res* 12: 340-344, 2006.