

# Molecular pathophysiology of cerebral edema

Jesse A Stokum<sup>1</sup>, Volodymyr Gerzanich<sup>1</sup> and J Marc Simard<sup>1,2,3</sup>

Journal of Cerebral Blood Flow & Metabolism

2016, Vol. 36(3) 513–538

© Author(s) 2015

Reprints and permissions:

sagepub.co.uk/

journalsPermissions.nav

DOI: 10.1177/0271678X15617172

jcbfm.sagepub.com



## Abstract

Advancements in molecular biology have led to a greater understanding of the individual proteins responsible for generating cerebral edema. In large part, the study of cerebral edema is the study of maladaptive ion transport. Following acute CNS injury, cells of the neurovascular unit, particularly brain endothelial cells and astrocytes, undergo a program of pre- and post-transcriptional changes in the activity of ion channels and transporters. These changes can result in maladaptive ion transport and the generation of abnormal osmotic forces that, ultimately, manifest as cerebral edema. This review discusses past models and current knowledge regarding the molecular and cellular pathophysiology of cerebral edema.

## Keywords

Astrocytes, brain edema, capillaries, cerebrospinal fluid, endothelium

Received 11 September 2015; Revised 21 October 2015; Accepted 22 October 2015

## Introduction

Historically, the goal of brain protection following injury has been to reduce neuronal damage. Edema, the inevitable accompaniment, was considered a secondary event. A renewed interest in edema, its molecular antecedents and its importance in all but the smallest ischemic insults has shifted this paradigm. Strikingly, recent advancements in understanding molecular mechanisms of edema formation suggest that the translation of novel treatments for edema may be closer at hand than the translation of treatments for neuronal demise.

Cerebral edema is a pressing clinical problem. Cerebral edema and brain swelling inevitably accompany ischemic infarcts and intracerebral hemorrhages and, when severe, may increase mortality to nearly 80%.<sup>1</sup> Even in non-life-threatening stroke, the magnitude of brain swelling is strongly predictive of patients' functional outcome.<sup>2</sup> Cerebral edema and brain swelling occur in 20–30% of patients with acute liver failure, and increase mortality to ~55%.<sup>3</sup> Cerebral edema and brain swelling after traumatic brain injury are estimated to account for up to 50% of patient mortality.<sup>4</sup>

Currently approved treatments for cerebral edema—decompressive craniectomy and osmotherapy—were developed prior to any knowledge of modern cerebral edema pathophysiology. These therapies

attempt to manage downstream end-stage events without directly attenuating the underlying molecular mechanisms of cerebral edema. New advances have shed light on heretofore poorly understood cellular and molecular pathophysiology of cerebral edema, and have led to clinical trials of antagonists of key molecular events in cerebral edema formation. It is now understood that cerebral edema evolves in stages, where each stage is characterized by distinct morphological and molecular changes. Cytotoxic edema, or cellular swelling, manifests minutes after acute central nervous system (CNS) injuries. Ionic edema, an extracellular edema that occurs in the presence of an intact blood brain barrier (BBB), forms immediately following cytotoxic edema. Vasogenic edema, an extracellular

<sup>1</sup>Department of Neurosurgery, University of Maryland School of Medicine, Baltimore, USA

<sup>2</sup>Department of Pathology, University of Maryland School of Medicine, Baltimore, USA

<sup>3</sup>Department of Physiology, University of Maryland School of Medicine, Baltimore, USA

### Corresponding author:

Jesse A Stokum, Department of Neurosurgery, University of Maryland School of Medicine, 10 S. Pine St., MSTF 634, Baltimore, MD 21201-1595, USA.

Email: Jesse.Stokum@som.umaryland.edu

edema that includes extravasation of plasma proteins, manifests hours after the initial insult.

This review is intended to serve as a foundation and a reference from which researchers and clinicians might extend their molecular understanding of cerebral edema formation and clearance. Here, the most current cellular and molecular models of cerebral edema formation, transport, and clearance are discussed.

## Models of the cerebral vasculature

### *The blood–brain barrier*

The term BBB refers to a complex of cells that separates the brain interstitium from the luminal contents of the cerebral vasculature (Figure 1). In addition, the term can be used to describe the functional consequence of these cells, namely, the relative independence of brain interstitial fluid (ISF) composition from that of blood plasma, an arrangement that was likely evolved to tightly regulate perisynaptic electrolyte homeostasis, thereby ensuring accurate synaptic integration.<sup>5</sup> Brain ISF, which freely communicates with cerebrospinal fluid (CSF), is optimized for neuronal activity and differs from blood serum in that it contains higher concentrations of  $\text{Cl}^-$  and  $\text{Mg}^{2+}$  and lower concentrations of  $\text{K}^+$ ,  $\text{Ca}^{2+}$ , and  $\text{HCO}_3^-$  (Table 1).<sup>6</sup> The solute composition of plasma and brain ISF in adult humans is shown in Table 1.

Various aspects of the BBB functionality are distributed among its constituent cell types. The innermost layer of the BBB is comprised of a monolayer of endothelial cells that directly contacts the circulating blood. Excepting select fenestrated capillary beds in the circumventricular organs,<sup>7</sup> vertebrate cerebral endothelial cells are interconnected by tight junctions that collectively form the “physical barrier” of the BBB. Notably, interendothelial tight junctions physically divide the brain endothelium plasmalemma into luminal and abluminal membrane faces, which allows for polarized localization of transporters and channels<sup>8</sup> akin to secretory epithelium.<sup>9</sup>

External to the endothelium is a layer of basement membrane, a connective tissue composed of extracellular matrix proteins including collagens, laminins, heparin sulfate proteoglycans, fibronectin, vitronectin, nidogens, perlecan, and agrin.<sup>10</sup> Pericytes, a mesenchymal cell type that contributes to cerebral blood flow regulation,<sup>11</sup> and increases BBB “tightness”,<sup>12,13</sup> are embedded within the endothelial basement membrane at varying intervals along the vessel.

In vessels larger than capillaries, the endothelial basement membrane is bounded externally by the perivascular Virchow Robin space. The Virchow Robin space is a CSF filled extension of the subarachnoid

space that is internally bounded by the endothelial basement membrane and externally bounded by a second, glial, basement membrane. The Virchow Robin space follows penetrating arterioles into the brain parenchyma, becomes fenestrated close to the capillary bed, and eventually disappears at the level of brain capillaries, where astrocyte endfeet directly contact the vessel wall.<sup>14,15</sup>


Astrocyte endfeet, the terminal pads of large astrocyte processes, comprise the outermost layer of the BBB at all levels of the vasculature, including capillaries. Astrocytes are supportive cells that completely fill the brain parenchyma, and in grey matter, are arranged in a three-dimensional matrix with nonoverlapping spatial domains.<sup>16</sup> Nearly all astrocytes extend at least one process that contacts a vessel with an astrocyte endfoot.<sup>16</sup> The parenchymal surface of cerebral vessels are completely covered by a mosaic of astrocyte endfeet that are separated by gaps of approximately 20  $\mu\text{m}$ .<sup>17</sup> The astrocyte endfoot is a specialized membrane domain that is enriched in transporters and channels involved with brain ISF homeostasis.

Recently, the Virchow Robin spaces and the astrocyte endfeet were identified as key anatomical components of the so-called “cerebral glymphatic system”.<sup>18</sup> This system was conceived to account for CSF movements observed in the healthy brain that may function to clear solutes such as amyloid  $\beta$  from the parenchyma and facilitate transport of small lipophilic molecules, particularly during sleep.<sup>18–20</sup> While some controversy exists regarding details of this model,<sup>21</sup> these observations are clearly impactful, in that they give potential insight into the function of the astrocyte endfoot syncytium with regards to brain fluid movements.

### *The neurovascular unit*

The BBB is not a static barrier, but rather it dynamically alters its properties in response to neuronal activity. The term “neurovascular unit” is used to reflect the communication between components of the BBB and cells in the greater brain parenchyma.<sup>22</sup> The neurovascular unit is a system composed of neurons, glia, endothelial cells, vascular smooth muscle, and immune cells that functions in part to trigger the hemodynamic responses to neuronal activity,<sup>23,24</sup> regulate nutrient influx to support neuronal metabolism,<sup>25</sup> and modulate neuronal remodeling.

The term “neurovascular unit” highlights the dependence of neurons upon other central CNS cell types. One example of this dependency is the astrocyte-neuron lactate shuttle.<sup>25</sup> Neurons also rely on astrocytes for neurotransmitter recycling<sup>26</sup> and maintenance of neurons’ antioxidant capabilities through the production of ascorbic acid.<sup>27</sup>



**Figure 1.** Anatomy of cerebral arterioles (top) and capillary (bottom). The innermost layer of arterioles and capillaries is composed of a continuous layer of endothelial cells, linked by tight junctions, and bounded externally by a layer of basement membrane that contains pericytes; arterioles, but not capillaries, travel inside the perivascular Virchow Robin Space (VRS); the outermost layer of the blood brain barrier is composed of astrocyte endfeet, the terminal pads of large astrocyte processes.

## Historical perspectives

### *Historical models of cerebral edema*

In the mid-to-late 1700s, cerebral edema was beginning to be recognized as an entity distinct from acute hydrocephalus, which theretofore was believed to etiologically underlie all cases of excess intracranial water.

This recognition was driven by the observations of Robert Whytt (1714–1766) and George Cheyne (1671–1743) that excess intracranial fluid can occur without enlarged ventricles, a presentation typically accompanied by a “soft” appearing brain.<sup>28,29</sup> Some observers speculated that the brain itself held the excess fluid.

**Table 1.** Mean concentrations of select solutes in plasma and CSF of healthy adult humans.

	Blood plasma <i>a,b</i>	Interstitial fluid/CSF <sup>a,b</sup>
Na <sup>+</sup>	150	147
Cl <sup>-</sup>	99	113
K <sup>+</sup>	4.63	2.86
Mg <sup>2+</sup>	1.61	2.23
Ca <sup>2+</sup>	2.25	1.25
HCO <sub>3</sub> <sup>-</sup>	26.8	23.3
Total protein (mg/dL)	6987.2	39.2
Osmolality (mosm/kg)	289	289
pH	7.397	7.3
pCO <sub>2</sub>	41.1	50.5

<sup>a</sup>Unless otherwise specified, concentrations are shown as mEq/kg.

<sup>b</sup>Adapted from Irani.<sup>261</sup>

Soon after these observations, writings and experiments by Alexander Monro (1733–1817), George Kellie (1720–1779), and John Abercrombie (1780–1844) led to the formation and rise in popularity of the Monro-Kellie axiom, which states that during health, the volume occupied by the contents of the cranium must remain in dynamic equilibrium, the implication being that the fluid influx rate must equal the efflux rate.<sup>30,31</sup> This axiom was an important precursor to models of brain swelling. Unfortunately, excepting a few articles describing brains with a soggy gross appearance, interest in cerebral edema waned until the late 1800s.<sup>32</sup>

Interest in cerebral edema was renewed due to Paul Ehrlich's (1854–1915) identification of the BBB with aniline dyes in 1886, and the revival of the Monro-Kellie axiom. The latter was partially driven by the writings of Harvey Cushing (1869–1939).<sup>33</sup> The growing popularity of these concepts drew attention to the unique anatomy and physiology of cerebral circulation and indicated to contemporaneous researchers that mechanisms of cerebral edema formation were unique from those that drive edema formation in other body regions.

Based upon the gross appearance of brain tissue and data from a newly developed technique to quantify brain swelling, in the late 1910s and early 1920s cerebral edema was subdivided into “brain swelling” and “cerebral edema”, characterized by wet and dry tissue, respectively.<sup>34,35</sup> While debate arose regarding the exclusivity of these subtypes, the controversy remained unresolved until the advent and optimization of electron microscopy techniques.

In 1965, Bakay and Lee applied electron microscopy to describe two different types of cerebral edema in their text *Cerebral Edema*.<sup>36</sup> In 1967, Igor Klatzo (1916–2007) termed these subtypes “cytotoxic” and

“vasogenic”.<sup>37</sup> Vasogenic edema was defined by extravasation of fluid that contained plasma proteins and was attributed to vascular injury. In contrast, cytotoxic edema was characterized by cellular swelling and was attributed to inhibited cell volume regulation. While the latter edema subtype was termed cytotoxic to reflect its occurrence following toxicant exposure, such as triethyltin (TET) poisoning, water intoxication, or cyanide poisoning, cytotoxic edema was also known to occur following ischemia. Importantly, in Klatzo's schema, cytotoxic edema *could* include extravasation of ions and water, but by definition did not include extravasation of plasma protein.

While reviewing these historical models, it is important to note that they are purely phenomenological and offer little in terms of mechanistic explanation. The development of molecular biology in the 1950s and 1960s allowed researchers to probe the molecular drivers of edema formation. Findings from studies utilizing these techniques indicated that all subtypes of cerebral edema, as well as hemorrhagic transformation, share common molecular antecedents.<sup>38</sup> Thus, subtypes of cerebral edema are best viewed as the manifestations of a program of pre- and post-transcriptional molecular events that is ultimately triggered by a brain insult.<sup>38</sup>

#### *Historical approaches to post-ischemic therapeutic intervention*

Excepting neurons in specialized regions, neurons in the adult mammalian brain are arrested in the G<sub>0</sub>-phase of cell-cycle and can be considered to be essentially irreplaceable. Therefore, over the past few decades, acute CNS research has attempted to mediate direct neuroprotection through strategies such as attenuation of excitotoxicity, apoptosis, or oxidative stress. During this time, preclinical work in animal models of acute CNS injury led to the identification of over 1000 new potential neuroprotectants.<sup>39,40</sup> However, this great expenditure of effort, time, and money has essentially failed, as none of these agents has shown effectiveness in clinical trials.<sup>39</sup>

Possible explanations have been offered for the failure to translate promising preclinical findings into the clinic. Some have criticized the commonly used animal models of acute CNS injury, arguing that they do not accurately reflect human disease.<sup>41</sup> Others find fault with the experimental design used in many preclinical studies, arguing that methods like blinding would have prevented many of said false positives.<sup>42</sup> Yet others point out that clinical trials often do not replicate the experimental preclinical studies that appeared so promising.

While model validity and experimental design are clearly important, a more fundamental issue might be

that agents designed to specifically salvage neurons may not abort the death or dysregulation of other components of the neurovascular unit. Neurons are fragile cells and cannot survive without the support of other cell types. Therefore, in addition to direct neuroprotection, a new goal for acute brain injury research is to investigate and attenuate mechanisms of endothelial, astrocytic, and microglial dysfunction and, thereby, create an environment permissible to neuronal survival. It follows that cerebral edema, a phenomenon that arises from dysfunction of astrocytes and endothelium, represents an important target for basic research and therapeutic intervention.

## Core concepts of cerebral edema

### *Cerebral edema and swelling*

The cranial contents are divided into a series of fluid compartments, which are spaces separated by barriers that are relatively impermeable to water and are maintained at homeostatic volumes. Examples of fluid compartments include the vasculature (~100 mL), CSF (~100 mL), brain interstitial space (~100 mL), and brain intracellular space (~1.1 L) (volumes refer to the human brain).<sup>43</sup> The water masses contained by these compartments are dynamic during health; for example, neuronal activity precipitates an increase in the intracellular water mass of local astrocytes.<sup>44,45</sup>

Cerebral edema is a pathological increase in the water mass contained by the brain interstitial space. Incidentally, although cytotoxic edema (oncotic cell swelling) is referred to as “edema” for purely historical reasons, it results in intracellular, rather than extracellular, fluid accumulation, it does not include a swelling component, and it is best regarded as a premorbid precursor to extracellular ionic edema. Transvascular cerebral edema (ionic edema and vasogenic edema) is detrimental because it manifests as brain tissue swelling. Swelling refers to a volumetric expansion of a given mass of tissue and can be generated by the accumulation of tumor, edema, or blood, although here, the focus is on edema.

Brain swelling causes a mass effect that exerts pressure on the surrounding shell of tissue. This pressure increase is magnified by the rigid enclosure of the skull, which places an upper limit on the volume that the brain might expand to. As the brain swells, it exerts mechanical forces on the skull interior, thereby increasing intracranial tissue pressure. When tissue pressure exceeds capillary pressure, capillary lumens collapse, precipitating a feedforward process wherein ischemia of the surrounding shell triggers further edema formation and further swelling in the next shell.<sup>46</sup>

### *Cerebral edema requires perfusion*

For cerebral edema and swelling to occur, the brain tissue must be perfused by an external fluid source. To illustrate this concept, imagine that a fresh biopsy of brain tissue is placed upon a dry surface. As the tissue is completely ischemic, cytotoxic edema will form and water will be redistributed from the interstitium to the intracellular compartment. However, as the tissue is isolated from any possible source of new ions or water mass, the tissue will not become heavier and will not swell. For *in vivo* brain tissue, blood or CSF might serve as the source of this new water, although the relative contributions of these sources are in debate, as is described in the next section.

### *Water sources of ionic cerebral edema*

While the water that drives ionic edema originates ultimately from the vasculature, there exist two major hypotheses regarding the immediate source of the new water mass that is required for brain swelling. In one hypothesis, water moves from the capillary lumen into the parenchyma, driven by osmotic forces, and is conveyed across capillary endothelial cells by mechanisms discussed in later sections of this review. In the context of ischemic stroke, the requirement for perfusion by an outside water source is fulfilled by post-ischemic reperfusion of the core and/or the ischemic penumbra.

The recent description of the glymphatic system led to the formulation of a second hypothesis, whereby CSF serves as the immediate source of ions and water. In this hypothesis, swelling occurs when influx of CSF into the parenchyma is enhanced and/or efflux of ISF is impaired, a situation that precipitates relative accumulation of ISF within the parenchyma.<sup>47</sup> Interestingly, both of these major hypotheses were alluded to in an essay on cerebral edema published in 1894.<sup>32</sup>

In support of the first hypothesis, ionic edema formation is intimately associated with the local blood perfusion status. The post-ischemic reperfusion flow magnitude is positively correlated with edema load.<sup>48</sup> Furthermore, regional brain perfusion is correlated with the spatial magnitude of edema influx: In a rat model of malignant cerebral edema 8 h after permanent middle cerebral artery occlusion, edema fluid is located mostly in peri-infarct regions, with minimal edema fluid in the poorly-perfused core.<sup>49</sup> In human stroke, magnetic resonance imaging (MRI) shows that edema is first found in peri-infarct regions that are actively perfused.<sup>50</sup> In addition, following acute liver failure, cerebral blood flow is positively associated with edema load.<sup>51</sup> However, acceptance of this hypothesis is not universal, as there are uncertainties regarding the

expression level of commonly cited molecular mechanisms for ion and water influx across brain endothelium.<sup>47</sup>

As the second hypothesis was only recently formulated, relatively little literature exists to support or refute it. However, as acute CNS injury does appear to trigger chronic dysregulation of glymphatic clearance of interstitial solutes, this hypothesis merits additional scrutiny.<sup>52</sup>

Two salient features of the BBB indicate that these hypotheses need not be mutually exclusive. Firstly, since brain endothelial cells in the healthy brain mediate high rates of water flux between the vascular compartment and interstitium without net water movements,<sup>53</sup> and thus exhibit high water permeability, the two aforementioned hypotheses could represent steps in a sequential process of water movements from blood to parenchyma. Secondly, the absence of a perivascular CSF-filled space surrounding brain capillaries suggests that their contribution to the glymphatic system may be minimal.<sup>14,15</sup> Thus, these two hypotheses could explain ionic edema formation at different levels of the vascular tree.

Interestingly, the absence of a perivascular CSF-filled space surrounding brain capillaries does not prevent intracisternally-delivered fluorescent tracers from appearing in the capillary basement membrane.<sup>18</sup> In the glymphatic hypothesis, this presumably occurs due to the marker migrating first along the periarteriolar Virchow Robin spaces and then ostensibly by migrating longitudinally through the capillary basement membrane itself. However, it is unclear how the relatively dense pericapillary basement membrane is able to serve as a high-capacity channel for CSF flux, especially since the basement membrane is usually thought to serve as a physical barrier to solute movements.<sup>54</sup> It has been suggested that nonspecific binding of the fluorescent dextran tracer to the capillary basement membrane could underlie the pericapillary fluorescence reported in the abovementioned study.<sup>21</sup>

Importantly, although the aforementioned points accurately reflect our current understanding of the formation of *ionic* edema, the two hypotheses do not account as equally well for the formation of *vasogenic* edema, which is best understood as reflecting the transcapillary flow of plasma proteins.

### Diffusion versus bulk flow

The phenomenon of diffusion, namely, the tendency for molecules to spread from a point of high concentration to surrounding points of lower concentration, arises from random Brownian motions exhibited by particles when suspended in fluid. One property of diffusion is that the probability of finding a molecule at a given

distance from its starting position within a given period of time, essentially the molecule's speed of diffusion, is inversely proportional to the hydrodynamic radius of the molecule. Larger molecules will diffuse more slowly than smaller molecules.

Solutes may also move by bulk flow. In contrast to diffusion, where molecular migration is driven by movement of the molecule within a static substrate, bulk flow migration is driven by the movement of the fluid substrate itself, which in turn is driven by hydrodynamic or osmotic forces. Essentially, suspended particles are "swept along" by the fluid substrate. An important property of bulk flow is that the speed of migration of a suspended particle is dictated solely by the forces driving the migration of the fluid substrate and is therefore independent of the solute's hydrodynamic radius. This property can be exploited to probe whether observed movements are due to diffusion or bulk flow. Parenthetically, bulk flow is utilized in convection-enhanced delivery, an intraparenchymal drug delivery technique that allows for much wider distribution of a given agent than diffusion-driven delivery.<sup>55</sup>

### Cerebral edema moves by bulk flow

The movement of ISF through the brain parenchyma was recognized as early as 1865, when His observed that material injected into the brain parenchyma spreads from the initial point of delivery.<sup>56</sup> In the early 1980s, Cserr et al. observed that radiotracers of varying molecular weights were cleared from the parenchyma with identical rates, indicating that brain ISF in the healthy brain moves by bulk flow rather than diffusion, with white matter tracts exhibiting higher fluid transport.<sup>57,58</sup> Experiments using a cold lesion model of extracellular edema indicated that edema fluid also spreads through the parenchyma through bulk flow rather than diffusion.<sup>59</sup> Interestingly, extracellular fluid appears to take both pericellular and transcellular routes through the parenchyma.<sup>60</sup> Bulk flow of edema fluid is driven by hydrostatic and osmotic forces, produced by mass effect and derangements in ion transport, respectively. Better models of how these forces are generated in vivo are needed to improve our understanding of directional flux of ISF and cerebral edema in vivo.

### Clearance of cerebral edema

The clearance routes of extracellular edema are incompletely understood. Studies examining clearance of ISF indicate that a sizable portion is removed from the parenchyma along perivascular spaces and is deposited into the subarachnoid space, whereupon it is cleared by

mechanisms of CSF absorption, a topic that itself is not without controversy.<sup>18,21,61,62</sup> Historically, the arachnoid villi were considered to mediate most of the absorption of CSF. However, a significant portion of CSF might be cleared through the cervical lymphatics, either through the perineuronal subarachnoid spaces that surround cranial nerves or by passage through the olfactory submucosa.<sup>63</sup> In addition to efflux into the subarachnoid space, a portion of brain ISF is cleared by cerebral capillaries, though their precise contribution remains controversial.

The clearance pathway taken by ISF depends upon its location in the brain: In rats, CSF in the cisterna magna accounts for efflux of 60–70% of midbrain ISF but only 10–15% of caudate nucleus ISF.<sup>64</sup>

### *The transcapillary mechanism and Starling's principle*

In the late 1800s, Starling formulated the basic model for the transcapillary driving forces during edema formation.<sup>65</sup> In his model, edema formation requires two factors: (i) a driving force that either “pushes” or “pulls” substances into or out of tissues; (ii) a “permeability pore” that mediates the transcapillary flux of these substances. In its original formulation, Starling's principle stated that the transendothelial flux of fluid depends upon the net osmotic and hydrostatic pressure and a single permeability coefficient,  $K$ . To reflect mechanisms of cerebral edema formation, Starling's principle was reformulated in 2007, and the permeability coefficient  $K$ , was separated into two constants, the net hydraulic conductivity,  $K_O$ , and the net osmotic conductivity,  $K_H$ , to account for the transcapillary efflux of water ( $J_v$ ).<sup>38</sup>

$$J_v = K_O(\pi_i - \pi_c) + K_H(P_c - P_i)$$

Starling's principle states that the driving force is the sum of the hydrostatic and osmotic pressure gradients. Capillary hydrostatic pressure ( $P_c$ ) is dictated by the precapillary arteriolar pressure, the postcapillary venular pressure, and the capillary resistance, while the tissue pressure ( $P_i$ ) is a function of the volume of the ISF and the tissue compliance. The osmotic pressures of blood ( $\pi_c$ ) and of ISF ( $\pi_i$ ) are a function of the number of particles suspended in each. The ability of the driving forces to actually produce fluid flux then depends on the net hydraulic ( $K_H$ ) and net osmotic ( $K_O$ ) conductivities of the BBB, i.e. the “permeability pores”.

In healthy tissues, both the osmotic term [ $K_O(\pi_i - \pi_c)$ ] and hydrostatic term [ $K_H(P_c - P_i)$ ] are near zero and net water flux is minimal. For the osmotic term, although  $K_O$  is nonzero due to passive endothelial water transport, the vascular and interstitial compartments normally exhibit equal osmolality

(Table 1). For the hydrostatic term, while the hydrostatic gradient is nonzero (for example, in rats, the brain interstitial pressure is approximately 3.43 ± 0.65 mmHg while the brain capillary pressure is approximately 20 mmHg, resulting in a hydrostatic gradient of approximately 16.57 mmHg),<sup>66</sup>  $K_H$  is near zero due to brain endothelium tight junctions.

## **Driving forces for cerebral edema formation**

### *Cytotoxic edema*

Cytotoxic edema (oncotic cell swelling) is a premorbid process whereby cells swell due to influx of osmolites (mainly  $\text{Na}^+$  and  $\text{Cl}^-$ ) and water from the interstitial spaces into the intracellular compartment. This process takes place in all CNS cell types following CNS injury, and is particularly prominent in astrocytes. Astrocyte swelling appears to be a general response of astrocytes to injury and occurs quickly following a variety of types of CNS injury, including ischemia, trauma, hypoglycemia, status epilepticus, and fulminant hepatic failure, though importantly, the mechanisms driving the swelling may differ among etiologies.<sup>67</sup> Importantly, cytotoxic (cellular) edema does *not* generate tissue swelling, as it simply represents a rearrangement of parenchymal water mass and does not involve the addition of new water mass to the brain. Nevertheless, cytotoxic edema is an important initial step in the formation of cerebral edema and swelling, as it generates the driving force for influx of ionic and vasogenic edema, which do cause swelling.

Cellular influx of osmolites may occur due to primary active transport or secondary active transport. Primary active transport requires a continuous supply of adenosine triphosphate (ATP) to provide energy for “pumps” such as the  $\text{Na}^+/\text{K}^+$ -ATPase and  $\text{Ca}^{2+}$ -ATPase. Secondary active transport harnesses the potential energy stored in transmembrane ionic gradients previously generated through primary active transport. Examples of secondary active transporters include ion channels and cotransporters such as the  $\text{Na}^+/\text{K}^+/\text{Cl}^-$  co-transporter (NKCC1) and the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. Following many types of CNS injury, intracellular ATP becomes depleted and thus, mechanisms that are independent of intracellular ATP, like secondary active transport, are more likely to be relevant to the formation of ionic edema.

Ions involved in cytotoxic edema can be conceptually divided into primary drivers and secondary participants. Primary drivers are substances that, through extrusion by primary active transport, are more concentrated outside of the cell than inside. Secondary participants exhibit no pre-existing electrochemical


gradient. However, rearrangement of primary drivers stimulates secondary participants to flux. For example, if  $\text{Na}^+$  is the primary driver,  $\text{Cl}^-$  and water are secondary participants that move in order to maintain electrical and osmotic neutrality.

Per its namesake, astrocytic cytotoxic edema is usually triggered by exposure to endogenous toxins ( $\text{K}^+$ , glutamate,  $\text{H}^+$ ), exogenous toxins (ammonia, TET, cyanide)<sup>68,69</sup> or, unique to the Sur1-Trpm4 (sulfonyleurea receptor 1 – transient receptor potential melastatin 4) channel, intracellular ATP depletion. Maladaptive ion influx may ensue, generating a transmembrane osmotic gradient that drives water influx and causes cell swelling. Cytotoxic edema is one instance of the more general category of astrocyte swelling; the latter includes forms of astrocyte swelling that are not strictly pathological. For example, while hypotonicity, i.e. water intoxication, may trigger astrocyte swelling, this swelling is a purely osmotic phenomenon and lacks the maladaptive ion transport characteristic of cytotoxic edema.

### Constitutively expressed drivers of cytotoxic edema

Swelling due to endogenous brain toxin exposure occurs because astrocytes possess strong homeostatic mechanisms that evolved to maintain the extracellular fluid composition within a range of acceptable values. Following injury, certain molecules normally present in ISF greatly increase in concentration, whereupon astrocytes attempt to maintain ISF homeostasis by activating a variety of normally beneficial secondary active transporters that drive solute transport. In extreme conditions, excessive activation of these secondary active transport mechanisms occurs, leading to massive  $\text{Na}^+$  and water influx and cytotoxic edema (Figure 2).<sup>70</sup> All mechanisms of cytotoxic edema involve  $\text{Na}^+$  overload which, interestingly, is sufficient to impair astrocyte volume regulation<sup>71</sup> perhaps indicative of why astrocytes normally exhibit relatively strong volume regulation,<sup>47,72–74</sup> yet swell so markedly following injury.

The normal ISF  $\text{K}^+$  concentration ranges between 2.7 and 3.5 mM and is maintained by astrocytic  $\text{Na}^+/\text{K}^+$ -ATPase, NKCC1, and Kir4.1.<sup>44,75,76</sup> Following many types of CNS injury, extracellular  $\text{K}^+$  accumulates to dangerous levels, sometimes in excess of 60 mM, due to energy depletion and  $\text{Na}^+/\text{K}^+$ -ATPase failure, cell membrane rupture, or as a byproduct of glutamate excitotoxicity.<sup>77</sup> In the healthy brain, astrocytes function to clear excess extracellular  $\text{K}^+$ , a function that is associated with benign astrocyte swelling.<sup>44,45,78</sup> However, in conditions of greatly increased extracellular  $\text{K}^+$ ,  $\text{K}^+$  clearance triggers cytotoxic edema formation. In the range of 25–117 mM, the



**Figure 2.** Major routes for influx of ions and water in astrocytic cytotoxic edema. Schematic depiction of the major astrocytic transporters and channels that are implicated in the formation of cytotoxic edema; in regards to water transport, single-headed arrows denote water co-transport, while double-headed arrows denote passive water transport.

magnitude of astrocyte swelling becomes linearly related to extracellular  $[\text{K}^+]$ .<sup>79</sup> The bumetanide-sensitive electroneutral NKCC1 transporter, a member of the  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  transporter family, is particularly important to  $\text{K}^+$ -induced astrocyte swelling. NKCC1 is constitutively expressed by astrocytes in all regions of the adult brain and its activity is enhanced after ischemia and acute liver failure due to increased protein expression and phosphorylation.<sup>80–82</sup> NKCC1 carries a net of four osmolites inward per turnover and is capable of water co-transport.<sup>83</sup> In vitro experiments using cultured primary astrocytes demonstrated that NKCC1 contributes to cell swelling in conditions of high extracellular potassium.<sup>84–86</sup> In vivo, swelling is attenuated with bumetanide, an NKCC1 inhibitor, following trauma and ischemia.<sup>87–90</sup>

ISF glutamate in the healthy brain is typically maintained around 10  $\mu\text{M}$ , depending on the region sampled. Following CNS injury such as stroke or trauma, extracellular glutamate can accumulate to greater than 200  $\mu\text{M}$  due to spreading depolarization, synaptic release, or neuronal lysis.<sup>91–93</sup> Astrocyte swelling occurs when extracellular glutamate ranges between 50  $\mu\text{M}$  and 5 mM.<sup>94</sup> Parenthetically, while brain extracellular glutamate is increased after acute liver failure,<sup>95</sup> glutamate is thought to play a minor role in astrocyte swelling after acute liver failure, a process that is driven primarily by ammonia. Glutamate can induce astrocyte swelling through two mechanisms. Firstly, the human excitatory amino acid transporter



(EAAT) glia-specific family members EAAT1 (a.k.a. GLAST in rat) and EAAT2 (a.k.a. Glt-1 in rat) mediate influx of glutamate with co-transport of  $\text{Na}^+$  and water.<sup>96</sup> Both transporters are constitutively expressed by astrocytes in the adult brain.<sup>97</sup> EAAT2 is the major (>90%) contributor to CNS glutamate uptake and homeostasis during health, and forms a multiprotein complex with aquaporin-4.<sup>98–100</sup> Secondly, the metabotropic glutamate receptor 5 (mGluR5),<sup>101</sup> which forms a multiprotein complex with  $\text{Na}^+/\text{K}^+$ -ATPase and aquaporin-4, also has been implicated in glutamate-induced astrocyte swelling.<sup>102,103</sup> Notably, mGluR5 is minimally expressed by resting (nonreactive) adult astrocytes.<sup>104</sup> In line with the notion that many of the mechanisms that drive cytotoxic edema were evolved due to their beneficial actions, if glutamate uptake and glial swelling are inhibited, glutamate mediated neurotoxicity is worsened.<sup>105</sup>

ISF pH is also tightly regulated by astrocytes. Many types of CNS injury interrupt oxygen delivery and lead to ATP depletion, triggering anaerobic metabolism and lactic acid generation that can precipitate a drop in extracellular pH. If ISF pH drops below 6.8, compensatory ion fluxes in astrocytes are sufficient to induce cytotoxic edema.<sup>106,107</sup> Two general classes of astrocyte transporters are involved in pH homeostasis and acidosis-induced astrocyte swelling. Firstly, the constitutively expressed bicarbonate-independent and amiloride sensitive  $\text{Na}^+/\text{H}^+$  exchanger (NHE),<sup>108</sup> which facilitates a 1:1 exchange of intracellular  $\text{H}^+$  for extracellular  $\text{Na}^+$ , mediates acidosis-induced astrocyte swelling in vitro and in vivo.<sup>109–116</sup> Secondly, the constitutively expressed bicarbonate dependent  $\text{Na}^+/\text{HCO}_3^-$  transporter family (NBC) may also contribute to acid-induced swelling.<sup>107,108,117</sup> However, compared to the NHE family, the contribution of the NBC family to cytotoxic edema in vivo is less well understood.

Brain interstitial ammonia is normally  $\sim 100 \mu\text{M}$ ,<sup>118</sup> and can rise above 5 mM during acute liver failure.<sup>119</sup> Astrocyte swelling occurs when cells are exposed to millimolar concentrations of ammonia.<sup>120</sup> Ammonia is internalized by astrocytes and converted to glutamine via glutamine synthetase. It has been hypothesized that glutamine is shuttled to the mitochondria, whereupon mitochondrial phosphate-activated glutaminase (PAG) converts glutamine to glutamate and ammonia, which can trigger production of reactive oxygen species (ROS) and activation of the mitochondrial permeability transition, culminating in astrocyte swelling, i.e. the “Trojan horse hypothesis”.<sup>121</sup> Treatment of astrocytes with ammonia also triggers upregulation and phosphorylation of NKCC1, which mediates ammonia-induced osmolyte influx and astrocyte swelling.<sup>122</sup>

### *De novo expressed drivers of cytotoxic edema*

In contrast to constitutively expressed mechanisms of cytotoxic edema, which can be viewed as maladaptive versions of normally beneficial processes, the Sur1-Trpm4 channel is expressed in the CNS only following injury, and is absent from healthy brain. Trpm4, a constitutively expressed monovalent cation channel that opens in response to intracellular  $\text{Ca}^{2+}$ , is the pore forming subunit of Sur1-Trpm4.<sup>123–125</sup> In all CNS cells, CNS injury triggers activation of the hypoxia-inducible factor 1 (HIF1) transcription factor, which induces *de novo* expression of Sur1,<sup>126</sup> an ATP-binding cassette, which associates with Trpm4 and doubles its  $\text{Ca}^{2+}$  sensitivity and sensitizes the channel to ATP depletion.<sup>123,125,127</sup> It is hypothesized that this channel evolved to protect cells against  $\text{Ca}^{2+}$  influx in less extreme types of CNS injury by allowing  $\text{Na}^+$  influx, membrane depolarization, and thus a reduction of the inward driving force for  $\text{Ca}^{2+}$ . However, in the context of severe CNS injury and ATP depletion, excessive influx of  $\text{Na}^+$  through Sur1-Trpm4 drives maladaptive cell swelling and cytotoxic edema (Figure 2). Inhibition of Sur1-Trpm4 with sulfonyleurea drugs prevents cytotoxic edema in vitro and reduces edema in vivo following ischemia and trauma.<sup>49,123,128,129</sup> Evidence also suggests that the Sur1-Trpm4 channel contributes to ammonia-induced astrocyte swelling in vitro and in vivo.<sup>130</sup>

### *Routes for transmembrane water flow during cytotoxic edema*

The influx of ions generates a transmembrane osmotic gradient that favors the influx of water. Water might flow into astrocytes through three routes. Firstly, simple diffusion through the lipid bilayer can account for significant water influx; however, this route is low capacity and not thermodynamically favored. Secondly, transmembrane water channels, including the aquaporin family as well as certain astrocytic transporters such as SGLT1, GLUT1, and GLUT2 possess passive water permeable pores, where water flux is driven by an osmotic gradient.<sup>83</sup> Thirdly, certain ion transporters expressed by astrocytes that drive ion fluxes during cytotoxic edema also mediate secondary water co-transport by carrying a fixed number of water molecules with their ionic cargo per turnover. Examples of these transporters include NKCC1 and the glutamate transporters EAAT1 and EAAT2.<sup>83</sup>

### *Aquaporin channels in the CNS*

Recently, aquaporin channels have been recognized as important mediators of plasmalemmal water fluxes in cerebral edema. In 1992, Peter Agre described a novel

molecular conduit for bulk water flow through the plasma membrane, a protein later christened aquaporin-1.<sup>131</sup> Aquaporin monomers constitute the functional water channel subunit of aquaporin channels and are exquisitely selective to water by virtue of a dumb-bell-shaped pore with an amphipathic bottleneck.<sup>132</sup> Aquaporin water transport is passive and bidirectional; the rate and directionality of aquaporin mediated water flux is determined exclusively by the transmembrane osmotic pressure gradient.

While there are 14 known aquaporin channels, only aquaporin-1, aquaporin-4, aquaporin-9, and aquaporin-11 are expressed in the CNS.<sup>133,134</sup> Of these, aquaporin-4 is the major aquaporin expressed by astrocytes, and is the dominant contributor to cerebral edema formation and clearance. In the prefrontal cerebrum under normal conditions, aquaporin-4 expression is strongly localized to the perivascular astrocyte endfoot; astrocyte soma and main processes do not exhibit notable aquaporin-4 immunoreactivity.<sup>135</sup> Aquaporin-4 also is localized to the submeningeal astrocyte endfeet and in the leading lamellipodia of migrating astrocytes.<sup>136</sup> While aquaporin-4 is mostly astrocyte-specific, it has been reported to be upregulated in microglia following LPS injection.<sup>137</sup>

Aquaporin-4 is expressed in two major N-terminal splice variants, the M1 (323 amino acids) and M23 (301 amino acids), as well as four other alternative isoforms whose functional significance is still being investigated.<sup>138–141</sup> The M23 isoform is able to multimerize on the plasmalemma into large complexes called orthogonal arrays of intramembraneous particles (OAPs) that, on the luminal face of the astrocyte endfoot, exhibit a density of 500–600/ $\mu\text{m}^2$  and occupy approximately 50% of its surface area.<sup>135,142,143</sup>

The precise contribution of the astrocyte endfeet and aquaporin-4 to brain ISF during health is currently being investigated. Aquaporin-4 represents a high capacity route of water flux past the astrocyte endfoot layer of the BBB; other routes include paracellular water flux, co-transport of water, or simple diffusion.<sup>70</sup> Aquaporin-4 may then be necessary to rapidly neutralize osmotic gradients that are generated during ion transport, akin to its role in skeletal muscle during contraction.<sup>144</sup> This might explain the dependence of the aforementioned glymphatic system on aquaporin-4.<sup>18</sup> In addition, aquaporin-4 knockout animals exhibit CNS abnormalities linked with altered ion transport, such as expansion of the extracellular space, increased brain water content, cochlear deafness, and increased seizure threshold.<sup>145–148</sup>

### *Aquaporin channels rely on ion transport*

As passive channels, aquaporins are completely dependent upon the activity of ion transporters for

water flux. Therefore, the study of aquaporin-dependent cerebral edema is essentially the study of ion transport.<sup>70</sup> While aquaporins alone are undoubtedly important in the generation of cerebral edema, future work will address the interaction between ion transport and aquaporin water flux.<sup>149</sup>

### *Role of aquaporin-4 in cytotoxic edema*

Aquaporin-4 worsens subtypes of cerebral edema that form in the context of an intact BBB.<sup>150–154</sup> Mice with aquaporin-4 knockout or mislocalized aquaporin-4 exhibit reduced astrocyte swelling during water intoxication, suggesting a role for aquaporin-4 in cytotoxic edema of astrocytes.<sup>153,155,156</sup> Notably, the alternative routes of astrocyte plasmalemma water influx are physiologically and pathologically important, as even with knockout of aquaporin-4, astrocytes still swell quickly in response to hypoosmotic challenge.<sup>157</sup>

### *Cytotoxic edema generates driving forces*


The influx of primary drivers like  $\text{Na}^+$  and secondary participants like  $\text{Cl}^-$  and water into cells during cytotoxic edema depletes these constituents from the extracellular space.<sup>158,159</sup> Sequestration of these constituents is possible because the intracellular compartment is much larger than the extracellular compartment, which only comprises 12–19% of total brain volume.<sup>160</sup> Cytotoxic edema thereby generates a new  $\text{Na}^+$  gradient across the BBB, where  $\text{Na}^+$  concentration becomes higher in the vascular compartment compared to the interstitial compartment. For example, in one study that used a rat model of global ischemia, extracellular  $[\text{Na}^+]$  declined from a baseline of 141 mM to 74 mM; as plasma  $[\text{Na}^+]$  was 134 mM, cytotoxic edema generated a transendothelial  $\text{Na}^+$  concentration differential of approximately 60 mM.<sup>158</sup>

This newly formed  $\text{Na}^+$  gradient is preserved even if cells lyse and release their intracellular contents because the extracellular space is much smaller than the intracellular space and because  $\text{K}^+$  will remain mostly bound to intracellular proteins and macromolecules.<sup>161</sup> The  $\text{Na}^+$  gradient generated by cytotoxic edema serves as a source of potential energy that drives subsequent influx of ionic edema fluid.

## **Endothelial dysfunction**

### *The endothelial permeability pore*

Acute CNS injury triggers a program of pre- and post-transcriptional molecular changes in the neurovascular unit that results in the formation of endothelial “permeability pores” and subsequent loss of BBB integrity.



**Figure 3.** Phases and select mechanisms of endothelial dysfunction. In ionic edema, water flux (blue arrows) and ion flux (grey arrows) are mediated by plasmalemma channels and transporters; vasogenic edema, which includes extravasation of plasma proteins, but not erythrocytes, is mediated by transcellular channels, MMP degradation of tight junctions, and endothelial retraction, phenomena that are, in part, triggered by VEGF, Ang2, and CCL2 signaling; hemorrhagic transformation occurs due to structural failure of the vessel, driven by either complete degradation of tight junctions or Sur1-Trpm4-mediated oncotic cell death of endothelial cells.

Progressive endothelial dysfunction can be organized into three phases (ionic edema, vasogenic edema, and hemorrhagic transformation), based upon the principle substances that undergo transcapillary movement (Figure 3).

The three phases of endothelial dysregulation are thought to occur sequentially, although the rapidity of transition between phases probably depends on injury type and severity. Furthermore, as many etiologies of brain endothelial dysregulation and cerebral edema are focal in nature, brain tissue usually exhibits a complex spatiotemporal pattern of the different phases of endothelial dysregulation.

### First phase: Ionic edema


During ionic edema, the potential energy contained in the transendothelial  $\text{Na}^+$  gradient generated by cytotoxic edema drives extravasation of osmolites and water.  $\text{Na}^+$  is transported inward along its concentration gradient by brain endothelial cells, resulting in the accumulation of  $\text{Na}^+$  in the brain parenchyma and the generation of a nonzero osmotic driving force (the  $(\pi_c - \pi_i)$  term in Starling's principle).<sup>162</sup> This influx of  $\text{Na}^+$  requires blood perfusion, supportive of the hypothesis that the vascular compartment serves as the source of ionic edema.<sup>163</sup> As secondary participants,  $\text{Cl}^-$  and water follow  $\text{Na}^+$  inward to maintain electrical and osmotic neutrality, resulting in the formation of ionic edema.<sup>164</sup> While the relative contributions of brain arterioles, venules, and capillaries to ionic edema formation has not been experimentally tested, ionic edema may form

primarily at the capillary level due to the relatively thin walls and large surface area of the capillary bed.

Ionic edema and vasogenic edema are forms of extracellular edema that differ in an important way. Vasogenic edema, but not ionic edema, includes extravasated serum proteins. The preservation of BBB integrity has two important implications for mechanisms of ionic edema. Firstly, ion influx during ionic edema is exclusively mediated by endothelial ion channels and transporters; physical disruptions to the BBB, such as occurs with reverse pinocytosis or degradation of tight junctions, do not represent viable pathways for ion flow, as they would inevitably include plasma proteins. Secondly, preservation of the BBB integrity implies that  $K_H$  remains zero. The dynamics of ionic edema are governed only by the osmotic term in Starling's principle.

Parenthetically, while a subtype of extracellular edema that lacked serum protein extravasation has long been recognized, ionic edema was only recently defined as a distinct cerebral edema subtype.<sup>38</sup> Prior to this distinction, all edema subtypes not involving serum protein extravasation were grouped under the common term "cytotoxic edema", which therefore encompassed cytotoxic edema, as the term is used presently, as well as ionic edema.

For ionic edema to form,  $\text{Na}^+$ ,  $\text{Cl}^-$ , and water must first be transported inward through the luminal membrane and then transported through the abluminal membrane of brain capillary endothelial cells (Figure 4). Therefore, ionic edema is essentially a two-step transport process. Given that many brain endothelial channels and transporters exhibit a



**Figure 4.** Major routes for influx of ions and water in ionic edema. Schematic depiction of the major endothelial transporters and channels that have been implicated in the formation of ionic edema; in regards to water transport, single-headed arrows denote water co-transport, while double-headed arrows denote passive water transport.

polarized distribution at these membrane faces, the transmembrane routes taken by ions and water differ between the luminal and abluminal membrane. Similar to cytotoxic edema, ion flux is driven by either primary active transport or secondary active transport.

As ionic edema involves, in part, the uptake of ions and water by brain endothelial cells, it is analogous to cytotoxic edema, but differs in that the ion and water fluxes are polarized. Specifically, channels on the luminal membrane of brain endothelial cells drive the cellular uptake of vascular ions and water, which may manifest as endothelial cell swelling. Endothelial swelling is then “relieved” by channels on the abluminal membrane, which permit efflux of ions and water into the brain interstitium, thereby completing the transcapillary flux of ions and water.

### Constitutively expressed drivers of ionic edema

The sodium-hydrogen antiporter (NHE) family members, NHE1 and NHE2, are constitutively expressed on both luminal and abluminal membranes of brain endothelium. During ionic edema, NHE family members contribute to  $\text{Na}^+$  influx across the luminal membrane.<sup>165</sup>  $\text{Na}^+/\text{H}^+$  exchange in vitro is stimulated by hypoxia and hypoglycemia, conditions that occur following cerebral ischemia.<sup>165</sup> Intravenous delivery of  $\text{Na}^+/\text{H}^+$  exchange inhibitors reduce cerebral edema formation in a rat stroke model, putatively by attenuating luminal NHE activity.<sup>113,166,167</sup>

The cation-chloride transport family member, NKCC1, is constitutively expressed on the luminal

membrane of brain endothelium, is upregulated and activated via phosphorylation in response to ischemia, and contributes to  $\text{Na}^+$  influx across the luminal membrane during ionic edema formation.<sup>90,168</sup> In addition to mediating solute influx that osmotically drives water influx, NKCC1 also mediates secondary active transport of water at 590 molecules of  $\text{H}_2\text{O}$  per transporter turnover, thereby allowing it to pump water up an osmotic gradient.<sup>83</sup> Influx of ionic edema is attenuated with intravenous delivery of bumetanide, an NKCC1 inhibitor, thus supporting the role of NKCC1 in ionic edema formation.<sup>87,88</sup>

Under conditions of adequate energy,  $\text{Na}^+$  efflux across the abluminal membrane is primarily mediated by the  $\text{Na}^+-\text{K}^+-\text{ATPase}$ , a primary active transporter that is selectively expressed on the abluminal membrane of brain endothelial cells.<sup>9</sup> Notably, as this efflux route depends on ATP, its contribution is likely minimal during severe energy depletion such as during ischemia, although it may become relevant with timely reperfusion. In addition, the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger exists on the abluminal membrane and might contribute to  $\text{Na}^+$  efflux by virtue of its ability to operate in reverse-mode where  $\text{Na}^+$  is expelled in exchange for  $\text{Ca}^{2+}$  influx.<sup>169,170</sup> As this efflux route is ATP-independent,  $\text{Na}^+/\text{Ca}^{2+}$  mediated  $\text{Na}^+$  efflux might be particularly relevant during total ischemia.

### De novo expressed drivers of ionic edema

The Sur1-Trpm4 channel is upregulated by capillary, arteriole, and venule endothelial cells in response to

CNS injury,<sup>127</sup> and contributes to the formation of ionic edema by mediating  $\text{Na}^+$  influx at the luminal membrane and  $\text{Na}^+$  efflux at the abluminal membrane. Post-ischemic blockade of the channel by glibenclamide, a potent Sur1 antagonist,<sup>171–173</sup> reduces edema formation by 50%, indicating that Sur-Trpm4 plays a key role in transcapillary  $\text{Na}^+$  influx that occurs during ionic edema. Parenthetically, as glybenclamide penetrates the BBB in ischemic brain tissue,<sup>49</sup> and Sur1 is expressed after injury by all CNS cells, its effects upon edema formation may not be solely due to inhibition of luminal endothelial Sur1.<sup>174</sup> As Sur1 is not constitutively expressed, this mechanism is injury-specific. Furthermore, as this mechanism relies on transcriptional gene expression and hence ATP, it is most relevant in ischemic, but still perfused vessels.

A disease-specific molecular driver of cytotoxic and ionic edema, the Sur1-Trpm4 channel is a highly promising target for pharmacological inhibition. Notably, the glyburide advantage in malignant edema and stroke (GAMES-RP) trial, a randomized phase II trial that evaluated the efficacy of an intravenous formulation of glybenclamide (CIRARA or RP-1127) in patients with large territory ischemic stroke,<sup>175</sup> was recently concluded. Adverse drug reactions are most commonly due to unwanted effects upon normal function that are mediated by the drug's primary pharmacological mechanism. For example, inhibition of the *N*-methyl-D-aspartate receptor (NMDAR), a protein that is constitutively expressed throughout the brain, is associated with adverse CNS drug effects such as dizziness, sedation, agitation, hallucination, and confusion.<sup>176</sup> These adverse effects have led many researchers to abandon NMDAR antagonism as a strategy to combat excitotoxic injury following ischemia. However, as Sur1-Trpm4 is only expressed in the injured brain, CNS effects resulting from the drug's primary pharmacology will be specific to injured brain tissue, and adverse drug effects will be minimized.

### *Transendothelial routes for $\text{Cl}^-$ and water during ionic edema*

$\text{Na}^+$ , the primary driver of ionic edema, drives the influx of secondary participants like  $\text{Cl}^-$  and water in order to equalize electrical and osmotic gradients. Transendothelial  $\text{Cl}^-$  flux is likely mediated by  $\text{Cl}^-$  channels and  $\text{Cl}^-$  co-transporters such as NKCC1 and KCC.<sup>177</sup>

As during cytotoxic edema, there are three possible routes for water transport across the plasmalemma; however, some of the molecular details are specific to brain endothelium (Table 2). Firstly, simple diffusion can occur across the endothelial plasma membrane. While the water permeability of a small patch of

endothelium is low, the surface area of the brain capillary bed is quite large, and could permit nontrivial net flux. Secondly, secondary co-transport of water can be mediated by NKCC1 at the luminal face,<sup>83,84</sup> and by the  $\text{K}^+/\text{Cl}^-$  co-transporter KCC at the abluminal face.<sup>178</sup> In addition to NKCC1 and KCC, all solute transporters that co-transport water can be driven to operate in response to an osmotic gradient. Thus, endothelial transporters such as MCT1, GAT-1 and EAAT-1, which, unlike NKCC1, do not contribute to ionic edema related  $\text{Na}^+$  influx, are able to contribute to water influx during ionic edema.<sup>83</sup> Thirdly, brain endothelial cells express certain membrane proteins that can mediate passive water transport. Some studies report that brain endothelial cells weakly express certain members of the aquaporin family such as aquaporin-1.<sup>179</sup> However, given their low abundance in endothelial cells, aquaporins are unlikely to mediate the bulk of endothelial water transport across the endothelium. Rather, the highly abundant glucose transporters, sodium-glucose linked transporter 1 (SGLT1) and glucose transporter 1 (GLUT1) might represent a major route for transendothelial water flux, as they have the ability to passively transport water independently of glucose, analogously to aquaporin channels.<sup>83,180,181</sup> While the single protein water permeability of GLUT1 is less than 1% than that of aquaporin-4,<sup>83</sup> GLUT1 is highly expressed by brain endothelium. Interestingly, the estimated net contribution of GLUT1 to passive water permeability of endothelium ( $0.5 \times 10^{-3}$  cm/s) is remarkably close to the measured total passive water permeability of the BBB ( $1 \times 10^{-3}$  cm/s).<sup>83,182</sup> GLUT1 and GLUT2 are expressed at both luminal and abluminal faces, while SGLT1 is expressed at the luminal face.<sup>180,181</sup>

### *Role of the astrocyte endfoot and aquaporin-4 in ionic edema*

Aquaporin-4 worsens ionic edema, a subtype of cerebral edema that occurs in the context of an intact BBB.<sup>150–154</sup> Conversely, overexpression of aquaporin-4 enhances ionic edema formation.<sup>183</sup> Interestingly, following cerebral ischemia, aquaporin-4 is upregulated primarily by white matter astrocytes.<sup>184</sup> Given that white matter can exhibit greater swelling after ischemia than grey matter,<sup>184–186</sup> these data indicate that white matter may play an underappreciated role in ionic edema formation and brain swelling.

Recently, it was postulated that a primary function of the astrocyte endfoot syncytium, and of aquaporin-4, is to mediate transglial water flux.<sup>18</sup> It was suggested that dysregulation of this function occurs following CNS injury, and results in reduced glymphatic flow, which is thought to represent a primary mechanism

**Table 2.** Routes for transendothelial water influx during cerebral edema formation.<sup>a</sup>

Endothelial dysfunction phase	Route	Mediator	Single-channel $L_p$ <sup>b</sup>	Oocyte $P_f$ <sup>c</sup>	Water co-transport per turnover <sup>d</sup>	
Ionic edema	Transmembrane diffusion	Lipid bilayer	– <sup>e</sup>	$\sim 1 \times 10^{-4}$ <sup>262</sup>	–	
	Channel mediated passive diffusion and water co-transport	Aquaporin-1 <sup>f</sup>	$4.9 \times 10^{-14}$ <sup>263</sup>	$1.9 \times 10^{-2}$ <sup>264</sup>	–	
		Aquaporin-4 <sup>f</sup>	$1.5 \times 10^{-13}$ <sup>263</sup>	$2.9 \times 10^{-2}$ <sup>264</sup>	–	
		SGLT1	$1.5 \times 10^{-15}$ <sup>265</sup>	$1.6 \times 10^{-4}$ <sup>265</sup>	260 <sup>266</sup>	
		GLUT1	$2 \times 10^{-15}$ <sup>83</sup>	$2.8 \times 10^{-3}$ <sup>267</sup>	40 <sup>83</sup>	
		GLUT2	ND <sup>g</sup>	$4.8 \times 10^{-4}$ <sup>268</sup>	35 <sup>268</sup>	
		GAT-1	$1.4 \times 10^{-14}$ <sup>269</sup>	$1.002 \times 10^{-4}$ <sup>270</sup>	330 <sup>271</sup>	
		EAAT1	$7 \times 10^{-15}$ <sup>272</sup>	$1.006 \times 10^{-4}$ <sup>272</sup>	436 <sup>272</sup>	
		NKCC1	–	–	590 <sup>273</sup>	
		KCC4	–	–	500 <sup>178</sup>	
		MCT1	–	–	500 <sup>83</sup>	
	Vasogenic edema	Uncoupling of tight junctions	Claudins, Occludins, Zo-1, Actin			
		Endothelial retraction	Actin			
Pinocytic vesicle fusion		ND				
Hemorrhagic conversion	Endothelial oncotic cell death	Sur I-Trpm4				
		Reactive oxygen species				
		Leukocyte transmigration				
	Tight junction degradation	Endothelial activation				
		MMPs				

<sup>a</sup>Portions of Table 2 are adapted from MacAulay and Zeuthen.<sup>83</sup> <sup>b</sup>Expressed as the hydraulic permeability coefficient ( $L_p$  with units of  $\text{cm}^3/\text{s}$ ). <sup>c</sup>Expressed as the diffusive water permeability coefficient ( $P_f$  with units of  $\text{cm sec}^{-1}$ ) of unmodified *Xenopus* oocytes (first row), or *Xenopus* oocytes expressing a particular water-permeable channel. Note that  $P_f$  is dependent upon the abundance of the expressed water channel and thus measurements of single-channel  $L_p$  is not directly translatable to plasmalemma  $P_f$  measurements. In addition, note that  $P_f$  is surface-area independent and is related to  $L_p$  through the equation:

$$P_f = \frac{L_p RT}{\bar{V}_w A}$$

where  $A$  is the membrane area,  $\bar{V}_w$  is the partial molar volume of water,  $R$  is the gas constant, and  $T$  is the temperature.<sup>274</sup> <sup>d</sup>Expressed as number of water molecules co-transported per transporter turnover. <sup>e</sup>Not applicable (–). <sup>f</sup>Aquaporin channels are minimally expressed by brain endothelium. <sup>g</sup>Not determined (ND).

driving ionic edema formation.<sup>47</sup> While this hypothesis might account in part for ionic edema, a strict adherence to this model would imply that the brain endothelium has little, if any, contribution to cerebral edema formation. Omitting a key role for endothelium is implausible, given the many endothelial transporters and channels implicated in the formation of edema. It is more likely that, in addition to its hypothesized role in glymphatic-mediated edema, aquaporin-4 interacts with and potentiates the endothelial water fluxes that drive ionic edema formation. Unfortunately, without a greater understanding of how aquaporin-4 controls water and solute flux through the endfoot layer of the BBB, it is unclear how aquaporin-4 water transport might affect endothelial ion transport.

### Second phase: Vasogenic edema

Vasogenic edema is a form of extracellular edema characterized by breakdown of the BBB, wherein a transendothelial permeability pore forms that permits extravasation of water and plasma proteins such as albumin and IgG into the brain interstitial compartment. Unlike hemorrhage, capillary structural integrity is maintained during vasogenic edema such that passage of erythrocytes is prohibited. Therefore, vasogenic edema is best viewed as a cell-free blood ultrafiltrate, i.e. plasma.<sup>37,187</sup> The permeability pore that allows the passage of solutes during vasogenic edema likely has contributions from more than one mechanism (see below). While all levels of the vascular tree contribute

to vasogenic edema formation, brain capillaries are a particularly major contributor.<sup>188</sup>

Once physical communication between the vascular and interstitial compartments is established, microvessels behave like fenestrated capillaries and therefore both hydrostatic and osmotic pressure gradients can affect edema formation, although hydrostatic pressure represents the primary driving force for vasogenic edema formation.<sup>189</sup> Determinants of the hydrostatic pressure gradient, such as intracranial pressure, systemic blood pressure, capillary occlusion, and vasospasm, are important to vasogenic edema dynamics. Determinants of the osmotic pressure gradient, which now include all osmotically active molecules such as  $\text{Na}^+$  and proteins, also can influence water flux.

The influence of hydrostatic pressure on vasogenic edema has direct clinical implications. For example, systemic blood pressure must be kept high enough to maintain brain perfusion, but in excess will promote edema formation.<sup>189,190</sup> In addition, intracranial pressure must be kept low enough to maintain tissue perfusion, but high enough to counteract edema influx.<sup>189</sup> Optimization of these parameters is a difficult, multifactorial problem.

The concept that only the osmotic forces influence ionic edema, while both osmotic and hydrostatic gradients influence vasogenic edema may help to explain the mixed outcomes that occur following decompressive craniectomy, a procedure that abruptly lowers intraparenchymal pressure.<sup>191,192</sup> Decompression is safe if done early, during the ionic edema stage, as it aids in the restoration of tissue perfusion. However, if done later during vasogenic edema, decompression will decrease tissue pressure, thereby increasing the hydrostatic gradient and driving edema influx.<sup>193,194</sup>

### *Mechanisms of vasogenic edema*

Protein and water may passage from the vascular compartment to the interstitial compartment through transendothelial channels formed by dysregulation of pinocytosis. Pinocytosis, a process whereby blood solutes are enveloped by luminal endothelial membrane ruffles,<sup>195</sup> are transported across the cytoplasm, and are released at the abluminal membrane.<sup>196</sup> Following CNS injury, pinocytic vesicles that are capable of carrying solutes and water have been observed to fuse and form transendothelial channels that span from the luminal to the abluminal membrane.<sup>197,198</sup> Controversy still exists regarding this mechanism.<sup>199</sup>

It is generally agreed that vasogenic edema can form via paracellular transport past endothelial cells. Inflammation and cerebral ischemia can trigger actin-dependent endothelial cell rounding or retraction and increased endothelial permeability.<sup>200</sup> Endothelial

retraction is an ATP-dependent process that can be triggered by thrombin,<sup>200–203</sup> a protease that is upregulated following ischemia and is highly abundant in the brain parenchyma following intracerebral hemorrhage. However, as some evidence suggests that in lieu of tight junction disruption, endothelial retraction is not sufficient to impair barrier resistance,<sup>204</sup> this mechanism might serve to enhance rather than initiate vasogenic edema formation. It has been speculated that endothelial cell retraction might have been evolved to facilitate transmigration of leukocytes that contribute to the beneficial clearance of necrotic debris produced by many types of CNS injury.<sup>205</sup>

Paracellular permeability pores also can be generated by vascular endothelial growth factor (VEGF) signaling. Brain injury triggers expression of VEGF,<sup>206–208</sup> which triggers decreased expression of tight junction proteins,<sup>209,210</sup> uncoupling of interendothelial tight junctions, increased hydraulic permeability of vessels<sup>211</sup> and promotes edema formation.<sup>212</sup> Inhibition of VEGF reduces edema associated with post-ischemia reperfusion and brain tumors.<sup>213,214</sup> Like many mechanisms that produce cerebral edema, VEGF signaling is not purely maladaptive, but rather is linked to angiogenesis. If administered early following experimental stroke in rats, recombinant VEGF increases edema formation, but if given at later times, VEGF stimulates angiogenesis in the penumbra and improves neurological recovery.<sup>215</sup> As some studies have shown that the extent of angiogenesis is correlated with survival following ischemic stroke, angiogenesis might be an important factor in recovery following ischemia.<sup>216</sup> In addition to VEGF, a host of signaling molecules including CCL2, angiopoietin 2 (Ang2), and nitric oxide are released following injury, and can inhibit expression of tight junction proteins and thereby exacerbate vasogenic edema formation.<sup>217–221</sup>

Endothelial basement membrane proteins and tight junction proteins also can be lost following CNS injury through protease degradation. Following injury, matrix metalloproteinase (MMP) activity increases through *de novo* expression and activation of latent MMPs, resulting in degradation of basement membrane and tight junction proteins.<sup>222–227</sup> MMP inhibitors reduce ischemia- and reperfusion-associated cerebral edema,<sup>36,228,229</sup> partially by preventing degradation of tight junction proteins.<sup>230</sup>

### *Role of the astrocyte endfoot and aquaporin-4 in vasogenic edema*

In contrast to its role in cytotoxic edema or ionic edema, knockout of aquaporin-4 is associated with *worsened* edema following injuries that precipitate vasogenic edema formation, such as trauma or cold

lesion.<sup>148</sup> These data suggest that the astrocyte endfoot and aquaporin-4 contributes to the clearance of vasogenic edema, and likely mediates the clearance of other forms of extracellular edema, such as ionic edema.

### *Third phase: Intracerebral hemorrhage*

Intracerebral hemorrhage can occur as a primary injury, as in the context of traumatic brain injury, or as secondary injury, where its formation is due to downstream injury-related mechanisms. The latter, also referred to as hemorrhagic conversion or hemorrhagic transformation, represents the third and final phase of endothelial dysfunction, where the structural integrity of the capillary is lost, allowing extravasation of all constituents of blood, including erythrocytes and other cells. Up to 30–40% of ischemic strokes undergo hemorrhagic transformation, a phenomenon that accounts for approximately 26–154 additional deaths per 1000 patients.<sup>231–233</sup>

Like vasogenic edema, hydrostatic pressure is the primary driving force for hemorrhagic transformation. Interestingly, extravasated blood increases the local tissue hydrostatic pressure and should thus impair further hemorrhage. However, this meager benefit is outweighed by the mass effect and tissue distortion created, as well as the robust inflammatory response triggered by blood products such as methemoglobin. Implications for clinical management of hemorrhage are similar to vasogenic edema, but this challenge is much more difficult, given the inflammatory milieu created by hemorrhage.

### *Mechanisms of hemorrhagic transformation*

Mechanisms that drive hemorrhagic transformation are complicated, multifactorial, and incompletely understood. The aforementioned mechanisms of vasogenic edema formation are likely also relevant to hemorrhagic transformation. For example, exogenous VEGF administration following reperfusion worsens hemorrhagic transformation. MMP-driven extracellular proteolysis appears to play a major role in hemorrhagic transformation as its inhibition reduces hemorrhage.<sup>224,234–237</sup> In addition to these shared mechanisms, there exist several mechanisms that are specific to hemorrhagic transformation. Oncotic death of endothelial cells mediated by Surl-Trpm4 is likely an important factor in hemorrhagic transformation after a variety of CNS injuries.<sup>49,238,239</sup> Other mechanisms might include damage mediated by ROS, basement membrane degradation, endothelial cell activation, and transmigration of leukocytes.<sup>231,235</sup>

### *Perihematomal cerebral edema*

As blood is exquisitely toxic to brain tissue, hemorrhage by itself is a form of focal CNS injury that triggers formation of cerebral edema in the shell of tissue immediately surrounding the hemorrhage, i.e. the perihematomal space, a phenomenon referred to as perihematomal edema.<sup>240</sup> Perihematomal edema occurs in three stages: ionic edema, vasogenic edema, and delayed vasogenic edema. While the aforementioned core concepts that govern the formation of edema during phases of endothelial dysregulation (e.g. Starling's principle) apply also to perihematomal edema, some mechanistic details are unique to the latter.

Perihematomal ionic edema, the first stage of perihematomal edema, is driven by transendothelial osmotic forces generated by two processes. Firstly, cytotoxic edema forms in the perihematoma space, putatively because glutamate tends to accumulate in this region.<sup>241,242</sup> As described above, cytotoxic edema generates a strong driving force for the influx of ionic edema. Secondly, a hemorrhage-specific phenomenon called clot retraction, where activation of the coagulation cascade in the hematoma results in exudation of serum proteins and increased colloidal pressure of the perihematomal space, drives influx of water.<sup>243,244</sup>

Perihematomal vasogenic edema, the second stage of perihematomal edema, occurs when extravasation of blood products triggers changes in brain endothelium that manifest as extravasation of serum proteins without extravasation of erythrocytes. Thrombin, a protein that is extravasated with hemorrhaged blood and is produced at the site of injury, is a major contributor to the formation of perihematomal vasogenic edema.<sup>240</sup> Thrombin activates microglia primarily through PAR-1 receptors,<sup>245–247</sup> resulting in secretion of tumor necrosis factor (TNF) and IL-1 $\beta$ ,<sup>246,248,249</sup> cytokines that elicits downregulation of tight-junction proteins in endothelial cells and BBB opening.<sup>250,251</sup> In addition, thrombin enables the transmigration of circulating leukocytes by triggering endothelial retraction (as discussed above) and endothelial upregulation of chemokines and adhesion molecules.<sup>246,252</sup> Infiltrating leukocytes contribute to perihematomal edema through the secretion of mediators such as ROS. In addition to thrombin, the complement cascade is an important mediator of perihematomal vasogenic edema. Activation of the complement cascade results in the production of anaphylatoxins, membrane attack complex (MAC)-mediated lysis of red blood cells and iron-induced edema, as well as infiltration of neutrophils.<sup>253</sup>

The third and final stage of perihematomal edema, delayed vasogenic edema, is principally mediated by



hemoglobin degradation products that originate from extravasated and lysed erythrocytes. In the interstitium, hemoglobin is quickly oxidized to methemoglobin; the latter can spontaneously release its heme moiety, which may be further degraded by heme oxygenase enzymes to free iron.<sup>254</sup> Erythrocyte lysis and hemoglobin degradation is apparently a relatively slow process, as free iron reaches maximal tissue levels at approximately 3 days following a hemorrhage event.<sup>246,255</sup> Perihematomal delayed vasogenic edema exhibits similar temporality: Infusion of free iron increases brain water content by 24 h, whereas infusion of packed red blood cells increases brain water content only after 3 days.<sup>255,256</sup> Many hemoglobin degradation products can independently contribute to perihematomal delayed vasogenic edema. Free iron triggers ROS generation, MMP-9 activation, and BBB breakdown, while iron chelation reduces edema influx in models of intracerebral hemorrhage.<sup>257–259</sup> In addition, extracellular methemoglobin is a potent TLR-4 ligand that can trigger microglial TNF secretion and neuroinflammation.<sup>260</sup>

## Conclusions

While historical models have focused on the gross or ultrastructural appearance of edematous brain tissue, cerebral edema is better understood in a cellular and molecular context. The water movements involved in cerebral edema are dependent upon ionic fluxes, which are ultimately mediated by individual channels and transporters. The study of cerebral edema is essentially the study of maladaptive ion transport. While significant gaps still remain in our understanding of how specific proteins contribute to cerebral edema, the fields of cerebral edema and brain ISF dynamics are robust and productive. Doubtlessly, the next few years will yield new knowledge of how particular proteins drive edema influx, paving the way for rationally designed therapeutics that directly target key steps in cerebral edema formation, thereby achieving what currently approved therapies do not.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by grants to JMS from the National Institute of Neurological Disorders and Stroke (NINDS) (NS060801; NS061808) and the National Heart, Lung and Blood Institute (HL082517), and to VG from NINDS (NS061934; NS072501).

## Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Author's contributions

JS conceived of and wrote the manuscript. JMS and VG contributed intellectually and provided critical feedback on the manuscript.

## References

1. Kochanek KD, Xu J, Murphy SL, et al. Deaths: Final data for 2009. *Natl Vital Stat Rep* 2011; 60: 1–116.
2. Battley TW, Karki M, Singhal AB, et al. Brain edema predicts outcome after nonlacunar ischemic stroke. *Stroke* 2014; 45: 3643–3648.
3. Bernal W, Hyyrylainen A, Gera A, et al. Lessons from look-back in acute liver failure? A single centre experience of 3300 patients. *J Hepatol* 2013; 59: 74–80.
4. Donkin JJ and Vink R. Mechanisms of cerebral edema in traumatic brain injury: Therapeutic developments. *Curr Opin Neurol* 2010; 23: 293–299.
5. Abbott NJ. Comparative physiology of the blood-brain barrier. In: Bradbury MWB (ed.) *Physiology and pharmacology of the blood-brain barrier*. Berlin: Springer-Verlag, 1992, pp.171–196.
6. Di TR and Platt S. The function, composition and analysis of cerebrospinal fluid in companion animals: Part I - function and composition. *Vet J* 2006; 172: 422–431.
7. Ballabh P, Braun A and Nedergaard M. The blood-brain barrier: An overview: structure, regulation, and clinical implications. *Neurobiol Dis* 2004; 16: 1–13.
8. Cornford EM and Hyman S. Localization of brain endothelial luminal and abluminal transporters with immunogold electron microscopy. *NeuroRx* 2005; 2: 27–43.
9. Betz AL, Firth JA and Goldstein GW. Polarity of the blood-brain barrier: Distribution of enzymes between the luminal and antiluminal membranes of brain capillary endothelial cells. *Brain Res* 1980; 192: 17–28.
10. Yao Y and Tsirka SE. Monocyte chemoattractant protein-1 and the blood-brain barrier. *Cell Mol Life Sci* 2014; 71: 683–697.
11. Bell RD, Winkler EA, Sagare AP, et al. Pericytes control key neurovascular functions and neuronal phenotype in the adult brain and during brain aging. *Neuron* 2010; 68: 409–427.
12. Nakagawa S, Deli MA, Nakao S, et al. Pericytes from brain microvessels strengthen the barrier integrity in primary cultures of rat brain endothelial cells. *Cell Mol Neurobiol* 2007; 27: 687–694.
13. Armulik A, Genove G, Mae M, et al. Pericytes regulate the blood-brain barrier. *Nature* 2010; 468: 557–561.
14. Jones EG. On the mode of entry of blood vessels into the cerebral cortex. *J Anat* 1970; 106: 507–520.
15. Zhang ET, Inman CB and Weller RO. Interrelationships of the pia mater and the perivascular (Virchow-Robin) spaces in the human cerebrum. *J Anat* 1990; 170: 111–123.
16. Bushong EA, Martone ME, Jones YZ, et al. Protoplasmic astrocytes in CA1 stratum radiatum occupy separate anatomical domains. *J Neurosci* 2002; 22: 183–192.

17. Mathiisen TM, Lehre KP, Danbolt NC, et al. The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction. *Glia* 2010; 58: 1094–1103.
18. Iliff JJ, Wang M, Liao Y, et al. A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid beta. *Sci Transl Med* 2012; 4: 147ra111.
19. Thrane VR, Thrane AS, Plog BA, et al. Paravascular microcirculation facilitates rapid lipid transport and astrocyte signaling in the brain. *Sci Rep* 2013; 3: 2582.
20. Iliff JJ and Nedergaard M. Is there a cerebral lymphatic system? *Stroke* 2013; 44: S93–S95.
21. Brinker T, Stopa E, Morrison J, et al. A new look at cerebrospinal fluid circulation. *Fluids Barriers CNS* 2014; 11: 10.
22. Harder DR, Zhang C and Gebremedhin D. Astrocytes function in matching blood flow to metabolic activity. *News Physiol Sci* 2002; 17: 27–31.
23. Gordon GR, Mulligan SJ and MacVicar BA. Astrocyte control of the cerebrovasculature. *Glia* 2007; 55: 1214–1221.
24. Attwell D, Buchan AM, Charpak S, et al. Glial and neuronal control of brain blood flow. *Nature* 2010; 468: 232–243.
25. Belanger M, Allaman I and Magistretti PJ. Brain energy metabolism: Focus on astrocyte-neuron metabolic cooperation. *Cell Metab* 2011; 14: 724–738.
26. Bak LK, Schousboe A and Waagepetersen HS. The glutamate/GABA-glutamine cycle: Aspects of transport, neurotransmitter homeostasis and ammonia transfer. *J Neurochem* 2006; 98: 641–653.
27. Castro MA, Beltran FA, Brauchi S, et al. A metabolic switch in brain: Glucose and lactate metabolism modulation by ascorbic acid. *J Neurochem* 2009; 110: 423–440.
28. Cheyne J. *An essay on hydrocephalus acutus*. Philadelphia, PA: Finley, 1814.
29. Balfour . *Observations on the Dropsy in the brain*. Edinburgh, 1768.
30. *Pathological and practical researches on diseases of the brain and the spinal cord*. Philadelphia, PA: Carey & Lea, 1831.
31. Monro SA. *Observations on the structure and function of the nervous system*. Edinburgh: W Creech, 1783.
32. Preston G. Cerebral oedema. *J Nerv Ment Dis* 1894; 19: 494–503.
33. Torack RM. Historical aspects of normal and abnormal brain fluids. III. Cerebral edema. *Arch Neurol* 1982; 39: 355–357.
34. Reichardt M. Hirnschwellung. *Allg Z Psychiatr* 1919; 75: 34–103.
35. Spatz H. Die bedeutun der symptomatischen hirnschwellung fur die hirntumoren und fur andere raumbeengende prozesse in der scadelgrube. *Arch Psychiatr* 1929; 88: 790–794.
36. Bakay L and Lee JC. *Cerebral edema*. Springfield, IL: Thomas, 1965.
37. Klatzo I. Presidential address. Neuropathological aspects of brain edema. *J Neuropathol Exp Neurol* 1967; 26: 1–14.
38. Simard JM, Kent TA, Chen M, et al. Brain oedema in focal ischaemia: Molecular pathophysiology and theoretical implications. *Lancet Neurol* 2007; 6: 258–268.
39. Minnerup J, Sutherland BA, Buchan AM, et al. Neuroprotection for stroke: Current status and future perspectives. *Int J Mol Sci* 2012; 13: 11753–11772.
40. O'Collins VE, Macleod MR, Donnan GA, et al. 1,026 experimental treatments in acute stroke. *Ann Neurol* 2006; 59: 467–477.
41. Hussain MS and Shuaib A. Research into neuroprotection must continue... But with a different approach. *Stroke* 2008; 39: 521–522.
42. Fisher M, Feuerstein G, Howells DW, et al. Update of the stroke therapy academic industry roundtable preclinical recommendations. *Stroke* 2009; 40: 2244–2250.
43. Tait MJ, Saadoun S, Bell BA, et al. Water movements in the brain: Role of aquaporins. *Trends Neurosci* 2008; 31: 37–43.
44. Larsen BR, Assentoft M, Cotrina ML, et al. Contributions of the Na(+)/K(+)-ATPase, NKCC1, and Kir4.1 to hippocampal K(+) clearance and volume responses. *Glia* 2014; 62: 608–622.
45. Ransom BR, Yamate CL and Connors BW. Activity-dependent shrinkage of extracellular space in rat optic nerve: A developmental study. *J Neurosci* 1985; 5: 532–535.
46. Hossmann KA and Schuier FJ. Experimental brain infarcts in cats. I. Pathophysiological observations. *Stroke* 1980; 11: 583–592.
47. Thrane AS, Rangroo TV and Nedergaard M. Drowning stars: Reassessing the role of astrocytes in brain edema. *Trends Neurosci* 2014; 37: 620–628.
48. Bell BA, Symon L and Branston NM. CBF and time thresholds for the formation of ischemic cerebral edema, and effect of reperfusion in baboons. *J Neurosurg* 1985; 62: 31–41.
49. Simard JM, Chen M, Tarasov KV, et al. Newly expressed SUR1-regulated NC(Ca-ATP) channel mediates cerebral edema after ischemic stroke. *Nat Med* 2006; 12: 433–440.
50. Quast MJ, Huang NC, Hillman GR, et al. The evolution of acute stroke recorded by multimodal magnetic resonance imaging. *Magn Reson Imaging* 1993; 11: 465–471.
51. Jalan R, Olde Damink SW, Deutz NE, et al. Moderate hypothermia prevents cerebral hyperemia and increase in intracranial pressure in patients undergoing liver transplantation for acute liver failure. *Transplantation* 2003; 75: 2034–2039.
52. Iliff JJ, Chen MJ, Plog BA, et al. Impairment of glymphatic pathway function promotes tau pathology after traumatic brain injury. *J Neurosci* 2014; 34: 16180–16193.
53. Bering EA. Water exchange of central nervous system and cerebrospinal fluid. *J Neurosurg* 1952; 9: 275–287.
54. Barber AJ and Lieth E. Agrin accumulates in the brain microvascular basal lamina during development of the blood-brain barrier. *Dev Dyn* 1997; 208: 62–74.
55. Raghavan R, Brady ML, Rodriguez-Ponce MI, et al. Convection-enhanced delivery of therapeutics for brain disease, and its optimization. *Neurosurg Focus* 2006; 20: E12.

56. His W. Uber ein perivascuales kanalsystem in den nervosen central-organen und uber dessen beziehungen zum lymphsystem. *Zeitschrift Wissens Zoo* 1865; 15: 127–141.
57. Cserr HF, Cooper DN, Suri PK, et al. Efflux of radiolabeled polyethylene glycols and albumin from rat brain. *Am J Physiol* 1981; 240: F319–F328.
58. Geer CP and Grossman SA. Interstitial fluid flow along white matter tracts: A potentially important mechanism for the dissemination of primary brain tumors. *J Neurooncol* 1997; 32: 193–201.
59. Reulen HJ, Graham R, Spatz M, et al. Role of pressure gradients and bulk flow in dynamics of vasogenic brain edema. *J Neurosurg* 1977; 46: 24–35.
60. Badaut J, Ashwal S, Adami A, et al. Brain water mobility decreases after astrocytic aquaporin-4 inhibition using RNA interference. *J Cereb Blood Flow Metab* 2011; 31: 819–831.
61. Ohata K and Marmarou A. Clearance of brain edema and macromolecules through the cortical extracellular space. *J Neurosurg* 1992; 77: 387–396.
62. Reulen HJ, Tsuyumu M, Tack A, et al. Clearance of edema fluid into cerebrospinal fluid. A mechanism for resolution of vasogenic brain edema. *J Neurosurg* 1978; 48: 754–764.
63. Bradbury MW, Cserr HF and Westrop RJ. Drainage of cerebral interstitial fluid into deep cervical lymph of the rabbit. *Am J Physiol* 1981; 240: F329–F336.
64. Szentistvanyi I, Patlak CS, Ellis RA, et al. Drainage of interstitial fluid from different regions of rat brain. *Am J Physiol* 1984; 246: F835–F844.
65. Starling EH. On the absorption of fluids from the connective tissue spaces. *J Physiol* 1896; 19: 312–326.
66. Wiig H and Reed RK. Rat brain interstitial fluid pressure measured with micropipettes. *Am J Physiol* 1983; 244: H239–H246.
67. Norenberg MD. Astrocyte responses to CNS injury. *J Neuropathol Exp Neurol* 1994; 53: 213–220.
68. Norenberg MD, Jayakumar AR, Rama Rao KV, et al. New concepts in the mechanism of ammonia-induced astrocyte swelling. *Metab Brain Dis* 2007; 22: 219–234.
69. Magee PN, Stoner HB and Barnes JM. The experimental production in oedema in the central nervous system of the rat by triethyltin compounds. *J Pathol* 1957; 73: 107–124.
70. Stokum JA, Kurland DB, Gerzanich V, et al. Mechanisms of astrocyte-mediated cerebral edema. *Neurochem Res* 2015; 40: 317–328.
71. Minieri L, Pivonkova H, Harantova L, et al. Intracellular Na(+) inhibits volume-regulated anion channel in rat cortical astrocytes. *J Neurochem* 2015; 132: 286–300.
72. Takano T, Tian GF, Peng W, et al. Cortical spreading depression causes and coincides with tissue hypoxia. *Nat Neurosci* 2007; 10: 754–762.
73. Thrane AS, Rappold PM, Fujita T, et al. Critical role of aquaporin-4 (AQP4) in astrocytic Ca<sup>2+</sup> signaling events elicited by cerebral edema. *Proc Natl Acad Sci USA* 2011; 108: 846–851.
74. Zhou N, Gordon GR, Feighan D, et al. Transient swelling, acidification, and mitochondrial depolarization occurs in neurons but not astrocytes during spreading depression. *Cereb Cortex* 2010; 20: 2614–2624.
75. Rose CR and Ransom BR. Intracellular sodium homeostasis in rat hippocampal astrocytes. *J Physiol* 1996; 491: 291–305.
76. Somjen GG. Ion regulation in the brain: implications for pathophysiology. *Neuroscientist* 2002; 8: 254–267.
77. Gido G, Kristian T and Siesjo BK. Extracellular potassium in a neocortical core area after transient focal ischemia. *Stroke* 1997; 28: 206–210.
78. Florence CM, Baillie LD and Mulligan SJ. Dynamic volume changes in astrocytes are an intrinsic phenomenon mediated by bicarbonate ion flux. *PLoS One* 2012; 7: e51124.
79. Bourke RS and Nelson KM. Further studies on the K<sup>+</sup>-dependent swelling of primate cerebral cortex in vivo: The enzymatic basis of the K<sup>+</sup>-dependent transport of chloride. *J Neurochem* 1972; 19: 663–685.
80. Yan Y, Dempsey RJ and Sun D. Expression of Na(+)-K(+)-Cl(-) cotransporter in rat brain during development and its localization in mature astrocytes. *Brain Res* 2001; 911: 43–55.
81. Yan Y, Dempsey RJ and Sun D. Na<sup>+</sup>-K<sup>+</sup>-Cl<sup>-</sup> cotransporter in rat focal cerebral ischemia. *J Cereb Blood Flow Metab* 2001; 21: 711–721.
82. Jayakumar AR and Norenberg MD. The Na-K-Cl Cotransporter in astrocyte swelling. *Metab Brain Dis* 2010; 25: 31–38.
83. MacAulay N and Zeuthen T. Water transport between CNS compartments: Contributions of aquaporins and cotransporters. *Neuroscience* 2010; 168: 941–956.
84. Hamann S, Herrera-Perez JJ, Zeuthen T, et al. Cotransport of water by the Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransporter NKCC1 in mammalian epithelial cells. *J Physiol* 2010; 588: 4089–4101.
85. Su G, Kintner DB and Sun D. Contribution of Na(+)-K(+)-Cl(-) cotransporter to high-[K(+)](o)- induced swelling and EAA release in astrocytes. *Am J Physiol Cell Physiol* 2002; 282: C1136–C1146.
86. Su G, Kintner DB, Flagella M, et al. Astrocytes from Na(+)-K(+)-Cl(-) cotransporter-null mice exhibit absence of swelling and decrease in EAA release. *Am J Physiol Cell Physiol* 2002; 282: C1147–C1160.
87. Lu KT, Cheng NC, Wu CY, et al. NKCC1-mediated traumatic brain injury-induced brain edema and neuron death via Raf/MEK/MAPK cascade. *Crit Care Med* 2008; 36: 917–922.
88. O'Donnell ME, Tran L, Lam TI, et al. Bumetanide inhibition of the blood-brain barrier Na-K-Cl cotransporter reduces edema formation in the rat middle cerebral artery occlusion model of stroke. *J Cereb Blood Flow Metab* 2004; 24: 1046–1056.
89. O'Donnell ME, Lam TI, Tran L, et al. The role of the blood-brain barrier Na-K-2Cl cotransporter in stroke. *Adv Exp Med Biol* 2004; 559: 67–75.
90. Yan Y, Dempsey RJ, Flemmer A, et al. Inhibition of Na(+)-K(+)-Cl(-) cotransporter during focal cerebral ischemia decreases edema and neuronal damage. *Brain Res* 2003; 961: 22–31.
91. Wahl F, Obrenovitch TP, Hardy AM, et al. Extracellular glutamate during focal cerebral ischaemia in rats: Time course and calcium dependency. *J Neurochem* 1994; 63: 1003–1011.

92. Guyot LL, Diaz FG, O'Regan MH, et al. Real-time measurement of glutamate release from the ischemic penumbra of the rat cerebral cortex using a focal middle cerebral artery occlusion model. *Neurosci Lett* 2001; 299: 37–40.
93. Obrenovitch TP and Urenjak J. Is high extracellular glutamate the key to excitotoxicity in traumatic brain injury? *J Neurotrauma* 1997; 14: 677–698.
94. Schneider GH, Baethmann A and Kempfski O. Mechanisms of glial swelling induced by glutamate. *Can J Physiol Pharmacol* 1992; 70: S334–S343.
95. de Knegt RJ, Schalm SW, van der Rijt CC, et al. Extracellular brain glutamate during acute liver failure and during acute hyperammonemia simulating acute liver failure: An experimental study based on in vivo brain dialysis. *J Hepatol* 1994; 20: 19–26.
96. Hansson E, Muyderman H, Leonova J, et al. Astroglia and glutamate in physiology and pathology: Aspects on glutamate transport, glutamate-induced cell swelling and gap-junction communication. *Neurochem Int* 2000; 37: 317–329.
97. Schmitt A, Asan E, Puschel B, et al. Cellular and regional distribution of the glutamate transporter GLAST in the CNS of rats: Nonradioactive in situ hybridization and comparative immunocytochemistry. *J Neurosci* 1997; 17: 1–10.
98. Haugeto O, Ullensvang K, Levy LM, et al. Brain glutamate transporter proteins form homomultimers. *J Biol Chem* 1996; 271: 27715–27722.
99. Tanaka K, Watase K, Manabe T, et al. Epilepsy and exacerbation of brain injury in mice lacking the glutamate transporter GLT-1. *Science* 1997; 276: 1699–1702.
100. Hinson SR, Roemer SF, Lucchinetti CF, et al. Aquaporin-4-binding autoantibodies in patients with neuromyelitis optica impair glutamate transport by down-regulating EAAT2. *J Exp Med* 2008; 205: 2473–2481.
101. Aronica E, Gorter JA, Jansen GH, et al. Expression and cell distribution of group I and group II metabotropic glutamate receptor subtypes in Taylor-type focal cortical dysplasia. *Epilepsia* 2003; 44: 785–795.
102. Bender AS, Schousboe A, Reichelt W, et al. Ionic mechanisms in glutamate-induced astrocyte swelling: Role of K<sup>+</sup> influx. *J Neurosci Res* 1998; 52: 307–321.
103. Illarionova NB, Gunnarson E, Li Y, et al. Functional and molecular interactions between aquaporins and Na,K-ATPase. *Neuroscience* 2010; 168: 915–925.
104. Sun W, McConnell E, Pare JF, et al. Glutamate-dependent neuroglial calcium signaling differs between young and adult brain. *Science* 2013; 339: 197–200.
105. Izumi Y, Kirby CO, Benz AM, et al. Muller cell swelling, glutamate uptake, and excitotoxic neurodegeneration in the isolated rat retina. *Glia* 1999; 25: 379–389.
106. Kempfski O, Staub F, Jansen M, et al. Glial swelling during extracellular acidosis in vitro. *Stroke* 1988; 19: 385–392.
107. Staub F, Baethmann A, Peters J, et al. Effects of lactic acidosis on glial cell volume and viability. *J Cereb Blood Flow Metab* 1990; 10: 866–876.
108. Douglas RM, Schmitt BM, Xia Y, et al. Sodium-hydrogen exchangers and sodium-bicarbonate cotransporters: ontogeny of protein expression in the rat brain. *Neuroscience* 2001; 102: 217–228.
109. Xue J and Haddad GG. The Na<sup>+</sup>/H<sup>+</sup> exchanger: A target for therapeutic intervention in cerebral ischemia. In: Annunziato L (ed.) *New strategies in stroke intervention: Ionic transporters, pumps, and new channels*. New York: Humana Press, 2010, pp.113–128.
110. Ferrazzano P, Shi Y, Manhas N, et al. Inhibiting the Na<sup>+</sup>/H<sup>+</sup> exchanger reduces reperfusion injury: A small animal MRI study. *Front Biosci (Elite Ed)* 2011; 3: 81–88.
111. Jakubovicz DE and Klip A. Lactic acid-induced swelling in C6 glial cells via Na<sup>+</sup>/H<sup>+</sup> exchange. *Brain Res* 1989; 485: 215–224.
112. Kintner DB, Su G, Lenart B, et al. Increased tolerance to oxygen and glucose deprivation in astrocytes from Na(+)/H(+) exchanger isoform 1 null mice. *Am J Physiol Cell Physiol* 2004; 287: C12–C21.
113. Kitayama J, Kitazono T, Yao H, et al. Inhibition of Na<sup>+</sup>/H<sup>+</sup> exchanger reduces infarct volume of focal cerebral ischemia in rats. *Brain Res* 2001; 922: 223–228.
114. Kuribayashi Y, Itoh N, Kitano M, et al. Cerebroprotective properties of SM-20220, a potent Na(+)/H(+) exchange inhibitor, in transient cerebral ischemia in rats. *Eur J Pharmacol* 1999; 383: 163–168.
115. Park HS, Lee BK, Park S, et al. Effects of sabiporide, a specific Na<sup>+</sup>/H<sup>+</sup> exchanger inhibitor, on neuronal cell death and brain ischemia. *Brain Res* 2005; 1061: 67–71.
116. Wang Y, Luo J, Chen X, et al. Gene inactivation of Na<sup>+</sup>/H<sup>+</sup> exchanger isoform 1 attenuates apoptosis and mitochondrial damage following transient focal cerebral ischemia. *Eur J Neurosci* 2008; 28: 51–61.
117. Bevensee MO, Weed RA and Boron WF. Intracellular pH regulation in cultured astrocytes from rat hippocampus. I. Role Of HCO<sub>3</sub><sup>-</sup>. *J Gen Physiol* 1997; 110: 453–465.
118. Cooper AJ and Plum F. Biochemistry and physiology of brain ammonia. *Physiol Rev* 1987; 67: 440–519.
119. Butterworth RF. Pathophysiology of hepatic encephalopathy: A new look at ammonia. *Metab Brain Dis* 2002; 17: 221–227.
120. Norenberg MD, Baker L, Norenberg LO, et al. Ammonia-induced astrocyte swelling in primary culture. *Neurochem Res* 1991; 16: 833–836.
121. Albrecht J and Norenberg MD. Glutamine: a Trojan horse in ammonia neurotoxicity. *Hepatology* 2006; 44: 788–794.
122. Jayakumar AR, Liu M, Moriyama M, et al. Na-K-Cl Cotransporter-1 in the mechanism of ammonia-induced astrocyte swelling. *J Biol Chem* 2008; 283: 33874–33882.
123. Chen M, Dong Y and Simard JM. Functional coupling between sulfonyleurea receptor type 1 and a nonselective cation channel in reactive astrocytes from adult rat brain. *J Neurosci* 2003; 23: 8568–8577.
124. Nilius B, Prenen J, Tang J, et al. Regulation of the Ca<sup>2+</sup>-sensitivity of the nonselective cation channel TRPM4. *J Biol Chem* 2005; 280: 6423–6433.

125. Woo SK, Kwon MS, Ivanov A, et al. The sulfonylurea receptor 1 (Sur1)-transient receptor potential melastatin 4 (Trpm4) channel. *J Biol Chem* 2013; 288: 3655–3667.
126. Woo SK, Kwon MS, Geng Z, et al. Sequential activation of hypoxia-inducible factor 1 and specificity protein 1 is required for hypoxia-induced transcriptional stimulation of Abcc8. *J Cereb Blood Flow Metab* 2012; 32: 525–536.
127. Mehta RI, Ivanova S, Tosun C, et al. Sulfonylurea receptor 1 expression in human cerebral infarcts. *J Neuropathol Exp Neurol* 2013; 72: 871–883.
128. Chen M and Simard JM. Cell swelling and a nonselective cation channel regulated by internal Ca<sup>2+</sup> and ATP in native reactive astrocytes from adult rat brain. *J Neurosci* 2001; 21: 6512–6521.
129. Simard JM, Yurovsky V, Tsybalyuk N, et al. Protective effect of delayed treatment with low-dose glibenclamide in three models of ischemic stroke. *Stroke* 2009; 40: 604–609.
130. Jayakumar AR, Valdes V, Tong XY, et al. Sulfonylurea receptor 1 contributes to the astrocyte swelling and brain edema in acute liver failure. *Transl Stroke Res* 2014; 5: 28–37.
131. Preston GM, Carroll TP, Guggino WB, et al. Appearance of water channels in *Xenopus* oocytes expressing red cell CHIP28 protein. *Science* 1992; 256: 385–387.
132. Pohl P. Combined transport of water and ions through membrane channels. *Biol Chem* 2004; 385: 921–926.
133. Gorelick DA, Praetorius J, Tsunenari T, et al. Aquaporin-11: a channel protein lacking apparent transport function expressed in brain. *BMC Biochem* 2006; 7: 14.
134. Badaut J, Petit JM, Brunet JF, et al. Distribution of Aquaporin 9 in the adult rat brain: preferential expression in catecholaminergic neurons and in glial cells. *Neuroscience* 2004; 128: 27–38.
135. Nielsen S, Nagelhus EA, Amiry-Moghaddam M, et al. Specialized membrane domains for water transport in glial cells: High-resolution immunogold cytochemistry of aquaporin-4 in rat brain. *J Neurosci* 1997; 17: 171–180.
136. Saadoun S, Papadopoulos MC, Watanabe H, et al. Involvement of aquaporin-4 in astroglial cell migration and glial scar formation. *J Cell Sci* 2005; 118: 5691–5698.
137. Tomas-Camardiel M, Venero JL, de Pablos RM, et al. In vivo expression of aquaporin-4 by reactive microglia. *J Neurochem* 2004; 91: 891–899.
138. Jung JS, Bhat RV, Preston GM, et al. Molecular characterization of an aquaporin cDNA from brain: Candidate osmoreceptor and regulator of water balance. *Proc Natl Acad Sci USA* 1994; 91: 13052–13056.
139. Strand L, Moe SE, Solbu TT, et al. Roles of aquaporin-4 isoforms and amino acids in square array assembly. *Biochemistry* 2009; 48: 5785–5793.
140. Moe SE, Sorbo JG, Sogaard R, et al. New isoforms of rat Aquaporin-4. *Genomics* 2008; 91: 367–377.
141. Potokar M, Stenovec M, Jorgacevski J, et al. Regulation of AQP4 surface expression via vesicle mobility in astrocytes. *Glia* 2013; 61: 917–928.
142. Silberstein C, Bouley R, Huang Y, et al. Membrane organization and function of M1 and M23 isoforms of aquaporin-4 in epithelial cells. *Am J Physiol Renal Physiol* 2004; 287: F501–F511.
143. Wolburg H. Orthogonal arrays of intramembranous particles: A review with special reference to astrocytes. *J Hirnforsch* 1995; 36: 239–258.
144. Basco D, Blaauw B, Pisani F, et al. AQP4-dependent water transport plays a functional role in exercise-induced skeletal muscle adaptations. *PLoS One* 2013; 8: e58712.
145. Binder DK, Papadopoulos MC, Haggie PM, et al. In vivo measurement of brain extracellular space diffusion by cortical surface photobleaching. *J Neurosci* 2004; 24: 8049–8056.
146. Binder DK, Oshio K, Ma T, et al. Increased seizure threshold in mice lacking aquaporin-4 water channels. *Neuroreport* 2004; 15: 259–262.
147. Li J and Verkman AS. Impaired hearing in mice lacking aquaporin-4 water channels. *J Biol Chem* 2001; 276: 31233–31237.
148. Papadopoulos MC, Manley GT, Krishna S, et al. Aquaporin-4 facilitates reabsorption of excess fluid in vasogenic brain edema. *FASEB J* 2004; 18: 1291–1293.
149. Benfenati V, Caprini M, Dovizio M, et al. An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. *Proc Natl Acad Sci USA* 2011; 108: 2563–2568.
150. Da T and Verkman AS. Aquaporin-4 gene disruption in mice protects against impaired retinal function and cell death after ischemia. *Invest Ophthalmol Vis Sci* 2004; 45: 4477–4483.
151. Haj-Yasein NN, Vindedal GF, Eilert-Olsen M, et al. Glial-conditional deletion of aquaporin-4 (Aqp4) reduces blood-brain water uptake and confers barrier function on perivascular astrocyte endfeet. *Proc Natl Acad Sci USA* 2011; 108: 17815–17820.
152. Higashida T, Kreipke CW, Rafols JA, et al. The role of hypoxia-inducible factor-1alpha, aquaporin-4, and matrix metalloproteinase-9 in blood-brain barrier disruption and brain edema after traumatic brain injury. *J Neurosurg* 2011; 114: 92–101.
153. Manley GT, Fujimura M, Ma T, et al. Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke. *Nat Med* 2000; 6: 159–163.
154. Shenaq M, Kassem H, Peng C, et al. Neuronal damage and functional deficits are ameliorated by inhibition of aquaporin and HIF1alpha after traumatic brain injury (TBI). *J Neurol Sci* 2012; 323: 134–140.
155. Amiry-Moghaddam M, Xue R, Haug FM, et al. Alpha-syntrophin deletion removes the perivascular but not endothelial pool of aquaporin-4 at the blood-brain barrier and delays the development of brain edema in an experimental model of acute hyponatremia. *FASEB J* 2004; 18: 542–544.

156. Vajda Z, Pedersen M, Fuchtbauer EM, et al. Delayed onset of brain edema and mislocalization of aquaporin-4 in dystrophin-null transgenic mice. *Proc Natl Acad Sci USA* 2002; 99: 13131–13136.
157. Solenov E, Watanabe H, Manley GT, et al. Sevenfold-reduced osmotic water permeability in primary astrocyte cultures from AQP-4-deficient mice, measured by a fluorescence quenching method. *Am J Physiol Cell Physiol* 2004; 286: C426–C432.
158. Mori K, Miyazaki M, Iwase H, et al. Temporal profile of changes in brain tissue extracellular space and extracellular ion (Na(+), K(+)) concentrations after cerebral ischemia and the effects of mild cerebral hypothermia. *J Neurotrauma* 2002; 19: 1261–1270.
159. Stiefel MF and Marmarou A. Cation dysfunction associated with cerebral ischemia followed by reperfusion: a comparison of microdialysis and ion-selective electrode methods. *J Neurosurg* 2002; 97: 97–103.
160. Go KG. The normal and pathological physiology of brain water. *Adv Tech Stand Neurosurg* 1997; 23: 47–142.
161. Young W and Constantini S. Ionic and water shifts in injured central nervous tissues. In: Salzman S (ed.) *The neurobiology of central nervous system trauma*. New York: Oxford University Press, 2015, pp.123–130.
162. Lo WD, Betz AL, Schielke GP, et al. Transport of sodium from blood to brain in ischemic brain edema. *Stroke* 1987; 18: 150–157.
163. Hossmann KA. Development and resolution of ischemic brain swelling. In: Pappius HM and Feindel W (eds) *Dynamics of brain edema*. New York: Springer-Verlag, 1976, pp.219–227.
164. Ito U, Ohno K, Nakamura R, et al. Brain edema during ischemia and after restoration of blood flow. Measurement of water, sodium, potassium content and plasma protein permeability. *Stroke* 1979; 10: 542–547.
165. Lam TI, Wise PM and O'Donnell ME. Cerebral microvascular endothelial cell Na/H exchange: evidence for the presence of NHE1 and NHE2 isoforms and regulation by arginine vasopressin. *Am J Physiol Cell Physiol* 2009; 297: C278–C289.
166. O'Donnell ME, Chen YJ, Lam TI, et al. Intravenous HOE-642 reduces brain edema and Na uptake in the rat permanent middle cerebral artery occlusion model of stroke: Evidence for participation of the blood-brain barrier Na/H exchanger. *J Cereb Blood Flow Metab* 2013; 33: 225–234.
167. Suzuki Y, Matsumoto Y, Ikeda Y, et al. SM-20220, a Na(+)/H(+) exchanger inhibitor: Effects on ischemic brain damage through edema and neutrophil accumulation in a rat middle cerebral artery occlusion model. *Brain Res* 2002; 945: 242–248.
168. Foroutan S, Brillault J, Forbush B, et al. Moderate-to-severe ischemic conditions increase activity and phosphorylation of the cerebral microvascular endothelial cell Na<sup>+</sup>-K<sup>+</sup>-Cl<sup>-</sup> cotransporter. *Am J Physiol Cell Physiol* 2005; 289: C1492–C1501.
169. Betz AL. Sodium transport in capillaries isolated from rat brain. *J Neurochem* 1983; 41: 1150–1157.
170. Domotor E, Abbott NJ and Adam-Vizi V. Na<sup>+</sup>-Ca<sup>2+</sup> exchange and its implications for calcium homeostasis in primary cultured rat brain microvascular endothelial cells. *J Physiol* 1999; 515: 147–155.
171. Proks P, Reimann F, Green N, et al. Sulfonylurea stimulation of insulin secretion. *Diabetes* 2002; 51: S368–S376.
172. Gribble FM and Reimann F. Sulphonylurea action revisited: the post-cloning era. *Diabetologia* 2003; 46: 875–891.
173. Fujita A and Kurachi Y. Molecular aspects of ATP-sensitive K<sup>+</sup> channels in the cardiovascular system and K<sup>+</sup> channel openers. *Pharmacol Ther* 2000; 85: 39–53.
174. Simard JM, Sheth KN, Kimberly WT, et al. Glibenclamide in cerebral ischemia and stroke. *Neurocrit Care* 2014; 20: 319–333.
175. Sheth KN, Elm JJ, Beslow LA, et al. Glyburide advantage in malignant edema and stroke (GAMES-RP) trial: Rationale and design. *Neurocrit Care* 2015.
176. Davis SM, Lees KR, Albers GW, et al. Selfotel in acute ischemic stroke: Possible neurotoxic effects of an NMDA antagonist. *Stroke* 2000; 31: 347–354.
177. Nilius B and Droogmans G. Ion channels and their functional role in vascular endothelium. *Physiol Rev* 2001; 81: 1415–1459.
178. Zeuthen T. Cotransport of K<sup>+</sup>, Cl<sup>-</sup> and H<sub>2</sub>O by membrane proteins from choroid plexus epithelium of *Necturus maculosus*. *J Physiol* 1994; 478: 203–219.
179. Dolman D, Drndarski S, Abbott NJ, et al. Induction of aquaporin 1 but not aquaporin 4 messenger RNA in rat primary brain microvessel endothelial cells in culture. *J Neurochem* 2005; 93: 825–833.
180. Elfeber K, Kohler A, Lutzenburg M, et al. Localization of the Na<sup>+</sup>-D-glucose cotransporter SGLT1 in the blood-brain barrier. *Histochem Cell Biol* 2004; 121: 201–207.
181. Farrell CL and Pardridge WM. Blood-brain barrier glucose transporter is asymmetrically distributed on brain capillary endothelial luminal and abluminal membranes: An electron microscopic immunogold study. *Proc Natl Acad Sci USA* 1991; 88: 5779–5783.
182. Paulson OB, Hertz MM, Bolwig TG, et al. Filtration and diffusion of water across the blood-brain barrier in man. *Microvasc Res* 1977; 13: 113–124.
183. Yang B, Zador Z and Verkman AS. Glial cell aquaporin-4 overexpression in transgenic mice accelerates cytotoxic brain swelling. *J Biol Chem* 2008; 283: 15280–15286.
184. Stokum JA, Mehta RI, Ivanova S, et al. Heterogeneity of aquaporin-4 localization and expression after focal cerebral ischemia underlies differences in white versus grey matter swelling. *Acta Neuropathol Commun* 2015; 3: 61.
185. Aarabi B and Long DM. Dynamics of cerebral edema. The role of an intact vascular bed in the production and propagation of vasogenic brain edema. *J Neurosurg* 1979; 51: 779–784.
186. Fenske A, Samii M, Reulen HJ, et al. Extracellular space and electrolyte distribution in cortex and white

- matter of dog brain in cold induced oedema. *Acta Neurochir (Wien)* 1973; 28: 81–94.
187. Vorbrodth AW, Lossinsky AS, Wisniewski HM, et al. Ultrastructural observations on the transvascular route of protein removal in vasogenic brain edema. *Acta Neuropathol* 1985; 66: 265–273.
188. Hauck EF, Apostel S, Hoffmann JF, et al. Capillary flow and diameter changes during reperfusion after global cerebral ischemia studied by intravital video microscopy. *J Cereb Blood Flow Metab* 2004; 24: 383–391.
189. Durward QJ, Del Maestro RF, Amacher AL, et al. The influence of systemic arterial pressure and intracranial pressure on the development of cerebral vasogenic edema. *J Neurosurg* 1983; 59: 803–809.
190. Kogure K, Busto R and Scheinberg P. The role of hydrostatic pressure in ischemic brain edema. *Ann Neurol* 1981; 9: 273–282.
191. Kilincer C, Asil T, Utku U, et al. Factors affecting the outcome of decompressive craniectomy for large hemispheric infarctions: A prospective cohort study. *Acta Neurochir (Wien)* 2005; 147: 587–594.
192. Mori K, Nakao Y, Yamamoto T, et al. Early external decompressive craniectomy with duroplasty improves functional recovery in patients with massive hemispheric embolic infarction: Timing and indication of decompressive surgery for malignant cerebral infarction. *Surg Neurol* 2004; 62: 420–429.
193. Cooper PR, Hagler H, Clark WK, et al. Enhancement of experimental cerebral edema after decompressive craniectomy: Implications for the management of severe head injuries. *Neurosurgery* 1979; 4: 296–300.
194. Hofmeijer J, Schepers J, Veldhuis WB, et al. Delayed decompressive surgery increases apparent diffusion coefficient and improves peri-infarct perfusion in rats with space-occupying cerebral infarction. *Stroke* 2004; 35: 1476–1481.
195. Swanson JA and Watts C. Macropinocytosis. *Trends Cell Biol* 1995; 5: 424–428.
196. Broadwell RD and Salzman M. Expanding the definition of the blood-brain barrier to protein. *Proc Natl Acad Sci USA* 1981; 78: 7820–7824.
197. Castejon OJ. Formation of transendothelial channels in traumatic human brain edema. *Pathol Res Pract* 1984; 179: 7–12.
198. Nag S. Presence of transendothelial channels in cerebral endothelium in chronic hypertension. *Acta Neurochir Suppl (Wien)* 1990; 51: 335–337.
199. Balin BJ, Broadwell RD and Salzman M. Tubular profiles do not form transendothelial channels through the blood-brain barrier. *J Neurocytol* 1987; 16: 721–735.
200. Garcia JG, Siflinger-Birnboim A, Bizios R, et al. Thrombin-induced increase in albumin permeability across the endothelium. *J Cell Physiol* 1986; 128: 96–104.
201. Ochoa CD and Stevens T. Studies on the cell biology of interendothelial cell gaps. *Am J Physiol Lung Cell Mol Physiol* 2012; 302: L275–L286.
202. McDonald RI, Shepro D, Rosenthal M, et al. Properties of cultured endothelial cells. *Ser Haematol* 1973; 6: 469–478.
203. Laposata M, Dohnarsky DK and Shin HS. Thrombin-induced gap formation in confluent endothelial cell monolayers in vitro. *Blood* 1983; 62: 549–556.
204. Moy AB, Van EJ, Bodmer J, et al. Histamine and thrombin modulate endothelial focal adhesion through centripetal and centrifugal forces. *J Clin Invest* 1996; 97: 1020–1027.
205. Muller WA. Leukocyte-endothelial-cell interactions in leukocyte transmigration and the inflammatory response. *Trends Immunol* 2003; 24: 327–334.
206. Kovacs Z, Ikezaki K, Samoto K, et al. VEGF andflt. Expression time kinetics in rat brain infarct. *Stroke* 1996; 27: 1865–1872.
207. Dore-Duffy P, Wang X, Mehedi A, et al. Differential expression of capillary VEGF isoforms following traumatic brain injury. *Neurol Res* 2007; 29: 395–403.
208. Skold MK, von GC, Sandberg-Nordqvist AC, et al. VEGF and VEGF receptor expression after experimental brain contusion in rat. *J Neurotrauma* 2005; 22: 353–367.
209. Wang W, Dentler WL and Borchardt RT. VEGF increases BMEC monolayer permeability by affecting occludin expression and tight junction assembly. *Am J Physiol Heart Circ Physiol* 2001; 280: H434–H440.
210. Fischer S, Wobben M, Marti HH, et al. Hypoxia-induced hyperpermeability in brain microvessel endothelial cells involves VEGF-mediated changes in the expression of zonula occludens-1. *Microvasc Res* 2002; 63: 70–80.
211. Bates DO and Curry FE. Vascular endothelial growth factor increases hydraulic conductivity of isolated perfused microvessels. *Am J Physiol* 1996; 271: H2520–H2528.
212. Weis SM and Cheresh DA. Pathophysiological consequences of VEGF-induced vascular permeability. *Nature* 2005; 437: 497–504.
213. van BN, Thibodeaux H, Palmer JT, et al. VEGF antagonism reduces edema formation and tissue damage after ischemia/reperfusion injury in the mouse brain. *J Clin Invest* 1999; 104: 1613–1620.
214. Batchelor TT, Sorensen AG, di TE, et al. AZD2171, a pan-VEGF receptor tyrosine kinase inhibitor, normalizes tumor vasculature and alleviates edema in glioblastoma patients. *Cancer Cell* 2007; 11: 83–95.
215. Zhang ZG, Zhang L, Jiang Q, et al. VEGF enhances angiogenesis and promotes blood-brain barrier leakage in the ischemic brain. *J Clin Invest* 2000; 106: 829–838.
216. Krupinski J, Kaluza J, Kumar P, et al. Role of angiogenesis in patients with cerebral ischemic stroke. *Stroke* 1994; 25: 1794–1798.
217. Song L and Pachter JS. Monocyte chemoattractant protein-1 alters expression of tight junction-associated proteins in brain microvascular endothelial cells. *Microvasc Res* 2004; 67: 78–89.
218. Yamagata K, Tagami M, Takenaga F, et al. Hypoxia-induced changes in tight junction permeability of brain capillary endothelial cells are associated with IL-1beta and nitric oxide. *Neurobiol Dis* 2004; 17: 491–499.
219. Iadecola C, Zhang F, Casey R, et al. Inducible nitric oxide synthase gene expression in vascular cells after

- transient focal cerebral ischemia. *Stroke* 1996; 27: 1373–1380.
220. Sharma HS, Drieu K, Alm P, et al. Role of nitric oxide in blood-brain barrier permeability, brain edema and cell damage following hyperthermic brain injury. An experimental study using EGB-761 and Ginkgolide B pretreatment in the rat. *Acta Neurochir Suppl* 2000; 76: 81–86.
221. Nag S, Kapadia A and Stewart DJ. Molecular pathogenesis of blood-brain barrier breakdown in acute brain injury. *Neuropathol Appl Neurobiol* 2011; 37: 3–23.
222. Asahi M, Wang X, Mori T, et al. Effects of matrix metalloproteinase-9 gene knock-out on the proteolysis of blood-brain barrier and white matter components after cerebral ischemia. *J Neurosci* 2001; 21: 7724–7732.
223. Asahi M, Asahi K, Jung JC, et al. Role for matrix metalloproteinase 9 after focal cerebral ischemia: Effects of gene knockout and enzyme inhibition with BB-94. *J Cereb Blood Flow Metab* 2000; 20: 1681–1689.
224. Fukuda S, Fini CA, Mabuchi T, et al. Focal cerebral ischemia induces active proteases that degrade microvascular matrix. *Stroke* 2004; 35: 998–1004.
225. Kolev K, Skopal J, Simon L, et al. Matrix metalloproteinase-9 expression in post-hypoxic human brain capillary endothelial cells: H<sub>2</sub>O<sub>2</sub> as a trigger and NF- $\kappa$ B as a signal transducer. *Thromb Haemost* 2003; 90: 528–537.
226. Mun-Bryce S and Rosenberg GA. Matrix metalloproteinases in cerebrovascular disease. *J Cereb Blood Flow Metab* 1998; 18: 1163–1172.
227. Romanic AM, White RF, Arleth AJ, et al. Matrix metalloproteinase expression increases after cerebral focal ischemia in rats: inhibition of matrix metalloproteinase-9 reduces infarct size. *Stroke* 1998; 29: 1020–1030.
228. Lapchak PA, Chapman DF and Zivin JA. Metalloproteinase inhibition reduces thrombolytic (tissue plasminogen activator)-induced hemorrhage after thromboembolic stroke. *Stroke* 2000; 31: 3034–3040.
229. Pfefferkorn T and Rosenberg GA. Closure of the blood-brain barrier by matrix metalloproteinase inhibition reduces rtPA-mediated mortality in cerebral ischemia with delayed reperfusion. *Stroke* 2003; 34: 2025–2030.
230. Yang Y, Estrada EY, Thompson JF, et al. Matrix metalloproteinase-mediated disruption of tight junction proteins in cerebral vessels is reversed by synthetic matrix metalloproteinase inhibitor in focal ischemia in rat. *J Cereb Blood Flow Metab* 2007; 27: 697–709.
231. Hamann GF, del Zoppo GJ and von KR. Hemorrhagic transformation of cerebral infarction—possible mechanisms. *Thromb Haemost* 1999; 82: 92–94.
232. Jaillard A, Cornu C, Durieux A, et al. Hemorrhagic transformation in acute ischemic stroke. The MAST-E study. MAST-E Group. *Stroke* 1999; 30: 1326–1332.
233. Larrue V, von KR, del ZG, et al. Hemorrhagic transformation in acute ischemic stroke. Potential contributing factors in the European Cooperative Acute Stroke Study. *Stroke* 1997; 28: 957–960.
234. Abumiya T, Yokota C, Kuge Y, et al. Aggravation of hemorrhagic transformation by early intraarterial infusion of low-dose vascular endothelial growth factor after transient focal cerebral ischemia in rats. *Brain Res* 2005; 1049: 95–103.
235. Wang X and Lo EH. Triggers and mediators of hemorrhagic transformation in cerebral ischemia. *Mol Neurobiol* 2003; 28: 229–244.
236. Heo JH, Lucero J, Abumiya T, et al. Matrix metalloproteinases increase very early during experimental focal cerebral ischemia. *J Cereb Blood Flow Metab* 1999; 19: 624–633.
237. Sumii T and Lo EH. Involvement of matrix metalloproteinase in thrombolysis-associated hemorrhagic transformation after embolic focal ischemia in rats. *Stroke* 2002; 33: 831–836.
238. Gerzanich V, Woo SK, Vennekens R, et al. De novo expression of Trpm4 initiates secondary hemorrhage in spinal cord injury. *Nat Med* 2009; 15: 185–191.
239. Simard JM, Tsymbalyuk O, Ivanov A, et al. Endothelial sulfonylurea receptor 1-regulated NC Ca-ATP channels mediate progressive hemorrhagic necrosis following spinal cord injury. *J Clin Invest* 2007; 117: 2105–2113.
240. Urday S, Kimberly WT, Beslow LA, et al. Targeting secondary injury in intracerebral haemorrhage-perihematomal oedema. *Nat Rev Neurol* 2015; 11: 111–122.
241. Brunswick AS, Hwang BY, Appelboom G, et al. Serum biomarkers of spontaneous intracerebral hemorrhage induced secondary brain injury. *J Neurol Sci* 2012; 321: 1–10.
242. Li N, Liu YF, Ma L, et al. Association of molecular markers with perihematomal edema and clinical outcome in intracerebral hemorrhage. *Stroke* 2013; 44: 658–663.
243. Wagner KR, Xi G, Hua Y, et al. Lobar intracerebral hemorrhage model in pigs: Rapid edema development in perihematomal white matter. *Stroke* 1996; 27: 490–497.
244. Xi G, Keep RF and Hoff JT. Pathophysiology of brain edema formation. *Neurosurg Clin N Am* 2002; 13: 371–383.
245. Wang J. Preclinical and clinical research on inflammation after intracerebral hemorrhage. *Prog Neurobiol* 2010; 92: 463–477.
246. Bodmer D, Vaughan KA, Zacharia BE, et al. The molecular mechanisms that promote edema after intracerebral hemorrhage. *Transl Stroke Res* 2012; 3: 52–61.
247. Suo Z, Wu M, Citron BA, et al. Persistent protease-activated receptor 4 signaling mediates thrombin-induced microglial activation. *J Biol Chem* 2003; 278: 31177–31183.
248. Masada T, Hua Y, Xi G, et al. Attenuation of intracerebral hemorrhage and thrombin-induced brain edema by overexpression of interleukin-1 receptor antagonist. *J Neurosurg* 2001; 95: 680–686.
249. Lei B, Dawson HN, Roulhac-Wilson B, et al. Tumor necrosis factor alpha antagonism improves neurological recovery in murine intracerebral hemorrhage. *J Neuroinflammation* 2013; 10: 103.
250. Aslam M, Ahmad N, Srivastava R, et al. TNF-alpha induced NF $\kappa$ B signaling and p65 (RelA)



- overexpression repress Cldn5 promoter in mouse brain endothelial cells. *Cytokine* 2012; 57: 269–275.
251. Megyeri P, Abraham CS, Temesvari P, et al. Recombinant human tumor necrosis factor alpha constricts pial arterioles and increases blood-brain barrier permeability in newborn piglets. *Neurosci Lett* 1992; 148: 137–140.
252. Bijli KM, Minhajuddin M, Fazal F, et al. c-Src interacts with and phosphorylates RelA/p65 to promote thrombin-induced ICAM-1 expression in endothelial cells. *Am J Physiol Lung Cell Mol Physiol* 2007; 292: L396–L404.
253. Garrett MC, Otten ML, Starke RM, et al. Synergistic neuroprotective effects of C3a and C5a receptor blockade following intracerebral hemorrhage. *Brain Res* 2009; 1298: 171–177.
254. Chen-Roetling J, Lu X and Regan RF. Targeting heme oxygenase after intracerebral hemorrhage. *Ther Targets Neurol Dis* 2015; 2.
255. Xi G, Keep RF and Hoff JT. Erythrocytes and delayed brain edema formation following intracerebral hemorrhage in rats. *J Neurosurg* 1998; 89: 991–996.
256. Huang FP, Xi G, Keep RF, et al. Brain edema after experimental intracerebral hemorrhage: Role of hemoglobin degradation products. *J Neurosurg* 2002; 96: 287–293.
257. Keep RF, Hua Y and Xi G. Intracerebral haemorrhage: Mechanisms of injury and therapeutic targets. *Lancet Neurol* 2012; 11: 720–731.
258. Nakamura T, Keep RF, Hua Y, et al. Deferoxamine-induced attenuation of brain edema and neurological deficits in a rat model of intracerebral hemorrhage. *J Neurosurg* 2004; 100: 672–678.
259. Xie Q, Gu Y, Hua Y, et al. Deferoxamine attenuates white matter injury in a piglet intracerebral hemorrhage model. *Stroke* 2014; 45: 290–292.
260. Kwon MS, Woo SK, Kurland DB, et al. Methemoglobin is an endogenous toll-like receptor 4 ligand-relevance to subarachnoid hemorrhage. *Int J Mol Sci* 2015; 16: 5028–5046.
261. Irani DN. *Cerebrospinal fluid in clinical practice*. Philadelphia, PA: Elsevier, 2015.
262. Zhang RB and Verkman AS. Water and urea permeability properties of Xenopus oocytes: Expression of mRNA from toad urinary bladder. *Am J Physiol* 1991; 260: C26–C34.
263. Yang B, Van Hoek AN and Verkman AS. Very high single channel water permeability of aquaporin-4 in baculovirus-infected insect cells and liposomes reconstituted with purified aquaporin-4. *Biochemistry* 1997; 36: 7625–7632.
264. Yang B and Verkman AS. Water and glycerol permeabilities of aquaporins 1-5 and MIP determined quantitatively by expression of epitope-tagged constructs in Xenopus oocytes. *J Biol Chem* 1997; 272: 16140–16146.
265. Duquette PP, Bissonnette P and Lapointe JY. Local osmotic gradients drive the water flux associated with Na(+)/glucose cotransport. *Proc Natl Acad Sci USA* 2001; 98: 3796–3801.
266. Loo DD, Zeuthen T, Chandy G, et al. Cotransport of water by the Na<sup>+</sup>/glucose cotransporter. *Proc Natl Acad Sci USA* 1996; 93: 13367–13370.
267. Iserovich P, Wang D, Ma L, et al. Changes in glucose transport and water permeability resulting from the T310I pathogenic mutation in Glut1 are consistent with two transport channels per monomer. *J Biol Chem* 2002; 277: 30991–30997.
268. Zeuthen T, Zeuthen E and MacAulay N. Water transport by GLUT2 expressed in Xenopus laevis oocytes. *J Physiol* 2007; 579: 345–361.
269. Loo DD, Hirayama BA, Meinild AK, et al. Passive water and ion transport by cotransporters. *J Physiol* 1999; 518: 195–202.
270. MacAulay N, Gether U, Klaeke DA, et al. Passive water and urea permeability of a human Na(+)-glutamate cotransporter expressed in Xenopus oocytes. *J Physiol* 2002; 542: 817–828.
271. MacAulay N, Zeuthen T and Gether U. Conformational basis for the Li(+)-induced leak current in the rat gamma-aminobutyric acid (GABA) transporter-1. *J Physiol* 2002; 544: 447–458.
272. MacAulay N, Gether U, Klaeke DA, et al. Water transport by the human Na<sup>+</sup>-coupled glutamate cotransporter expressed in Xenopus oocytes. *J Physiol* 2001; 530: 367–378.
273. Hamann S, Herrera-Perez JJ, Bundgaard M, et al. Water permeability of Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup>-cotransporters in mammalian epithelial cells. *J Physiol* 2005; 568: 123–135.
274. *Water movement through lipid bilayers, pores and plasma membranes: Theory and reality*. New York: John Wiley and Sons, 2015.

## Appendix

### Anatomical abbreviations

BBB	blood–brain barrier
CSF	cerebrospinal fluid
ISF	interstitial fluid
CNS	central nervous system

### Protein abbreviations

Pfkfb3	6-phosphofructose-2-kinase/fructose-2,6-biphosphatase-3 ()
NKCC1	Na <sup>+</sup> / K <sup>+</sup> / Cl <sup>-</sup> co-transporter
Sur1-Trpm4	sulfonylurea receptor 1 – transient receptor potential melastatin 4 ()
EAAT	excitatory amino acid transporter
mGluR5	metabotropic glutamate receptor 5
NHE	Na <sup>+</sup> / H <sup>+</sup> exchanger
NCX	Na <sup>+</sup> /Ca <sup>2+</sup> exchanger
KCC	K <sup>+</sup> /Cl <sup>-</sup> co-transporter
NBC	bicarbonate dependent Na <sup>+</sup> /HCO <sub>3</sub> <sup>-</sup> transporter family
PAG	phosphate-activated glutaminase

HIF1	hypoxia-inducible factor 1
SGLT-1	sodium-glucose linked transporter 1
VEGF	vascular endothelial growth factor
Ang2	angiopoietin 2
MMP	matrix metalloproteinase
ROS	reactive oxygen species
TNF	tumor necrosis factor
MAC	membrane attack complex
Aqp4	aquaporin-4
OAPs	orthogonal arrays of intramembraneous particles

#### *Small molecule abbreviations*

ATP	adenosine triphosphate
TET	triethyltin

#### *Physiological term abbreviations*

$P_c$	capillary hydrostatic pressure ()
$P_i$	tissue hydrostatic pressure ()
$\pi_c$	blood osmotic pressure
$\pi_i$	interstitial osmotic pressure
$K_H$	BBB net hydraulic conductivity
$K_O$	BBB net osmotic conductivity

#### *Miscellaneous abbreviations*

MRI	magnetic resonance imaging
-----	----------------------------