

Isolation of a Novel Insect-Specific Flavivirus from *Culiseta melanura* in the Northeastern United States

Michael J. Misencik,¹ Nathan D. Grubaugh,² Theodore G. Andreadis,¹
Gregory D. Ebel,² and Philip M. Armstrong¹

Abstract

The genus *Flavivirus* includes a number of newly recognized viruses that infect and replicate only within mosquitoes. To determine whether insect-specific flaviviruses (ISFs) may infect *Culiseta (Cs.) melanura* mosquitoes, we screened pools of field-collected mosquitoes for virus infection by RT-PCR targeting conserved regions of the *NS5* gene. *NS5* nucleotide sequences amplified from *Cs. melanura* pools were genetically similar to other ISFs and most closely matched Calbertado virus from *Culex tarsalis*, sharing 68.7% nucleotide and 76.1% amino acid sequence identity. The complete genome of one virus isolate was sequenced to reveal a primary open reading frame (ORF) encoding a viral polyprotein characteristic of the genus *Flavivirus*. Phylogenetic analysis showed that this virus represents a distinct evolutionary lineage that belongs to the classical ISF group. The virus was detected solely in *Cs. melanura* pools, occurred in sampled populations from Connecticut, New York, New Hampshire, and Maine, and infected both adult and larval stages of the mosquito. Maximum likelihood estimate infection rates (MLE-IR) were relatively stable in overwintering *Cs. melanura* larvae collected monthly from November of 2012 through May of 2013 (MLE-IR = 0.7–2.1/100 mosquitoes) and in host-seeking females collected weekly from June through October of 2013 (MLE-IR = 3.8–11.5/100 mosquitoes). Phylogenetic analysis of viral sequences revealed limited genetic variation that lacked obvious geographic structure among strains in the northeastern United States. This new virus is provisionally named Culiseta flavivirus on the basis of its host association with *Cs. melanura*.

Key Words: *Culiseta melanura*—Flavivirus—Insect viruses—Mosquito-only flavivirus—Genomic analysis—Phylogenetic analysis.

Introduction

THE *FLAVIVIRUS* GENUS REPRESENTS a diverse group of viruses that includes a number of important human pathogens, such as dengue virus, Japanese encephalitis virus, tick-borne encephalitis virus, and West Nile virus (WNV). These viruses are maintained in transmission cycles between arthropod vectors and vertebrate hosts, whereas other flaviviruses appear to be limited to infecting insects (Blitvich and Firth 2015). Cell fusing agent was the first insect-specific flavivirus (ISF) discovered 40 years ago from an *Aedes (Ae.) aegypti* cell line (Stollar and Thomas 1975). Since then, a multitude of ISFs have been isolated and characterized from a wide range of mosquito species worldwide. Some of these include Kamiti River virus (KRV) from *Ae. macintoshi* in

Kenya (Crabtree et al. 2003), *Culex flavivirus* (CxFV) from *Culex (Cx.) pipiens* in Japan (Hoshino et al. 2007), *Aedes flavivirus* (AeFV) from *Ae. albopictus* and *Ae. flavopictus* in Japan (Hoshino et al. 2009), Calbertado virus (CLBOV) from *Cx. tarsalis* and *Cx. pipiens* in Canada and the United States (Bolling et al. 2011, Tyler et al. 2011), and Palm Creek virus (PCV) from *Coquillettidia xanthogaster* in Australia (Hobson-Peters et al. 2013). These ISFs form a monophyletic group and are distantly related to the dual-host mosquito-borne and tick-borne flaviviruses and to viruses with no known vector. However, more recently, a second group of ISF-like viruses has been isolated from mosquitoes that replicate solely within arthropod cells but cluster phylogenetically with the dual-host mosquito-borne flaviviruses. This group includes Lammi virus (Huhtamo et al. 2009), Nounane virus

¹Center for Vector Biology and Zoonotic Diseases, The Connecticut Agricultural Experiment Station, New Haven, Connecticut.

²Arthropod-Borne and Infectious Diseases Laboratory, Department of Microbiology, Immunology, and Pathology, Colorado State University, Fort Collins, Colorado.

(Junglen et al. 2009), Barkedji virus (Kolodziejek et al. 2013), Chaoyang virus (Lee et al. 2013), and Nhumirim virus (Kenney et al. 2014).

CxFV represents the most extensively studied ISF and has been detected in multiple mosquito species from around the world. The virus was found to be highly prevalent in *Cx. pipiens* and *Cx. quinquefasciatus* sampled from East Asia, Africa, and the Americas (Hoshino et al. 2007, Morales-Betoulle et al. 2008, Blitvich et al. 2009, Cook et al. 2009, Kim et al. 2009, Farfan-Ale et al. 2010, Bolling et al. 2011, Newman et al. 2011, Machado et al. 2012), and it was infrequently detected in *Cx. tarsalis* from the United States (Blitvich et al. 2009, Bolling et al. 2011) and *Cx. tritaeniorhynchus* from Japan (Obara-Nagoya et al. 2013). CxFV appears to perpetuate in mosquito populations by vertical transmission, the passing of the virus from an infected female to her progeny (Saiyasombat et al. 2011, Bolling et al. 2012), and venereal transmission may also play a minor role in viral maintenance (Bolling et al. 2012). Evidence for vertical transmission has also been observed for other ISFs, including KRV (Lutomiah et al. 2007), AeFV (Haddow et al. 2013), and CLBOV (Bolling et al. 2011), on the basis of virus detection in immature mosquitoes.

ISFs may inhibit medically important flaviviruses from establishing infection in mosquitoes through “superinfection exclusion.” This phenomenon occurs when a viral infection establishes itself in a cell, and this infection interferes with the establishment of a secondary infection (Randolph and Hardy 1988, Zou et al. 2009). There are conflicting reports on CxFV-infected mosquitoes’ vector competence for WNV. Laboratory studies have suggested that CxFV infection in *Cx. pipiens* may suppress a secondary infection of WNV up to 7 days postinfection (Bolling et al. 2012). However, prior infection of CxFV was not shown to impact WNV transmission rates in *Cx. quinquefasciatus*, and simultaneous co-infection actually increased the rate of WNV transmission (Kent et al. 2010). Observations from field studies also fail to support the concept of superinfection exclusion of WNV by CxFV (Newman et al. 2011, Crockett et al. 2012). Nonetheless, other studies in Australia have shown that PCV infection suppresses WNV and Murray Valley encephalitis virus replication in mosquito cell culture (Hobson-Peters et al. 2013). These studies show that although ISFs cannot infect humans, they may impact public health by suppressing or enhancing infection of pathogenic flaviviruses in the mosquito vector.

In this study, we describe a novel ISF infecting *Culiseta* (*Cs.*) *melanura* mosquitoes collected from the northeastern United States. *Cs. melanura* is a bird-biting mosquito species that serves as the main vector of eastern equine encephalitis virus (EEEV) and a secondary vector of WNV in eastern North America (Scott and Weaver 1989, Andreadis et al. 2004). To determine whether *Cs. melanura* may be infected by ISFs, we surveyed mosquito populations in the northeastern United States for flavivirus infection using generic and specific PCR primers. The virus was isolated in mosquito cell culture and detected by electron microscopy, and the complete genome was sequenced and compared to other flaviviruses by phylogenetic analysis. We provisionally named it *Culiseta flavivirus* (CsFV) and describe here the annual cycle of CsFV infection in *Cs. melanura* females and overwintering mosquito larvae.

Materials and Methods

Mosquito collections

Adult mosquitoes were collected in 91 trapping locations throughout Connecticut as part of the statewide mosquito and arbovirus surveillance program (Andreadis et al. 2004). A dry ice-baited CDC light trap and gravid trap baited with hay-infusion water were set overnight at each site on a weekly rotation from June through October. Additional *Cs. melanura* were collected in Cheshire, Hillsborough, and Rockingham Counties in New Hampshire and York County, Maine, during June, 2011, and in Oswego County, New York, during May, 2012. Mosquitoes were identified to species on a chill table under a dissecting microscope using diagnostic keys (Andreadis et al. 2005). Female mosquitoes from each trap were pooled by species with a maximum of 50 individuals per pool, and placed into 2.0-mL microcentrifuge tubes containing a copper BB. Pools were stored at -80°C until tested.

Virus detection and partial sequencing

In preparation for virus testing, 1.0 mL of phosphate-buffered saline (PBS) containing 0.5% gelatin, 30% rabbit serum, and $1\times$ antibiotic + antimycotic was added to each mosquito pool. Mosquito pools were homogenized using a MM 300 Mixer-Mill (Retsch Laboratory, Irvine, CA) set at 25 cycles per second for 4 min. Samples were centrifuged at 4°C for 7 min at 7000 rpm, and RNA was extracted from a 70- μL aliquot of the supernatant using the Viral RNA Kit (Qiagen, Valencia, CA). RT-PCR was performed using the Titan One Tube RT-PCR System (Roche Diagnostics, Indianapolis, IN) in 25- μL reactions. Primer sequences are given in Table 1. Primer pairs FU2new/cFD3 and ISFfwd/CSFLAVrev were used to detect flavivirus infection initially in *Cs. melanura* by amplifying portions of the *NS5* gene (Kuno et al. 1998). Amplification was as follows: 1 cycle of 50°C for 30 min and 94°C for 2 min, 10 cycles of 94°C for 15 s, 50°C for 30 s, and 68°C for 1 min, followed by 25 cycles of 94°C for 15 s, 50°C for 30 s, and 68°C for 1 min + 5 s per cycle, and 1 cycle of 68°C for 7 min. RT-PCR amplification products were purified using the QIAquick PCR Purification Kit (Qiagen, Valencia, CA) and sequenced at the DNA Analysis Facility (Yale University, New Haven, CT).

To detect CsFV in all other mosquito pools, primers CSFLAVfwd and CSFLAVrev were used to target specifically and amplify a 255-nucleotide fragment of the *NS5* gene of CsFV. The thermal cycling protocol was as follows: 1 cycle of 50°C for 30 min and 94°C for 2 min, 10 cycles of 94°C for 15 s, 58°C for 30 s, and 68°C for 30 s, followed by 25 cycles of 94°C for 15 s, 58°C for 30 s, and 68°C for 30 s + 5 s per cycle, and one cycle of 68°C for 7 min. Primer set CSLfwd/CSLrev was used to retest samples that had produced ambiguous or faint bands with CSFLAVfwd/CSFLAVrev primers and for sequencing and phylogenetic analyses of selected CsFV strains. These primers amplify a 913 bp portion of the *NS5* gene under the following parameters: One cycle of 50°C for 30 min and 94°C for 2 min, 10 cycles of 94°C for 15 s, 60°C for 30 s, and 68°C for 1 min, followed by 25 cycles of 94°C for 15 s, 60°C for 30 s, and 68°C for 1 min + 5 s per cycle, and one cycle of 68°C for 7 min. Infection rates per 100 individuals were calculated as bias-corrected maximum likelihood estimates using the Excel Add-In PooledInfRate, version 4.0 (Biggerstaff 2006).

TABLE 1. PRIMERS USED FOR RT-PCR AND NUCLEOTIDE SEQUENCING

Primer name	Primer sequence (5' → 3')	Target gene, virus	Amplicon length (bp)	Used for
FU2new	GCCGATGACGTAGCCGGTTGGGATAC	NS5, Flavivirus genus	888	Sequencing
cFD3	AGCATGTCTTCCGTGGTCATCCA	NS5, Flavivirus genus		
ISFfwd	GGATTATCTGGTACATGTGG	NS5, ISFs	518	Sequencing
CSFLAVrev	TAAGCCAGGCGTCAATCTCT	NS5, CsFV		
CSFLAVfwd	CCGTACCACAGGAAACTCGT	NS5, CsFV	255	Virus detection
CSFLAVrev	TAAGCCAGGCGTCAATCTCT	NS5, CsFV		
CSLfwd	CGAGTACGAGGCTCTAGGGT	NS5, CsFV	913	Sequencing, virus confirmation
CSLrev	GACCAGTTGGCATCCACTCA	NS5, CsFV		

ISF, insect-specific flavivirus; CsFV, *Culiseta flavivirus*.

Virus isolation and propagation

Ae. albopictus (C6/36) cells were grown in 25-cm² flasks containing minimum essential medium (MEM), 5% fetal bovine serum (FBS), 1× nonessential amino acids, and antibiotics + antimycotics, and held at 28°C, 5% CO₂. African green monkey (Vero E6) cells and baby hamster kidney (BHK-21) cells were maintained in 25-cm² flasks containing MEM, 5% FBS, and antibiotics + antimycotics at 37°C, 5% CO₂. CsFV was initially isolated in C6/36 cell culture by inoculating 100 μL of mosquito pool homogenate onto confluent cells and by splitting cell cultures for three passages at a 1:5 dilution every 7 days. CsFV isolate 502-13, passage 3, was used to assess its ability to propagate in mosquito (C6/36) and mammalian (Vero E6 and BHK-21) cells by inoculating 100 μL of virus into each cell line in triplicate. Cell cultures were maintained for 7 days, and then 100 μL of the culture medium was used for each subsequent passage for a total of three passages.

Electron microscopy

Five-day-old C6/36 cell cultures previously inoculated with CsFV and cultures that were not inoculated (negative controls) were centrifuged, infused with a 5% agar solution, and fixed overnight at 4°C in a 2.5% (vol/vol) glutaraldehyde + 2% paraformaldehyde solution containing 0.1% (wt/vol) CaCl₂ and 1% (wt/vol) sucrose in 100 mM sodium cacodylate buffer (pH 7.4). Samples were postfixed in aqueous 1% (wt/vol) OsO₄, dehydrated through an ascending ethanol and acetone series, and embedded in LX-112/Araldite (Ladd Research Industries, Williston, VT). Thin sections were poststained with 5% (wt/vol) uranyl acetate in 50% (vol/vol) methanol followed by Reynold's lead citrate and examined in a Zeiss EM 10C electron microscope at an accelerating voltage of 100 kV.

Genomic sequencing

Total RNA (15 μL) from virus-infected C6/36 cell culture supernatants (virus isolate 502-13) were prepared for whole genome sequencing, as previously described (Grubaugh et al. 2015). In brief, the samples were treated with DNase, amplified using the Ovation RNA-Seq System V2 (NuGEN, San Carlos, CA), sheared using the Covaris S2 Focused-ultrasonicator (Covaris, Woburn, MA), and prepared using the Ovation Ultralow Library Kit (NuGEN, San Carlos, CA). Agencourt RNAClean XP beads (Beckman Coulter, Brea,

CA) were used for all purification steps. Finished libraries were analyzed for correct size distribution using the 2100 Bioanalyzer (Agilent, Santa Clara, CA) and were quantified using the SYBR Green I-based qPCR Library Quantification Kit (KAPA Biosystems, Wilmington, MA). Paired-end reads of 100 nucleotides were generated using the Illumina HiSeq 2500 platform at Beckman Coulter Genomics (Danvers, MA). De-multiplexed paired-end reads were assembled *de novo* using Trinity (Grabherr et al. 2011), and the putative viral contigs were aligned and edited manually using Geneious version 7.0.6 (Kearse et al. 2012). The reference-guided aligner MOSAIK (Lee et al. 2014) was used to verify the accuracy of the assembled viral genome.

Phylogenetic analysis

Sequence alignments were generated in MEGA 6.0 based on nucleotide or translated amino acid sequences. Full-length polyprotein amino acid sequences were aligned using the ClustalW algorithm. Ambiguous positions in the alignment were removed using Gblocks (Castresana 2000), and 1449 amino acid positions were retained in the final alignment. A second alignment was generated from an 863-nucleotide portion of the NS5 gene from 22 CsFV strains. The optimal substitution model for each dataset was identified after performing maximum likelihood fits of 24 different models in MEGA. Phylogenetic trees were reconstructed by maximum likelihood analysis in MEGA and support for individual nodes was obtained by performing 1000 bootstrap replicates.

Results

Virus isolation and characterization

To determine whether *Cs. melanura* mosquitoes may be infected by ISFs, we initially screened seven pools consisting of 249 mosquitoes collected in Connecticut using PCR primers (FU2new and cFD3) targeting a highly conserved region of the *Flavivirus NS5* gene. Three *Cs. melanura* pools yielded bands of the expected size (888 bp). The amplification product from one pool was sequenced to design new primers (ISFfwd and CSFLAVrev) to amplify an overlapping region of the NS5 gene of 518 bp. A partial NS5 nucleotide sequence (960 bp) was compared to sequences on GenBank and most closely matched CLBOV from *Cx. tarsalis*, sharing 68.7% nucleotide and 76.1% amino acid sequence identity. This level of sequence divergence is consistent with distinct viral species for the *Flavivirus* genus (Kuno et al. 1998). The

virus was provisionally named *Culiseta flavivirus* (CsFV) on the basis of its host association with *Cs. melanura*.

To isolate and propagate CsFV, positive mosquito pools were inoculated into C6/36 cell cultures and blind-passaged three times. Two virus isolates (50-13 and 502-13) were identified when screened by RT-PCR using virus-specific primers (CSFLAVfwd and CSFLAVrev); however, cytopathic effect was not apparent in virus-positive cell cultures. CsFV isolate 50-13 was then inoculated and serially passaged in mosquito (C6/36) and mammalian (Vero E6 and BHK-21) cell lines. CsFV appeared to replicate only within the mosquito cell line. CsFV RNA was detected by RT-PCR after three passages in C6/36 cells, but was not detected after the second and third passage in Vero and BHK cells.

When examined by transmission electron microscopy, infected C6/36 cell cultures revealed spherical virus particles surrounded by a faint outer envelope membrane measuring 40 nm in diameter. These were located within the cisternae of the endoplasmic reticulum (Fig. 1A) and were not observed in noninoculated cell cultures. Naked nucleocapsid precursors were not detected. Dense accumulations of mature virus particles associated with distended cisternae of endoplasmic reticulum characterized by intercisternal invaginations of tubular membranes were further seen in disrupted cells (Fig. 1B). These characteristics are consistent with insect cell lines infected with a flavivirus (Grimley 1991).

Genomic sequence and phylogenetic analysis

The complete genome of CxFV was sequenced and assembled (GenBank acc. no. KT599442). The sequence was based on a total of 6.8 million reads, of which 58.8% aligned to CsFV. The genome consists of 10,864 nucleotides with a primary open reading frame (ORF) from nucleotides 92 to 10,282 that encodes a 3396-amino acid polyprotein. The CsFV coding region was most similar to that of Naikwogo virus, PCV, and CxFV (49.1–49.4% nucleotide identity);

however, the complete polyprotein sequence of the aforementioned CLBOV was not included in this analysis because it was not available on GenBank. Common to other ISFs, a “slippery” programmed –1 ribosomal frameshift motif (GGAUUUU) was detected at nucleotide position 3343 near the expected NS1 and NS2A junction (Blitvich and Firth 2015). This generated a second predicted overlapping ORF encoding a novel 247-amino-acid protein, previously termed *fifo*, from nucleotides 3508 to 4248 (Firth et al. 2010).

Phylogenetic relationships of 42 flaviviruses were evaluated by maximum likelihood analysis of polyprotein amino acid sequences, and the resulting phylogenetic tree is shown in Figure 2. The overall topology of the tree was similar to that of other analyses, with viruses clustering into groups defined by host range—ISFs, tick-borne flaviviruses, mosquito-borne flaviviruses, and no known vector flaviviruses (Cook et al. 2012). The recently discovered ISF-like viruses (ISF-2) fell within the mosquito-borne flavivirus group, as previously described (Kenney et al. 2014, Blitvich and Firth 2015). CsFV belonged to the classical ISF group but represents a distinct evolutionary lineage that grouped with ISFs associated with *Culex*, *Mansonia*, and *Coquillettidia* spp. mosquitoes (Naikwogo virus, PCV, Nienokoue virus, CxFV, Quang Binh virus, and *Culex theileri* flavivirus).

Prevalence of CsFV in field-collected mosquitoes

To evaluate the host range of CsFV in field-collected mosquitoes, we tested 3620 mosquitoes representing 20 species that were collected in Connecticut from 2012 to 2013 (Table 2). CsFV was detected in 24 pools of *Cs. melanura* ($n = 584$, 33 pools) by RT-PCR, and the maximum likelihood estimate-infection rate (MLE-IR) was 9.2/100 mosquitoes. All other mosquito species were negative by RT-PCR. CsFV was then detected in additional pools of *Cs. melanura* collected from Maine, New Hampshire, and New York (Table 3). The MLE-IR was similar among states and ranged




FIG. 1. Section of 5-day-old C6/36 cell cultures infected with CsFV. (A) Spherical virus particles associated with cisternae of the endoplasmic reticulum. N, cell nucleus. Magnification, 31,000 \times . (B) Dense accumulations of mature virus particles associated with distended cisternae of endoplasmic reticulum and intercisternal tubular membranes. Magnification 48,000 \times . (Inset) Higher magnification of mature enveloped virions. Magnification, 155,000 \times .




FIG. 2. Maximum likelihood tree based on polyprotein amino acid sequences. Labels include virus names and corresponding GenBank accession numbers. Numbers at nodes indicate bootstrap support values. MBF, mosquito-borne flaviviruses; NKVF, no known vector flaviviruses; TBF, tick-borne flaviviruses; ISF, insect-specific flaviviruses.

TABLE 2. DETECTION OF *CULISETA FLAVIVIRUS* IN FEMALE MOSQUITOES COLLECTED IN CONNECTICUT 2012–2013

<i>Mosquito species</i>	<i>No. mosquitoes</i>	<i>No. mosquito pools</i>	<i>No. pools positive</i>	<i>MLE-IR/100 mosquitoes (95% CI)</i>
<i>Aedes cinereus</i>	182	7	0	
<i>Aedes vexans</i>	421	9	0	
<i>Anopheles punctipennis</i>	153	5	0	
<i>Anopheles walker</i>	115	4	0	
<i>Coquillettidia perturbans</i>	239	8	0	
<i>Culex pipiens</i>	366	20	0	
<i>Culex salinarius</i>	272	6	0	
<i>Culex restuans</i>	124	6	0	
<i>Culiseta melanura</i>	584	33	24	9.2 (6.0–14.5)
<i>Culiseta minnesotae</i>	5	5	0	
<i>Culiseta morsitans</i>	23	16	0	
<i>Ochlerotatus canadensis</i>	260	9	0	
<i>Ochlerotatus cantator</i>	16	1	0	
<i>Ochlerotatus japonicus</i>	15	2	0	
<i>Ochlerotatus sollicitans</i>	26	1	0	
<i>Ochlerotatus taeniorhynchus</i>	135	3	0	
<i>Ochlerotatus triseriatus</i>	128	4	0	
<i>Ochlerotatus trivittatus</i>	122	4	0	
<i>Psorophora ferox</i>	188	6	0	
<i>Uranotaenia sapphirina</i>	246	9	0	
Total	3620	158	24	

TABLE 3. DETECTION OF CULISETA FLAVIVIRUS IN *Cs. MELANURA* COLLECTED FROM MAINE, NEW HAMPSHIRE, AND NEW YORK DURING 2011–2012

State	Sex	No. mosquitoes	No. mosquito pools	No. pools positive	MLE-IR/100 mosquitoes (95% CI)
Maine	F	164	12	8	9.6 (4.5–22.2)
New Hampshire	F	251	26	12	8.7 (4.7–15.9)
New York	F	90	9	6	9.5 (4.1–20.8)
New York	M	102	11	7	9.4 (4.4–18.8)

MLE-IR, maximum likelihood estimate-infection rate; CI, confidence interval.

from 8.7 to 9.6/100 mosquitoes. CsFV was detected in male and female mosquitoes from New York and both sexes yielded similar MLE-IRs (9.4–9.5/100 mosquitoes). Culiseta virus was detected solely in *Cs. melanura* mosquitoes and was found in all of the northeastern states surveyed.

Seasonal and spatial dynamics of CsFV in Connecticut

To characterize the seasonal and spatial dynamics of CsFV in Connecticut, additional *Cs. melanura* were collected and tested from 32 sites from June 3 through October 17, 2013. Each site was trapped on a 7- to 10-day rotation as part of the statewide mosquito surveillance program. A total of 7178 *Cs. melanura* females were processed as 474 mosquito pools for virus testing by RT-PCR, and 274 of these pools tested positive for CsFV. The MLE-IR for all of the tested *Cs. melanura* adults was 8.5 (per 100 mosquitoes). For the temporal and spatial analyses, two of 32 sites, OL22 and FR13, were omitted due to the low number of *Cs. melanura* trapped at each site (<20 mosquitoes). CsFV was detected during all 20 weeks of the study (Fig. 3), and although MLE-IRs fluctuated from week to week, they remained fairly stable, ranging from 3.8 to 11.5. In addition, the MLE-IR was calculated for each site and plotted on a map of Connecticut

(Fig. 4). CsFV was present at all 30 sites, and the site infection rates ranged from 3.4 to 50.9.

To investigate the overwintering portion of the annual virus cycle, pools of *Cs. melanura* larvae were collected from Chester, Connecticut (site CH23), during the collection period of November 6, 2012 to May 3, 2013. A total of 5822 mosquito larvae were processed as 144 pools for virus testing, and 69 of these pools tested positive for CsFV. The MLE-IR for all of the tested *Cs. melanura* larvae was 1.7 (per 100 larvae), and monthly infection rates were relatively stable throughout this time period, ranging from 0.7 to 2.1 (Fig. 5).

Genetic diversity of CsFV

To evaluate the genetic relationships of CsFV strains circulating in the northeastern United States, we sequenced part of the NS5 gene (863 nucleotides) from 22 viral strains and compared these sequences by phylogenetic analysis (Fig. 6). We also included three CsFV strains that were detected retrospectively in historical pools of *Cs. melanura* collected during 1998–2001, in addition to more recent strains from Connecticut, Maine, New Hampshire, and New York. Mean nucleotide distances were 1.03% among all CsFV strains





FIG. 3. Weekly infection rates of Culiseta flavivirus (CsFV) in *Cs. melanura* females collected at 30 sites in Connecticut between June and October of 2013. Error bars show 95% confidence intervals. Numbers in parentheses indicate infection rates for each month.

FIG. 4. County map of Connecticut showing geographic location of *Cs. melanura* infected with Culiseta flavivirus (CsFV) during 2013. Pie charts show the estimated proportion of infected *Cs. melanura* (indicated by white sector) per 100 individuals at each site. Pie chart size indicates the total number of *Cs. melanura* trapped at each site during the study period. Infection rates are not shown for sites (OL22 and FR13) that produced fewer than 20 *Cs. melanura*.



analyzed. Viruses segregated into two major clades by maximum likelihood analysis with 70% and 88% bootstrap support, but without distinct geographic or temporal patterns. Phylogenetic analysis indicates that viral genetic variation is limited in the northeastern United States and lacks obvious temporal and spatial definition.

Discussion

In this study, we describe a new ISF, termed CsFV, that was detected solely in *Cs. melanura* mosquitoes and found in all northeastern US states surveyed. CsFV represents the first ISF known to infect mosquitoes of the genus *Culiseta*, indicating that these single-host flaviviruses occur in diverse mosquito genera. CsFV was successfully isolated and visualized in mosquito cells, but failed to propagate in mammalian cell lines. The genome comprised a primary ORF

encoding a viral polyprotein characteristic of the genus *Flavivirus* and a second overlapping ORF expressing a novel gene *fifo* due to a ribosomal frameshift that is common to other ISFs. Our phylogenetic analyses showed that CsFV represents a distinct evolutionary lineage that belongs to the classical ISF group.

CsFV was found to be very prevalent in northeastern US populations of *Cs. melanura*. The estimated MLE-IR for pools of adult females was 8.7–9.6 individuals per 100 mosquitoes collected in Connecticut, Maine, New Hampshire, and New York. This infection rate is at the lower end of the range of annual MLE-IRs calculated for CxFV in *Cx. pipiens* from Colorado, which ran from 7.2 to 46.2, but greater than ranges of annual infection rates of CxFV in *Cx. tarsalis*, 0.08–0.13, and of CLBOV in both *Cx. pipiens* (0–0.3) and *Cx. tarsalis* (0.3–4.0) from Colorado (Bolling et al. 2011), and similar to MLE-IRs for CxFV infection in *Culex*




FIG. 5. Monthly infection rates of Culiseta flavivirus (CsFV) in overwintering *Cs. melanura* larvae collected in Chester, Connecticut, between November of 2012 and May of 2013. Error bars indicate 95% confidence interval (CIs).




FIG. 6. Maximum likelihood tree of *Culiseta flavivirus* (CsFV) strains based on partial *NS5* nucleotide sequences. Labels specify GenBank accession number, geographic origin, and year of virus isolation of virus strains. Numbers at nodes indicate bootstrap support values >60%.

pools from Illinois (10.0–11.2) (Newman et al. 2011). The estimated CsFV infection rate falls well within the range of infection rates for other ISFs described in the literature.

To characterize spatial and temporal patterns of CsFV infection, *Cs. melanura* females were collected and tested from 30 sites in Connecticut between June and October, 2013. During the study period, weekly MLE-IRs per 100 mosquitoes ranged from 3.8 to 11.5 (Fig. 3). Although there were fluctuations in the MLE-IRs from week to week, overlapping 95% confidence intervals (CIs) indicate that the infection rate did not differ significantly during the study period. This differs from the weekly infection rates observed for another flavivirus, WNV, which cycles between bird and mosquito populations. In Connecticut, WNV is typically absent in mosquito populations until early July, at which point it begins to increase in prevalence before peaking in August, and then declining into early fall (Andreadis et al. 2004). This suggests that CsFV does not undergo seasonal amplification via horizontal transmission, as WNV does, and that a different mechanism acts to maintain CsFV in *Cs. melanura* populations.

Geographically, CsFV was found at all test sites in Connecticut with a sample size of >20 *Cs. melanura* per site (Fig. 4). The calculated MLE-IRs at these sites ranged between 3.4 and 50.9 infected individuals per 100 mosquitoes. Similar infection rates were found throughout the state, with the exception of site BB17 (MLE-IR = 50.9/100 mosquitoes), which was more than twice the next highest infection rate. We cannot explain this result either by a low sample size or by characteristics of the mosquito trapping site. BB17 is characterized as a woodland depression bordered by a forested wetland complex that is typical for this region. CsFV appears to be widespread and was found at all sites with a sufficient sample of more than 20 *Cs. melanura* mosquitoes.

Monthly MLE-IRs were calculated for *Cs. melanura* larvae collected from Chester, Connecticut, between November, 2012, and May, 2013 (Fig. 5). As with the MLE-IRs of CsFV in adults, the monthly larvae infection rates were very stable. For the most part, the infection rate per 100 larvae ranged from 1.1 to 2.1. The only month when the MLE-IR was outside of this range was March of 2013 (MLE-IR = 0.7). However, this anomaly is likely due to the small sample size for that month. A large snowstorm in February interfered with collection efforts, and only four pools containing 148 larvae were collected in March, of which one pool tested positive for CsFV. The infection rate for CsFV in overwintering larval *Cs. melanura* appears to be stable over the winter months. CsFV detection in larvae suggests that vertical transmission is occurring, and stable infection rates suggest that horizontal transmission is not occurring between larvae during the overwintering period. This reinforces observations of other ISFs, which indicate that vertical transmission is the major mode of transmission (Lutomiah et al. 2007, Bolling et al. 2012).

An attempt was made to compare the MLE-IRs of overwintering larval *Cs. melanura* from Chester, Connecticut (1.7/100, 95% CI = 1.3–2.1), and adults trapped at the same site during the summer and fall of 2013 (4.0/100, 95% CI = 1.7–9.2). While a higher MLE-IR was calculated for the adult *Cs. melanura*, these differences were not significantly different on the basis of overlapping 95% CIs. Even if the larval and adult infection rates differed significantly, this is a difficult comparison to make because the larvae and adults were collected during different time periods. We attempted to collect *Cs. melanura* larvae during the fall of 2013, so that larvae and adults from the same location and time period could be compared, but this was prevented by extremely dry conditions.

Phylogenetic analyses were also performed to evaluate the genetic diversity and relationships of CsFV strains circulating in the northeastern United States (Fig. 6). Viral sequences proved to be highly conserved and grouped into two well-supported clades, but without evidence of spatial or temporal clustering. This indicates that CsFV has not undergone local isolation and differentiation among sampled locations in the northeastern United States. However, further testing of *Cs. melanura* from the mid-Atlantic and southeastern regions of the United States is needed to determine the prevalence of infection and phylogeography of this virus. CsFV could also potentially serve as a surrogate marker to identify geographically isolated *Cs. melanura* populations, given that this RNA virus likely mutates and differentiates faster than its mosquito host and appears to be stably maintained by vertical transmission. Future studies are warranted to evaluate the geographic range and population structure of CsFV.

In conclusion, we discovered and characterized CsFV that was found at high infection rates in *Cs. melanura* populations in the northeastern United States. Many of the findings, including the presence of the virus in larvae and the stable infection rates throughout the study period, suggest that CsFV is maintained in *Cs. melanura* populations through vertical transmission, and that it does not amplify through horizontal transmission. Studies need to be performed to determine whether the high prevalence of CsFV infection throughout Connecticut may be interfering with the transmission of WNV, a widespread flavivirus and human pathogen, through superinfection exclusion. Further studies are needed to determine the impact of CsFV on WNV infections in *Cs. melanura*.

Acknowledgments

We thank our support staff, specifically Angela Bransfield, John Shepard, and Michael Thomas, who provided technical assistance in the collection, identification, and testing of mosquitoes. This work was supported in part by grants from the Centers for Disease Control and Prevention (U50/CCU116806-01-1), the US Department of Agriculture Hatch Funds (CONH00773), the Multistate Research Project (NE1043), and the National Institutes of Health (AI067380).

Author Disclosure Statement

No competing financial interests exist.

References

- Andreadis TG, Anderson JF, Vossbrinck CR, Main AJ. Epidemiology of West Nile virus in Connecticut: A five-year analysis of mosquito data 1999–2003. *Vector Borne Zoonotic Dis* 2004; 4:360–378.
- Andreadis TG, Thomas MC, Shepard JJ. *Identification Guide to the Mosquitoes of Connecticut*. Bulletin of The Connecticut Agricultural Experiment Station, No. 966, 2005:1–173.
- Biggerstaff BL. PooledInfRate, Version 3.0: A Microsoft Excel Add-In to Compute Prevalence Estimates from Pooled Samples. Centers for Disease Control and Prevention, Fort Collins, CO, 2006.
- Blitvich BJ, Firth AE. Insect-specific flaviviruses: A systematic review of their discovery, host range, mode of transmission, superinfection exclusion potential and genomic organization. *Viruses* 2015; 7:1927–1959.
- Blitvich BJ, Lin M, Dorman KS, Soto V, et al. Genomic sequence and phylogenetic analysis of *Culex* flavivirus, an insect-specific flavivirus, isolated from *Culex pipiens* (Diptera: Culicidae) in Iowa. *J Med Entomol* 2009; 46:934–941.
- Bolling BG, Eisen L, Moore CG, Blair CD. Insect-specific flaviviruses from *Culex* mosquitoes in Colorado, with evidence of vertical transmission. *Am J Trop Med Hyg* 2011; 85:169–177.
- Bolling BG, Olea-Popelka FJ, Eisen L, Moore CG, et al. Transmission dynamics of an insect-specific flavivirus in a naturally infected *Culex pipiens* laboratory colony and effects of co-infection on vector competence for West Nile virus. *Virology* 2012; 427:90–97.
- Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 2000; 17:540–552.
- Cook S, Moureau G, Harbach RE, Mukwaya L, et al. Isolation of a novel species of flavivirus and a new strain of *Culex* flavivirus (Flaviviridae) from a natural mosquito population in Uganda. *J Gen Virol* 2009; 90:2669–2678.
- Cook S, Moureau G, Kitchen A, Gould EA, et al. Molecular evolution of the insect-specific flaviviruses. *J Gen Virol* 2012; 93:223–234.
- Crabtree MB, Sang RC, Stollar V, Dunster LM, et al. Genetic and phenotypic characterization of the newly described insect flavivirus, Kamiti River virus. *Arch Virol* 2003; 148:1095–1118.
- Crockett RK, Burkhalter K, Mead D, Kelly R, et al. *Culex* flavivirus and West Nile virus in *Culex quinquefasciatus* populations in the southeastern United States. *J Med Entomol* 2012; 49:165–174.
- Farfan-Ale JA, Lorono-Pino MA, Garcia-Rejon JE, Soto V, et al. Detection of flaviviruses and orthobunyaviruses in mosquitoes in the Yucatan Peninsula of Mexico in 2008. *Vector Borne Zoonotic Dis* 2010; 10:777–783.
- Firth AE, Blitvich BJ, Wills NM, Miller CL, et al. Evidence for ribosomal frameshifting and a novel overlapping gene in the genomes of insect-specific flaviviruses. *Virology* 2010; 399:153–166.
- Grabherr MG, Haas BJ, Yassour M, Levin JZ, et al. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol* 2011; 29:644–652.
- Grimley PM. Togaviridae and Flaviviridae. In: Adams JR, Bonami JR, eds. *Atlas of Invertebrate Viruses*. Boca Raton, FL: CRC Press, 1991:461–497.
- Grubaugh ND, Smith DR, Brackney DE, Bosco-Lauth AM, et al. Experimental evolution of an RNA virus in wild birds: evidence for host-dependent impacts on population structure and competitive fitness. *PLoS Pathog* 2015; 11:e1004874.
- Haddow AD, Guzman H, Popov VL, Wood TG, et al. First isolation of *Aedes* flavivirus in the Western Hemisphere and evidence of vertical transmission in the mosquito *Aedes* (*Stegomyia*) *albopictus* (Diptera: Culicidae). *Virology* 2013; 440:134–139.
- Hobson-Peters J, Yam AW, Lu JW, Setoh YX, et al. A new insect-specific flavivirus from northern Australia suppresses replication of West Nile virus and Murray Valley encephalitis virus in co-infected mosquito cells. *PLoS One* 2013; 8:e56534.
- Hoshino K, Isawa H, Tsuda Y, Yano K, et al. Genetic characterization of a new insect flavivirus isolated from *Culex pipiens* mosquito in Japan. *Virology* 2007; 359:405–414.
- Hoshino K, Isawa H, Tsuda Y, Sawabe K, et al. Isolation and characterization of a new insect flavivirus from *Aedes albopictus* and *Aedes flavopictus* mosquitoes in Japan. *Virology* 2009; 391:119–129.

- Huhtamo E, Putkuri N, Kurkela S, Manni T, et al. Characterization of a novel flavivirus from mosquitoes in northern Europe that is related to mosquito-borne flaviviruses of the tropics. *J Virol* 2009; 83:9532–9540.
- Junglen S, Kopp A, Kurth A, Pauli G, et al. A new flavivirus and a new vector: Characterization of a novel flavivirus isolated from *Uranotaenia* mosquitoes from a tropical rain forest. *J Virol* 2009; 83:4462–4468.
- Kearse M, Moir R, Wilson A, Stones-Havas S, et al. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 2012; 28:1647–1649.
- Kenney JL, Solberg OD, Langevin SA, Brault AC. Characterization of a novel insect-specific flavivirus from Brazil: Potential for inhibition of infection of arthropod cells with medically important flaviviruses. *J Gen Virol* 2014; 95:2796–2808.
- Kent RJ, Crabtree MB, Miller, BR. Transmission of West Nile virus by *Culex quinquefasciatus* say infected with *Culex* Flavivirus Izabal. *PLoS Negl Trop Dis* 2010; 4:e671.
- Kim DY, Guzman H, Bueno R, Dennett JA, et al. Characterization of *Culex* flavivirus (Flaviviridae) strains isolated from mosquitoes in the United States and Trinidad. *Virology* 2009; 386:154–159.
- Kolodziejek J, Pachler K, Bin H, Mendelson E, et al. Barkedji virus, a novel mosquito-borne flavivirus identified in *Culex perexiguus* mosquitoes, Israel, 2011. *J Gen Virol* 2013; 94:2449–2457.
- Kuno G, Chang GJ, Tsuchiya KR, Karabatsos N, et al. Phylogeny of the genus *Flavivirus*. *J Virol* 1998; 72:73–83.
- Lee JS, Grubaugh ND, Kondig JP, Turell MJ, et al. Isolation and genomic characterization of Chaoyang virus strain ROK144 from *Aedes vexans nipponii* from the Republic of Korea. *Virology* 2013; 435:220–224.
- Lee WP, Stromberg MP, Ward A, Stewart C, et al. MOSAIK: A hash-based algorithm for accurate next-generation sequencing short-read mapping. *PLoS One* 2014; 9:e90581.
- Lutumiah JJ, Mwandawiro C, Magambo J, Sang RC. Infection and vertical transmission of Kamiti River virus in laboratory bred *Aedes aegypti* mosquitoes. *J Insect Sci* 2007; 7:1–7.
- Machado DC, Mondini A, dos Santos Santana V, Yonamine PT, et al. First identification of *Culex* flavivirus (Flaviviridae) in Brazil. *Intervirology* 2012; 55:475–483.
- Morales-Betoulle ME, Monzon Pineda ML, Sosa SM, Panella N, et al. *Culex* flavivirus isolates from mosquitoes in Guatemala. *J Med Entomol* 2008; 45:1187–1190.
- Newman CM, Cerutti F, Anderson TK, Hamer GL, et al. *Culex* flavivirus and West Nile virus mosquito coinfection and positive ecological association in Chicago, United States. *Vector Borne Zoonotic Dis* 2011; 11:1099–1105.
- Obara-Nagoya M, Yamauchi T, Watanabe M, Hasegawa S, et al. Ecological and genetic analyses of the complete genomes of *Culex* flavivirus strains isolated from *Culex tritaeniorhynchus* and *Culex pipiens* (Diptera: Culicidae) group mosquitoes. *J Med Entomol* 2013; 50:300–309.
- Randolph VB, Hardy JL. Phenotypes of St Louis encephalitis virus mutants produced in persistently infected mosquito cell cultures. *J Gen Virol* 1988; 69:2199–2207.
- Saiyasombat R, Bolling BG, Brault AC, Bartholomay LC, et al. Evidence of efficient transovarial transmission of *Culex* flavivirus by *Culex pipiens* (Diptera: Culicidae). *J Med Entomol* 2011; 48:1031–1038.
- Scott TW, Weaver SC. Eastern equine encephalomyelitis virus: Epidemiology and evolution of mosquito transmission. *Adv Virus Res* 1989; 37:277–328.
- Stollar V, Thomas VL. An agent in the *Aedes aegypti* cell line (Peleg) which causes fusion of *Aedes albopictus* cells. *Virology* 1975; 64:367–377.
- Tyler S, Bolling BG, Blair CD, Brault AC, et al. Distribution and phylogenetic comparisons of a novel mosquito flavivirus sequence present in *Culex tarsalis* mosquitoes from western Canada with viruses isolated in California and Colorado. *Am J Trop Med Hyg* 2011; 85:162–168.
- Zou G, Zhang B, Lim PY, Yuan Z, et al. Exclusion of West Nile virus superinfection through RNA replication. *J Virol* 2009; 83:11765–11776.

Address correspondence to:

Philip M. Armstrong

Center for Vector Biology and Zoonotic Diseases

The Connecticut Agricultural Experiment Station

123 Huntington Street

PO Box 1106

New Haven, CT 06504

E-mail: philip.armstrong@ct.gov