

# Tristetraprolin mediates the anti-proliferative effects of metformin in breast cancer cells

Indira Pandiri<sup>1</sup> · Yingqing Chen<sup>1</sup> · Yeonsoo Joe<sup>1</sup> · Hyo Jeong Kim<sup>1</sup> · Jeongmin Park<sup>1</sup> · Hun Taeg Chung<sup>1</sup> · Jeong Woo Park<sup>1</sup>

Received: 12 October 2015 / Accepted: 2 March 2016 / Published online: 8 March 2016  
© The Author(s) 2016. This article is published with open access at Springerlink.com

**Abstract** Metformin, which is a drug commonly prescribed to treat type 2 diabetes, has anti-proliferative effects in cancer cells; however, the molecular mechanisms underlying this effect remain largely unknown. The aim is to investigate the role of tristetraprolin (TTP), an AU-rich element-binding protein, in anti-proliferative effects of metformin in cancer cells. p53 wild-type and p53 mutant breast cancer cells were treated with metformin, and expression of TTP and c-Myc was analyzed by semi-quantitative RT-PCR, Western blots, and promoter activity assay. Breast cancer cells were transfected with siRNA against TTP to inhibit TTP expression or c-Myc and, after metformin treatment, analyzed for cell proliferation by MTS assay. Metformin induces the expression of tristetraprolin (*TTP*) in breast cancer cells in a *p53*-independent manner. Importantly, inhibition of *TTP* abrogated the anti-proliferation effect of metformin. We observed that metformin decreased *c-Myc* levels, and ectopic expression of *c-Myc* blocked the effect of metformin on *TTP* expression and cell proliferation. Our data indicate that metformin induces *TTP* expression by reducing the expression of *c-Myc*, suggesting a new model whereby TTP acts as a mediator of metformin's anti-proliferative activity in cancer cells.

**Keywords** Metformin · Tristetraprolin · p53 · c-Myc · Anti-proliferation

## Introduction

Metformin is a first-line hypoglycemic agent used for the treatment of type 2 diabetes (T2D) [1]. Recently, metformin has received increased attention because of its potential anti-tumorigenic effects [2–4]. Metformin exerts its effects by disrupting mitochondrial respiratory chain 1, which leads to decreased ATP synthesis and increased AMP associated with AMPK, ultimately leading to AMPK activation [5]. This regulation of AMPK by metformin leads to its anti-proliferative effects due to subsequent modification of the activity of mammalian target of rapamycin (mTOR) [6, 7] and p53 [8]. On the other hand, synthetic AMPK agonists provide a proliferative advantage to the cells [9], and there is emerging evidence to suggest that metformin can arrest cell proliferation in an AMPK-independent manner [9–15]. Specifically, metformin has been reported to affect several other intracellular pathways in tumor cells including HER1/HER2, Src, S6K1, c-Myc, and STAT3 [8, 16–19] and is also able to overcome dietary restriction resistance in cancer cells [20]. Overall, however, the mechanistic aspects of metformin action with respect to its anti-proliferative functions remain ill-defined.

AU-rich elements (AREs) post-transcriptionally regulate the expression of a variety of short-lived mRNAs such as cytokines and proto-oncogenes [21]. The stability of ARE-containing mRNAs is regulated by ARE-binding proteins [22]. One of the best-characterized ARE-binding proteins is tristetraprolin (TTP, ZFP36), which promotes the degradation of ARE-containing transcripts [23, 24]. *TTP* expression is significantly decreased in various cancers

Indira Pandiri and Yingqing Chen authors have been contributed equally to this work.

✉ Hun Taeg Chung  
chung@ulsan.ac.kr

✉ Jeong Woo Park  
jwpark@ulsan.ac.kr

<sup>1</sup> Department of Biological Sciences, University of Ulsan, Ulsan 680-749, Korea

[25], which correlates with increased expression of proto-oncogenes and may contribute to cancer processes. Likewise, re-expression of *TTP* has a growth inhibitory effect [26–28]. The expression of *TTP* in cancer cells is induced by *p53* [29] but inhibited by *Myc* [30]. However, nearly all types of cancers have abnormalities in the *p53* pathway [31]. Furthermore, *c-Myc* is often activated in human cancers [32]. Together, these features may lead to a widespread decrease in the expression of *TTP* in human cancers.

We show here for the first time that metformin induces the expression of *TTP* in a *p53*-independent manner, and also that *TTP* mediates the anti-proliferative effect of metformin in both *p53* wild-type and *p53* mutant cancer cells. Specifically, metformin decreased the expression of *c-Myc* and increased the expression of *TTP* in both *p53* wild-type and *p53* mutant cells. Ectopic expression of *c-Myc* abrogated the effects of metformin with respect to *TTP* induction, while siRNA-mediated inhibition of *TTP* attenuated the anti-proliferative effects of metformin. Together, these studies identify a novel signaling pathway by which metformin induces *TTP* expression in a *p53*-independent manner, representing a possible novel pharmacological approach to treat *p53* mutant cancer cells.

## Methods and materials

### Cell culture

The human MCF7 and MDA-MB-231 breast cancer cell lines were purchased from the Korean Cell Line Bank (Seoul, Korea). Cells were cultured in RPMI 1640 media supplemented with 10 % heat-inactivated fetal bovine serum (FBS) (Welgene, Korea) and were maintained at 37 °C in a humidified 5 % CO<sub>2</sub> atmosphere. To investigate the induction of *TTP*, cells were treated with metformin (D150959 Sigma) in the presence or absence of 20 μM compound C (P5499 Sigma). Cells were harvested at the indicated length of time and analyzed for mRNA by RT-PCR, protein by Western blotting, and cell viability by MTS assay.

### Cell viability/proliferation

For the MTS cell proliferation assay, cells were plated in triplicate at  $1.0 \times 10^4$  cells/well in 96-well culture plates in culture media. At 24 h after plating, CellTiter 96<sup>®</sup> Aqueous One Solution reagent (Promega) was added to each well according to the manufacturer's instructions, and absorbance at 490 nm was determined for each well using a Victor 1420 Multilabel Counter (EG&G Wallac, Turku, Finland).

### Plasmids, small interfering RNAs, transfections, and dual-luciferase assay

The pcDNA6/V5-TTP containing full-length ORF of human *TTP* [33] and the pGL3/TTPp-1343 containing human *TTP* promoter [29] were described previously. The pcDNA3-cMyc vector was purchased from Addgene.

For luciferase assays, cells were co-transfected with a pGL3/TTPp-1343-luciferase reporter construct and pRL-SV40 Renilla luciferase construct using TurboFect<sup>™</sup> in vitro transfection reagent (Fermentas). Transfected cells were lysed with lysis buffer and mixed with luciferase assay reagent (Promega). The chemiluminescent signal was measured using a SpectraMax L Microplate (Molecular Devices, Sunnyvale, CA, USA). Firefly luciferase was normalized to Renilla luciferase in each sample. All luciferase assays reported in this study represent at least three independent experiments, each consisting of three wells per transfection.

Small interfering RNAs (siRNAs) against human *TTP* (*TTP*-siRNA, sc-36761), human *c-Myc* (*c-Myc*-siRNA, sc-29226), and control siRNA [scrambled siRNA (scRNA), sc-37007] were purchased from Santa Cruz Biotechnology (Santa Cruz). Cells were transfected 24 h after plating using Lipofectamine<sup>™</sup> RNAiMAX (Invitrogen) and were harvested at 48 h after transfection. The expression levels of *TTP* or *c-Myc* mRNA and protein were analyzed by RT-PCR and Western blotting, respectively.

### SDS-PAGE analysis and immunoblotting

Proteins were resolved by SDS-PAGE, transferred onto Hybond-P membranes (Amersham Biosciences Inc.), and probed with appropriate dilutions of the following antibodies: rabbit anti-human *TTP* (T5327, Sigma), anti-human *c-Myc* (sc-40, Santa Cruz), anti-p53 (1026-1, Epitomics), anti-phospho-p53 (#9284, Cell Signaling), anti-AMPK (#2603, Cell Signaling), anti-phospho-AMPK (#2535, Cell Signaling), anti-STAT3 (#12640, Cell Signaling), anti-phospho-STAT3 (#9134, Cell Signaling), and anti-β-actin (A2228, Sigma). Immunoreactivity was detected using an ECL detection system (Amersham Biosciences Inc.). Films were exposed at multiple time points to ensure that the images were not saturated.

### Quantitative real-time PCR and semi-qRT-PCR

DNase I-treated total RNA (3 mg) was reverse transcribed using oligo-dT and Superscript II reverse transcriptase (Invitrogen) according to the manufacturer's instructions. qRT-PCR was performed by real-time monitoring of the increase in fluorescence of SYBR Green dye (QIAGEN, Hilden, Germany) using a StepOnePlus<sup>™</sup> real-time PCR system

(Applied Biosystems). Semi-qRT-PCR was performed using Taq polymerase (Solgent, Daejeon, Korea). The PCR primer pairs were as follows: *TTP*: 5'-CGCTACAA-GACTGAGCTAT-3', 5'-GAGGTAGAAGTTGTGACAG A-3'; *c-Myc*: 5'-ACAGCATACATCCTGTCCGTCCAA-3', 5'-TGTTCTCGTCGTTTCCGCAACAAG-3'; GAPDH: 5'-ACATCAAGAAGGTGGTGAAG-3', 5'-CTGTTGCTGTAGCCAAATTC-3'.

### Statistical analysis

For statistical comparisons, *p* values were determined using Student's *t* test.

## Results

### Metformin induces tristetrapiolin expression in both p53 wild-type and p53 mutant breast cancer cells

We previously reported that *p53* is required for *TTP* induction in cancer cells [29]. In addition, metformin can enhance *p53* activity [8], suggesting the possibility that metformin can induce *TTP* expression in *p53* wild-type cancer cells. To evaluate this possibility, we treated *p53* wild-type MCF7 breast cancer cells with metformin. As expected, treatment with metformin increased the mRNA and protein levels of *TTP* in MCF7 cells in a dose- (Fig. 1a) and time-dependent (Fig. 1b) manner. Metformin treatment enhanced *TTP* promoter activity (Fig. 1c), indicating that metformin enhances *TTP* gene expression in MCF7 cells. To test whether *p53* activity is required for *TTP* induction by metformin, we treated MDA-MB-231 breast cancer cells, which have a mutant *p53* status, with metformin and analyzed the induction of *TTP*. We confirmed that, while metformin increased phosphorylation of *p53* in MCF7 cells, it did not do so in MDA-MB-231 cells (Fig. 1d). Unexpectedly, we found that metformin treatment also increased the mRNA and protein levels of *TTP* in MDA-MB-231 cells in a dose-(Fig. 1e) and time-(Fig. 1f) dependent manner., as well as promoter activity (Fig. 1g). Together, these data suggest that metformin can induce *TTP* expression in breast cancer cells in a *p53*-independent manner.


### TTP mediates metformin's anti-proliferative function

Previously metformin has been reported to exert anti-tumorigenic effect [2–4]. To confirm whether metformin shows anti-proliferative effect on MCF7 and MDA-MB-231 cells, we incubated these cells in the presence of different concentration of metformin for 24 and 48 h and

analyzed the cell proliferation using MTS assay. Consistent with previous reports [2–4], metformin treatment significantly inhibited the growth of both MCF7 and MDA-MB-231 cells in a dose-dependent (Fig. 2a, b) and time-dependent manner (Fig. 2c, d). Since incubation of both MCF7 and MDA-MB-231 cells with 6 mM metformin for 24 h induced *TTP* expression (Fig. 1a, e, f) and showed anti-proliferative effect (Fig. 2a–d), we conducted further experiment under conditions of incubating cells with 6 mM metformin for 24 h. Ectopic expression of *TTP* has been reported to inhibit cancer cell growth [26–28], and we confirmed that overexpression of *TTP* (Fig. 2e) inhibited cell proliferation in both MCF7 (Fig. 2g) and MDA-MB-231 cells (Fig. 2h). We next tested whether *TTP* is required for the anti-proliferative activity of metformin. To this end, we inhibited the expression of *TTP* using siRNA (Fig. 2f) and examined the effects of *TTP* knock-down on the anti-proliferative activity of metformin in MCF7 and MDA-MB-231 cells. Importantly, treatment of cells with siRNA against *TTP* (*TTP*-siRNA) but not scRNA attenuated the inhibitory effects of metformin on the growth of both MCF7 (Fig. 2g) and MDA-MB-231 cells (Fig. 2h) These results strongly suggested that *TTP* mediates the anti-proliferative functions of metformin in breast cancer cells.

### Metformin induces TTP expression in a c-Myc-dependent manner

We previously reported that *TTP* expression is induced by STAT3 in LPS-stimulated macrophages [34]. However, it is unlikely that STAT3 mediates the induction of *TTP* expression by metformin in breast cancer cells, since metformin decreased STAT3 phosphorylation (Fig. 3a). It has been reported that metformin decreases c-Myc levels in an AMPK-dependent manner [18] and also that c-Myc suppresses *TTP* expression [30]. Consistently, we found that metformin treatment increased phosphorylation of AMPK and decreased c-Myc levels in a time-dependent manner in both MCF7 (Fig. 3a, left) and MDA-MB-231 cells (Fig. 3a, right). In addition, siRNA-mediated inhibition of *c-Myc* increased *TTP* promoter activity (Fig. 3b, c) in the absence of metformin in both MCF7 and MDA-MB-231 cells (Fig. 3b, c). These results suggest that down-regulation of c-Myc level by metformin induces *TTP* expression. However, it is not likely that, in MDA-MB-231 cells, down-regulation of c-Myc is the only mechanism for the metformin-induced *TTP* expression, since metformin further increased the *TTP* promoter activity in *c-Myc*-depleted MDA-MB-231 cells (Fig. 3c). These results indicate that while metformin induces *TTP* expression in c-Myc-dependent manner in MCF7 cells, it induces *TTP* expression through both c-Myc-dependent and c-Myc-independent manner in MDA-MB-231 cells.




**Fig. 1** Metformin induces expression of *TTP* in both *p53* wild-type and *p53* mutant breast cancer cells. **a, b** Metformin increases *TTP* levels in *p53* wild-type human breast cancer MCF7 cells. MCF7 cells were treated **a** with the indicated concentrations of metformin for 24 h or **b** with 6 mM metformin for the indicated length of time. The levels of *TTP* were measured by semi-qRT-PCR (**a, b, top**) and Western blotting (**a, b, bottom**). **c** Metformin induces *TTP* promoter activity in *p53* wild-type MCF7 cells. MCF7 cells were transfected with pGL3/TTPp-1343 containing the *TTP* promoter. After treatment with 6 mM metformin for 24 h, luciferase activity was measured. The expression levels obtained from pGL3-transfected cells without metformin treatment were set to 1. Data are presented as the mean  $\pm$  SD ( $n = 3$ ).  $***p < 0.001$ . **d** Metformin induces p53 phosphorylation in *p53* wild-type MCF7 cells but not *p53* mutant MDA-MB-231 cells. MCF7 and MDA-MB-231 cells were treated


with 6 mM metformin for the indicated length of time. The levels of p53 and phospho-p53 (p-p53) were measured by Western blotting. **e, f** Metformin increases *TTP* levels in *p53* mutant human breast cancer MDA-MB-231 cells. MDA-MB-231 cells were treated **e** with the indicated concentrations of metformin for 24 h or **f** with 6 mM metformin for the indicated length of time. The levels of *TTP* were measured by semi-qRT-PCR (**e, f, top**) and Western blotting (**e, f, bottom**). **g** Metformin induces *TTP* promoter activity in MDA-MB-231 cells. MDA-MB-231 cells were transfected with pGL3/TTPp-1343 containing the *TTP* promoter. After treatment with 6 mM metformin for 24 h, luciferase activity was measured. The expression levels obtained from pGL3-transfected cells without metformin treatment were set to 1. Data are presented as the mean  $\pm$  SD ( $n = 3$ ).  $***p < 0.001$

We next tested whether metformin induces *TTP* expression in an AMPK-dependent manner. We treated MCF7 and MDA-MB-231 cells with 6 mM metformin in the presence or absence of AMPK inhibitor compound C for 24 h and analyzed the effect of compound C on the metformin-induced *TTP* expression. The compound C inhibited AMPK phosphorylation and blocked metformin-mediated induction of *TTP* in both MCF7 and MDA-MB-231 cells (Fig. 3d), indicating that AMPK phosphorylation is required for metformin-induced *TTP* expression. We next tested whether overexpression of *c-Myc* could block

the effect of metformin on the induction of *TTP* expression in cancer cells. Specifically, we transfected MCF7 and MDA-MB-231 cells with *c-Myc* (Fig. 3e) and analyzed the extent of *TTP* induction by real-time PCR (Fig. 3f) and cell proliferation by MTS assay (Fig. 3g, h) after treatment of the cells with metformin. Cells transfected with pcDNA3 empty vector were used as controls. Metformin treatment increased *TTP* expression (Fig. 3f) and inhibited cell proliferation in both pcDNA3-transfected MCF7 (Fig. 3g) and MDA-MB-231 cells (Fig. 3h). However, in cells over-expressing *c-Myc*, metformin failed to induce *TTP*

**Fig. 2** *TTP* mediates the anti-proliferative function of metformin in both *p53* wild-type and *p53* mutant breast cancer cells. **a–d** Metformin inhibits proliferation of both *p53* wild-type MCF7 cells and *p53* mutant MDA-MB-231 cells. **a, c** MCF7 and **b, d** MDA-MB-231 cells were treated (**a, b**) with the indicated concentrations of metformin for 24 h and (**c, d**) with 6 mM metformin for 24 and 48 h. Cell viability was assessed by measuring absorbance at 490 nm using an MTS cell proliferation assay. The values obtained with mock-treated cells were set to 100. Values are the mean  $\pm$  SD ( $n = 3$ ). \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ . **e–h** Inhibition of *TTP* attenuates the anti-proliferative effects of metformin in both MCF7 and MDA-MB-231 cells. MCF7 and MDA-MB-231 cells were transfected with **e, g, h** pcDNA6/*TTP* or **f, g, h** *TTP*-specific siRNA (*TTP*-siRNA). scRNA and pcDNA6 were used as negative controls. After treatment with 6 mM metformin for 24 h, cell viability was assessed by measuring the absorbance at 490 nm using an MTS cell proliferation assay. The values obtained with mock-treated cells were set to 100. Values are the mean  $\pm$  SD ( $n = 3$ ). \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ . *ns* not significant





**Fig. 3** Metformin induces *TTP* expression through down-regulation of *c-Myc*. **a** Metformin treatment increases phospho-AMPK (pAMPK) but decreases *c-Myc* and phospho-STAT3 (pSTAT3) in MCF7 and MDA-MB-231 cells. MCF7 and MDA-MB-231 cells were treated with 6 mM metformin for the indicated length of time, and the levels of *TTP*, *STAT3*, p*STAT3*, *AMPK*, p*AMPK*, and *c-Myc* were measured by Western blotting. **b, c** Inhibition of *c-Myc* by siRNA enhances *TTP* promoter activity in MCF7 and MDA-MB-231 cells. **b** MCF7 and **c** MDA-MB-231 cells were transfected with pGL3/*TTP*<sub>p-1343</sub> containing the *TTP* promoter. After treatment with 6 mM metformin for 24 h, luciferase activity was measured. The expression levels obtained from pGL3-transfected cells without metformin treatment were set to 1. Data are presented as the mean  $\pm$  SD ( $n = 3$ ). \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ . **d** Metformin increases *TTP* expression in an AMPK-dependent manner. MCF7 and MDA-MB-231 cells were treated with 6 mM metformin with or

without 20  $\mu$ M Compound C for 24 h. The levels of *TTP*, *AMPK*, and p*AMPK* were measured by Western blotting. **e, f** Overexpression of *c-Myc* blocks the effect of metformin on *TTP* induction. MDA-MB-231 cells were transfected with pcDNA3/*c-Myc* or control pcDNA3. Cells were treated with 6 mM metformin for 24 h. **e** *c-Myc* and **f** *TTP* levels were measured by quantitative RT-PCR. The values obtained with pcDNA-transfected and mock-treated cells were set to 1. Data are presented as the mean  $\pm$  SD ( $n = 3$ ). \*\*\* $p < 0.001$ . ns not significant. **g, h** Overexpression of *c-Myc* attenuates the anti-proliferative effect of metformin in MCF7 and MDA-MB-231 cells. **g** MCF7 and **h** MDA-MB-231 cells were transfected with pcDNA3/*c-Myc* or control pcDNA3. Cells were treated with 6 mM metformin for 24 h. Cell viability was assessed by measuring the absorbance at 490 nm using an MTS cell proliferation assay. The values obtained with mock-treated cells were set to 100. Data are presented as the mean  $\pm$  SD ( $n = 3$ ). \*\* $p < 0.01$ , \*\*\* $p < 0.001$ . ns not significant

expression (Fig. 3f) and did not show anti-proliferative effect in both MCF7 (Fig. 3g) and MDA-MB-231 cells (Fig. 3h). Taken together, these data suggested that

metformin induces *TTP* expression by down-regulating *c-Myc* and also that *TTP* mediates metformin's anti-proliferative function in breast cancer cells.



## Discussion

Metformin, a first-line drug for type 2 diabetes, has recently received increased attention because of its anti-proliferative effects in cancer cells [2–4]. However, the mechanisms underlying the anti-proliferative effects of metformin remain unclear. Here, we describe a role for *TTP* in mediating metformin's anti-proliferative effect in cancer cells. Specifically, we found that metformin increased expression level of *TTP* in cancer cells in an AMPK-dependent manner and also that down-regulation of *TTP* by siRNA attenuated the anti-proliferative effect of metformin. Thus, our data indicate that *TTP* induction is required for the anti-proliferative activity of metformin in cancer cells.

*TTP* can inhibit the growth of cancer cells by down-regulating expression of oncogenes [26, 35, 36]. However, a significant decrease in the expression of *TTP* has been observed in many cancer cells [25, 26, 33]. Thus, we hypothesized that induction of *TTP* in cancer cells may lead to an inhibition of growth. In this study, we found that *TTP* expression was induced by metformin in breast cancer cells. Metformin enhanced *TTP* promoter activity, indicating that this regulation occurs at the level of transcription.

We previously reported that activation of p53 increases *TTP* transcription in cancer cells [29]. Furthermore, Metformin can activate p53 in an AMPK-dependent manner [8, 37]. Thus, we considered the possibility that metformin may induce the expression of *TTP* through p53. Consistently, we found that metformin increased *TTP* expression level in an AMPK-dependent manner. However, considering the abnormalities in the p53 pathway in nearly all types of cancers [31], if metformin induces *TTP* expression through p53 pathway, *TTP* induction by metformin would be limited to only a small portion of cancer cells containing with an intact p53 signaling pathway. Indeed, it is unlikely that p53 is essential for *TTP* induction by metformin, since metformin induced the expression of *TTP* in p53 mutant cells as well as p53 wild-type cancer cells as indicated by MDA-MB-231 and MCF7 cells in this study, respectively. However, we found that, in p53 mutant MDA-MB-231 cells, *TTP* protein increased at high concentration of metformin and to less extent compared with p53 wild-type MCF7 cells. These results suggest the possibility that p53 may be involved in the induction of *TTP* expression by metformin.

What would be the p53-independent mechanism of *TTP* induction by metformin in cancer cells? It has been reported that c-Myc acts as a negative regulator of *TTP* expression [30]. The induction of *TTP* by metformin is likely the result of inhibition of c-Myc. In this study, we

obtained strong evidence in support of this hypothesis: inhibition of c-Myc increased *TTP* expression level; metformin decreased the expression of c-Myc; and ectopic expression of c-Myc abrogated the effects of metformin with respect to induction of *TTP*. Metformin has been reported to decrease c-Myc expression via the AMPK pathway [18]. If metformin induces *TTP* expression through inhibition of c-Myc, *TTP* induction by metformin would depend on AMPK activity. Consistently, we found that metformin induced *TTP* expression in an AMPK-dependent manner. However, the transcription factors acting as positive regulators for *TTP* induction by metformin remain elusive. Thus, further investigation into the specific transcription factors required for *TTP* induction will reveal the mechanisms underlying how metformin induces *TTP* expression.

In conclusion, we identified *TTP* as a down-stream target of metformin and a mediator of the anti-proliferative effects of metformin in cancer cells. Specifically, metformin induced *TTP* expression by down-regulating c-Myc, a negative regulator of *TTP* expression [30]. Importantly, our study provides a molecular basis for the anti-proliferative effects of metformin in cancer cells. Our finding that *TTP* is induced by metformin in both p53 wild-type and p53 mutant cancer cells and mediates the anti-proliferative effect of metformin further highlights the important role of *TTP* in human cancer cells. Metformin showed only modest anti-proliferative effect on cancer cells used in this study. However, since the mechanism of action of metformin is unique compared with that of other chemotherapeutic agents, metformin in combination with other chemotherapeutic agents may trigger significant tumor growth inhibition in vivo. Given the emerging evidence supporting the anti-proliferative effects of metformin in various types of cancer cells [2–4], it will be of interest to explore whether the regulatory mechanisms described here are relevant to other types of cancers.

**Acknowledgments** This work was supported by Priority Research Centers Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2014R1A6A1030318) and NRF-2014R1A1A2007525.

### Compliance with ethical standard

**Conflict of interest** The authors declare that they have no conflict of interest.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Bailey CJ, Turner RC (1996) Metformin. *N Engl J Med* 334:574–579
- Kourelis TV, Siegel RD (2012) Metformin and cancer: new applications for an old drug. *Med Oncol* 29:1314–1327
- Jiralerspong S, Palla SL, Giordano SH et al (2009) Metformin and pathologic complete responses to neoadjuvant chemotherapy in diabetic patients with breast cancer. *J Clin Oncol* 27:3297–3302
- Quinn BJ, Kitagawa H, Memmott RM, Gills JJ, Dennis PA (2013) Repositioning metformin for cancer prevention and treatment. *Trends Endocrinol Metab* 24:469–480
- Dowling RJ, Goodwin PJ, Stambolic V (2011) Understanding the benefit of metformin use in cancer treatment. *BMC Med* 9:33
- Javeshghani S, Zakikhani M, Austin S et al (2012) Carbon source and myc expression influence the antiproliferative actions of metformin. *Cancer Res* 72:6257–6267
- Russo GL, Russo M, Ungaro P (2013) AMP-activated protein kinase: a target for old drugs against diabetes and cancer. *Biochem Pharmacol* 86:339–350
- Vazquez-Martin A, Oliveras-Ferreros C, Del Barco S, Martin-Castillo B, Menendez JA (2009) If mammalian target of metformin indirectly is mammalian target of rapamycin, then the insulin-like growth factor-1 receptor axis will audit the efficacy of metformin in cancer clinical trials. *J Clin Oncol* 27:e207–209
- Vincent EE, Coelho PP, Blagih J, Griss T, Viollet B, Jones RG (2015) Differential effects of AMPK agonists on cell growth and metabolism. *Oncogene* 34:3627–3639
- Cufi S, Corominas-Faja B, Vazquez-Martin A et al (2012) Metformin-induced preferential killing of breast cancer initiating CD44+ CD24–/low cells is sufficient to overcome primary resistance to trastuzumab in HER2+ human breast cancer xenografts. *Oncotarget* 3:395–398
- Hirsch HA, Iliopoulos D, Tsiachlis PN, Struhl K (2009) Metformin selectively targets cancer stem cells, and acts together with chemotherapy to block tumor growth and prolong remission. *Cancer Res* 69:7507–7511
- Wurth R, Pattarozzi A, Gatti M et al (2013) Metformin selectively affects human glioblastoma tumor-initiating cell viability: a role for metformin-induced inhibition of Akt. *Cell Cycle* 12:145–156
- Sato A, Sunayama J, Okada M et al (2012) Glioma-initiating cell elimination by metformin activation of FOXO3 via AMPK. *Stem Cells Transl Med* 1:811–824
- Memmott RM, Mercado JR, Maier CR, Kawabata S, Fox SD, Dennis PA (2010) Metformin prevents tobacco carcinogen-induced lung tumorigenesis. *Cancer Prev Res (Phila)* 3:1066–1076
- Liu X, Chhipa RR, Pooya S et al (2014) Discrete mechanisms of mTOR and cell cycle regulation by AMPK agonists independent of AMPK. *Proc Natl Acad Sci USA* 111:E435–444
- Liu B, Fan Z, Edgerton SM, Yang X, Lind SE, Thor AD (2011) Potent anti-proliferative effects of metformin on trastuzumab-resistant breast cancer cells via inhibition of erbB2/IGF-1 receptor interactions. *Cell Cycle* 10:2959–2966
- Martin-Castillo B, Vazquez-Martin A, Oliveras-Ferreros C, Menendez JA (2010) Metformin and cancer: doses, mechanisms and the dandelion and hornetic phenomena. *Cell Cycle* 9:1057–1064
- Akinyeke T, Matsumura S, Wang X et al (2013) Metformin targets c-MYC oncogene to prevent prostate cancer. *Carcinogenesis* 34:2823–2832
- Deng XS, Wang S, Deng A et al (2012) Metformin targets Stat3 to inhibit cell growth and induce apoptosis in triple-negative breast cancers. *Cell Cycle* 11:367–376
- Cufi S, Corominas-Faja B, Lopez-Bonet E et al (2013) Dietary restriction-resistant human tumors harboring the PIK3CA-activating mutation H1047R are sensitive to metformin. *Oncotarget* 4:1484–1495
- Shaw G, Kamen R (1986) A conserved AU sequence from the 3′ untranslated region of GM-CSF mRNA mediates selective mRNA degradation. *Cell* 46:659–667
- Shyu AB, Wilkinson MF (2000) The double lives of shuttling mRNA binding proteins. *Cell* 102:135–138
- Carballo E, Lai WS, Blackshear PJ (1998) Feedback inhibition of macrophage tumor necrosis factor- $\alpha$  production by tristetraprolin. *Science* 281:1001–1005
- Brooks SA, Blackshear PJ (2013) Tristetraprolin (TTP): interactions with mRNA and proteins, and current thoughts on mechanisms of action. *Biochim Biophys Acta* 1829:666–679
- Brennan SE, Kuwano Y, Alkharouf N, Blackshear PJ, Gorospe M, Wilson GM (2009) The mRNA-destabilizing protein tristetraprolin is suppressed in many cancers, altering tumorigenic phenotypes and patient prognosis. *Cancer Res* 69:5168–5176
- Lee HH, Son YJ, Lee WH et al (2010) Tristetraprolin regulates expression of VEGF and tumorigenesis in human colon cancer. *Int J Cancer* 126:1817–1827
- Marderosian M, Sharma A, Funk AP et al (2006) Tristetraprolin regulates Cyclin D1 and c-Myc mRNA stability in response to rapamycin in an Akt-dependent manner via p38 MAPK signaling. *Oncogene* 25:6277–6290
- Young LE, Sanduja S, Bemis-Standoli K, Pena EA, Price RL, Dixon DA (2009) The mRNA binding proteins HuR and tristetraprolin regulate cyclooxygenase 2 expression during colon carcinogenesis. *Gastroenterology* 136:1669–1679
- Lee JY, Kim HJ, Yoon NA et al (2013) Tumor suppressor p53 plays a key role in induction of both tristetraprolin and let-7 in human cancer cells. *Nucleic Acids Res* 41:5614–5625
- Rounbehler RJ, Fallahi M, Yang C et al (2012) Tristetraprolin impairs myc-induced lymphoma and abolishes the malignant state. *Cell* 150:563–574
- Soussi T, Beroud C (2001) Assessing TP53 status in human tumours to evaluate clinical outcome. *Nat Rev Cancer* 1:233–240
- Cole MD (1986) The myc oncogene: its role in transformation and differentiation. *Annu Rev Genet* 20:361–384
- Kim HK, Kim CW, Vo MT et al (2012) Expression of proviral integration site for Moloney murine leukemia virus 1 (Pim-1) is post-transcriptionally regulated by tristetraprolin in cancer cells. *J Biol Chem* 287:28770–28778
- Joe Y, Kim HJ, Kim S et al (2011) Tristetraprolin mediates anti-inflammatory effects of nicotine in lipopolysaccharide-stimulated macrophages. *J Biol Chem* 286:24735–24742
- Lykke-Andersen J, Wagner E (2005) Recruitment and activation of mRNA decay enzymes by two ARE-mediated decay activation domains in the proteins TTP and BRF-1. *Genes Dev* 19:351–361
- Kim CW, Vo MT, Kim HK et al (2012) Ectopic over-expression of tristetraprolin in human cancer cells promotes biogenesis of let-7 by down-regulation of Lin28. *Nucleic Acids Res* 40:3856–3869
- He G, Zhang YW, Lee JH et al (2014) AMP-activated protein kinase induces p53 by phosphorylating MDMX and inhibiting its activity. *Mol Cell Biol* 34:148–157