## **Corrections**

## Forces Applied by Cilia Measured on Explants from Mucociliary Tissue

Zvi Teff, Zvi Priel, and Levi A. Ghebery\*

(Biophysical Journal 92, 1813–1823; March 1, 2007)

In Eq. 3 of this article,  $L_{\text{eff}}$  should have read (L-P). The calculations of results were based on the correct equation, and thus the published results remain valid. The correct equation appears below.

$$\tan \alpha = \frac{(L-P)\tan \beta + D'}{\tan \beta (D+D')}$$

\*Correspondence: glevi@bgu.ac.il

http://dx.doi.org/10.1016/j.bpj.2016.02.004

## Label-free Imaging of Microtubules with Sub-nm Precision Using Interferometric Scattering Microscopy

Joanna Andrecka, Jaime Ortega Arroyo, Katie Lewis, Robert A. Cross, and Philipp Kukura\*

(Biophysical Journal 110, 214–217; January 5, 2016)

In Fig. 2 b, the scale bar at the bottom of the figure should have read 4 nm instead of 8 nm. The corrected figure is shown here and in the online article.

FIGURE 2 Analysis of MT motion driven by surface-bound kinesin. (a) Representative time traces showing 8 nm steps for MTs bound to single kinesins, including a representative xy trajectory acquired at 1000 frames/s. (b) Tracking results for higher motor densities exhibiting smaller but distinct fractional steps, including representative xy trajectories acquired at 100 frames/s. To see this figure in color, go online. (corrected)







FIGURE 2 Analysis of MT motion driven by surface-bound kinesin. (a) Representative time traces showing 8 nm steps for MTs bound to single kinesins, including a representative xy trajectory acquired at 1000 frames/s. (b) Tracking results for higher motor densities exhibiting smaller but distinct fractional steps, including representative xy trajectories acquired at 100 frames/s. To see this figure in color, go online. (original)



<sup>\*</sup>Correspondence: philipp.kukura@chem.ox.ac.uk http://dx.doi.org/10.1016/j.bpj.2016.02.005