

SCIENTIFIC REPORTS



OPEN

Rationalising the role of Keratin 9 as a biomarker for Alzheimer's disease

Received: 21 September 2015

Accepted: 19 February 2016

Published: 14 March 2016

Joanna L. Richens¹, Hannah L. Spencer¹, Molly Butler¹, Fiona Cantlay¹, Kelly-Ann Vere¹, Nin Bajaj², Kevin Morgan³ & Paul O'Shea^{1,*}

Keratin 9 was recently identified as an important component of a biomarker panel which demonstrated a high diagnostic accuracy (87%) for Alzheimer's disease (AD). Understanding how a protein which is predominantly expressed in palmoplantar epidermis is implicated in AD may shed new light on the mechanisms underlying the disease. Here we use immunoassays to examine blood plasma expression patterns of Keratin 9 and its relationship to other AD-associated proteins. We correlate this with the use of an *in silico* analysis tool VisANT to elucidate possible pathways through which the involvement of Keratin 9 may take place. We identify possible links with Dickkopf-1, a negative regulator of the wnt pathway, and propose that the abnormal expression of Keratin 9 in AD blood and cerebrospinal fluid may be a result of blood brain barrier dysregulation and disruption of the ubiquitin proteasome system. Our findings suggest that dysregulated Keratin 9 expression is a consequence of AD pathology but, as it interacts with a broad range of proteins, it may have other, as yet uncharacterized, downstream effects which could contribute to AD onset and progression.

Keratin 9 is a 623 amino acid type 1 cytokeratin expressed predominantly in the skin¹. Mutations within the KRT9 gene (Keratin 9) have been strongly associated with the skin disorder Epidermolytic palmoplantar keratoderma^{2–4}. More recently however, evidence has emerged demonstrating alternative expression sites of Keratin 9 including natural killer cell membranes⁵, intestinal extracellular exosomes⁶, sperm nuclei⁷, tears⁸, follicular fluid⁹ and blood serum¹⁰ with several of these studies implicating the molecule in disease pathology. Dysregulated expression of Keratin 9 has, for example, been implicated in polycystic ovary syndrome⁹ and suggested as a metastatic marker of hepatocellular carcinoma¹⁰.

Of particular interest is recent evidence suggesting that Keratin 9 may be implicated in neurological disorders including dementia. Cerebrospinal fluid (CSF), a fluid produced by brain ventricles and which surrounds the central nervous systems, has been a target for studies into neurological disorders as it is contiguous with brain interstitial fluid and is, therefore, indicative of brain pathology¹¹. A review of the CSF proteome found evidence of Keratin 9 expression in CSF using both ESI Ion Trap/FT-ICR (18 peptide sequences) and MALDI TOF/TOF (23 peptide sequences)¹² mass spectrometry. These findings were further substantiated by a 2012 study in which 2D electrophoresis examination of CSF demonstrated that levels of Keratin 9 expression in both Multiple Sclerosis and Neuromyelitis Optica patients were twice that of control individuals¹³.

Alzheimer's disease (AD), the most common form of dementia¹⁴, is posing a huge societal problem and indicators of disease that can be utilised in identification and monitoring of disease progression are vital¹⁴. A proteomic study of CSF undertaken by Vafadar-Isfahani *et al.* identified Keratin 9 as an important component of a biomarker panel for diagnosing AD¹⁵. The power of these markers resided in their combined use as a complete biomarker panel, but the existence of any marker within the panel is suggestive of some involvement in the mechanisms underlying AD pathology. A subsequent study undertaken within our laboratory utilised immunoassays to examine Keratin 9 expression in 'matched' CSF and blood plasma samples collected simultaneously from individual donors¹⁶. This study provided further validation of the presence of Keratin 9 in both CSF and

¹Cell Biophysics Group, School of Life Sciences, University of Nottingham, University Park, Nottingham, United Kingdom. ²Department of Neurology, Nottingham University Hospitals NHS Trust, Queen's Medical Centre, Nottingham, United Kingdom. ³School of Life Sciences, University of Nottingham, Queen's Medical Centre, Nottingham, United Kingdom. *Present address: Faculty of Pharmaceutical Sciences, The University of British Columbia, Vancouver Campus, 2405 Wesbrook Mall, Vancouver, BC Canada. Correspondence and requests for materials should be addressed to P.O. (email: paul.oshea@nottingham.ac.uk or paul.oshea@ubc.ca)

	Healthy Cohort	AD Cohort
Sample number	60	58
Gender (M/F)	25/35	31/27
Age at entry (years \pm SD)	80.33 \pm 6.64	79.03 \pm 6.96
Time between study entry and recalled onset (years \pm SD)	—	4.32 \pm 2.71

Table 1. Patient demographics of the blood plasma samples used within this study.

blood plasma. Interestingly, Keratin 9 was detected exclusively in CSF collected from patients with AD but not in healthy individuals. Whilst the majority of the other components of the biomarker panel¹⁵ had previously been linked to AD^{17–21}, it was thought that these studies were the first to implicate Keratin 9 (and also Tetranectin) in AD pathology. It has since emerged, however, that a previous study by Mueller and colleagues in 2010 used mass spectrometry to demonstrate dysregulation of Keratin 9 expression levels in AD²² substantiating our previous findings^{15,16}. In addition to this, a recent study undertaken by Li *et al.* identified Keratin 9 amongst a number of potential molecules and pathways involved in AD pathology, although its involvement within these mechanisms was not expanded upon²³.

Whilst the understandable focus of many biomarker studies, including those mentioned above, is to identify molecules with high diagnostic utility they do also offer the often overlooked opportunity to obtain a further understanding of the mechanistic origins of any disease. As evidence lends strength to the potential importance of Keratin 9 in AD, it also raises the question of how a protein typically associated with the skin can be intimately involved in a neurodegenerative disorder. The question of what has led to dysregulation of its normal expression pattern therefore requires attention and particularly whether it is a cause or consequence of the disease process.

In the present study we aim to shed light on the potential involvement of Keratin 9 in AD. Expression patterns of Keratin 9 in blood plasma samples from AD and healthy patient cohorts are examined to determine its diagnostic utility as a stand-alone biomarker and its relationship with other AD-associated protein targets. This is followed by a detailed *in silico* investigation of potential mechanisms and pathways through which Keratin 9 could influence disease pathology. These *in silico* studies were undertaken using the VisANT pathways analysis package developed at Boston University and based upon the Predictome database (<http://visant.bu.edu>)²⁴. This software enables examination of the biological interactions of an experimentally determined biomarker, facilitating construction of molecular pathways and networks that are associated with the disease condition. In a previous study we applied VisANT analysis protocols to AD-related gene and protein panels in order to explore and redefine the mechanistic basis of the disease²⁵. Other studies that utilised VisANT include investigations into the mammalian 14-3-3-phosphoproteome²⁶, the involvement of the Stat5a network in prostate cancer²⁷ and the role of FoxP1 in the regulation of autism-related pathways²⁸. Identification of molecular pathways of which Keratin 9 is a potential component allows discussion on how Keratin 9 expression could influence or be influenced by AD pathology. Understanding the mechanisms which are associated with these pathways therefore could lead to the identification of novel drug targets or additional diagnostic biomarkers.

Methods

Patient samples. This study was approved by the local Ethics Committee, NRES Committee East Midlands, Ref 12/EM/0052. All methods were carried out in accordance with the approved guidelines and all participants provided written informed consent. Case samples were diagnosed as either confirmed or probable AD according to the Consortium to Establish a Registry for Alzheimer's disease (CERAD) guidelines. All patients were diagnosed as Late-Onset Alzheimer's disease (LOAD). Patients with significant cognitive comorbidity, including but not limited to head trauma, alcoholism, learning disability or Parkinson's disease, were excluded from the study. The demographics of the sample cohorts analysed within this study are identified in Table 1.

Protein detection by Immunoassay: ELISAs. Keratin 9 was detected according to the manufacturer's protocol using an ELISA kit purchased from antibodies-online GmbH (ABIN417500). Plasma and CSF samples (diluted 1:1) were incubated in wells pre-coated with a capture antibody for 2 hr followed by incubation with a biotinylated detector antibody for 1 hr and a 30 min incubation with horseradish peroxidase-conjugated avidin. The reaction was developed using TMB substrate solution, stopped with 1 M HCl and read at 450 nm. All assay steps were undertaken at 37 °C. All samples and standard curves were measured in duplicate and concentrations of Keratin 9 in the clinical samples were determined from the standard curve generated using the standards provided in the kit.

Amyloid beta peptide 42 (A β 42) was detected using a commercially available ELISA kit processed according to manufacturer's instructions (Wako Chemicals GmbH; 296–64401). All samples and standard curves were measured in duplicate and concentrations of A β 42 in the clinical samples were determined from the standard curve generated using the reagents provided in the kit.

SPARC-like 1 (SPARCL1) was measured by ELISA as previously described¹⁶. Briefly, ELISA plates were coated overnight with 5 μ g/ml capture antibody (CAB-701MH in PBS; Creative Biomart). Following this and all subsequent incubations wells were washed three times with PBS-0.05% Tween20 (PBST). A blocking solution of PBST-1% BSA was applied to antibody-coated wells for 1 hr prior to addition of 50 μ l sample (either clinical sample or protein standard) for 2 hr. Biotinylated detection antibody (BAF2728; R&D Systems) was added at 500 ng/ml for 2 hr and the reaction was developed using streptavidin-HRP and TMB substrate, stopped with 1 M HCl. Samples




Figure 1. Keratin 9 expression in blood plasma in healthy and AD patient cohorts (A) and the correlation of these concentrations with the age of healthy (B(i)) and AD (B(ii)) individuals.

were read at 450 nm with concentrations of SPARCL1 determined from a standard curve generated with recombinant SPARCL1 (R&D Systems; 2728-SL).

Protein detection by Immunoassay: Luminex assays. Apolipoprotein E (ApoE), Clusterin, Fibrinogen and Tau concentrations were determined using Luminex assays developed in-house as previously described¹⁶. Briefly, monoclonal capture antibodies to the target proteins were coupled to COOH-coated fluorescently dyed microspheres (Bio-Rad, Hercules, CA) using the Bio-Plex Amine Coupling Kit (Bio-Rad, 171-406001) according to manufacturer's instructions and as previously validated¹⁶. A bead suspension containing 5000 of each antibody-conjugated bead set was added to designated wells of a filter plate (pre-wetted with wash buffer (PBST)), washed twice with PBST and resuspended in incubation buffer (PBS-1% BSA). To this, a sample of appropriately diluted clinical sample or protein standard mixture was added, and the plate was incubated for 2 hr at 25 °C on a rotating plate shaker (600 rpm). Wells were washed, incubated with a cocktail of biotinylated detector antibodies (containing each antibody at a pre-determined optimal concentration) for 1 hr, washed again and incubated for 30 mins with streptavidin-RPE. Wells were then washed to remove unbound streptavidin-RPE prior to analysis. Data was acquired on a Bio-Plex 200 system and analysed with associated software (Bio-Rad). All samples and standard curves were performed in duplicate and in each well, a minimum of 100 beads per target molecule were analysed for both bead designation and R-phycoerythrin fluorescence. Concentrations of the target proteins within the clinical samples were determined from standard curves generated using recombinant proteins.

In silico analysis. *In silico* protein association studies were undertaken using the free licence software VisANT (<http://visant.bu.edu>)²⁴. This network analysis tool is constructed around the Predictome database²⁹ which contains details of experimentally-derived biological interactions collated from a variety of data sources²⁴. An initial search of Keratin 9 interactions was undertaken to identify binding partners of the molecule. The potential contribution of Keratin 9 to AD pathology was then examined through investigation of the biological interactions of Keratin 9 with the well-established AD-associated molecules ApoE, amyloid precursor protein (APP) and Tau^{30–32}. Searches were undertaken using the primary UniProt accession number of the appropriate proteins; P35527 (Keratin 9), P05067 (APP), P02649 (ApoE) and P10636 (Tau). An interactome was defined as the network of interactions that occurs between the target molecules under consideration. Interactions were restricted to those which occur no more than one degree away from the inputted molecules. The components identified within the derived interactome were further analysed using the free, open access pathway database Reactome (www.reactome.org)³³ to identify potential pathways through which they may contribute to AD pathology.

Statistics. Data were analysed using GraphPad Prism Version 6.04. Cohorts were compared using the Mann-Whitney test and Spearman correlation coefficients were determined as appropriate. A p-value ≤ 0.05 was considered to be a practical level of clinical significance. Significance values are denoted by asterisks as defined by GraphPad Prism version 6.04: ****extremely significant ($p < 0.0001$); ***extremely significant ($0.001 \leq p \leq 0.0001$); **very significant ($0.01 \leq p \leq 0.001$); *significant ($0.05 \leq p \leq 0.01$).

Results & Discussion

Keratin 9 Expression and Correlations with Other AD-associated Proteins. Keratin 9 expression levels were examined in blood plasma samples from a cohort comprising 60 healthy individuals and 58 AD patients, as shown in Fig. 1A. The actual concentrations observed in this cohort were lower than those obtained for an alternative cohort in a previous study¹⁶, however, the trends observed were analogous, with expression levels decreasing in healthy individuals (Mean = 34.7 ± 47.6 pg/ml) when compared to AD patients (Mean = 64.4 ± 88.0 pg/ml). The variation in concentrations observed between the different studies may be due to the use of different anticoagulants during blood collection, a factor which is known to affect metabolite concentrations³⁴. Whilst still not reaching significance, the p-value of the change due to AD decreased for the larger cohort ($p = 0.1399$; $n = 58$ AD/60 Healthy) indicating that it may continue to improve as the sample size increases. In Fig. 1B, there appears to be some level of association between blood plasma Keratin 9 concentration and age in healthy individuals, ($r = -0.2316$; $p = 0.075$) which is dysregulated in the AD cohort ($r = 0.1646$; $p = 0.2168$).

Molecule	Healthy Cohort			AD Cohort		
	Concentration	Correlation with age		Concentration	Correlation with age	
		Correlation coefficient (r)	p-value		Correlation coefficient (r)	p-value
A β 42	6.49 \pm 1.25 pmol/L	0.0755	0.6432	6.32 \pm 1.74 pmol/L	0.3802	0.0185*
ApoE	4776 \pm 3238 ng/ml	-0.3833	0.0025**	4607 \pm 2689 ng/ml	0.1369	0.3056
Clusterin	2534 \pm 1397 ng/ml	-0.2536	0.0505	2516 \pm 1336 ng/ml	0.0824	0.5386
Fibrinogen	49.59 \pm 20.72 μ g/ml	0.0570	0.6652	41.57 \pm 21.72 μ g/ml	0.2234	0.0918
Keratin 9	34.7 \pm 47.6 pg/ml	-0.2316	0.0750	64.4 \pm 88.0 pg/ml	0.1646	0.2168
SPARCL1	1126 \pm 533 ng/ml	0.4619	0.0002***	1204 \pm 757 ng/ml	0.1711	0.1992
Tau	43.90 \pm 13.78 ng/ml	-0.4082	0.0012**	43.10 \pm 14.84 ng/ml	0.1062	0.4275

Table 2. Blood plasma concentrations of putative AD biomarker targets as measured by immunoassay. Spearman correlation coefficients for protein concentration and patient age were determined using GraphPad Prism Version 5.02. A p -value \leq 0.05 was considered statistically significant.

Molecule 1	Molecule 2	Healthy Cohort		AD Cohort	
		Spearman correlation coefficient (r)	p-value	Spearman correlation coefficient (r)	p-value
Keratin 9	A β 42	0.1045	p = 0.5209	-0.05246	p = 0.7478
Keratin 9	ApoE	0.3829	p = 0.0025**	0.6380	p < 0.0001****
Keratin 9	Clusterin	0.2707	p = 0.0364*	0.5276	p < 0.0001****
Keratin 9	Fibrinogen	0.1107	p = 0.3998	0.03512	p = 0.7899
Keratin 9	SPARCL1	-0.3026	p = 0.0188*	-0.3102	p = 0.0158*
Keratin 9	Tau	0.4338	p = 0.0005***	0.6256	p < 0.0001****

Table 3. Correlations between protein concentrations of Keratin 9 and other AD-associated molecules in blood plasma samples. Protein concentrations were determined using the appropriate ELISA of Luminex assay. Spearman correlation coefficients were determined for all protein pairings in healthy and AD patient cohorts using GraphPad Prism Version 5.02. A p -value \leq 0.05 was considered statistically significant.

In order to gain an understanding of the role of Keratin 9 in AD, interactions with other AD-associated proteins were sought. Immunoassays of A β 42, ApoE, Clusterin, Fibrinogen, SPARCL1 and Tau^{15,16,35} were undertaken on blood plasma samples from the two patient cohorts. The concentrations of these proteins are detailed in Table 2. It should be noted that, whilst not the focus of this study, a significant difference ($p = 0.0063$) between the concentration of fibrinogen in the healthy (49.59 \pm 20.72 μ g/ml) and AD (41.57 \pm 21.72 μ g/ml) patient samples was observed lending weight to previous studies which identify it as a potential biomarker of AD³⁶. Additionally, the correlations between age and concentration of ApoE, SPARCL1 and Tau were found to be disrupted upon onset of AD whilst the correlation between A β 42 and age strengthened in the AD cohort when compared to healthy individuals (Table 2).

When the relationship between concentrations of Keratin 9 and A β 42, ApoE, Clusterin, Fibrinogen, SPARCL1 and Tau were examined (Table 3), significant correlations were demonstrated between blood plasma concentrations of Keratin 9 and ApoE, Clusterin or Tau. The strength of all these correlations was found to increase in the AD cohort when compared to the healthy individuals. A correlation with SPARCL1 was observed, but the strength of this association remained unchanged between the two cohorts. No correlations were identified between blood plasma concentrations of Keratin 9 and A β 42 or Fibrinogen in either the healthy or AD cohorts (Table 3).

Before speculating on the meaning of these results, the possibility of contamination was also addressed. Due to the expression of Keratin 9 in the outer, terminally differentiated epidermis, it is present in shed skin, which is a major component of dust. However unlikely, it is conceivable, that results from some previous studies could have attributed dust contamination to the detection of Keratin 9³⁷. Furthermore, taking blood and CSF samples involves puncturing the skin, and as such may be a route through which keratinocyte components could enter the clinical samples. ELISA measurements of Keratin 9 in both AD patients and healthy controls collected within this study suggest that there is some level of expression in both cohorts (Fig. 1A). Whilst the possibility of contamination causing these baseline levels of expression cannot be disregarded, it cannot account for studies where differences in expression levels have been identified in identically handled samples. Work in our laboratory has previously identified the presence of Keratin 9 in AD CSF but not in normal CSF¹⁶ substantiating the significance of our findings and suggesting that there is a biological basis behind the results as opposed to contamination.

Initial *in silico* Analysis. The data outlined above suggests that Keratin 9 may be implicated in the mechanistic pathways underlying AD. To further ascertain the nature of association, *in silico* protein association studies were undertaken using VisANT software, as described in Methods. Initially, the known interacting partners of Keratin 9 were examined with the interactome produced by these molecules illustrated in Fig. 2. This identified 54 proteins that have been experimentally proven to interact with Keratin 9, which are listed in Table 4, along with




Figure 2. A basic Keratin 9 interactome determined using VisANT software. The interactome comprises the 54 proteins demonstrated to interact directly with Keratin 9 with the lines linking each node colour coded according to method used to establish the interaction. Further information on the nodes and interactions can be found in Table 4.

the experimental methods used to ascertain these interactions. Current knowledge of the function of Keratin 9 is limited; it has been shown to provide scaffolding to cells³ and act as an epidermal differentiation marker³⁸. This VisANT analysis implies it may have further complex roles as it interacts with proteins that possess a wide range of functions (Table 4). The names and details of these proteins are listed in Table 4. Additionally, the interactome produced substantiates recent findings that Keratin 9 may have alternate expression sites to its well-characterized expression in palms and soles. For example, Keratin 9 has been shown to interact with IQCB1³⁹, which is localized in cilia of renal epithelial cells and, interestingly photoreceptor cells⁴⁰. This adds weight to the recent discovery that Keratin 9 can be found in the proteome of tears⁸. One of the most important implications of this interactome is that it provides further validation of the presence of Keratin 9 in blood plasma^{10,16} as it can interact with five of the most abundant proteins in blood serum: Albumin, Apolipoprotein A-I and the heavy chain constant regions of Immunoglobins A, G and M⁴¹.

Further *in silico* Analysis: Expanding the Keratin 9 Interactome. Although this interactome provides a useful start point from which to understand the involvement of Keratin 9 in AD, it only details direct molecular interactions; expanding the interactome would highlight further biochemical pathways that may influence/be influenced by variations in Keratin 9 expression. With this in mind, we examined the additional interactions of all 54 identified Keratin 9 interacting proteins. However, simply expanding every node resulted in high levels of complexity in the outputted data. We proceeded, therefore, to examine the interactions between Keratin 9 and other molecules known to be involved in AD pathology directly rather than indirectly. VisANT was utilised to produce an interactome detailing the interactions of Keratin 9 with APP, Tau (MAPT) and ApoE, three molecules with a longstanding association with AD (together, referred to as the AMA interactome). Molecules found to have two or more interacting partners within the derived network (i.e. they interact with at least two out of ApoE, APP, Keratin 9 and Tau) are included in the resultant interactome (Fig. 3). Correlations between expression levels of Keratin 9 and those of ApoE and Tau were earlier found to be disrupted in blood plasma samples from AD patients (Table 3). Whilst Keratin 9 has not yet been demonstrated to interact directly with APP, ApoE or Tau, Fig. 3 demonstrates that it is closely connected to all three molecules, requiring only one bridging molecule. The molecules found to interact with Keratin 9 and the AMA interactome are summarised in Table 4. In the case of ApoE Keratin 9 is linked via either VCAM1, ALB or UBC whilst for Tau it is linked via UBC and YWHAQ and for APP, it is linked via ALB, APOA1, COPS5, CUL3, CUL4B, GABARAPL2, GRB2, IGHM, MAP1LC3A, MDM2, NEDD8, PIK3R2, SHC1, TANK, TRAF3IP1, UBC and UCHL5. These molecular associations fail, however, to shed much more light on the problem since it is unlikely that Keratin 9 levels would affect APOE, MAPT or APP and vice versa through such a short pathway. Furthermore, many of the linking molecules, such as UBC, are ubiquitously expressed and interact with a broad range of proteins so any effect manifesting through these molecules would not explain the specificity of AD symptoms. It is therefore necessary to expand the interactome network: a series of several, more specific interactions that link these proteins is potentially the key to solving the problem as opposed to searching for the shortest connection paths. One way to solve this problem, whilst avoiding generating too much complexity in the interactome, is to place Keratin 9 in the context of pathways that are already thought to play a role in AD.

In previous work we have used Reactome (<http://www.reactome.org/>)^{25,42} to identify pathways in which AD biomarkers may be participating²⁵. Initially, the same approach was implemented here whereby Keratin 9 and all 54 of its interacting molecules that were identified through VisANT were inputted into Reactome. However 10 out of these 55 molecules, including Keratin 9, were not found in the Reactome database, and as a consequence

KRT9 interacting partner	Experimental Evidence of Interaction	Reference	Direct Link to AMA Interactome		
			APP	APOE	MAPT
ADAM metallopeptidase domain 3B (ADAM3B)	Two-hybrid Test	72	—	—	—
Adrenoceptor beta 2, surface (ADRB2)	Affinity Capture-MS	73	—	—	—
Albumin (ALB)	Affinity Capture-MS; Anti-Bait Coimmunoprecipitation	41	✓	✓	—
Adaptor-related protein complex 2, mu 1 subunit (AP2M1)	Affinity Capture-MS	67	—	—	—
Adenomatous polyposis coli (APC)	Surface Plasmon Resonance	74	—	—	—
Apolipoprotein A-I (APOA1)	Affinity Capture-MS; Anti-Bait Coimmunoprecipitation	41	✓	—	—
Cullin-associated and neddylation-dissociated 1 (CAND1)	Affinity Capture-MS	66	—	—	—
Cbl proto-oncogene, E3 ubiquitin protein ligase (CBL)	Affinity Capture-MS	67	—	—	—
Cadherin 1, type 1, E-cadherin (epithelial) (CDH1)	Affinity Capture-MS; Surface Plasmon Resonance	74	—	—	—
Cystic fibrosis transmembrane conductance regulator (CFTR)	Anti-Bait Coimmunoprecipitation	75	—	—	—
COP9 signalosome subunit 5 (COPS5)	Affinity Capture-MS	66	✓	—	—
COP9 signalosome subunit 6 (COPS6)	Affinity Capture-MS	66	—	—	—
V-crk avian sarcoma virus CT10 oncogene homolog (CRK)	Affinity Capture-MS	67	—	—	—
Cullin 1 (CUL1)	Affinity Capture-MS	66	—	—	—
Cullin 2 (CUL2)	Affinity Capture-MS	66	—	—	—
Cullin 3 (CUL3)	Affinity Capture-MS	66	✓	—	—
Cullin 4A (CUL4A)	Affinity Capture-MS	66	—	—	—
Cullin 4B (CUL4B)	Affinity Capture-MS	66	✓	—	—
Cullin 5 (CUL5)	Affinity Capture-MS	66	—	—	—
DCN1, defective in cullin neddylation 1, domain containing 1 (DCUN1D)	Affinity Capture-MS	66	—	—	—
E2F transcription factor 3 (E2F3)	Pull Down	76	—	—	—
Eukaryotic translation initiation factor 4A3 (EIF4A3)	Affinity Capture-MS	77	—	—	—
Epidermal growth factor receptor pathway substrate 15 (EPS15)	Affinity Capture-MS; Anti-Tag Coimmunoprecipitation	67	—	—	—
Fibronectin 1 (FN1)	Affinity Capture-MS	78	—	—	—
GABA(A) receptor-associated protein (GABARAP)	Anti-Tag Coimmunoprecipitation	79	—	—	—
GABA(A) receptor-associated protein like 1 (GABARAPL1)	Anti-Tag Coimmunoprecipitation	79	—	—	—
GABA(A) receptor-associated protein like 2 (GABARAPL2)	Anti-Tag Coimmunoprecipitation	79	✓	—	—
Growth factor receptor-bound protein 2 (GRB2)	Affinity Capture-MS	67	✓	—	—
Immunoglobulin heavy constant alpha 1 (IGHA1)	Anti-Bait Coimmunoprecipitation	41	—	—	—
Immunoglobulin heavy constant gamma 1 (IGHG1)	Anti-Bait Coimmunoprecipitation	41	—	—	—
Immunoglobulin heavy constant mu (IGHM)	Anti-Bait Coimmunoprecipitation	41	✓	—	—
Inositol polyphosphate phosphatase-like 1 (INPPL1)	Affinity Capture-MS	67	—	—	—
IQ motif containing B1 (IQCB1)	Affinity Capture-MS	39	—	—	—
Integrin, alpha 4 (ITGA4)	Affinity Capture-MS	80	—	—	—
Keratin 1 (KRT1)	Co-Fractionation	81	—	—	—
Leucine-rich repeat containing G protein-coupled receptor 4 (LGR4)	Affinity Capture-MS	82	—	—	—
Mago-nashi homolog, proliferation-associated (MAGOH)	Affinity Capture-MS	77	—	—	—
Microtubule-associated protein 1 light chain 3 alpha (MAP1LC3A)	Anti-Tag Coimmunoprecipitation	79	✓	—	—
Microtubule-associated protein 1 light chain 3 beta (MAP1LC3B)	Anti-Tag Coimmunoprecipitation	79	—	—	—
MDM2 proto-oncogene, E3 ubiquitin protein ligase (MDM2)	Affinity Capture-MS	69	✓	—	—
Neural precursor cell expressed, developmentally down-regulated 8 (NEDD8)	Affinity Capture-MS	66	✓	—	—
Phosphoinositide-3-kinase, regulatory subunit 2 (beta) (PIK3R2)	Affinity Capture-MS	67	✓	—	—
POU class 5 homeobox 1 (POU5F1)	Affinity Capture-MS	83	—	—	—
Protein phosphatase 2, regulatory subunit B, beta (PPP2R2B)	Pull Down; Anti-Tag Coimmunoprecipitation	84	—	—	—
SH3-domain GRB2-like 3 (SH3GL3)	Two Hybrid Test	72	—	—	—
Src homology 2 domain containing transforming protein 1 (SHC1)	Affinity Capture-MS	67	✓	—	—
TRAF family member-associated NFKB activator (TANK)	Functional Linkage Network; Anti-Tag Coimmunoprecipitation	85	✓	—	—
TNF receptor-associated factor 3 interacting protein 1 (TRAF3IP1)	Anti-Bait Coimmunoprecipitation; Affinity Capture-MS	86	✓	—	—
Ubiquitin associated and SH3 domain containing B (UBASH3B)	Affinity Capture-MS	67	—	—	—
Continued					

KRT9 interacting partner	Experimental Evidence of Interaction	Reference	Direct Link to AMA Interactome		
			APP	APOE	MAPT
Ubiquitin C (UBC)	Affinity Capture-MS	68	✓	✓	✓
Ubiquitin carboxyl-terminal hydrolase L5 (UCHL5)	Affinity Capture-MS	87	✓	—	—
PAN2 poly (A) specific ribonuclease subunit (USP52)	Affinity Capture-MS	88	—	—	—
Vascular cell adhesion molecule 1 (VCAM1)	Cross-linking Studies; Affinity Capture-MS	78,80	—	✓	—
Tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, theta (YWHAQ)	<i>In vivo</i> ; Coimmunoprecipitation; Affinity Capture-MS	89	—	—	✓

Table 4. Components of the Keratin 9 interactome determined using VisANT software. The means of determining their interaction with Keratin 9 and their links to the AMA Interactome are displayed.

this method was deemed inappropriate for determining the pathways associated with Keratin 9. Instead, the functions of each of the Keratin 9 interacting proteins were evaluated using the UniProtKB database, which displays the complete Gene Ontology annotation as well as a full list of Reactome entries for each protein. Additionally, since VisANT only displays proteins that share physical interactions, we searched the literature for evidence of transcriptional regulation of Keratin 9 to establish whether Keratin 9 regulators play a role in AD. As a result of this, several pathways linked to these interacting proteins and regulatory proteins were highlighted, including wnt signalling and the ubiquitin proteasome system which have already been linked to AD^{43–45}.

Wnt Signalling in AD. The canonical wnt signalling pathway regulates the expression of a wide range of genes throughout the body and has been strongly implicated in AD^{43,44}. The cascade is initiated when wnt ligands complex with LRP5/6 and Frizzled (Fz) receptors. Fz then activates Dishevelled protein, which forms a complex with GSK3 β , adenomatous polyposis coli (APC), Axin and beta-catenin resulting in inactivation of GSK3 β and stabilization of beta-catenin. Beta catenin can then bind to T cell receptor (Tcf)/Lymphoid enhancer-binding factor (LEF) transcription factors by displacement of the repressor protein, Groucho leading to activation or repression of target genes. In the absence of wnt, or in the presence of a wnt antagonist, GSK3 β remains active, facilitating phosphorylation of beta-catenin, which is targeted to the ubiquitin-proteasome degradation pathway⁴⁴. This prevents beta-catenin from entering the nucleus and interacting with transcription factors. It is thought that downregulation of components of this cascade can result in many of the key changes that occur in the AD brain: repression of wnt signalling has so far been linked to the processing of APP, A β peptide neurotoxicity and tau phosphorylation⁴³. For example beta secretase (BACE1), the APP cleaving enzyme, is suppressed following wnt activation via beta-catenin binding to TCF4, a BACE1 promoter⁴⁶. Therefore, downregulation of the wnt pathway can result in increased cleavage of APP and therefore increased A β levels.

VisANT analysis suggests that Keratin 9 may be linked to this cascade via interactions with APC and Cadherin 1 (CDH1), which interacts with β -catenin in adherens junctions. It is unlikely however, that Keratin 9 could affect this pathway through binding to CDH1 or APC as there is no evidence of Keratin 9 acting as a signalling molecule. It is more likely that Keratin 9 levels could be influenced by this pathway as it has been suggested that wnt signalling may be involved in keratinocyte development. Dickkopf 1 (DKK1), an antagonist of the wnt pathway, has been shown to induce the expression of Keratin 9, resulting in a thicker epidermis⁴⁷. Furthermore, increased expression of DKK1 has been seen in brain tissue of AD patients and this has been causally linked to the neurodegeneration associated with the disease⁴⁸. It is thought that expression of A β induces expression of DKK1 in a P53 dependent manner and conversely, repressing DKK1 reduces A β toxicity⁴⁹. Increased expression of DKK1 could therefore not only help explain the characteristic amyloid beta toxicity, but also the presence of Keratin 9 in the CSF of AD patients. Although, to our knowledge, this is the first time a link has been proposed between DKK1 and Keratin 9 in non-epidermal tissue, both DKK1 and Keratin 9 have separately been shown to have diagnostic potential for hepatocellular carcinoma due to their dysregulation in blood serum^{10,50}. This suggests that, in certain disease conditions, DKK1 could induce the expression of Keratin 9 in non-epidermal tissue.

Interestingly, our results also show a correlation between changes in Keratin 9 and Clusterin levels in AD (Table 3). This could be explained by previous research by Killick and colleagues⁴⁹, which suggests linkages between Clusterin, DKK1 and A β . They found that knockdown of Clusterin expression results in reduced A β toxicity and DKK1 expression, suggesting that A β induced DKK1 expression is dependent on Clusterin, as well as P53. They also highlighted a potential mechanism, thought to be the amyloid cascade, whereby Clusterin mediates neurotoxicity in an A β induced clusterin/P53/DKK1/wnt-PCP-JNK pathway. Since our immunoassay detection did not include DKK1 or P53, we can only speculate that these may be linked to Clusterin and Keratin 9 based on previous research. It would therefore be of interest to include DKK1 and P53 in our biomarker panel in future in order to establish for ourselves whether these correlate with Clusterin, A β or Keratin 9.

Keratin 9 and Blood Brain Barrier Dysregulation in AD. The presence of Keratin 9 in CSF has previously been demonstrated using various mass spectrometry techniques^{12,15} and 2D electrophoresis¹³, but these techniques gave no indication of the concentration at which it existed. To our knowledge, the first immunoassay validation of the presence of Keratin 9 in CSF was obtained during our previous study which demonstrated the presence of Keratin 9 in the CSF of AD patients but not in healthy individuals¹⁶. Keratin 9 has also been identified in the proteome of blood plasma multiple times^{41,51–62}. Although sample contamination cannot be ruled out, and




Figure 3. The Interactome of Keratin 9, Apolipoprotein E (APOE), Tau (MAPT) and Amyloid Precursor Protein (APP) (nodes shown in red). VisANT software was used to identify all interacting molecules. Interacting molecules are identified as green (interacting with 2 proteins), yellow (interacting with 3 proteins), or blue (interacting with 4 proteins). Proteins with only a single interaction were excluded from the Interactome. Only the nodes with direct links to Keratin 9 have been labelled.

we can only speculate on why Keratin 9 is present in blood, the sheer number of studies documenting this result adds plausibility to our finding of Keratin 9 in blood plasma in both healthy and AD patients. Furthermore, our finding that Keratin 9 was present in the CSF of AD individuals but not in healthy individuals¹⁶, could suggest that there are differences in the blood brain barrier (BBB), enabling Keratin 9 to pass from the blood into CSF in AD individuals.

Dysfunction of the BBB has already been strongly implicated in AD. The integrity of the BBB is compromised in AD due to down-regulation of Claudin proteins, resulting in destabilisation of tight junctions between endothelial cells⁶³. Evidence also suggests that p-glycoprotein, an A β transporter at the BBB, is down-regulated in AD, resulting in compromised A β clearance and accumulation of A β in the brain⁶⁴. Interestingly, expression of p-glycoprotein and maintenance of tight junctions in the BBB are known to be regulated by the wnt signalling pathway^{63,65}. Increased expression of DKK1, as mentioned previously, inhibits this pathway and therefore contributes to increased BBB permeability, decreased p-glycoprotein expression and A β accumulation. Increased permeability of the BBB could therefore enable Keratin 9, which is up-regulated by DKK1⁴⁷, to pass from blood to CSF in AD patients.

Keratin 9 and the Ubiquitin Proteasome System in AD. The interactome produced using VisANT and subsequent analysis revealed that at least 12 of the Keratin 9 interacting partners: CAND1, CBL, CUL1, CUL2, CUL3, CUL4A, CUL4B, CUL5, DCUN1D, MDM2, NEDD8 and UBC, are components of the Ubiquitin Proteasome System (UPS)^{66–69}. This suggests that Keratin 9 may be ubiquitinated and targeted for degradation by the proteasome. However, the abnormal presence of Keratin 9 in CSF and the elevated levels of Keratin 9 in blood of AD patients indicate a failure in this system, which prevents Keratin 9 degradation. Dysregulation of the UPS has previously been linked to AD as a significant decrease in proteasome activity has been observed in various regions of the AD brain and this is thought to result in the aggregation of A β into plaques⁴⁵. This may also explain the build-up of Keratin 9 and therefore Keratin 9 may be a useful biomarker to detect the UPS dysregulation associated with AD.

Keratin 9: More than just a structural molecule. In addition to functioning as scaffolding proteins within cellular cytoskeletons, some keratins are known to be involved in cell signalling, communicating with extracellular matrix (ECM) components and other cells through desmosomes and hemidesmosomes^{70,71}. Although Keratin 9 has not yet been implicated in any signalling pathways, VisANT analysis reveals interactions with several cytoskeletal and ECM components, implying a potential signalling role. Specifically, Keratin 9 interacts with CDH1 and ITGA4, which are involved in cell-cell junctions and cell-matrix interactions, and FN1, a component of the ECM. Keratin 9 also interacts directly with other cytoskeletal components, including KRT1 and indirectly with microtubules through interactions with MAP1LC3A and MAP1LC3B. There is a wide range of other proteins with which Keratin 9 also interacts (Table 4) for which the implications of the interactions are currently unclear. As these proteins have a variety of functions it could be speculated that the interactions may have unforeseen consequences for other downstream pathways. If this were to be proven and if the affected pathways influence AD it would mean that dysregulation of Keratin 9 expression, which currently appears to be a consequence of AD, may have consequences of its own that may contribute to AD onset and progression.




Figure 4. Keratin 9 in Alzheimer's disease pathology. The diagram illustrates how Keratin 9 may interact with some of the major known factors in Alzheimer's disease including those pathways which implicate ApoE, APP, Clusterin and Tau. Dotted lines indicate unknown mechanisms.

Conclusion

Rapidly emerging evidence suggests that Alzheimer's disease is multifactorial and therefore, finding a single marker for the disease is not the key to early diagnosis. With this in mind, we do not present Keratin 9 as a stand-alone biomarker for the disease but rather, a useful component of a biomarker panel. The use of Keratin 9 as a biomarker seems controversial and counterintuitive considering that it was until recently believed to be solely expressed in terminally differentiated layers of the epidermis. However, biochemical analysis of expression patterns and *in silico* mapping of keratin 9 interactions has proved to be useful in understanding potential mechanisms that may explain the involvement of keratin 9 in the disease. These pathways, alongside other proposed AD risk factors and their likely role in AD pathology are summarized in Fig. 4. We can speculate from our research that the abnormal expression of keratin 9 in AD is likely the consequence of dysregulated signalling pathways, notably wnt signalling pathways and potentially the UPS, in addition to a break-down in the BBB. Based on this information, we suggest that P53 and DKK1 should also be considered as potential biomarkers in addition to the panel of markers we identified previously. Furthermore, the interactome of keratin 9 also suggests that its abnormal expression may have other downstream consequences resulting from its interactions with a broad range of other molecules, so this may be a potential area for future research.

References

- Langbein, L., Heid, H. W., Moll, I. & Franke, W. W. Molecular characterization of the body site-specific human epidermal cytokeratin 9: cDNA cloning, amino acid sequence, and tissue specificity of gene expression. *Differentiation* **55**, 57–71 (1993).
- Du, Z. F. *et al.* A novel mutation within the 2B rod domain of keratin 9 in a Chinese pedigree with epidermolytic palmoplantar keratoderma combined with knuckle pads and camptodactyly. *Eur J Dermatol* **21**, 675–679 (2011).
- Reis, A. *et al.* Keratin 9 gene mutations in epidermolytic palmoplantar keratoderma (EPPK). *Nat Genet* **6**, 174–179 (1994).
- Shimomura, Y., Wajid, M., Weiser, J., Kraemer, L. & Christiano, A. M. Mutations in the keratin 9 gene in Pakistani families with epidermolytic palmoplantar keratoderma. *Clin Exp Dermatol* **35**, 759–764 (2010).
- Ghosh, D., Lippert, D., Krokhin, O., Cortens, J. P. & Wilkins, J. A. Defining the membrane proteome of NK cells. *J Mass Spectrom* **45**, 1–25 (2010).
- van Niel, G. *et al.* Intestinal epithelial cells secrete exosome-like vesicles. *Gastroenterology* **121**, 337–349 (2001).
- de Mateo, S., Castillo, J., Estanyol, J. M., Balleca, J. L. & Oliva, R. Proteomic characterization of the human sperm nucleus. *Proteomics* **11**, 2714–2726 (2011).
- Bohm, D. *et al.* Comparison of tear protein levels in breast cancer patients and healthy controls using a *de novo* proteomic approach. *Oncol Rep* **28**, 429–438 (2012).
- Kim, Y. S. *et al.* Apolipoprotein A-IV as a novel gene associated with polycystic ovary syndrome. *Int J Mol Med* **31**, 707–716 (2013).
- Fu, B. S. *et al.* [Serum proteomic analysis on metastasis-associated proteins of hepatocellular carcinoma]. *Nan Fang Yi Ke Da Xue Xue Bao* **29**, 1775–1778 (2009).
- Veening, J. G. & Barendregt, H. P. The regulation of brain states by neuroactive substances distributed via the cerebrospinal fluid; a review. *Cerebrospinal Fluid Res* **7**, 1 (2010).
- Pan, S. *et al.* A combined dataset of human cerebrospinal fluid proteins identified by multi-dimensional chromatography and tandem mass spectrometry. *Proteomics* **7**, 469–473 (2007).
- Jiang, S. *et al.* Proteomic analysis of the cerebrospinal fluid in multiple sclerosis and neuromyelitis optica patients. *Mol Med Rep* **6**, 1081–1086 (2012).
- Ballard, C. *et al.* Alzheimer's disease. *Lancet* **377**, 1019–1031 (2011).
- Vafadar-Isfahani, B. *et al.* Identification of SPARC-like 1 protein as part of a biomarker panel for Alzheimer's disease in cerebrospinal fluid. *J Alzheimers Dis* **28**, 625–636 (2012).
- Richens, J. L. *et al.* Practical detection of a definitive biomarker panel for Alzheimer's disease; comparisons between matched plasma and cerebrospinal fluid. *Int J Mol Epidemiol Genet* **5**, 53–70 (2014).
- Alafuzoff, I., Adolfsson, R., Bucht, G. & Winblad, B. Albumin and immunoglobulin in plasma and cerebrospinal fluid, and blood-cerebrospinal fluid barrier function in patients with dementia of Alzheimer type and multi-infarct dementia. *J Neurol Sci* **60**, 465–472 (1983).
- Buchhave, P. *et al.* Cerebrospinal fluid levels of beta-amyloid 1-42, but not of tau, are fully changed already 5 to 10 years before the onset of Alzheimer dementia. *Arch Gen Psychiatry* **69**, 98–106 (2012).
- Cortes-Canteli, M. *et al.* Fibrinogen and beta-amyloid association alters thrombosis and fibrinolysis: a possible contributing factor to Alzheimer's disease. *Neuron* **66**, 695–709 (2010).
- Soares, H. D. *et al.* Plasma Biomarkers Associated With the Apolipoprotein E Genotype and Alzheimer Disease. *Arch Neurol*, 1–8 (2012).
- Yin, G. N., Lee, H. W., Cho, J. Y. & Suk, K. Neuronal pentraxin receptor in cerebrospinal fluid as a potential biomarker for neurodegenerative diseases. *Brain Res* **1265**, 158–170 (2009).
- Mueller, C. *et al.* The heme degradation pathway is a promising serum biomarker source for the early detection of Alzheimer's disease. *J Alzheimers Dis* **19**, 1081–1091 (2010).
- Li, X., Long, J., He, T., Belshaw, R. & Scott, J. Integrated genomic approaches identify major pathways and upstream regulators in late onset Alzheimer's disease. *Sci Rep* **5**, 12393 (2015).
- Hu, Z. *et al.* VisANT 4.0: Integrative network platform to connect genes, drugs, diseases and therapies. *Nucleic Acids Res* **41**, W225–231 (2013).
- Richens, J. L., Morgan, K. & O'Shea, P. Reverse engineering of Alzheimer's disease based on biomarker pathways analysis. *Neurobiol Aging* **35**, 2029–2038 (2014).
- Johnson, C. *et al.* Visualization and biochemical analyses of the emerging mammalian 14-3-3-phosphoproteome. *Mol Cell Proteomics* **10**, M110 005751 (2011).
- Reddy, A. *et al.* Robust gene network analysis reveals alteration of the STAT5a network as a hallmark of prostate cancer. *Genome Inform* **24**, 139–153 (2010).
- Araujo, D. J. *et al.* FoxP1 orchestration of ASD-relevant signaling pathways in the striatum. *Genes Dev* **29**, 2081–2096 (2015).
- Mellor, J. C., Yanai, I., Clodfelter, K. H., Mintseris, J. & DeLisi, C. Predictome: a database of putative functional links between proteins. *Nucleic Acids Res* **30**, 306–309 (2002).
- Bohm, C. *et al.* Current and future implications of basic and translational research on amyloid-beta peptide production and removal pathways. *Mol Cell Neurosci* **66**, 3–11 (2015).
- Kim, J., Yoon, H., Basak, J. & Kim, J. Apolipoprotein E in synaptic plasticity and Alzheimer's disease: potential cellular and molecular mechanisms. *Mol Cells* **37**, 767–776 (2014).
- Sato, N. & Morishita, R. The roles of lipid and glucose metabolism in modulation of beta-amyloid, tau, and neurodegeneration in the pathogenesis of Alzheimer disease. *Front Aging Neurosci* **7**, 199 (2015).
- Croft, D. *et al.* The Reactome pathway knowledgebase. *Nucleic Acids Res* **42**, D472–477 (2014).
- Denery, J. R., Nunes, A. A. & Dickerson, T. J. Characterization of differences between blood sample matrices in untargeted metabolomics. *Anal Chem* **83**, 1040–1047 (2011).
- Thambisetty, M. *et al.* Association of plasma clusterin concentration with severity, pathology, and progression in Alzheimer disease. *Arch Gen Psychiatry* **67**, 739–748 (2010).
- Noguchi, M. *et al.* Roles of serum fibrinogen alpha chain-derived peptides in Alzheimer's disease. *Int J Geriatr Psychiatry* **29**, 808–818 (2014).
- Moll, R., Divo, M. & Langbein, L. The human keratins: biology and pathology. *Histochem Cell Biol* **129**, 705–733 (2008).
- Fu, D. J. *et al.* Keratin 9 is required for the structural integrity and terminal differentiation of the palmoplantar epidermis. *J Invest Dermatol* **134**, 754–763 (2014).
- Sang, L. *et al.* Mapping the NPHP-JBTS-MKS protein network reveals ciliopathy disease genes and pathways. *Cell* **145**, 513–528 (2011).
- Otto, E. A. *et al.* Nephrocystin-5, a ciliary IQ domain protein, is mutated in Senior-Loken syndrome and interacts with RPGR and calmodulin. *Nat Genet* **37**, 282–288 (2005).
- Zhou, M. *et al.* An investigation into the human serum “interactome”. *Electrophoresis* **25**, 1289–1298 (2004).
- Croft, D. *et al.* Reactome: a database of reactions, pathways and biological processes. *Nucleic Acids Res* **39**, D691–697 (2011).
- De Ferrari, G. V. *et al.* Wnt/beta-catenin signaling in Alzheimer's disease. *CNS Neurol Disord Drug Targets* **13**, 745–754 (2014).
- De Ferrari, G. V. & Inestrosa, N. C. Wnt signaling function in Alzheimer's disease. *Brain Res Brain Res Rev* **33**, 1–12 (2000).
- Riederer, B. M., Leuba, G., Vernay, A. & Riederer, I. M. The role of the ubiquitin proteasome system in Alzheimer's disease. *Exp Biol Med (Maywood)* **236**, 268–276 (2011).

46. Parr, C., Mirzaei, N., Christian, M. & Sastre, M. Activation of the Wnt/beta-catenin pathway represses the transcription of the beta-amyloid precursor protein cleaving enzyme (BACE1) via binding of T-cell factor-4 to BACE1 promoter. *FASEB J* **29**, 623–635 (2015).
47. Yamaguchi, Y. *et al.* Dickkopf 1 (DKK1) regulates skin pigmentation and thickness by affecting Wnt/beta-catenin signaling in keratinocytes. *FASEB J* **22**, 1009–1020 (2008).
48. Caraci, F. *et al.* The Wnt antagonist, Dickkopf-1, as a target for the treatment of neurodegenerative disorders. *Neurochem Res* **33**, 2401–2406 (2008).
49. Killick, R. *et al.* Clusterin regulates beta-amyloid toxicity via Dickkopf-1-driven induction of the wnt-PCP-JNK pathway. *Mol Psychiatry* **19**, 88–98 (2014).
50. Fatima, S., Luk, J. M., Poon, R. T. & Lee, N. P. Dysregulated expression of dickkopfs for potential detection of hepatocellular carcinoma. *Expert Rev Mol Diagn* **14**, 535–548 (2014).
51. Bassani-Sternberg, M. *et al.* Soluble plasma HLA peptidome as a potential source for cancer biomarkers. *Proc Natl Acad Sci USA* **107**, 18769–18776 (2010).
52. Farrah, T. *et al.* A high-confidence human plasma proteome reference set with estimated concentrations in PeptideAtlas. *Mol Cell Proteomics* **10**, M110 006353 (2011).
53. Jeong, S. K. *et al.* Data management and functional annotation of the Korean reference plasma proteome. *Proteomics* **10**, 1250–1255 (2010).
54. Liu, X. *et al.* Mapping the human plasma proteome by SCX-LC-IMS-MS. *J Am Soc Mass Spectrom* **18**, 1249–1264 (2007).
55. Madian, A. G. & Regnier, F. E. Profiling carbonylated proteins in human plasma. *J Proteome Res* **9**, 1330–1343 (2010).
56. Manabe, T., Jin, Y. & Tani, O. Assignment of human plasma polypeptides on a nondenaturing 2-D gel using MALDI-MS and PMF and comparisons with the results of intact protein mapping. *Electrophoresis* **28**, 843–863 (2007).
57. Panchaud, A. *et al.* Precursor acquisition independent from ion count: how to dive deeper into the proteomics ocean. *Anal Chem* **81**, 6481–6488 (2009).
58. Prentice, R. L. *et al.* Novel proteins associated with risk for coronary heart disease or stroke among postmenopausal women identified by in-depth plasma proteome profiling. *Genome Med* **2**, 48 (2010).
59. Qian, W. J. *et al.* Quantitative proteome analysis of human plasma following *in vivo* lipopolysaccharide administration using ¹⁶O/¹⁸O labeling and the accurate mass and time tag approach. *Mol Cell Proteomics* **4**, 700–709 (2005).
60. Qian, W. J. *et al.* Plasma proteome response to severe burn injury revealed by ¹⁸O-labeled “universal” reference-based quantitative proteomics. *J Proteome Res* **9**, 4779–4789 (2010).
61. Sennels, L. *et al.* Proteomic analysis of human blood serum using peptide library beads. *J Proteome Res* **6**, 4055–4062 (2007).
62. Tu, C. J. *et al.* High-sensitivity analysis of human plasma proteome by immobilized isoelectric focusing fractionation coupled to mass spectrometry identification. *J Proteome Res* **4**, 1265–1273 (2005).
63. Liebner, S. *et al.* Wnt/beta-catenin signaling controls development of the blood-brain barrier. *J Cell Biol* **183**, 409–417 (2008).
64. Deo, A. K. *et al.* Activity of P-Glycoprotein, a beta-Amyloid Transporter at the Blood-Brain Barrier, Is Compromised in Patients with Mild Alzheimer Disease. *J Nucl Med* **55**, 1106–1111 (2014).
65. Liu, L., Wan, W., Xia, S., Kalionis, B. & Li, Y. Dysfunctional Wnt/beta-catenin signaling contributes to blood-brain barrier breakdown in Alzheimer’s disease. *Neurochem Int* **75**, 19–25 (2014).
66. Bennett, E. J., Rush, J., Gygi, S. P. & Harper, J. W. Dynamics of cullin-RING ubiquitin ligase network revealed by systematic quantitative proteomics. *Cell* **143**, 951–965 (2010).
67. Brehme, M. *et al.* Charting the molecular network of the drug target Bcr-Abl. *Proc Natl Acad Sci USA* **106**, 7414–7419 (2009).
68. Lopitz-Otsoa, F. *et al.* Integrative analysis of the ubiquitin proteome isolated using Tandem Ubiquitin Binding Entities (TUBEs). *J Proteomics* **75**, 2998–3014 (2012).
69. Nicholson, J. *et al.* A systems wide mass spectrometric based linear motif screen to identify dominant *in-vivo* interacting proteins for the ubiquitin ligase MDM2. *Cell Signal* **26**, 1243–1257 (2014).
70. Homberg, M. & Magin, T. M. Beyond expectations: novel insights into epidermal keratin function and regulation. *Int Rev Cell Mol Biol* **311**, 265–306 (2014).
71. Jones, J. C., Hopkinson, S. B. & Goldfinger, L. E. Structure and assembly of hemidesmosomes. *BioEssays* **20**, 488–494 (1998).
72. Stelzl, U. *et al.* A human protein-protein interaction network: a resource for annotating the proteome. *Cell* **122**, 957–968 (2005).
73. Roy, S. J. *et al.* Novel, gel-free proteomics approach identifies RNF5 and JAMP as modulators of GPCR stability. *Mol Endocrinol* **27**, 1245–1266 (2013).
74. Catimel, B. *et al.* Biosensor-based micro-affinity purification for the proteomic analysis of protein complexes. *J Proteome Res* **4**, 1646–1656 (2005).
75. Colas, J. *et al.* Disruption of cyokeratin-8 interaction with F508del-CFTR corrects its functional defect. *Hum Mol Genet* **21**, 623–634 (2012).
76. von Eyss, B. *et al.* The SNF2-like helicase HELLS mediates E2F3-dependent transcription and cellular transformation. *EMBO J* **31**, 972–985 (2012).
77. Singh, G. *et al.* The cellular EJC interactome reveals higher-order mRNP structure and an EJC-SR protein nexus. *Cell* **151**, 750–764 (2012).
78. Humphries, J. D. *et al.* Proteomic analysis of integrin-associated complexes identifies RCC2 as a dual regulator of Rac1 and Arf6. *Sci Signal* **2**, ra51 (2009).
79. Behrends, C., Sowa, M. E., Gygi, S. P. & Harper, J. W. Network organization of the human autophagy system. *Nature* **466**, 68–76 (2010).
80. Byron, A., Humphries, J. D., Craig, S. E., Knight, D. & Humphries, M. J. Proteomic analysis of alpha4beta1 integrin adhesion complexes reveals alpha-subunit-dependent protein recruitment. *Proteomics* **12**, 2107–2114 (2012).
81. Havugimana, P. C. *et al.* A census of human soluble protein complexes. *Cell* **150**, 1068–1081 (2012).
82. Carmon, K. S., Gong, X., Yi, J., Thomas, A. & Liu, Q. RSPO-LGR4 functions via IQGAP1 to potentiate Wnt signaling. *Proc Natl Acad Sci USA* **111**, E1221–1229 (2014).
83. Hawkins, P. G. & Morris, K. V. Transcriptional regulation of Oct4 by a long non-coding RNA antisense to Oct4-pseudogene 5. *Transcription* **1**, 165–175 (2010).
84. Glatter, T., Wepf, A., Aebersold, R. & Gstaiger, M. An integrated workflow for charting the human interaction proteome: insights into the PP2A system. *Mol Syst Biol* **5**, 237 (2009).
85. Bouwmeester, T. *et al.* A physical and functional map of the human TNF-alpha/NF-kappa B signal transduction pathway. *Nat Cell Biol* **6**, 97–105 (2004).
86. Guo, C. W. *et al.* Proteomic analysis reveals novel binding partners of MIP-T3 in human cells. *Proteomics* **10**, 2337–2347 (2010).
87. Fang, Y. *et al.* The interaction between ubiquitin C-terminal hydrolase 37 and glucose-regulated protein 78 in hepatocellular carcinoma. *Mol Cell Biochem* **359**, 59–66 (2012).
88. Bett, J. S. *et al.* The P-body component USP52/PAN2 is a novel regulator of HIF1A mRNA stability. *Biochem J* **451**, 185–194 (2013).
89. Jin, J. *et al.* Proteomic, functional, and domain-based analysis of *in vivo* 14-3-3 binding proteins involved in cytoskeletal regulation and cellular organization. *Curr Biol* **14**, 1436–1450 (2004).

Acknowledgements

This work was funded by a Medical Research Council Developmental Pathways Grant, grant number 1001500.

Author Contributions

J.L.R. and P.O. designed the research. J.L.R., H.L.S., M.B., F.C. and K.A.V. conducted the research. J.L.R., H.L.S. and P.O. wrote the manuscript. N.B. and K.M. contributed to the writing of the manuscript. All authors have reviewed and approved the final manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Richens, J. L. *et al.* Rationalising the role of Keratin 9 as a biomarker for Alzheimer's disease. *Sci. Rep.* **6**, 22962; doi: 10.1038/srep22962 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>