



Published in final edited form as:

Sex Transm Dis. 2010 September ; 37(9): 585–590.

Prevalence and Correlates of *Trichomonas vaginalis* Infection Among Female US Federal Prison Inmates

Siobhan Sutcliffe, PHD^{*}, Sara B. Newman, DrPH[†], Andrew Hardick, BA[‡], and Charlotte A. Gaydos, DrPH[‡]

^{*}Department of Surgery and The Alvin J. Siteman Cancer Center, Washington University School of Medicine, St. Louis, MO

[†]Division of Risk Management, National Park Service, Washington, DC

[‡]Division of Infectious Diseases, Department of Medicine, Johns Hopkins Medical Institutions, Baltimore, MD

Abstract

Background—Previous studies have observed high prevalences of *Trichomonas vaginalis* infection among women entering US jails and state prisons (22%–47%). We sought to determine the prevalence among women incarcerated in 2 US female-only federal prisons.

Methods—Female inmates were recruited at 2 prisons (n = 624). Participants completed a self-administered questionnaire and provided self-collected first-catch urine and vaginal swab specimens. Specimens were tested for *T. vaginalis* DNA.

Results—Approximately 8.5% of participants at the first prison, and 8.3% at the second prison had a positive urine result, vaginal swab result or both, for a combined prevalence of 8.5%. Using positivity in either specimen as the reference standard, urine polymerase chain reaction had a sensitivity of 66.7% and vaginal swab polymerase chain reaction had a sensitivity of 84.4%. The only significant positive correlate of *T. vaginalis* infection was lower household income before arrest. Other variables nonsignificantly positively correlated with *T. vaginalis* were being employed at the time of arrest, having experienced sexual, physical, or emotional abuse by a family member, having a parent who had not had a drug or alcohol addiction, never exchanging sex for money or drugs, ever being pregnant, having abnormal vaginal bleeding/spotting, and having concurrent chlamydia or gonorrhea.

Conclusions—Although not as high as in other studies of women entering US jails and state prisons, our observed *T. vaginalis* prevalence of 8.5% was much higher than in the general US population. Therefore, screening for *T. vaginalis* infection may be warranted at federal prison entry, as well as sexual health education during prison stay.

Previous studies have observed high prevalences of sexually transmitted infections (STIs) among women entering jail and juvenile corrections facilities.¹ In a study of adult women entering correctional facilities in several US states and Puerto Rico, the median prevalence

of chlamydia was 6.4%, and those of gonorrhea and syphilis were 2.9% and 2.1%, respectively.¹ Less is known about the prevalence of *Trichomonas vaginalis* infection among incarcerated women. *T. vaginalis* is a common, sexually transmitted protozoan known to cause vaginitis and adverse birth outcomes, such as premature rupture of the membranes, in women, and increased risk and transmission of human immunodeficiency virus infection.² In the 2 surveys of *T. vaginalis* conducted to date among incarcerated women, high prevalences of infection have been observed, ranging from 22% among nonpregnant inmates³ to 47% among newly incarcerated pregnant inmates.⁴

We previously investigated prevalences of chlamydia and gonorrhea in female US federal prison inmates, and found the prevalence of chlamydia to be high among women <30 years of age (prison 2: 3.5%), but low among all participants combined (prison 1: chlamydia = 1.2%, gonorrhea = 0.3%; prison 2: chlamydia = 2.3%, gonorrhea = 0.0%).⁵ This finding was not entirely unexpected because women at the first prison were screened for chlamydia and gonorrhea at entry, and women at the second prison may have been screened and treated before being transferred to federal prison, or treated earlier in their prison stay. In addition, federal prison inmates tend to be older than those in jails and corrections facilities, and are thus at possibly lesser risk for chlamydia and gonorrhea and greater risk for *T. vaginalis* infection, as the risk of chlamydia and gonorrhea typically decreases with age, and that of *T. vaginalis* infection typically increases with age beyond adolescence.⁶ Therefore, we investigated the prevalence and correlates of *T. vaginalis* infection in this older female incarcerated population.

MATERIALS AND METHODS

Study Population

Study participants were recruited from 2 US female-only federal prisons. The first prison, located in the Midwest, screened all inmates for chlamydia and gonorrhea at entry (screening prison, SP), whereas the second prison, located in New England, only tested inmates for STIs if they presented with signs or symptoms of infection (nonscreening prison, NSP). Female inmates learned about the study at gatherings for group announcements (“call-outs”) held between August and October, 2001 at each prison. Only women 18 to 45 years of age were invited to attend call-out. Women who consented to participate in the study completed a self-administered questionnaire and provided self-collected urine and vaginal swab specimens. No incentives were provided for participation. Further details of study procedures were described elsewhere.⁵

This study was approved by the institutional review boards of the Uniformed Services University of the Health Sciences, the Federal Bureau of Prisons, and the Johns Hopkins Medical Institutions.

Data Collection

On the questionnaire, women were asked to provide information on demographics; reason and length of incarceration; substance use, sexual and reproductive history; and current STI symptoms. This questionnaire was developed based on focus groups and group-based

cognitive interviews with female federal prison inmates,⁷ and was available in English and Spanish. After completion of the questionnaire, women were asked to provide approximately 20 mL of first-catch urine and a vaginal swab. Specimens were stored at 4°C and transported to the laboratory for processing within 4 days of collection. They were then maintained frozen at -80°C until testing.

Laboratory Methods

Urine and vaginal swabs were tested for *T. vaginalis* DNA by BTUB FRET real-time polymerase chain reaction (PCR). In a previous validation study, this assay had a sensitivity of 90.5% in male and female thawed urine specimens, and a specificity of 100%.⁸

Statistical Analysis

Proportions of women positive for *T. vaginalis* by urine PCR, swab PCR and combinations of these 2 methods were calculated separately for each prison, and for both prisons combined. *T. vaginalis* correlates were investigated by calculating medians or proportions of each covariate by *T. vaginalis* status, and comparing these estimates by Wilcoxon rank sum test or Chi-squared/Fisher's exact tests, as appropriate. *T. vaginalis* positivity was defined as a positive urine result, swab result, or both. Sensitivity analyses were performed using 2 alternate case definitions: (1) a positive urine result and (2) a positive swab result. Stratified analyses were performed by race (black, non-black) and duration of incarceration (<2, 2 years).

RESULTS

Of the approximately 1344 female inmates eligible for the study, 1230 (91.5%, 90% at the SP, and 93% at the NSP) attended call-out. Reasons for not attending call-out included confinement in a secure housing unit, illness, inability to be released from work, or personal choice. Nine hundred eighty-eight inmates (80.3%) who attended call-out volunteered to participate in the study and completed the self-administered questionnaire, 363 from the SP and 625 from the NSP. Of those who volunteered at the SP, 331 provided urine and swab specimens, 27 provided only urine, and 5 provided only swabs. Because of time constraints at the SP, 117 additional women were unable to complete the questionnaire, but did provide either urine (n = 17), swabs (n = 2), or both (n = 98) for testing. Thus, 473 urine and 436 swab specimens were available for testing at the SP. At the NSP, 614 participants originally provided either urine or swabs. However, several boxes of these specimens (77% of specimens) were discarded after chlamydia and gonorrhea testing to create additional freezer space before the decision was made to perform *T. vaginalis* testing. Boxes were originally filled in order of participation, and were discarded at random; therefore, it is unlikely that discarded specimens differed from nondiscarded specimens by *T. vaginalis* or correlate status. After discarding specimens, 142 urine and 79 swab specimens remained from the NSP, 77 urine and swab specimens from the same participant, 65 individual urine specimens, and 2 individual swabs.

At the SP, 27 (5.7%) participants had a positive urine result and 35 (8.0%) had a positive vaginal swab result, for a total of 41 (8.5%, 95% confidence interval [CI]: 6.0%–11.0%)

participants positive by either test. At the NSP, 11 (7.7%) participants had a positive urine result and 3 (3.8%) had a positive swab result, for a total of 12 (8.3%, 95% CI: 3.8%–12.8%) positive using either specimen. When information from both prisons was combined, 53 (8.5%, 95% CI: 6.3%–10.7%) participants had a positive urine result, swab result, or both. Considering only women who provided both specimens (n = 506), 7 (1.4%) were positive by urine results only, 15 (3%) by swab results only, and 23 (4.6%) by both specimens (Table 1). Among these women, the sensitivity of urine PCR was 66.7% (95% CI: 52.9%–80.4%) and that of swab PCR was 84.4% (95% CI: 73.9%–95.0%) using positivity in either specimen as the reference standard.

As the total number of positive participants was low and as similar prevalences were observed at each prison, *T. vaginalis* correlates were investigated using the combined data for both prisons. In this analysis, *T. vaginalis* positivity was defined as a positive urine result, swab result, or both. No differences were observed between *T. vaginalis*-positive and -negative women by age, race/ethnicity, education, or marital status. Positive women were nonsignificantly more likely to have been born outside the continental United States than negative women, although this finding was driven by urine results. No correlation was observed when infection was defined by a positive swab result (Table 2 and data not shown).

T. vaginalis-positive women were significantly more likely to have had a legal or illegal household income \geq \$15,000, and nonsignificantly more likely to have been legally employed before their arrest than negative women, irrespective of the case definition. No differences were observed by type of crime committed (only 1 participant reported incarceration for prostitution), or length of incarceration. In addition, no differences were observed by cigarette smoking or use of illegal drugs or controlled substances in the 12 months before arrest. Among women who reported using illegal drugs or controlled substances during this time, positive women tended to be more likely to have used marijuana, and less likely to have used methamphetamines or other drugs, such as LSD, PCP, or barbiturates. Positive women were nonsignificantly more likely to report sexual, physical, or emotional abuse by a family member, and less likely to report parental drug or alcohol addiction. No differences were observed by parental incarceration (Table 2).

About sexual history, no differences were observed between *T. vaginalis*-positive and -negative women by lifetime history of vaginal intercourse with a male partner, or recent history before prison entry. No differences were also observed by lifetime history of any form of sexual intercourse with a female partner. Null results were observed both before and after taking into consideration consistency of participant responses across related sexual history questions. Positive women were nonsignificantly less likely to have ever exchanged sexual activity for money or drugs, and nonsignificantly more likely to have ever been pregnant. No differences were observed by method of birth control or reported pelvic examination in the 12 months before arrest (Table 3).

About current symptoms, *T. vaginalis*-positive women were nonsignificantly more likely to report an abnormal/unusual discharge and vaginal bleeding/spotting other than their normal period than -negative women. The correlation between abnormal/unusual discharge and

infection was more pronounced for positive urine results. When only swab results were considered, positive women were more likely to report lower abdominal/pelvic pain. Finally, positive women were nonsignificantly more likely to have concurrent chlamydia or gonorrhea, a finding that was more pronounced for positive swab results. Generally similar correlates were observed when the analyses were stratified by race and 2 years' incarceration. Too few positive women were incarcerated for <1 year to stratify by 1 year's incarceration (Table 3 and data not shown).

DISCUSSION

The prevalence of *T. vaginalis* infection among female prison inmates at 2 US federal prisons was 8.5%. More infections were detected by self-collected vaginal swab than urine. The only significant positive correlate of *T. vaginalis* was lower income. Other variables that appeared to be positively correlated with infection, but that were not statistically significant, included being employed at the time of arrest, having experienced sexual, physical or emotional abuse by a family member, having a parent who had not had a drug or alcohol addiction, never exchanging sexual activity for money or drugs, ever being pregnant, having abnormal vaginal bleeding/spotting, and having concurrent chlamydia or gonorrhea. A positive urine result was additionally correlated with being born outside the continental United States and having an abnormal/unusual discharge, whereas a positive swab result was additionally correlated with a greater likelihood of lower abdominal/pelvic pain. Interestingly, length of incarceration and sexual history were not correlated with infection by any definition.

Relative to other incarcerated populations, in which prevalences as high as 22% to 47% have been observed,^{3,4} our observed prevalence of 8.5% is low. This finding is not surprising because a lower proportion of women in our federal prison population reported exchanging sexual activity for money/drugs than in other incarcerated populations (17% vs. 27%–29%^{3,4}), and because women in our population had been incarcerated for a longer period of time (median = 19 months vs. at entry^{3,4}), allowing for greater opportunity for screening, treatment for symptoms, or spontaneous cure. However, when compared to general population estimates (2.8%–3.1%^{9,10}), our observed prevalence of 8.5% is high. One possible explanation for this difference may be the greater likelihood of high-STI risk correlates/behaviors, such as black race,¹¹ lower education,^{10,12–14} lower income,^{10,13,14} greater sexual experience/history,^{10,12–15} illegal drug use,^{4,14–19} and previous or concomitant STIs^{4,9,12–14,17,20} in our prison population than in the general population. Within our study population, however, these high-STI risk correlates/behaviors were not correlated with *T. vaginalis*, with the exception of lower income and possibly concomitant STIs. The reasons for these differing correlations are unclear, but perhaps once incarceration is held constant, factors such as race/ethnicity, lower education, sexual experience/history, and illegal drug use are no longer as correlated with infection as in less highly selected populations, although correlations have been observed in other highly selected populations.^{4,12–18,20,21} The interpretation of the positive correlation with lower income in our study population is also difficult because no correlation was observed with education, which is typically correlated with income, and a nonsignificant positive correlation was observed with full- or part-time employment, which is typically correlated with higher

income. However, perhaps in our study population, full- or part-time employment was correlated with lower total family income because it represented the need for employment by female single heads of households, or a lesser likelihood of illegal contributions to household income.

As stated earlier, no correlation was observed for length of incarceration, which ranged from 1 month to almost 11 years among positive women (median = 18.5 months). This lack of correlation raises the question as to whether detected infections were long-term infections acquired before incarceration, short-term infections acquired in prison, or possibly misreported lengths of incarceration. *T. vaginalis* has been observed to persist for at least 12 weeks to 1 year in women^{16,22,23}; therefore, it is possible that some infections may have been acquired before incarceration and persisted until the study date. Alternatively, although no evidence of within-jail infection was observed at a county correctional center,²⁴ some infections may potentially have been acquired in our prison population through sexual relationships with prison staff members or fellow inmates, as infection has been observed following lesbian sexual activities.^{25,26} Therefore, detected *T. vaginalis* infections may represent a combination of infections acquired before prison and during prison stay, making it difficult to identify correlates of prevalent infection. We attempted to investigate possible differing correlates for infections acquired before or during prison stay by stratifying the analyses by length of incarceration, as participants incarcerated for a lesser amount of time may have been more likely to have been infected before prison entry, and participants incarcerated for a greater amount of time may have been more likely to have been infected in prison. However, no additional correlates were identified when we stratified the analyses by 2 years' incarceration, and too few positive women were incarcerated for <1 year to stratify by 1 year's incarceration.

Another factor that may have contributed to difficulties in identifying *T. vaginalis* correlates is some inconsistencies in participants' self-reported responses, particularly those related to sexual history. One possible reason for these inconsistencies, which emerged in focus group discussions, is a general mistrust of how study information might be used, particularly information on sensitive or illegal topics, such as sexual activity in prison.⁷ To reduce this concern, we limited sexual activity questions to the period before prison entry, or inquired about participants' entire sexual history. Nevertheless, some questions that incorporated information on sexual activity during prison stay may not have been answered truthfully. Another possible reason for inconsistencies is the distinction between consensual sexual relationships, and sexual abuse or rape, which was reported by a large proportion of inmates in group discussions. Whether participants included sexual abuse/rape in their responses to questions on sexual history is unclear. We attempted to address inconsistencies in participants' responses to sexual history questions by investigating both participants' original responses and those found to be consistent across related questions, neither of which was correlated with *T. vaginalis* in overall, race-specific, or duration of incarceration-specific analyses.

About assessment of the outcome, the PCR assay used had a high sensitivity and specificity for *T. vaginalis*,⁸ and use of 2 different specimens should have increased the sensitivity further. However, some infections may still have been missed because of using thawed

rather than fresh specimens, as *T. vaginalis* DNA is less stable in thawed specimens. Therefore, the actual prevalence of *T. vaginalis* in our prison population may have been even higher than observed. When we considered the sensitivity of each of the 2 specimens/methods separately, we found that vaginal swab PCR had a higher sensitivity than urine PCR. This finding is consistent with the results from several previous studies,^{27–30} and the more frequent localization of *T. vaginalis* to the vagina than the urethra.² It may also possibly be explained by lower stability of *T. vaginalis* DNA in urine than swabs both before and during storage.³¹ Despite these differences, most identified correlates were similar for urine and swab PCR. One exception was symptoms; women positive by urine PCR were nonsignificantly more likely to report an abnormal/unusual vaginal discharge, whereas women positive by swab PCR were nonsignificantly more likely to report lower abdominal/pelvic pain than negative women. The reasons for these differences are unclear. Perhaps, in the case of the stronger positive correlation between abnormal/unusual vaginal discharge and urine positivity, urine PCR is better at detecting infections associated with discharge because of contamination of the urethral area by vaginal discharge, whereas vaginal swab PCR may be equally good at detecting infections with or without discharge. For the positive correlation between abdominal/pelvic pain and vaginal swab positivity, possibly higher vaginal parasite load in women with *T. vaginalis*-associated abdominal/pelvic pain may make it easier to detect infections by swab than urine.

As a final consideration and possible limitation, our ability to detect statistically significant correlations may have been low because of the relatively small number of positive women in our study population. In addition, some observed correlations, particularly those that were not statistically significant and those that were not observed for both urine and swab PCR, may have been observed by chance.

In conclusion, although our observed *T. vaginalis* prevalence of 8.5% was not as high as in other studies of women entering US jails and state prisons, it was still considerably higher than in the general US population. It was also higher than prevalences of chlamydia and gonorrhea in the same study population, even among the subgroup of women with the highest prevalence, i.e., those <30 years of age (3.5% for chlamydia) for whom the Federal Bureau of Prisons modified their screening protocol to a more targeted age-based approach.⁵ Therefore, given the higher observed prevalence of *T. vaginalis* infection in this population, we believe that universal *T. vaginalis* screening at federal prison entry may also be warranted, as well as sexual health education during prison stay.

Acknowledgments

The authors thank Ratna Pakpahan for help with literature review, and Catherine G. Sutcliffe for critical review of this manuscript.

Supported by grants from the Centers for Disease Control and Prevention, the Federal Bureau of Prisons, the Uniformed Services University of the Health Sciences, and the HIV Prevention Trials Network (U01 A1068613).

References

1. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance, 2007. Atlanta, GA: US Department of Health and Human Services; 2008.

2. Hobbs, MM.; Sena, AC.; Swygard, H., et al. *Trichomonas vaginalis* and trichomoniasis. In: Sparling, PF.; Stamm, WE.; Piot, P., et al., editors. Sexually Transmitted Diseases. 4. New York, NY: McGraw-Hill Medical; 2008. p. 771-793.
3. Willers DM, Peipert JF, Allsworth JE, et al. Prevalence and predictors of sexually transmitted infection among newly incarcerated females. *Sex Transm Dis.* 2008; 35:68–72. [PubMed: 18090178]
4. Shuter J, Bell D, Graham D, et al. Rates of and risk factors for trichomoniasis among pregnant inmates in New York City. *Sex Transm Dis.* 1998; 25:303–307. [PubMed: 9662764]
5. Newman SB, Nelson MB, Friedman HB, et al. Should female federal inmates be screened for chlamydial and gonococcal infection? *J Correct Health Care.* 2005; 11:137–155.
6. Holmes, KK.; Sparling, PF.; Stamm, WE., et al. Sexually Transmitted Diseases. 4. New York, NY: McGraw-Hill Medical; 2008.
7. Newman SB, Girasek DC, Friedman H. Using qualitative methods to design an epidemiologic study on sexually transmitted diseases in female prisoners. *Sex Transm Dis.* 2003; 30:531–532. [PubMed: 12782959]
8. Hardick J, Yang S, Lin S, et al. Use of the Roche LightCycler instrument in a real-time PCR for *Trichomonas vaginalis* in urine samples from females and males. *J Clin Microbiol.* 2003; 41:5619–22. [PubMed: 14662951]
9. Miller WC, Swygard H, Hobbs MM, et al. The prevalence of trichomoniasis in young adults in the United States. *Sex Transm Dis.* 2005; 32:593–598. [PubMed: 16205299]
10. Sutton M, Sternberg M, Koumans EH, et al. The prevalence of *Trichomonas vaginalis* infection among reproductive-age women in the United States, 2001–2004. *Clin Infect Dis.* 2007; 45:1319–1326. [PubMed: 17968828]
11. Shafir SC, Sorvillo FJ, Smith L. Current issues and considerations regarding trichomoniasis and human immunodeficiency virus in African-Americans. *Clin Microbiol Rev.* 2009; 22:37–45. [PubMed: 19136432]
12. Helms DJ, Mosure DJ, Metcalf CA, et al. Risk factors for prevalent and incident *Trichomonas vaginalis* among women attending three sexually transmitted disease clinics. *Sex Transm Dis.* 2008; 35:484–488. [PubMed: 18360314]
13. Cotch MF, Pastorek JG II, Nugent RP, et al. The Vaginal Infections and Prematurity Study Group. Demographic and behavioral predictors of *Trichomonas vaginalis* infection among pregnant women. *Obstet Gynecol.* 1991; 78:1087–1092. [PubMed: 1945213]
14. Sorvillo F, Kovacs A, Kerndt P, et al. Risk factors for trichomoniasis among women with human immunodeficiency virus (HIV) infection at a public clinic in Los Angeles County, California: Implications for HIV prevention. *Am J Trop Med Hyg.* 1998; 58:495–500. [PubMed: 9574798]
15. Crosby R, DiClemente RJ, Wingood GM, et al. Predictors of infection with *Trichomonas vaginalis*: A prospective study of low income African-American adolescent females. *Sex Transm Infect.* 2002; 78:360–364. [PubMed: 12407241]
16. Miller M, Liao Y, Gomez AM, et al. Factors associated with the prevalence and incidence of *Trichomonas vaginalis* infection among African-American women in New York City who use drugs. *J Infect Dis.* 2008; 197:503–509. [PubMed: 18275272]
17. Wendel KA, Erbeding EJ, Gaydos CA, et al. *Trichomonas vaginalis* polymerase chain reaction compared with standard diagnostic and therapeutic protocols for detection and treatment of vaginal trichomoniasis. *Clin Infect Dis.* 2002; 35:576–580. [PubMed: 12173132]
18. Seth P, Wingood GM, DiClemente RJ. Exposure to alcohol problems and its association with sexual behaviour and biologically confirmed *Trichomonas vaginalis* among women living with HIV. *Sex Transm Infect.* 2008; 84:390–392. [PubMed: 18524841]
19. Cu-Uvin S, Ko H, Jamieson DJ, et al. Prevalence, incidence, and persistence or recurrence of trichomoniasis among human immunodeficiency virus (HIV)-positive women and among HIV-negative women at high risk for HIV infection. *Clin Infect Dis.* 2002; 34:1406–1411. [PubMed: 11981738]
20. Swygard H, Miller WC, Kaydos-Daniels SC, et al. Targeted screening for *Trichomonas vaginalis* with culture using a two-step method in women presenting for STD evaluation. *Sex Transm Dis.* 2004; 31:659–664. [PubMed: 15502673]

21. Plitt SS, Garfein RS, Gaydos CA, et al. Prevalence and correlates of *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, *Trichomonas vaginalis* infections, and bacterial vaginosis among a cohort of young injection drug users in Baltimore, Maryland. *Sex Transm Dis.* 2005; 32:446–453. [PubMed: 15976603]
22. Van Der Pol B, Williams JA, Orr DP, et al. Prevalence, incidence, natural history, and response to treatment of *Trichomonas vaginalis* infection among adolescent women. *J Infect Dis.* 2005; 192:2039–2044. [PubMed: 16288365]
23. Peterman TA, Tian LH, Metcalf CA, et al. Persistent, undetected *Trichomonas vaginalis* infections? *Clin Infect Dis.* 2009; 48:259–260. [PubMed: 19113985]
24. Klausner JD, Baer JT, Contento KM, et al. Investigation of a suspected outbreak of vaginal trichomoniasis among female inmates. *Sex Transm Dis.* 1999; 26:335–338. [PubMed: 10417021]
25. Sivakumar K, De Silva AH, Roy RB. *Trichomonas vaginalis* infection in a lesbian. *Genitourin Med.* 1989; 65:399–400. [PubMed: 2613220]
26. Kellock D, O'Mahony CP. Sexually acquired metronidazole-resistant trichomoniasis in a lesbian couple. *Genitourin Med.* 1996; 72:60–61. [PubMed: 8655171]
27. Garrow SC, Smith DW, Harnett GB. The diagnosis of chlamydia, gonorrhoea, and trichomonas infections by self obtained low vaginal swabs, in remote northern Australian clinical practice. *Sex Transm Infect.* 2002; 78:278–281. [PubMed: 12181467]
28. Smith KS, Tabrizi SN, Fethers KA, et al. Comparison of conventional testing to polymerase chain reaction in detection of *Trichomonas vaginalis* in indigenous women living in remote areas. *Int J STD AIDS.* 2005; 16:811–815. [PubMed: 16336764]
29. Simpson P, Higgins G, Qiao M, et al. Real-time PCRs for detection of *Trichomonas vaginalis* beta-tubulin and 18S rRNA genes in female genital specimens. *J Med Microbiol.* 2007; 56:772–777. [PubMed: 17510262]
30. Lawing LF, Hedges SR, Schwebke JR. Detection of trichomonosis in vaginal and urine specimens from women by culture and PCR. *J Clin Microbiol.* 2000; 38:3585–3588. [PubMed: 11015368]
31. Shafir SC, Sorvillo FJ. Viability of *Trichomonas vaginalis* in urine: Epidemiologic and clinical implications. *J Clin Microbiol.* 2006; 44:3787–3789. [PubMed: 16928970]

Detection of *Trichomonas vaginalis* by PCR in Urine and Vaginal Swabs From 624 Female Inmates at Two US Federal Prisons, 2001

TABLE 1

Urine	Screening Prison*			Non-Screening Prison [†]			Both Prisons				
	Swab	N	n (%)	Urine	Swab	N	n (%)	Urine	Swab	N	n (%)
Positive	‡	473	27 (5.7)	Positive	‡	142	11 (7.7)	Positive	‡	615	38 (6.2)
‡	Positive	436	35 (8.0)	‡	Positive	79	3 (3.8)	‡	Positive	515	38 (7.4)
Positive or	Positive	480	41 (8.5)	Positive or	Positive	144	12 (8.3)	Positive or	Positive	624	53 (8.5)
Negative	Negative	429	391 (91.1)	Negative	Negative	77	70 (90.9)	Negative	Negative	506	461 (91.1)
Positive	Negative	429	3 (0.7)	Positive	Negative	77	4 (5.2)	Positive	Negative	506	7 (1.4)
Negative	Positive	429	14 (3.3)	Negative	Positive	77	1 (1.3)	Negative	Positive	506	15 (3.0)
Positive	Positive	429	21 (4.9)	Positive	Positive	77	2 (2.6)	Positive	Positive	506	23 (4.6)

* All women were screened for chlamydia and gonorrhea upon entry at the screening prison.

† Women were tested for sexually transmitted infections if they presented with signs or symptoms of infection at the nonscreening prison.

‡ Positive, negative, or not tested.

TABLE 2

Demographic, Incarceration, Substance Use, and Familial Characteristics of Female Inmates at Two US Federal Prisons by *Trichomonas vaginalis* Status, 2001

	All Participants (n = 507 [*])	<i>T. vaginalis</i> -Positive (n = 48 [*])	<i>T. vaginalis</i> -Negative (n = 459 [*])	<i>P</i> [†]
Age (yr, %)				
<25	15.0	10.4	15.4	
25–34	41.7	47.9	41.1	0.53
35	43.3	41.7	43.5	
Race/ethnicity (%)				
Black	41.4	41.7	41.3	
Caucasian	38.8	33.3	39.3	
Hispanic	10.0	12.5	9.8	0.79
Other race/ethnicity	6.6	8.3	6.4	
Mixed race/ethnicity	3.2	4.2	3.1	
Education (%)				
Elementary school or less	7.3	6.2	7.4	
Some high school or high school degree	56.0	52.1	56.5	0.74
Some college or higher	36.6	41.7	36.1	
Marital status (%)				
Never married	42.6	42.6	42.6	
Married	34.6	27.7	35.4	0.40
Divorced, separated or widowed	22.8	29.8	22.0	
Birth outside the continental US (%)	14.4	23.4	13.5	0.07
Household income (%)				
\$15,000	30.4	47.9	28.5	
>\$15,000	52.3	37.5	53.8	0.02
Don't know or missing information	17.4	14.6	17.6	
Full- or part-time employment (%)	66.1	78.3	64.8	0.07
Type of crime committed (%)				
White collar	13.9	13.3	13.9	
Drug or violence-related	82.9	84.4	82.8	0.92
Other	3.2	2.2	3.3	
Median length of incarceration (range, mo)	19.0 (1–181)	18.5 (1–131)	19.0 (1–181)	0.92
Any cigarette smoking in the 12 mo before arrest (%)	65.0	66.0	64.9	0.88
Any drug use in the 12 mo before arrest (%)	70.3	63.8	71.0	0.30
Drugs used in the past 12 mo before arrest (%) [‡]				
Marijuana	74.4	86.7	73.2	0.11
Crack/cocaine	52.1	50.0	52.3	0.81
Heroin	12.2	6.7	12.8	0.56
Methamphetamines	19.9	10.0	20.9	0.15
Other drugs	25.1	13.3	26.2	0.12

	All Participants (n = 507 [*])	<i>T. vaginalis</i> -Positive (n = 48 [*])	<i>T. vaginalis</i> -Negative (n = 459 [*])	<i>P</i> [†]
Ever been a victim of sexual, physical or emotional abuse by a family member (%)	36.1	45.8	35.1	0.14
Either parent ever had a drug or alcohol addiction (%)	44.2	33.3	45.4	0.11
Either parent ever been incarcerated (%)	20.7	21.3	20.6	0.92

* Total number may vary slightly for each characteristic due to missing responses.

[†] *P* values were calculated by Wilcoxon rank sum tests for continuous variables, and Chi-squared or Fisher exact test tests for binary and categorical variables.

[‡] Among women who used drugs in the 12 months before their arrest.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

TABLE 3

Sexual and Reproductive Characteristics of Female Inmates at Two US Federal Prisons by *Trichomonas vaginalis* Status, 2001

	All Participants (n = 507 [*])	<i>T. vaginalis</i> -Positive (n = 48 [*])	<i>T. vaginalis</i> -Negative (n = 459 [*])	<i>P</i> [†]
Ever had vaginal sexual intercourse with a man (%)				
No	1.4 (1.4) [‡]	0.0 (0.0) [‡]	1.5 (1.5) [‡]	
Yes	64.3 (97.4) [‡]	58.3 (97.9) [‡]	64.9 (97.4) [‡]	0.39 (0.53) [‡]
Missing or inconsistent information	34.3 (1.2) [‡]	41.7 (2.1) [‡]	33.6 (1.1) [‡]	
No. lifetime male sexual partners (%)				
None	1.4 (34.7) [‡]	0.0 (39.6) [‡]	1.5 (34.2) [‡]	
1–4	28.6 (28.6) [‡]	29.2 (29.2) [‡]	28.5 (28.5) [‡]	0.51 (0.69) [‡]
5	33.9 (33.9) [‡]	27.1 (27.1) [‡]	34.6 (34.6) [‡]	
Missing or inconsistent information	36.1 (2.8) [‡]	43.8 (4.2) [‡]	35.3 (2.6) [‡]	
No. male sexual partners in the 3 mo before arrest (%)				
None	9.7 (15.4) [‡]	8.3 (18.8) [‡]	9.8 (15.0) [‡]	
1	32.2 (54.6) [‡]	33.3 (52.1) [‡]	32.0 (54.9) [‡]	0.82 (0.55) [‡]
2	21.1 (29.0) [‡]	16.7 (27.1) [‡]	21.6 (29.2) [‡]	
Missing or inconsistent information	37.1 (1.0) [‡]	41.7 (2.1) [‡]	36.6 (0.9) [‡]	
Ever had a non-regular (short-term or one time) male sexual partner (%)				
No	34.3 (34.3) [‡]	35.4 (35.4) [‡]	34.2 (34.2) [‡]	
Yes	34.5 (61.9) [‡]	35.4 (60.4) [‡]	34.4 (62.1) [‡]	0.95 (0.97) [‡]
Missing or inconsistent information	31.2 (3.8) [‡]	29.2 (4.2) [‡]	31.4 (3.7) [‡]	
Ever had vaginal sexual intercourse with a woman (%)				
No	66.9 (68.0) [‡]	72.9 (70.8) [‡]	66.2 (67.8) [‡]	
Yes	8.3 (28.0) [‡]	6.2 (25.0) [‡]	8.5 (28.3) [‡]	0.64 (0.89) [‡]
Missing or inconsistent information	24.8 (3.9) [‡]	20.8 (4.2) [‡]	25.3 (3.9) [‡]	
Ever exchanged sexual activity for money or drugs (%)	16.8	8.7	17.6	0.12
Ever pregnant (%)	88.0	93.8	87.4	0.20
Birth control in the 12 mo before prison entry (%)				
None	33.9	31.2	34.2	
Condoms	29.3	35.4	28.6	0.62
Other form of birth control	36.8	33.3	37.1	
Pelvic examination before prison entry (%)				
Never or 12 mo	54.2	54.2	54.2	
<12 mo	31.0	37.5	30.3	0.33
Don't know/missing information	14.8	8.3	15.5	
Current symptoms (%)				

	All Participants (n = 507 [*])	<i>T. vaginalis</i> -Positive (n = 48 [*])	<i>T. vaginalis</i> -Negative (n = 459 [*])	<i>P</i> [†]
Abnormal or unusual vaginal discharge	19.1	25.5	18.5	0.24
Vaginal irritation, itch or unusual odor	16.9	14.9	17.1	0.70
Lower abdominal or pelvic pain	18.5	21.3	18.2	0.61
Vaginal bleeding or spotting other than normal period	6.4	12.8	5.7	0.11
Pain during urination	4.8	4.3	4.8	1.00
Current chlamydia or gonorrhea (%)	1.8	4.2	1.5	0.21

* Total number may vary slightly for each characteristic due to missing responses.

[†] *P* values were calculated by chi squared or Fisher exact test tests.

[‡] Based on original participant responses without taking into consideration consistency of responses across related questions.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript