



Published in final edited form as:

Biochem Pharmacol. 2016 February 15; 102: 141.

Retraction notice to “Triptolide, histone acetyltransferase inhibitor, suppresses growth and chemosensitizes leukemic cells through inhibition of gene expression regulated by TNF-TNFR1-TRADD-TRAF2-NIK-TAK1-IKK pathway” [*Biochem. Pharmacol.* 82 (2011) 1134–1144]

Byoungduck Park^a, Bokyoung Sung^a, Vivek R. Yadav^a, Madan M. Chaturvedi^b, and Bharat B. Aggarwal^{a,*}

^aCytokine Research Laboratory, Department of Experimental Therapeutics, The University of Texas MD Anderson Cancer Center, Houston, TX, United States

^bLaboratory for Chromatin Biology, Department of Zoology, University of Delhi, Delhi, India

This article has been retracted: please see Elsevier Policy on Article Withdrawal (<http://www.elsevier.com/locate/withdrawalpolicy>).

This article has been retracted at the request of the Editor.

Fig. 4F, labeled Triptolide plus TNF in wild type mouse embryonic fibroblasts (Trp + TNF/MEF WT) was reused to represent TNF alone in knockout mouse embryonic fibroblasts (TNF/MEF p65^{-/-}). The image has been manipulated in contrast, brightness and size to represent the different experimental conditions.

The article has been retracted because the data integrity has become questionable.

*Corresponding author. Tel.: +1 713 794 1817; fax: +1 713 745 6339. aggarwal@mdanderson.org (B.B. Aggarwal).