

Erratum to: Choosing Wisely: Prevalence and Correlates of Low-Value Health Care Services in the United States

Carrie H. Colla, Ph.D.^{1,2}, Nancy E. Morden, M.D., M.P.H.^{1,2,3}, Thomas D. Sequist, M.D., M.P.H.^{4,5,6}, William L. Schpero, M.P.H.¹, and Meredith B. Rosenthal, Ph.D.⁷

¹The Dartmouth Institute for Health Policy & Clinical Practice, Geisel School of Medicine at Dartmouth, Lebanon, NH, USA; ²Norris Cotton Cancer Center, Dartmouth-Hitchcock Medical Center, Lebanon, NH, USA; ³Department of Community and Family Medicine, Geisel School of Medicine at Dartmouth, Lebanon, NH, USA; ⁴Department of Health Care Policy, Harvard Medical School, Boston, MA, USA; ⁵Division of General Internal Medicine, Brigham and Women's Hospital, Boston, MA, USA; ⁶Partners HealthCare, Boston, MA, USA; ⁷Department of Health Policy and Management, Harvard School of Public Health, Boston, MA, USA.

J Gen Intern Med 31(4):450

DOI: 10.1007/s11606-015-3420-5

© Society of General Internal Medicine 2015

Erratum to: J Gen Intern Med DOI 10.1007/s11606-014-3070-z

On page 225 of the initial publication, the results for cervical cancer screening in Table 2 mistakenly excluded HCPCS codes G0123, G0124, G0141, G0143–145, G0147, G0148, P3000, P3001 and Q0091. After incorporating these additional codes, the mean annual prevalence of cervical cancer screening increased to 9.8% from the reported 3.1%. The coefficients in Table 3 and the categorization of HRRs were not meaningfully affected. We regret this error.

Table 2 updates: Low-value diagnostic services: Annual Cervical Cancer Screening: Mean: 9.8%, Affected National Population (Millions): 14.3 Min/Max: 4.3–18.6%, IQR: 8.1%–11.0%, COV: 0.24, Annual Waste: \$23.4M

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11606-014-3070-z>.

Published online July 3, 2015