



Published in final edited form as:

*Angew Chem Int Ed Engl.* 2013 August 26; 52(35): 9153–9156. doi:10.1002/anie.201303994.

## Nickel-Catalyzed Enantioselective Arylation of Pyridinium Ions: Harnessing an Iminium Ion Activation Mode\*\*

Stephen T. Chau, J. Patrick Lutz, Kevin Wu, and Prof. Abigail G. Doyle

Department of Chemistry, Princeton University, Washington Rd, Princeton, NJ 08544-1009 (USA), Homepage: <http://www.princeton.edu/~doylegrp/>

Abigail G. Doyle: [agdoyle@princeton.edu](mailto:agdoyle@princeton.edu)

### Keywords

asymmetric catalysis; cross-coupling; Negishi reaction; nickel; nitrogen heterocycles

The ability to engage cationic substrates such as iminium and oxocarbenium ions in transition-metal-catalyzed C–C bond formation is a challenging task that has recently seen successful application.<sup>[1]</sup> In this context, we reported that a low-valent Ni catalyst facilitates unprecedented Suzuki–Miyaura cross-coupling reactions with allylic *N,O*- and *O,O*-acetal substrates.<sup>[2,3]</sup> Mechanistic studies revealed that boronic acids mediate allylic C–O activation, with oxidative addition occurring between the resulting iminium or oxocarbenium ion intermediate and the Ni catalyst (Scheme 1a).<sup>[4]</sup> The demonstration that a transition-metal catalyst can oxidatively insert into these prochiral intermediates offers a number of exciting possibilities for reaction design and enantioselective synthesis. Herein, we demonstrate that this activation mode enables the enantioselective synthesis of  $\alpha$ -substituted 2,3-dihydro-4-pyridones by Negishi cross-coupling with *N*-acyl pyridinium ions (Scheme 1b).

$\alpha$ -Substituted piperidines are among the most prevalent scaffolds in biologically active small molecules, and also serve as building blocks for natural product and pharmaceutical synthesis.<sup>[5]</sup> The stereoselective addition of carbon-centered nucleophiles to activated pyridines is a particularly attractive route to these core structures.<sup>[6]</sup> Seminal work from the groups of Comins and Charette established the possibility of using stoichiometric chiral acylating agents to control the stereo-chemical outcome of C–C bond formation.<sup>[7,8]</sup> However, only two examples have been described for the catalytic enantio-selective addition of organometallic reagents to prochiral *N*-acyl/alkyl pyridinium ions.<sup>[9]</sup> Both of these reactions likely proceed through the addition of a chiral [M]-R species into a prochiral pyridinium salt (M = Cu or Rh). As such, the strategy is limited to reactions with either highly nucleophilic R groups or highly electrophilic pyridinium ions. As the oxidative

\*\*We thank Phil Jeffrey for X-ray crystallographic structure determination of **2b** and **3**. Financial support provided by Princeton University and Boehringer Ingelheim is gratefully acknowledged. A.G.D. is an Alfred P. Sloan Foundation Fellow, an Eli Lilly Grantee, an Amgen Young Investigator, and a Roche Early Excellence in Chemistry Awardee.

Correspondence to: Abigail G. Doyle, [agdoyle@princeton.edu](mailto:agdoyle@princeton.edu).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201303994>.

addition/reductive elimination mechanism that we elucidated for cross-coupling with *N*-acyl quinolinium salts should not be subject to these same limitations, we envisioned that its application to enantioselective cross-coupling with pyridinium ions could provide a complementary approach to these methods. Furthermore, successful development of such a reaction would demonstrate for the first time that iminium ion activation by Ni<sup>0</sup> is subject to highly enantioselective C–C bond formation.


One challenge apparent at the outset of our endeavors was potential catalyst poisoning in the presence of free pyridine. To address this issue, we chose to use 4-methoxy-pyridine as a substrate, because it shows substantial formation of a pyridinium salt with chloroformates at –78°C.<sup>[10]</sup> To enable facile transmetalation at low temperature, a Negishi reaction platform was selected.<sup>[11]</sup> Notably, when we initiated the reaction at –78°C with warming to RT, the combination of [((methallyl)NiCl)<sub>2</sub>] (7.5 mol%) and (*R*)-Monophos (**L1**; 18 mol%) was found to promote arylation of 4-methoxypyridine in the presence of phenyl chloroformate and 4-FC<sub>6</sub>H<sub>4</sub>ZnBr with low but measurable *ee* (Table 1, entry 1).<sup>[12]</sup> Commercial phosphoramidite ligands **L2–L4**, which bear substituents at the 3 and 3' positions of the binaphthyl backbone, induced more promising levels of asymmetric induction (up to 91% *ee*, entries 2–4).<sup>[13]</sup> Accordingly, an extensive library of 3,3'-substituted ligands was prepared and evaluated, revealing that ligand **L7** was optimal (entry 7). With this ligand, the 2,3-dihydro-4-pyridone product **2a** was obtained in 95% *ee*, albeit with moderate reaction efficiency.

Upon selection of an optimal ligand (**L7**), we pursued further optimization of the reaction parameters. Among the Ni sources examined, NiBr<sub>2</sub>-diglyme was most promising, as its use increased the yield of the reaction while maintaining high levels of enantioselectivity (Table 2, entry 2). This result is particularly attractive because NiBr<sub>2</sub>-diglyme is air stable, thus permitting the reactions to be set up on the benchtop. In the absence of a Ni source, the reaction performs poorly, demonstrating that Ni induces substantial rate acceleration (entry 5). Additional changes to the reaction conditions, such as decreasing the amount of nucleophile or changing the identity of the acylating agent, had a negative effect on reactivity (entries 6–8). On the other hand, decreasing the Ni loading from 15 mol% to 10 mol% provided nearly identical levels of enantioselectivity and reaction efficiency (entry 9). A gram-scale reaction set up on the benchtop using a catalyst loading of 5 mol% delivered **2a** in 73% yield and 93% *ee* (entry 10).


Having identified conditions to prepare **2a** in high yield and enantiomeric excess, a survey of aryl zinc nucleophiles was conducted to evaluate the scope of the reaction (Scheme 2).<sup>[14]</sup> Phenyl zinc bromide afforded 2,3-dihydro-4-pyridone **2b** in 92% *ee*, which could be recrystallized to >99% *ee* in 78% yield. Generally, *ortho*-, *meta*-, and *para*-substituted zinc nucleophiles are well tolerated. Reactions with electron-withdrawing zinc reagents fared exceptionally well, delivering products bearing 3,5-difluoro (**2c**), 4-Cl (**2d**), 4-CF<sub>3</sub> (**2e**), 4-cyano (**2f**), and 4-sulfonamide (**2g**) groups in 96–99% *ee*. Notably, zinc reagents substituted with methyl ester and pivaloate groups afforded products (**2j**, **2k**) that are otherwise inaccessible by standard Grignard methods. Moreover, a reaction with a heteroaromatic nucleophile furnished **2l** in excellent *ee* and yield, highlighting the potential of this strategy to deliver medically relevant products. Whereas electron-neutral aryl zinc nucleophiles

such as 4-vinyl (**2m**), 4-Me (**2n**) and 2-naphthyl (**2o**) performed modestly in terms of enantioselectivity, electron-rich nucleophiles such as 4-OMe (**2p**) underwent reaction with no stereoselection owing to a competitive racemic background reaction (77% yield of isolated product without nickel).<sup>[15]</sup> An additional limitation is that other electrophile partners, including pyridine, 2-methoxy-pyridine, and 4-dimethylaminopyridine (DMAP), are not competent under the optimized conditions.

To understand the mechanism of this transformation, we studied the stoichiometric reaction of Ni<sup>0</sup> with the pyridinium salt derived from 4-methoxypyridine and phenyl chloroformate, which was generated in situ. For simplicity, we chose to conduct our studies with PPh<sub>3</sub>, as it is a competent ligand for the reaction of interest, providing (±)-**2a** in 87% yield under otherwise standard reaction conditions.<sup>[13]</sup> In the event, an air-sensitive allyl-Ni<sup>II</sup> adduct **3** was produced in 87% yield (Scheme 3), the structure of which was confirmed by single crystal X-ray diffraction (Figure 1). When subjected to 4-FC<sub>6</sub>H<sub>4</sub>ZnBr, allyl adduct **3** underwent C–C bond formation, providing **2a** in 25% yield [Eq. (1)]. Furthermore, **3** is catalytically competent, providing racemic **2a** in 68% yield at 10 mol% catalyst loading [Eq. (2)]. Taken together, these data provide compelling evidence for a redox mechanism distinct from that typically considered for transition-metal-catalyzed addition reactions to pyridinium ions.



(1)



(2)

With these data in mind, we propose the reaction mechanism shown in Scheme 4. Oxidative addition of Ni<sup>0</sup> into the C–N π bond of the pyridinium salt provides a Ni<sup>II</sup> allyl intermediate analogous to complex **3**.<sup>[16]</sup> Subsequent transmetalation with ArZnBr gives diorganonickel intermediate **4**, which can undergo reductive elimination to regenerate the Ni<sup>0</sup> catalyst and complete the catalytic cycle.<sup>[17]</sup> The presence of the methoxy substituent at C4 presumably serves as a blocking group, favoring the observed regioisomeric dihydropyridine **5**.<sup>[18]</sup> Acid hydrolysis of **5** then affords the 2,3-dihydro-4-pyridone product **2**.

In conclusion, we have described a novel nickel-catalyzed enantioselective Negishi cross-coupling reaction of 4-methoxypyridinium salts. A broad range of synthetically valuable enantioenriched 2,3-dihydro-4-pyridones can be obtained from commercially available or readily prepared starting materials and an air-stable, inexpensive Ni<sup>II</sup> precatalyst. Preliminary mechanistic data support the intermediacy of a Ni<sup>II</sup>  $\pi$ -allyl intermediate generated upon ionic oxidative addition of Ni<sup>0</sup> to the pyridinium electrophile. The study provides an exciting indication of the generality of this mode of activation and its amenability to asymmetric catalysis.

## Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

## References

1. For a review, see: Beller M, Eckert M. *Angew Chem.* 2000; 112:1026–1044. *Angew Chem Int Ed.* 2000; 39:1010–1027. for selected examples, see: Lu Y, Arndtsen BA. *Org Lett.* 2007; 9:4395–4397. [PubMed: 17887770] Davis JL, Dhawan R, Arndtsen BA. *Angew Chem.* 2004; 116:600–604. *Angew Chem Int Ed.* 2004; 43:590–594; Beveridge RE, Black DA, Arndtsen BA. *Eur J Org Chem.* 2010:3650–3656. Shack-lady-McAtee DM, Dasgupta S, Watson MP. *Org Lett.* 2011; 13:3490–3493; [PubMed: 21644568] Maity P, Srinivas HD, Watson MP. *J Am Chem Soc.* 2011; 133:17142–17145. [PubMed: 21988178] Gong H, Sinisi R, Gagné MR. *J Am Chem Soc.* 2007; 129:1908–1909. [PubMed: 17261000]
2. Graham TJA, Shields JD, Doyle AG. *Chem Sci.* 2011; 2:980–985.
3. Graham TJA, Doyle AG. *Org Lett.* 2012; 14:1616–1619. [PubMed: 22385385]
4. Sylvester KT, Wu K, Doyle AG. *J Am Chem Soc.* 2012; 134:16967–16970. [PubMed: 23030789]
5. For some recent examples of the use of 4-piperidones in the synthesis of biologically active 4-piperidines, see: Tawara JN, Lorenz P, Stermitz FR. *J Nat Prod.* 1999; 62:321–323. [PubMed: 10075774] Watson PS, Jiang B, Scott B. *Org Lett.* 2000; 2:3679–3681. [PubMed: 11073674] Brooks CA, Comins DL. *Tetrahedron Lett.* 2000; 41:3551–3553.
6. For a recent review, see: Bull JA, Mousseau JJ, Pelletier G, Charette AB. *Chem Rev.* 2012; 112:2642–2713. [PubMed: 22352938]
7. a) Comins DL, Goehring RR, Joseph SP, O'Connor S. *J Org Chem.* 1990; 55:2574–2576. b) Comins DL, Joseph SP, Goehring RR. *J Am Chem Soc.* 1994; 116:4719–4728. c) Comins DL, Kuethe JT, Hong H, Lakner FJ, Concolino TE, Rheingold AL. *J Am Chem Soc.* 1999; 121:2651–2652.
8. a) Legault C, Charette AB. *J Am Chem Soc.* 2003; 125:6360–6361. [PubMed: 12785763] b) Charette AB, Grenon M, Lemire A, Pourashraf M, Martel J. *J Am Chem Soc.* 2001; 123:11829–11830. [PubMed: 11716753] c) Barbe G, Pelletier G, Charette AB. *Org Lett.* 2009; 11:3398–3401. [PubMed: 19719187]
9. Fernández-Ibáñez MÁ, Maciá B, Pizzuti MG, Minnaard AJ, Feringa BL. *Angew Chem.* 2009; 121:9503–9505. *Angew Chem Int Ed.* 2009; 48:9339–9341; Nadeau C, Aly S, Belyk K. *J Am Chem Soc.* 2011; 133:2878–2880. [PubMed: 21309573] for the cross-coupling of pyridinium ions with copper acetylides generated in situ, see: Sun Z, Yu S, Ding Z, Ma D. *J Am Chem Soc.* 2007;


129:9300–9301. [PubMed: 17625864] Black DA, Beveridge RE, Arndtsen BA. *J Org Chem*. 2008; 73:1906–1910. [PubMed: 18215063]

10. Pabel J, Hçsl CE, Maurus M, Ege M, Wanner KT. *J Org Chem*. 2000; 65:9272–9275. [PubMed: 11149888]
11. a) Knochel P, Singer RD. *Chem Rev*. 1993; 93:2117–2188. b) Negishi, E-i; Liu, F. *Metal-Catalyzed Cross-Coupling Reactions*. Diederich, F.; Stang, P.J., editors. Wiley-VCH; Weinheim: 1998. p. 1-47.
12. For the synthesis of  $[(\text{methallyl})\text{NiCl}]_2$ , see: Shim CB, Kim YH, Lee BY, Dong Y, Yun H. *Organometallics*. 2003; 22:4272–4280. Smith CR, Zhang A, Mans DJ, Rajan-Babu TV. *Org Synth*. 2008; 85:248–266. [PubMed: 19672483]
13.  $\text{PPh}_3$  shares similar electronic properties to the phosphoramidite class of ligands; see: Teichert JF, Feringa BL. *Angew Chem*. 2010; 122:2538–2582. *Angew Chem Int Ed*. 2010; 49:2486–2528.
14. For the preparation of organozinc reagents, see: Jensen AE, Kneisel F, Knochel P. *Org Synth*. 2004 Coll. 10:391.
15. Alkyl zinc reagents also provided minimal enantioselectivity, owing to high rates of background reaction.
16. For a Ni-catalyzed vinylation of enones that proceeds through  $\text{Ni}^{\text{II}}$   $\pi$ -allyl intermediates resulting from oxidative addition to an activated electrophilic  $\pi$  system, see: Grisso BR, Johnson JR, Mackenzie PB. *J Am Chem Soc*. 1992; 114:5160–5165.; for the formation of  $\eta^3$ -1-methoxyallyl  $\text{Pd}^{\text{II}}$  and  $\text{Pt}^{\text{II}}$  complexes, see: Morita M, Inoue K, Ogoshi S, Kurosawa H. *Organometallics*. 2003; 22:5468–5472.
17. Whereas 4 may undergo  $\eta^3$  to  $\eta^1$  isomerization before reductive elimination, reductive elimination can occur directly from  $\text{Ni}^{\text{II}}$   $\pi$ -allyl complexes supported by one  $\text{PPh}_3$  ligand; see: Kurosawa H, Ohnishi H, Emoto M, Kawasaki Y, Murai S. *J Am Chem Soc*. 1988; 110:6272–6273. [PubMed: 22148827] Kurosawa H, Ohnishi H, Emoto M, Chatani N, Kawasaki Y, Murai S, Ikeda I. *Organometallics*. 1990; 9:3038–3042.
18. For a discussion of factors influencing the regioselectivity of nucleophilic addition to metal allyl complexes, see: Consiglio G, Waymouth RM. *Chem Rev*. 1989; 89:257–276.




**Figure 1.** Solid-state structure of allyl Ni<sup>II</sup> complex **3**. Ellipsoids set at 30% probability. Hydrogen atoms omitted for clarity.


a) Prior work: Elucidation of an ionic oxidative addition pathway



b) This work: Application to asymmetric cross coupling with pyridinium ions




**Scheme 1.**  
Nickel-catalyzed cross-coupling with iminium ions.


**Scheme 2.**

Scope of aryl zinc nucleophiles in the reaction. Yields and *ee* values are the average of two runs on a 0.5 mmol scale. [a] Yield and *ee* after recrystallization from ether. [b] Reaction warmed to RT. [c] Reaction warmed to -20°C. [d] Derived from an ArZnCl solution (0.3M in THF). [e] Reaction warmed to -50°C.






**Scheme 3.**  
Stoichiometric reaction between  $\text{Ni}^0$  and a pyridinium ion.



**Scheme 4.**  
Proposed catalytic cycle.

Table 1

Ligand optimization.<sup>[a]</sup>

Entry	Ligand	R	Yield [%] <sup>[b]</sup>	ee [%] <sup>[c]</sup>
1	L1	H	9	18
2	L2 <sup>[d]</sup>	Me	11	-40
3	L3	Ph	8	79
4	L4	OMe	52	91
5	L5	OEt	54	92
6	L6	O <i>i</i> Pr	61	96
7	L7	O <i>n</i> Pr	64	95
8	L8	O <i>i</i> Bu	57	96
9	L9	-	8	-40

<sup>[a]</sup> Reactions run on a 0.05 mmol scale; [(methallyl)NiCl]<sub>2</sub> (7.5 mol%), ligand (18 mol%), 4-FC<sub>6</sub>H<sub>4</sub>ZnBr (4 equiv; 0.3 M solution in THF), and phenyl chloroformate (4 equiv).

<sup>[b]</sup> Determined by <sup>19</sup>F NMR spectroscopy with 1,4-difluorobenzene as a quantitative internal standard.

<sup>[c]</sup> Determined by HPLC analysis on a chiral stationary phase.

<sup>[d]</sup> L2 was derived from (*S*)-1,1'-bi-2-naphthol.





Table 2

Optimization of reaction conditions.<sup>[a]</sup>

Entry	Deviation from standard conditions	Yield [%] <sup>[b]</sup>	ee [%] <sup>[c]</sup>
1	none (standard conditions)	64	95
2	NiBr <sub>2</sub> ·diglyme (15 mol%)	76	95
3	NiCl <sub>2</sub> ·glyme (15 mol%)	51	89
4	[Ni(cod) <sub>2</sub> ] (15 mol%)	32	72
5	No nickel or ligand	23	0
6	4-FC <sub>6</sub> H <sub>4</sub> ZnBr (2 equiv)	21	91
7	benzyl chloroformate	33	90
8	ethyl chloroformate	21	73
9	NiBr <sub>2</sub> ·diglyme (10 mol%), <b>L7</b> (12 mol%)	88 <sup>[d]</sup>	93
10	NiBr <sub>2</sub> ·diglyme (5 mol%), <b>L7</b> (6 mol%)	73 <sup>[e]</sup>	93

<sup>[a]</sup> Reactions run on a 0.05 mmol scale; [(methallyl)NiCl]<sub>2</sub> (7.5 mol%), ligand (**L7**; 18 mol%), 4-FC<sub>6</sub>H<sub>4</sub>ZnBr (4 equiv; 0.3 M solution in THF), and phenyl chloroformate (4 equiv).

<sup>[b]</sup> Determined by <sup>19</sup>F NMR spectroscopy with 1,4-difluorobenzene as a quantitative internal standard.

<sup>[c]</sup> Determined by HPLC analysis on a chiral stationary phase.

<sup>[d]</sup> Yield of isolated product on a 0.5 mmol scale.

<sup>[e]</sup> Yield of isolated product on a 5 mmol scale with the reaction warmed to -40°C. cod = 1,5-cyclo-octadiene.