

## ORIGINAL ARTICLE

# Multiple endosymbiont infections and reproductive manipulations in a linyphiid spider population

MM Curry<sup>1</sup>, LV Paliulis<sup>2</sup>, KD Welch<sup>1,3</sup>, JD Harwood<sup>1</sup> and JA White<sup>1</sup>

In many arthropods, maternally inherited endosymbiotic bacteria can increase infection frequency by manipulating host reproduction. Multiple infections of different bacteria in a single host population are common, yet few studies have documented concurrent endosymbiont phenotypes or explored their potential interactions. We hypothesized that spiders might be a particularly useful taxon for investigating endosymbiont interactions, because they are host to a plethora of endosymbiotic bacteria and frequently exhibit multiple infections. We established two matriline from the same population of the linyphiid spider *Mermessus fradeorum* and then used antibiotic curing and controlled mating assays to demonstrate that each matriline was subject to a distinct endosymbiotic reproductive manipulation. One matriline was co-infected with *Rickettsia* and *Wolbachia* and produced offspring with a radical female bias. Antibiotic treatment eliminated both endosymbionts and restored an even sex ratio to subsequent generations. Chromosomal and fecundity observations suggest a feminization mechanism. In the other matriline, a separate factorial mating assay of cured and infected spiders demonstrated strong cytoplasmic incompatibility (CI) induced by a different strain of *Wolbachia*. However, males with this *Wolbachia* induced only mild CI when mated with the *Rickettsia*–*Wolbachia* females. In a subsequent survey of a field population of *M. fradeorum*, we detected these same three endosymbionts infecting 55% of the spiders in almost all possible combinations, with nearly half of the infected spiders exhibiting multiple infection. Our results suggest that a dynamic network of endosymbionts may interact both within multiply infected hosts and within a population subject to multiple strong reproductive manipulations.

*Heredity* (2015) **115**, 146–152; doi:10.1038/hdy.2015.2; published online 22 April 2015

## INTRODUCTION

Maternally inherited endosymbiotic bacteria represent independent genetic elements within the cells and tissues of their arthropod hosts. Unlike virulent pathogens, the fate of these endosymbionts is wedded to their host's survival and reproduction (Engelstädter and Hurst, 2009), yet the interests of host and endosymbiont may not align. In particular, host and endosymbiont may come into conflict over offspring sex, as males are evolutionary dead ends for maternally inherited endosymbionts (O'Neill *et al.*, 1997; Werren *et al.*, 2008; Engelstädter and Hurst, 2009). Endosymbionts use a variety of mechanisms to manipulate host reproduction to promote the spread of infection within a host population, even if host fitness is reduced (Hoffmann and Turelli, 1997). The most common reproductive manipulation is cytoplasmic incompatibility (CI), which modifies the sperm of infected males such that it cannot produce a viable embryo unless the same or a closely related endosymbiont in the egg rescues sperm functionality (Werren *et al.*, 2008). Alternatively, endosymbionts can directly skew the sex ratio of hosts by inducing parthenogenesis (Huigens and Stouthamer, 2003), killing males (Hurst and Jiggins, 2000) or feminizing genetic males (Negri *et al.*, 2006).

Endosymbionts are very common; >30% of arthropods are infected with *Wolbachia* or other endosymbionts (for example, *Rickettsia*, *Cardinium*) known to cause reproductive anomalies (Duron *et al.*, 2008a). The extent of endosymbiont infection is likely

underestimated, as low frequency infections are easily missed in broad taxa screening studies (Hilgenboecker *et al.*, 2008) and low titer infections often go undetected (Hughes *et al.*, 2011). Rampant and assorted endosymbiont infections in arthropods suggest that endosymbionts employing different manipulative strategies will likely interact within the same host population or even the same host body. Although multiple endosymbiont infections have been documented across numerous host taxa (for example, Riegler and Stauffer, 2002; Duron *et al.*, 2010), relatively few studies have empirically determined the phenotypic effects of potentially interacting endosymbionts on their hosts (Hiroki *et al.*, 2004; Charlat *et al.*, 2006; White *et al.*, 2009; White *et al.*, 2011; Zhu *et al.*, 2012).

We suggest that spiders may be a particularly rich host taxon for investigating endosymbiont interactions. Though most endosymbiont investigation has focused on insects, recent surveys demonstrate that spiders harbor a wealth of endosymbionts (Goodacre *et al.*, 2006; Baldo *et al.*, 2008; Duron *et al.*, 2008b; Perlman *et al.*, 2010; Yun *et al.*, 2011; Stefanini and Duron, 2012), with multiple endosymbionts co-occurring within the same population or co-infecting the same individual (Goodacre, 2011; Vanthournout *et al.*, 2014). However, the functional role of these spider endosymbionts is not yet clearly defined. To date, only one reproductive manipulation has been reported in spiders: female bias thought to be male killing in *Oedothorax gibbosus* and *O. retusus* (Vanthournout *et al.*, 2011, 2014). However, two novel

<sup>1</sup>Department of Entomology, University of Kentucky, Lexington, KY, USA and <sup>2</sup>Department of Biology, Bucknell University, Lewisburg, PA, USA

Correspondence: M Curry, Department of Entomology, University of Kentucky, Lexington, KY 40546, USA.

E-mail: megmay111@gmail.com

<sup>3</sup>Current address: USDA-ARS, North Central Agricultural Research Laboratory, 2923 Medary Avenue, Brookings, SD 57006, USA.

Received 27 June 2014; revised 6 December 2014; accepted 19 December 2014; published online 22 April 2015

behavioral alterations have also been associated with facultative bacterial symbionts in spiders (Goodacre *et al.*, 2009; Gunnarsson *et al.*, 2009), implying that endosymbionts may have a different or expanded suite of manipulative tools in spiders than their insect relatives (Goodacre, 2011).

To better understand what endosymbiont interactions may arise in a diverse endosymbiont community, we characterized the phenotypic function of different infection types in a population of the linyphiid spider, *Mermessus fradeorum*. We established two infection types of *M. fradeorum* in the laboratory, cured each infection type with antibiotics and determined endosymbiont effects on sex ratio and fecundity by comparing infected and uninfected spiders. Endosymbiont effects on genetic sex were determined via chromosomal observation. We then conducted mating assays to test for CI. Finally, we determined the natural frequency of endosymbiont infection in a field population of *M. fradeorum* to evaluate the potential for interaction between endosymbionts within the host body and among differentially infected hosts in the same population.

## MATERIALS AND METHODS

### Establishment of laboratory matriline and determination of endosymbiont infection status

*M. fradeorum* is a small (average body length of 2.5 mm) sheet-weaving linyphiid spider common in Kentucky agroecosystems and generally found in agricultural fields and grasslands throughout North America (Welch, 2013). To investigate endosymbiont phenotypes in the laboratory, gravid *M. fradeorum* females were hand collected in February of 2012 from winter wheat at the Spindletop Research Farm in Lexington, KY, USA (38.1272°N, 84.5081°W). Each spider was placed in an individual plastic rearing cup (4.6 cm × 7.3 cm diameter) with a plaster base and imbedded plastic straws for web attachment, in a 24 h dark incubator with 98% relative humidity and a 16 h:8 h thermal cycle of 28 °C:18 °C. Egg masses were removed following deposition, and upon hatching, individual spiderlings were transferred to separate rearing cups. Spiders were fed twice a week: juveniles received *Proisotoma minuta* (Collembola), whereas adults received a combination of *P. minuta*, *Sinella curviseta* (Collembola) and *Drosophila melanogaster* (Diptera).

After egg mass deposition, mother spiders were screened for three endosymbionts common in spiders: *Wolbachia*, *Rickettsia*, and *Cardinium*. Prior to DNA extraction, spiders were surface sterilized with sequential 10 s rinses of ethanol, deionized water, 5% bleach solution and deionized water (adapted from Medina *et al.*, 2011). DNA was extracted using a Chelex extraction protocol adapted from White *et al.* (2009): abdomens of adults were pulverized in 4 µl of Proteinase K, incubated in a 20% Chelex bead solution for 60 min at 37 °C and then heat shocked for 10 min at 95 °C to deactivate Proteinase K. Extraction quality was assessed by measuring DNA concentration using a Nanodrop 2000 spectrophotometer (Thermo Scientific, Wilmington, DE, USA) and by PCR using universal eukaryotic primers NSF 18/19 or COI (Supplementary Table S1). To detect the endosymbionts, we used taxon-specific primers and protocols described in Supplementary Table S1 for segments of 16S rRNA of *Rickettsia* and *Cardinium* and the *Wolbachia* surface protein (*wsp*). PCR products were visualized under ultraviolet light on 1.5% agarose gels treated with Gel-Red (Biotium, Hayward, CA, USA). Endosymbiont identity for all positive PCR results was confirmed via Sanger Sequencing at the University of Kentucky's AGTC sequencing facility. Prey items were also screened for symbionts, none of which were detected in laboratory stocks. Morphological spider identification was confirmed by sequencing the COI barcoding region of each spider lineage (GenBank accession KJ561375).

Offspring of two females were used to establish two matriline of differential endosymbiont infection types: 'R<sub>1</sub>W<sub>1</sub>' bearing a double *Rickettsia*-*Wolbachia* infection, and 'W<sub>2</sub>' singly infected with a different *Wolbachia* strain. Vertical transmission of the endosymbionts was confirmed in each generation with diagnostic PCR using new primers (Ricklong and mfwp), which were designed using the Primer3 tool (Rozen and Skaletsky, 1998) in Geneious 5.6.4 (Drummond *et al.*, 2012) to specifically amplify the *Rickettsia* and *Wolbachia* detected in *M. fradeorum* (Supplementary Table S1). We previously

characterized the microbiomes of each matriline via 454-pyrosequencing of 16S bacterial diversity and did not find indication that other maternally inherited bacterial endosymbionts were present in the laboratory lines (Curry, 2013). Early rearing observation indicated that the R<sub>1</sub>W<sub>1</sub> line produced only female offspring, whereas the W<sub>2</sub> line produced equal numbers of male and female offspring. The R<sub>1</sub>W<sub>1</sub> line was therefore propagated by matings with W<sub>2</sub> line males. To prevent inbreeding depression, additional *M. fradeorum* infected with W<sub>2</sub> were collected from a different Lexington, KY population in 2012 and reared in the laboratory for several generations. Sons from three different field-collected W<sub>2</sub> mothers were introgressed equally into third generation R<sub>1</sub>W<sub>1</sub> and W<sub>2</sub> matriline prior to antibiotic treatment.

Experimentally cured lines were generated for both R<sub>1</sub>W<sub>1</sub> and W<sub>2</sub> matriline to investigate endosymbiont phenotype. To eliminate endosymbiotic bacteria in *M. fradeorum*, second and third instar spiderlings from each matriline were treated with antibiotics previously shown to be effective in spiders (Goodacre *et al.*, 2009; Vanthournout *et al.*, 2011). For 5 consecutive days, spiderlings (13 R<sub>1</sub>W<sub>1</sub> and 25 W<sub>2</sub>) were sprayed with a fine mist of 0.1% ampicillin solution daily. Following 1 week of no treatment, the same spiderlings were sprayed with a 0.1% tetracycline solution daily for 5 days. Treated spiders were mated, and curing was confirmed in the subsequent generation by screening offspring for *Rickettsia* and *Wolbachia*. Antibiotic treatment successfully eliminated all symbiotic bacteria from R<sub>1</sub>W<sub>1</sub> and W<sub>2</sub> lines, establishing R<sub>1</sub>-W<sub>1</sub>- and W<sub>2</sub>- cured lines. Experiments were conducted with spiders at least two generations removed from antibiotic treatment to alleviate any direct negative fitness effects of the antibiotics.

### Endosymbiont effects on sex ratio and fecundity

The effects of endosymbionts in the doubly infected R<sub>1</sub>W<sub>1</sub> line and the singly infected W<sub>2</sub> line were quantified by comparing life history parameters of naturally infected R<sub>1</sub>W<sub>1</sub> and W<sub>2</sub> lines with antibiotically cured R<sub>1</sub>-W<sub>1</sub>- or W<sub>2</sub>- lines. Females of the four lines were mated with cured males to eliminate the potential influence of paternal endosymbiont type (see Table 1 for sample sizes). Females were mated with males of either the R<sub>1</sub>-W<sub>1</sub>- or the W<sub>2</sub>- cured line. Mating was confirmed by visual observation, and males were removed 1 day after mating. Females were allowed to produce egg sacs for 10 days. We preserved all parents in 95% ethanol and stored them at -20 °C and then measured maternal size using patella-tibia length. All mothers and a subset of randomly selected fathers were subsequently screened for *Rickettsia* and *Wolbachia* to verify appropriate infection status.

Females typically deposited two to three egg sacs in total. One egg sac per mother was randomly selected for rearing; the remaining egg sacs were removed and stored in 95% ethanol for embryo counting. Upon hatching, spiderlings were transferred to individual rearing cups, and each egg sac was checked for undeveloped embryos. Offspring were individually reared to adulthood to determine survivorship and sex. Fecundity was calculated as the total number of offspring, hatched and unhatched, in all egg sacs for each mother.

Statistically, we first considered whether paternal lineage (R<sub>1</sub>-W<sub>1</sub>- and W<sub>2</sub>-) influenced any of the response variables. We found no difference in the total production of offspring (one-way analysis of variance  $F=0.043$ , d.f. = 1,42,  $P=0.836$ ), sex ratio (logistic regression  $\chi^2=1.556$ , d.f. = 1,  $P=0.212$ ) or survivorship (logistic regression  $\chi^2=0.525$ , d.f. = 1,  $P=0.217$ ) between R<sub>1</sub>-W<sub>1</sub>- and W<sub>2</sub>- fathers, therefore cured males of both matriline were pooled for subsequent analysis. We also did not find evidence that maternal size should be included in our models as a covariate, as there was no relationship between maternal size (patella-tibia length) and fecundity (regression  $R^2=0.002$ , d.f. = 42,  $P=0.7903$ ). Therefore fecundity per mother was compared among matriline categories (R<sub>1</sub>W<sub>1</sub>, R<sub>1</sub>-W<sub>1</sub>-, W<sub>2</sub>, W<sub>2</sub>-) using a simple one-way analysis of variance in JMP 10 (SAS Institute Inc., 2012) followed by Tukey's honestly significant difference test to separate means. Endosymbiont effect on sex ratio was determined using an exact binomial two-tailed test for goodness of fit (McDonald and Delaware, 2009), testing each category against the expected 1:1 sex ratio. Differences in embryo development (hatched versus unhatched eggs) and survivorship of hatched spiderlings to adulthood among lines were analyzed with logistic regression using Arc 1.06 (Cook and

**Table 1 Mean  $\pm$  s.e. for all life history parameters observed in maternal endosymbiont effects and cytoplasmic incompatibility experiments with *Mermessus fradeorum***

n Mothers	Cross type: (female $\times$ male)	Hatched	Unhatched	Fecundity	Ratio female	Proportion undeveloped	Survivorship of hatched spiderlings
<i>Maternal endosymbiont effects</i>							
12	R <sub>1</sub> W <sub>1</sub> $\times$ uninfected <sup>a</sup>	10.7 $\pm$ 1.1	1.0 $\pm$ 0.8	20.7 $\pm$ 2.4 <sup>b</sup>	1 $\pm$ 0	0.16 $\pm$ 0.11	0.97 $\pm$ 0.01
10	R <sub>1</sub> –W <sub>1</sub> – $\times$ uninfected	11.7 $\pm$ 1.4	0.1 $\pm$ 0.1	21.8 $\pm$ 2.2	0.62 $\pm$ 0.06	0.02 $\pm$ 0.02	0.97 $\pm$ 0.02
7	W <sub>2</sub> $\times$ uninfected	11.9 $\pm$ 0.8	1.2 $\pm$ 0.9	18.6 $\pm$ 3.4	0.54 $\pm$ 0.07	0.12 $\pm$ 0.08	0.97 $\pm$ 0.01
4	W <sub>2</sub> – $\times$ uninfected	12.0 $\pm$ 1.8	0 $\pm$ 0	16.4 $\pm$ 3.6	0.42 $\pm$ 0.17	0 $\pm$ 0	1 $\pm$ 0
<i>CI in W<sub>2</sub> females</i>							
12	W <sub>2</sub> – $\times$ W <sub>2</sub>	7.3 $\pm$ 2.4	14.0 $\pm$ 3.3	21.3 $\pm$ 2.5 <sup>c</sup>	0.53 $\pm$ 0.08	0.63 $\pm$ 0.063	1 $\pm$ 0
15	W <sub>2</sub> $\times$ W <sub>2</sub>	24.9 $\pm$ 1.9	1.5 $\pm$ 1.2	26.4 $\pm$ 1.6	0.54 $\pm$ 0.03	0.05 $\pm$ 0.052	0.87 $\pm$ 0.04
11	W <sub>2</sub> – $\times$ W <sub>2</sub> –	28.1 $\pm$ 1.9	1.0 $\pm$ 0.7	29.1 $\pm$ 1.5	0.40 $\pm$ 0.03	0.04 $\pm$ 0.063	0.94 $\pm$ 0.03
11	W <sub>2</sub> $\times$ W <sub>2</sub> –	27.6 $\pm$ 2.3	0.7 $\pm$ 0.6	28.3 $\pm$ 1.8	0.54 $\pm$ 0.07	0.05 $\pm$ 0.063	0.91 $\pm$ 0.03
<i>CI in R<sub>1</sub>W<sub>1</sub> females</i>							
4	R <sub>1</sub> W <sub>1</sub> $\times$ W <sub>2</sub>	25.5 $\pm$ 5.4	8.0 $\pm$ 4.3	33.5 $\pm$ 1.3 <sup>c</sup>	NA	0.25 $\pm$ 0.15	NA
4	R <sub>1</sub> W <sub>1</sub> $\times$ W <sub>2</sub> –	39.5 $\pm$ 3.8	0.5 $\pm$ 0.3	40.0 $\pm$ 3.7	NA	0.01 $\pm$ 0.01	NA

Abbreviations: CI, cytoplasmic incompatibility; NA, not applicable.

<sup>a</sup>Uninfected males are composed of R<sub>1</sub>–W<sub>1</sub>– and W<sub>2</sub>–.

<sup>b</sup>Fecundity for maternal endosymbiont effects experiment includes embryo counts from egg sacs not allowed to hatch.

<sup>c</sup>Fecundity is the sum of hatched and unhatched embryos; all egg sacs were allowed to hatch.

Weisberg, 1999). When overall significance of the model was established, Wald statistics were used to make comparisons among categories.

### Determination of genetic sex in R<sub>1</sub>W<sub>1</sub> offspring

To determine endosymbiont effect on genetic sex, infected R<sub>1</sub>W<sub>1</sub> eggs and uninfected eggs (both R<sub>1</sub>–W<sub>1</sub>– and W<sub>2</sub>–) at different stages of development were removed from their egg sacs using forceps and separately subjected to the staining protocol below. Eggs were pierced with forceps and soaked in hypotonic solution (0.075 M KCl) for 45 min at room temperature. The chorion was then removed, and egg contents were transferred to a freshly prepared solution of ethanol–acetic acid (3:1) and incubated for 45 min at room temperature. Egg contents were then transferred to a drop of 60% acetic acid on a clean glass slide and spread and dried on a slide drier at 45 °C. Dried slides were stained with either 5% giemsa in Sørensen buffer or aceto-orcein (2.5 g orcein per 100 ml 45% acetic acid) for 45 min at room temperature. Slides were then rinsed three times in distilled water, mounted in glycerol and sealed with nail polish. Slides were observed using a Zeiss Opton inverted microscope with phase contrast (Carl Zeiss Microscopy, Göttingen, Germany), and chromosomes from mitotically dividing cell spreads were counted. Although hundreds of spreads were prepared, spreads in which chromosomes were sufficiently dispersed so the total number could be reliably counted were rare. We counted chromosomes in spreads created from a total of 26 eggs: 12 eggs from 5 different egg sacs laid by uninfected females, and 14 eggs from 4 different egg sacs laid by R<sub>1</sub>W<sub>1</sub> infected females. Eggs were categorized as male or female based on chromosome number, and an exact binomial two-tailed goodness-of-fit test was used to determine whether the sex ratio differed from the expected 1:1 in either infection category (McDonald and Delaware, 2009).

### CI experiments

We tested whether W<sub>2</sub> *Wolbachia* induces CI by conducting a two-way factorial cross, mating W<sub>2</sub> and W<sub>2</sub>–females to W<sub>2</sub> or W<sub>2</sub>–males (see Table 1 for sample sizes). Following mating, mothers and fathers were screened for *Wolbachia*. Matings, husbandry methods and fitness parameters for the CI cross were the same as those used in the previously described experiment, except that all egg sacs were allowed to hatch. One egg sac from each pair was randomly selected for full individual rearing of offspring to adulthood. Emerging spiderlings from the other egg sacs were immediately preserved in ethanol. One day following hatching, egg sacs were preserved in ethanol to prevent embryo desiccation and subsequently dissected to quantify undeveloped embryos. Hatching success was calculated as the number of spiderlings that successfully hatched relative to the total number of embryos produced. Hatching success, spiderling survivorship

to adulthood and sex ratio were compared among the cross types using logistic regression in Arc 1.06 with Williams' correction (Williams, 1982) to address moderate overdispersion in the data when necessary.

We also investigated whether W<sub>2</sub> in males induces CI when mated with R<sub>1</sub>W<sub>1</sub> females; R<sub>1</sub>W<sub>1</sub> females were mated with either W<sub>2</sub> or W<sub>2</sub>– males (see Table 1 for sample sizes). For each mating pair, all conditions were identical to the previous CI cross, with the exception that no spiderlings were reared to adulthood. Hatching success was again analyzed using logistic regression in Arc v. 1.06.

### Endosymbiont infection frequency in *M. fradeorum*

In August 2013, live *M. fradeorum* adults were collected from winter wheat at the same location as the original matriline collections. To eliminate false positives due to residual gut contents (Greenstone *et al.*, 2013), spiders were reared on endosymbiont-free prey for at least 3 days and then starved for at least 2 additional days prior to DNA extraction. Upon morphological identification, specimens were dissected to remove epigynal and palpal vouchers. As described previously, adult spiders were surface sterilized and DNA extracted, and the 44 extractions that passed quality control were screened for *Rickettsia*, *Wolbachia* and *Cardinium* (Supplementary Table S1). Spider species (COI barcodes) and endosymbiont identity for all positive PCR results were confirmed via Sanger sequencing. Because two distinct *Wolbachia* strains were present in the laboratory *M. fradeorum* (W<sub>1</sub> or W<sub>2</sub>), we also amplified and sequenced the *fbpA* gene from each *Wolbachia*-positive field specimen and the laboratory lines. Upon inspection, a number of specimens had double peaks in their *Wolbachia* chromatograms that perfectly corresponded to the single-nucleotide polymorphisms between *Wolbachia* strains W<sub>1</sub> and W<sub>2</sub> (Riegler and Stauffer, 2002). These specimens were categorized as doubly infected with W<sub>1</sub> and W<sub>2</sub>; there was no evidence for additional *Wolbachia* strains. A few of the spiders had deposited eggs before being killed, and offspring of one *Rickettsia*-positive female were individually reared to adulthood to determine the sex ratio of offspring using the same rearing conditions described previously.

## RESULTS

### Endosymbiont strain characterization in laboratory matrilines

Diagnostic PCR and Sanger sequencing confirmed that each of the laboratory-reared matrilines carried a different *Wolbachia* strain and that one matriline also carried *Rickettsia*. R<sub>1</sub>W<sub>1</sub> and W<sub>2</sub> matrilines were reared in the laboratory for nine generations with nearly perfect vertical endosymbiont transmission rates. Transmission of *Rickettsia*

and *Wolbachia* in the  $R_1W_1$  line from mothers to offspring exhibited 100% fidelity (19 mothers, 51 offspring). In the  $W_2$  line, we did not detect maternal *Wolbachia* infection in 2/86 offspring (from 17 mothers), indicating 98% transmission efficiency under laboratory conditions. Sequencing of a 590-bp region of 16S rRNA indicated that the *Rickettsia* present in the  $R_1W_1$  line of *M. fradeorum* (KJ546642) is 99% similar to the *Rickettsia* of the leafhopper *Nephotettix cincticeps* (AB702995). The two *Wolbachia* strains in *M. fradeorum* diverge from one another by 8.8% for the *wsp* gene ( $W_1$ : KJ561379,  $W_2$ : KJ561380) and 2.8% for the *fbpA* gene ( $W_1$ : KJ561362,  $W_2$ : KJ561367). Sequences for both  $W_1$  and  $W_2$  *wsp* and *fbpA* genes were more similar to Supergroup A *Wolbachia* strains than strains from other supergroups.


### Endosymbiont effects on sex ratio and fecundity

The  $R_1W_1$  line offspring were entirely female and significantly differed from a 1:1 sex ratio ( $P < 0.001$ ). In contrast, the antibioticly cured  $R_1 - W_1 -$  line produced male offspring and showed a restoration to the expected 1:1 sex ratio ( $P = 0.064$ ; Figure 1, Table 1). Neither the *Wolbachia*-infected  $W_2$  line ( $P = 0.777$ ) nor its cured  $W_2 -$  counterpart ( $P = 0.618$ ) had sex ratios that differed from 1:1, indicating that this strain of *Wolbachia* does not affect sex ratio (Figure 1, Table 1). The extreme female bias in the  $R_1W_1$  line was observed not only in this experiment but also in nine generations of rearing. Only four males in total were produced out of the 328 non-experimental  $R_1W_1$  line spiders individually reared in the laboratory. Of the four  $R_1W_1$  males produced, all four were reproductively viable and successfully mated with females. Three of these males were screened for endosymbionts, and all three were positive for *Rickettsia* and *Wolbachia*.

No significant difference in fecundity was detected among the four infection types ( $F = 0.618$ , d.f. = 3,39,  $P = 0.608$ ; Table 1) and few undeveloped embryos were found in hatched egg sacs, yielding no significant differences in proportion of undeveloped embryos among groups ( $\chi^2 = 2.98$ , d.f. = 3,  $P = 0.395$ ; Table 1). Therefore, we found no evidence that endosymbiotic bacteria in the  $R_1W_1$  and  $W_2$  mothers affect fecundity or embryo development. Furthermore, survivorship to adulthood for hatched offspring did not significantly differ among infection categories ( $\chi^2 = 2.98$ , d.f. = 3,  $P = 0.395$ ; Table 1).

### Determination of genetic sex in $R_1W_1$ offspring

Counting of chromosomes from testes spreads and observation of chromosomes in living cells suggest that the male number of



**Figure 1** Mean  $\pm$  s.e. proportion female offspring from a controlled mating assay of each maternal infection type mated with uninfected males. Asterisk indicates a sex ratio that is significantly different from 1:1. R = *Rickettsia*, W = *Wolbachia*.

chromosomes in *M. fradeorum* is  $2n = 24$  and the female number is  $2n = 26$ , with a X1X1X2X2 (female)/X1X20 (male) sex determination system, similar to other spiders (Kořínková and Král, 2013). Examination of 12 uninfected eggs revealed 26 chromosomes in spreads from 7 eggs and 24 chromosomes in spreads from 5 eggs; indicating a sex ratio of 1.4:1 which does not differ significantly from 1:1 ( $P = 0.387$ ). Among the 14 eggs from  $R_1W_1$  mothers, 8 eggs had 26 chromosomes and 6 had 24 chromosomes; this sex ratio of 1.3:1 also does not differ significantly from 1:1 ( $P = 0.395$ ). These data suggest that  $R_1W_1$  females produce male eggs (24 chromosomes) but that these males are feminized into morphological females by bacterial endosymbionts in the all-female  $R_1W_1$  line.

### CI experiments

*Wolbachia*-infected  $W_2$  fathers induced CI when mated with cured  $W_2 -$  mothers, resulting in 73% fewer hatched spiderlings than any other mating combination ( $\chi^2 = 38.4$ , d.f. = 3,  $P < 0.001$ ; Table 1, Figure 2a) with no significant differences among the other cross types. Cross type did not influence the sex ratio of offspring ( $\chi^2 = 4.35$ , d.f. = 3,  $P = 0.22$ ; Table 1). Survivorship of hatched offspring to adulthood differed significantly among the four different cross types (Wald  $\chi^2 = 10.12$ , d.f. = 3,  $P = 0.017$ ; Table 1), with greater survival of offspring in the incompatible cross type than when  $W_2$ -infected mothers mated with  $W_2$ -infected fathers. Survivorship of hatched offspring in the other two crosses was intermediate and did not differ significantly from any other crosses. Anecdotally, visual observation of hatching spiderlings indicated that spiderlings from the incompatible cross often emerged later than other crosses, appeared larger and were observed consuming undeveloped embryos.


Though  $W_2$  induced strong CI when  $W_2 -$  females were crossed with  $W_2$  males, CI was relatively weak when  $R_1W_1$  females were crossed with  $W_2$  males. Egg hatch in  $R_1W_1 \times W_2$  matings was only 35% lower than  $R_1W_1 \times W_2 -$  matings ( $\chi^2 = 5.06$ , d.f. = 1,  $P = 0.025$ ; Figure 2b). This cross does not support the strong level of CI observed with the  $W_2 -$  mothers, suggesting that the endosymbionts or other elements present in  $R_1W_1$  mothers may partially rescue the CI induced by  $W_2$ .

### Endosymbiont infection frequency in a field population of *M. fradeorum*


Diagnostic PCR screening indicated that 55% of all field-collected *M. fradeorum* individuals bore an endosymbiont infection (Figure 3). All three endosymbiont strains present in the laboratory populations ( $R_1$ ,  $W_1$ ,  $W_2$ ) were detected in the field-collected *M. fradeorum* in six of the seven possible infection combinations;  $W_1$  was never found as a single infection but was always found as a co-infection with  $R_1$  and/or  $W_2$  (Figure 3). Males were more difficult to collect and represented only 25% of our sample, precluding tests of statistical association between endosymbiont infection and sex ratio. However, males corresponding to all female infection types were found in the field, with the exception of  $R_1W_1$  and  $W_2$ . One  $R_1$  female reproduced in the laboratory prior to being screened and produced male and female offspring. A subsequent F1 mating of  $R_1$  siblings produced 23 male and 23 female offspring under rearing conditions analogous to all other experimental matings.

### DISCUSSION

Multiple bacterially induced reproductive manipulations are simultaneously at work within a single population of the spider *M. fradeorum* (Figure 4). Some individuals, infected with *Rickettsia* and *Wolbachia* ( $R_1W_1$ ), exhibit an almost complete female bias, whereas other



**Figure 2** Mean  $\pm$  s.e. hatched spiderlings (black) and unhatched embryos (white) from experimental matings of (a)  $W_2$  infected and  $W_2^-$  uninfected females and males and (b)  $R_1W_1$  females with either  $W_2$  or  $W_2^-$  males. Asterisks indicate cross types that differed in proportion of hatched embryos from the other treatments in that experiment at  $\alpha=0.05$ . R = *Rickettsia*, W = *Wolbachia*.




**Figure 3** Percentage of specimens positive for three endosymbiont strains detected in the *Mermessus fradeorum* spider survey. Female ( $N=33$ ), male ( $N=11$ ) and total ( $N=44$ ) columns' order from left to right. Multiply infected individuals are indicated by more than one shaded region on the same horizontal plane. R = *Rickettsia* (dark grey), W = *Wolbachia* (light grey).

individuals, infected with a different strain of *Wolbachia* ( $W_2$ ), experience CI. Both CI-inducing (Hoffmann and Turelli, 1997) and female-biasing (Taylor, 1990) endosymbionts can invade and come to dominate uninfected host populations. However, theoretical models suggest that competing endosymbionts may disrupt one another's spread (Engelstädter *et al.*, 2004; Charlat *et al.*, 2006). Indeed, the few previous empirical studies documenting multiple manipulations within a host population have primarily confirmed the dominance of one or the other manipulation (Hiroki *et al.*, 2004; Charlat *et al.*, 2006; Narita *et al.*, 2006). In contrast, the presence of both

manipulations at both time points for *M. fradeorum* (initial laboratory line collection and field survey 2 years later) suggests that both manipulations may persist in this system. Linyphiid spiders such as *M. fradeorum* are often subject to frequent disturbance in agroecosystems, necessitating high rates of dispersal (Weyman *et al.*, 2002). It is possible that high migration rates may promote nonequilibrium coexistence of differentially infected hosts subject to conflicting reproductive manipulations (Engelstädter *et al.*, 2004).

In our field screen of *M. fradeorum*, we found these three endosymbiont strains in almost all possible combinations within hosts, indicating that endosymbiont infections in this host may be relatively dynamic, through imperfect vertical transmission and/or horizontal transmission of symbionts between hosts (Rowley *et al.*, 2004; Goodacre *et al.*, 2006). Vertical transmission rates in *M. fradeorum* were very high in the laboratory but may be lower in the field (Turelli and Hoffmann, 1995). High endosymbiont loss rates would be consistent with the relatively high proportion of uninfected individuals observed in the field (Figure 3). Horizontal transmission has previously been inferred among spiders, through hypothesized mechanisms such as shared vectors, cannibalism, paternal transmission (Baldo *et al.*, 2008) or a predatory lifestyle (Rowley *et al.*, 2004). Regardless of mechanism, the 'mix and match' among endosymbionts in this system may lead to conflicting selection on bacterial strains, depending on the presence or absence of co-infecting strains.

In  $R_1W_1$  spiders, the relative contribution of the two symbionts to the expressed phenotype is not entirely clear. Despite repeated attempts at low-dose antibiotics (for example, White *et al.* 2009), we were not able to differentially cure one or the other endosymbiont and therefore could not directly attribute feminization to either bacterial strain. However, 3 out of the 11 field males bore an  $R_1$  infection (Figure 3), and two generations of  $R_1$  rearing in the laboratory demonstrated an even sex ratio. These results indicate that  $R_1$  *Rickettsia* alone does not preclude production of males. It is possible that  $W_1$  alone is responsible for the female bias in  $R_1W_1$ , but no lone  $W_1$  infections were found in the field to allow evaluation of this hypothesis. It is also possible that both *Rickettsia* and *Wolbachia* are required to induce female bias in  $R_1W_1$  spiders; previous studies investigating the respective roles of co-infecting symbionts have sometimes found interactive effects among the symbionts (Zhu *et al.*, 2012) and sometimes have not (White *et al.*, 2009).



**Figure 4** Summary and implications of laboratory rearing and field collection for each endosymbiotic infection type for *Mermessus fradeorum*. FM=feminization, CI=cytoplasmic incompatibility, R = *Rickettsia*, W = *Wolbachia*.

This radical female bias induced by the  $R_1W_1$  infection type is likely the result of feminization, rather than male killing or parthenogenesis. Fecundity and the number of undeveloped embryos in  $R_1W_1$  versus  $R_1 - W_1$  spiders were similar (Table 1), whereas one would expect a substantial reduction in fecundity if male killing were at play in a population with a 1:1 ratio. Parthenogenetic reproduction is not supported, as we never observed virgin females to produce offspring during nine generations of laboratory rearing. Furthermore, the bimodal chromosome numbers in the  $R_1W_1$  line of *M. fradeorum* is consistent with a mixture of genetic females ( $n=26$  chromosomes) and feminized genetic males ( $n=24$  chromosomes). Feminization is suspected to be very common in isopods (Cordaux *et al.*, 2011) but has only rarely been reported in insects and mites (Engelstädter and Hurst, 2009). The present study is the first indication of feminization by endosymbionts in Araneae, but this mechanism of manipulation may be common in an order rife with female-biased populations (Eberhard, 2004).

On its own,  $W_2$  induces strong CI, reducing fecundity by 73% when  $W_2$  males were mated with uninfected  $W_2 -$  females. However,  $W_2$  CI strength may be undercut in two ways. First, the 27% of offspring that escaped CI in the incompatible cross enjoyed a higher chance of survival to adulthood, perhaps due to consumption of undeveloped siblings. Sibling cannibalism can provide a significant fitness benefit when resources are scarce, which is often the basis for the spread of male-killing endosymbionts in host populations (Freeland and McCabe, 1997). For a CI-inducing endosymbiont, however, a significant fitness boon for survivors of the incompatible cross is predicted to undermine its ability to spread or compete with other endosymbionts in the population (Freeland and McCabe, 1997). Second,  $W_2$  induced only weak CI in feminized  $R_1W_1$  *M. fradeorum*. In this instance, we were unable to perform reciprocal crosses that would allow us to distinguish whether this partial rescue was attributable to the  $R_1W_1$  endosymbionts; nevertheless, this lowered level of CI may help the  $R_1W_1$  infection type persist in the same host population as the CI-inducing  $W_2$  infection.

As a further indication of potential endosymbiont interactions within hosts, our field collection data suggest that  $R_1W_1$  feminization may be modulated by the additional presence of  $W_2$  within spiders. One of the three field-collected  $R_1W_1W_2$  spiders was male, as was one of the two field-collected  $W_1W_2$  spiders, ratios that are unlikely if these spiders show a similar female bias as the  $R_1W_1$  spiders in the laboratory ( $P < 0.05$ , Fisher's exact test). Future studies to explore the phenotypes of other symbiont combinations within this system

will provide important insights into the potential for conflict and collaboration among independent genetic elements within a shared host environment. Our results, in conjunction with a growing body of evidence for widespread multiple endosymbiont infection in spiders (Goodacre, 2011; Stefanini and Duron, 2012; Vanthournout *et al.*, 2014), indicate that these infections are common and dynamic, suggesting that spiders are an ideal taxon for the investigation of interactions among endosymbionts.

Importantly, our results also affirm that the same mechanisms of reproductive manipulation found in insects are active in spiders. To the best of our knowledge, this is the first demonstration of CI and feminization in a spider, signifying that the larger body of endosymbiont knowledge based upon insects is likely applicable to spiders as well. In one population of *M. fradeorum*, we have found multiple endosymbionts, a high prevalence of multiply infected individuals and two distinct reproductive manipulations. Though the picture is far from complete, the potential importance of interactions among endosymbionts in population dynamics should not be ignored.

#### DATA ARCHIVING

The data used in this study are available at the Dryad Digital Repository doi:10.5061/dryad.935hp.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### ACKNOWLEDGEMENTS

We thank A Dehnel, Z Helton and E Squires of the University of Kentucky for help with spider rearing and diagnostic PCR. We also thank all reviewers for their constructive comments on previous drafts of this manuscript. This research was supported by the University of Kentucky College of Agriculture and Department of Entomology and USDA National Institute of Food and Agriculture, Hatch/Multistate project KY00856. The information reported in this paper (15-08-016) is part of a project of the Kentucky Agricultural Experiment Station and is published with the approval of the Director.

Baldo L, Ayoub NA, Hayashi CY, Russell JA, Stahlhut JK, Werren JH (2008). Insight into the routes of *Wolbachia* invasion: high levels of horizontal transfer in the spider genus *Agelenopsis* revealed by *Wolbachia* strain and mitochondrial DNA diversity. *Mol Ecol* 17: 557–569.

- Charlat S, Engelstädter J, Dyson EA, Hornett EA, Duploux A, Tortosa P *et al.* (2006). Competing selfish genetic elements in the butterfly *Hypolimnas bolina*. *Curr Biol* **16**: 2453–2458.
- Cook RD, Weisberg S (1999). *Applied Regression Including Computing and Graphics*. John Wiley & Sons: New York, NY, USA.
- Cordaux R, Bouchon D, Grève P (2011). The impact of endosymbionts on the evolution of host sex-determination mechanisms. *Trends Genet* **27**: 332–341.
- Curry MM (2013). Endosymbiotic prevalence and reproductive manipulation of the spider *Mermessus fradeorum*. MSc thesis, University of Kentucky, Lexington, KY, USA.
- Duron O, Bouchon D, Boutin S, Bellamy L, Zhou LQ, Engelstädter J *et al.* (2008a). The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biol* **6**: 12.
- Duron O, Hurst GDD, Hornett EA, Josling JA, Engelstädter J (2008b). High incidence of the maternally inherited bacterium *Cardinium* in spiders. *Mol Ecol* **17**: 1427–1437.
- Duron O, Raymond M, Weill M (2010). Many compatible *Wolbachia* strains coexist within natural populations of *Culex pipiens* mosquito. *Heredity* **106**: 986–993.
- Drummond AJ, Ashton B, Buxton S, Cheung M, Cooper A, Duran C *et al.* (2012). Geneious v5.6. Available from <http://www.geneious.com>.
- Eberhard WG (2004). Why study spider sex: special traits of spiders facilitate studies of sperm competition and cryptic female choice. *J Arachnol* **32**: 545–556.
- Engelstädter J, Telschow A, Hammerstein P (2004). Infection dynamics of different *Wolbachia* types within one host population. *J Theor Biol* **231**: 345–355.
- Engelstädter J, Hurst GDD (2009). The ecology and evolution of microbes that manipulate host reproduction. *Annu Rev Ecol Syst* **40**: 127–149.
- Freeland SJ, McCabe BK (1997). Fitness compensation and the evolution of selfish cytoplasmic elements. *Heredity* **78**: 391–402.
- Goodacre SL, Martin OY, Thomas CFG, Hewitt GM (2006). *Wolbachia* and other endosymbiont infections in spiders. *Mol Ecol* **15**: 517–527.
- Goodacre SL, Martin OY, Bonte D, Hutchings L, Woolley C, Ibrahim K *et al.* (2009). Microbial modification of host long-distance dispersal capacity. *BMC Biol* **7**: 8.
- Goodacre SL (2011). Endosymbiont infections in spiders. Casas J (ed). *Advances in Insect Physiology*, Vol 60. Academic Press: London, UK. pp 137–153.
- Greenstone MH, Payton ME, Weber DC, Simmons AM (2013). The detectability half-life in arthropod predator-prey research: what it is, why we need it, how to measure it, and how to use it. *Mol Ecol* **23**: 3799–3813.
- Gunnarsson B, Goodacre SL, Hewitt GM (2009). Sex ratio, mating behaviour and *Wolbachia* infections in a sheetweb spider. *Biol J Linn Soc* **98**: 181–186.
- Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH (2008). How many species are infected with *Wolbachia*?—a statistical analysis of current data. *FEMS Microbiol Lett* **281**: 215–220.
- Hiroki M, Tagami Y, Miura K, Kato Y (2004). Multiple infection with *Wolbachia* inducing different reproductive manipulations in the butterfly *Eurema hecabe*. *Proc Soc Biol* **271**: 1751–1755.
- Hoffmann AA, Turelli M (1997). Cytoplasmic incompatibility in insects. O'Neill SL, Hoffmann AA, Werren JH (eds). *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*. Oxford University Press: Oxford, UK. pp 42–80.
- Hughes G, Allsopp P, Brumbley S, Woolfit M, McGraw E, O'Neill S (2011). Variable infection frequency and high diversity of multiple strains of *Wolbachia pipientis* in *Perkinsiella* planthoppers. *Appl Environ Microbiol* **77**: 2165–2168.
- Huigens M, Stouthamer R (2003). Parthenogenesis associated with *Wolbachia*. Bourtzis K, Miller TA (eds). *Insect Symbiosis*. CRC Press: Boca Raton, FL, USA. pp 247–266.
- Hurst GDD, Jiggins FM (2000). Male-killing bacteria in insects: mechanisms, incidence, and implications. *Emerg Infect Dis* **6**: 329–336.
- Kořínková T, Král J (2013). Karyotypes, sex chromosomes, and meiotic division in spiders. Nentwig W (ed). *Spider Ecophysiology*. Springer: Heidelberg, NY, USA. pp 159–171.
- McDonald JH, Delaware UO (2009). *Handbook of Biological Statistics*. Sparky House Publishing: Baltimore, MD, USA.
- Medina R, Nachappa P, Tamborindeguy C (2011). Differences in bacterial diversity of host-associated populations of *Phylloxera notabilis* Pergande (Hemiptera: Phylloxeridae) in pecan and water hickory. *J Evol Biol* **24**: 761–771.
- Narita S, Nomura M, Kato Y, Fukatsu T (2006). Genetic structure of sibling butterfly species affected by *Wolbachia* infection sweep: evolutionary and biogeographical implications. *Mol Ecol* **15**: 1095–1108.
- Negri I, Pellecchia M, Mazzoglio P, Patetta A, Alma A (2006). Feminizing *Wolbachia* in *Zyginidia pullula* (Insecta, Hemiptera), a leafhopper with an XX/X0 sex-determination system. *P Soc Biol* **273**: 2409–2416.
- O'Neill SL, Hoffmann AA, Werren JH (1997). *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*. Oxford University Press: Oxford, UK, 214 pp.
- Perlman SJ, Magnus SA, Copley CR (2010). Pervasive associations between *Cybaeus* spiders and the bacterial symbiont *Cardinium*. *J Invertebr Pathol* **103**: 150–155.
- Rowley SM, Raven RJ, McGraw EA (2004). *Wolbachia pipientis* in Australian spiders. *Curr Microbiol* **49**: 208–214.
- Rozen S, Skaletsky HJ (1998). *Primer3*. Whitehead Institute: Cambridge, MA, USA. Available from [http://www-genome.wi.mit.edu/genome\\_software/other/primer3.html](http://www-genome.wi.mit.edu/genome_software/other/primer3.html).
- Riegler M, Stauffer C (2002). *Wolbachia* infections and superinfections in cytoplasmically incompatible populations of the European cherry fruit fly *Rhagoletis cerasi* (Diptera, Tephritidae). *Mol Ecol* **11**: 2425–2434.
- SAS Institute Inc. (2012). *Using JMP 10*. SAS Institute Inc.: Cary, NC, USA.
- Stefanini A, Duron O (2012). Exploring the effect of the *Cardinium* endosymbiont on spiders. *J Evol Biol* **25**: 1521–1530.
- Taylor D (1990). Evolutionary consequences of cytoplasmic sex ratio distorters. *Evol Ecol* **4**: 235–248.
- Turelli M, Hoffmann AA (1995). Cytoplasmic incompatibility in *Drosophila simulans*: dynamics and parameter estimates from natural populations. *Genetics* **140**: 1319–1338.
- Vanhourntout B, Swaegers J, Hendrickx F (2011). Spiders do not escape reproductive manipulations by *Wolbachia*. *BMC Evol Biol* **11**: 9.
- Vanhourntout B, Vandomme V, Hendrickx F (2014). Sex ratio bias caused by endosymbiont infection in the dwarf spider *Oedothorax retusus*. *J Arachnol* **42**: 24–33.
- Welch KD (2013). Selective utilization of microhabitats by web-building spiders. PhDSc dissertation, University of Kentucky, Lexington, KY, USA.
- Werren JH, Baldo L, Clark ME (2008). *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* **6**: 741–751.
- Weyman G, Sunderland K, Jepson P (2002). A review of the evolution and mechanisms of ballooning by spiders inhabiting arable farmland. *Ethol Ecol Evol* **14**: 307–326.
- White JA, Kelly SE, Perlman SJ, Hunter MS (2009). Cytoplasmic incompatibility in the parasitic wasp *Encarsia inaron*: disentangling the roles of *Cardinium* and *Wolbachia* symbionts. *Heredity* **102**: 483–489.
- White JA, Kelly SE, Cockburn SN, Perlman SJ, Hunter MS (2011). Endosymbiont costs and benefits in a parasitoid infected with both *Wolbachia* and *Cardinium*. *Heredity* **106**: 585–591.
- Williams DA (1982). Extra-binomial variation in logistic linear models. *Appl Stat* **31**: 144–148.
- Yun YL, Lei CL, Peng Y, Liu FX, Chen JA, Chen LB (2011). *Wolbachia* strains typing in different geographic population spider, *Hylyphantes graminicola* (Linyphiidae). *Curr Microbiol* **62**: 139–145.
- Zhu L-Y, Zhang K-J, Zhang Y-K, Ge C, Gotoh T, Hong X-Y (2012). *Wolbachia* strengthens *Cardinium*-induced cytoplasmic incompatibility in the spider mite *Tetranychus piercei* McGregor. *Curr Microbiol* **65**: 516–523.

Supplementary Information accompanies this paper on Heredity website (<http://www.nature.com/hdy>)