



Published in final edited form as:

Dev Psychopathol. 2015 November ; 27(4 Pt 2): 1515–1526. doi:10.1017/S0954579415000917.

Maternal Elaborative Reminiscing Mediates the Effect of Child Maltreatment on Behavioral and Physiological Functioning

Kristin Valentino¹, Leah C Hibell², E. Mark Cummings¹, Amy K. Nuttall¹, Michelle Comas¹, and Christina G. McDonnell¹

¹Department of Psychology, University of Notre Dame

²Department of Human Ecology, University of California, Davis

Abstract

Theoretical and empirical evidence suggest that the way in which parents discuss everyday emotional experiences with their young children (i.e., elaborative reminiscing) has significant implications for child cognitive and socio-emotional functioning, and that maltreating parents have a particularly difficult time in engaging in this type of dialogue. This dyadic interactional exchange, therefore, has the potential to be an important process variable linking child maltreatment to developmental outcomes at multiple levels of analysis. The current investigation evaluated the role of maternal elaborative reminiscing in associations between maltreatment and child cognitive, emotional, and physiological functioning. Participants included 43 maltreated and 49 nonmaltreated children (aged 3–6) and their mothers. Dyads participated in a joint reminiscing task about four past emotional events, and children participated in assessments of receptive language and emotion knowledge. Child salivary cortisol was also collected from children three times a day (waking, midday, and bedtime) on two consecutive days to assess daily levels and diurnal decline. Results indicated that maltreating mothers engaged in significantly less elaborative reminiscing than nonmaltreating mothers. Maternal elaborative reminiscing mediated associations between child maltreatment and child receptive language and child emotion knowledge. Additionally, there was support for an indirect pathway between child maltreatment and child cortisol diurnal decline through maternal elaborative reminiscing. Directions for future research are discussed and potential clinical implications are addressed.

Keywords

child maltreatment; mother-child reminiscing; language; emotion knowledge; physiological regulation; salivary cortisol

Child maltreatment is a pathogenic relational experience characterized by parenting behavior that far exceeds acceptable disciplinary practices, fails to meet the child's basic physical and emotional needs, and is destructive to child development in multiple domains of functioning (Cicchetti & Toth, 2015; Cicchetti & Valentino, 2006). In 2013, approximately 3.9 million children were subjects of maltreatment reports to child welfare agencies; nearly 700,000

cases were substantiated leading to a national victimization rate of 9 per 1,000 children (US DHHS, 2015). Young children in particular are at increased risk for maltreatment such that nearly 50% of all victims are age 5 or younger. Among investigated cases, up to 50% exhibit clinically significant mental health symptoms (Burns et al., 2004), and approximately 68.1% of preschool-aged children exhibit substantial behavioral, emotional, and/or developmental problems (Stahmer et al., 2005). Though present across all demographics, child maltreatment tends to cluster in low income families and in 2013, more than 90% of child victims were maltreated by one or both parents (US DHHS, 2015).

Maltreatment, Parent-child Interactions, and Child Outcomes

Positive parenting is vital in supporting children's appropriate development of biological and psychological functioning (Gunnar, Fisch, & Malone, 1984; Sroufe, 2000; Tamis-LeMonda, Shannon, Cabrera, & Lamb, 2004), thus in the absence of such parental support, maltreated children are at heightened risk for a host of physical, mental, and cognitive impairments (e.g., Cicchetti & Toth, 2015; Cicchetti & Valentino, 2006). With nearly 1 million substantiated cases of child abuse and neglect in the United States each year, child maltreatment is clearly a public health concern requiring efforts to uncover the pathways to child maladjustment (US DHHS, 2015). Yet surprisingly little research has been conducted to identify specific parent-child interactions during early childhood that may link child maltreatment to maltreated children's developmental outcomes at multiple levels of analysis. Theoretical (Nelson & Fivush, 2004) and empirical (Fivush, Haden & Reese, 2006) evidence suggest that the way in which parents discuss everyday emotional experiences with their young children (i.e., elaborative reminiscing) has significant implications for child cognitive and socio-emotional functioning, and that maltreating parents have a particularly difficult time in engaging in this type of dialogue (Valentino, Comas, Nuttall, & Thomas, 2013). This dyadic interactional exchange therefore has the potential to be an important process variable linking child maltreatment to a host of developmental outcomes. Thus, the current investigation evaluates the role of maternal elaborative reminiscing in associations between maltreatment and child cognitive, emotional, and physiological functioning.

Parent-child interactions that are characterized by warmth, support, and engagement scaffold young children's cognitive development, and provide external emotional and physiological regulation for the child (e.g., Bowlby, 1969; Spangler, Schieche, Iig, Maier, & Ackermann, 1994; Sroufe, Carlson, Levy, & Egeland, 1999). In the long-term these skills produce well adjusted, socially competent individuals with the ability to form positive interpersonal relationships, understand others, articulate one's own feelings and needs, and manage conflict (Coster & Cicchetti, 1993; Parke, Cassidy, Burks, Carson, & Boyum, 1992; Rogosch, Cicchetti, & Aber, 1995; Stock & Fisher, 2006). For maltreated children, parent-child interactions often fail to foster these critical skills, with particular deficits in children's receptive and expressive language, emotional understanding, and ability to physiologically regulate stress (Cicchetti & Toth, 2015). Specifically, maltreated children use fewer words, have less complex sentence structure, and demonstrate poorer receptive language than nonmaltreated children matched on socioeconomic status and IQ (i.e., Coster & Cicchetti, 1993, Eigsti & Cicchetti, 2004; Pears & Fisher, 2005; Stacks, Beeghly, Partridge, & Dexter, 2011). Maltreated school-aged children demonstrate less emotional knowledge (i.e., ability

to express and recognize basic emotions, as well as to understand the causes and consequences of emotions) than nonmaltreated children (Edwards, Shipman, & Brown, 2005; Rogosch et al., 1995; Shipman & Zeaman, 1999; Sullivan, Bennett, Carpenter, & Lewis, 2008). Lastly, the hypothalamic-pituitary-adrenal (HPA) axis, with the end product cortisol, has the dual responsibility of being the body's primary stress response system, as well as the primary synchronizer of multiple physiological systems around the 24-hour dark/light cycle (i.e., a diurnal rhythm; Smyth, Hucklebridge, Thorn, Evans, & Clow, 2013). Maltreated children often exhibit dysregulation in this system characterized by a flattening of diurnal cortisol activity, including lower early morning cortisol, and less cortisol decline across the day (for review see Tarullo & Gunnar, 2006). Taken together these deficits and dysregulations place maltreated children at risk for poor school performance and academic success, as well as later emotion dysregulation and psychopathology (Rogosch et al., 1995; Ruttle et al., 2011; Shipman et al., 2007; Stock & Fisher, 2006). To successfully intervene and break this nefarious pathway to maladjustment, critical process variables translating maltreatment into child dysfunction must be uncovered.

Reminiscing as a Pathway to Child Behavioral and Physiological Outcomes

Associations among maltreatment, parenting, and developmental processes are dynamic and require multi-level, multi-systemed approaches to elucidate how these transactions unfold across the dyad, and over time (Cicchetti & Valentino, 2007). During the preschool years, regulation shifts from being primarily externally supported to more internally mediated processes, with the emergence of more effortful regulation of emotions and behavior (Calkins, 2009; Kopp, 1989). Likewise, during this time, parenting typically shifts to include more verbal and cognitive coping strategies such as reappraisal, redirection, or emotion coaching to help their children regulate distress and overcome challenge (Gottman, Katz, & Hooven, 1996; Sameroff, 2009; Sroufe, 2000). Compared to the extensive literature delineating the parenting patterns of maltreating mothers during infancy and toddlerhood (Barnett, Ganiban, & Cicchetti, 1999), and highlighting the salience of the mother-child attachment relationship for maltreated children's subsequent development (Toth, Gravener-Davis, Guild, & Cicchetti, 2013), relatively less work has focused on identifying these more verbal aspects of parenting behaviors among maltreating mothers with their preschool aged children. In particular, mothers' ability to co-construct elaborative and emotionally supportive narratives (i.e., elaborative reminiscing) about children's emotional experiences as children's verbal skills develop into the preschool years, becomes critical in shaping children's representational models, and in supporting cognitive and socio-emotional development (Fivush et al., 2006; Thompson, 2006). Whereas reminiscing has been identified as an important factor in associations between parenting and developmental processes among typically developing families, there are gaps in understanding the role of elaborative reminiscing in the context of child maltreatment.

There are several reasons to anticipate that the reminiscing style of maltreating mothers will diverge from nonmaltreating mothers when reminiscing with their preschool-aged children about past emotional events. There are clear individual differences in maternal reminiscing style (e.g., Fivush & Fromhoff, 1988), particularly with regard to elaboration during past-shared event discussion. Mothers possessing a high elaborative reminiscing style talk in rich,

detailed ways about past events with their preschool-aged children by asking more open-ended questions, providing elaborative details, and confirming children's contributions to the conversation (Fivush et al., 2006). In contrast, mothers from abusing families engage in fewer verbal interactions with their children throughout infancy (Valentino, Cicchetti, Toth, & Rogosch, 2006) and the preschool years (Alessandri, 1992; Eigsti & Cicchetti, 2004), at least during free play interactions. Furthermore, findings from the emotion socialization literature suggest that maltreating mothers engage in less discussion regarding emotional states with their school-aged children (Cicchetti, 1990; Shipman & Zeman, 1999). To date, there has been no research evaluating the elaborative reminiscing of maltreating mothers with their children during early childhood, the developmental period during which reminiscing may have the largest influence on children's emerging behavioral and regulatory development (Wareham & Salmon, 2006).

The way in which parents reminisce about past experiences with their young children may influence child cognitive functioning. An elaborative style of reminiscing benefits preschool-aged children's autobiographical memory, language and literacy development (Nelson & Fivush, 2004; Peterson, Jesso, & McCabe, 1999; Reese, 1995; Sparks & Reese, 2013). Evidence suggests that maltreating mothers engage in fewer verbal interactions with their children (Alessandri, 1992; Valentino et al., 2006), are less likely to use verbal means of instruction, and are less likely to respond to children's initiatives (Egeland & Sroufe, 1981). Thus, we examined poor elaborative reminiscing as a mechanism explaining associations between child maltreatment and child language development.

Moreover, when reminiscing focuses on children's past *emotional* events specifically, the parent's reminiscing style has important implications for children's emerging emotional development. In particular, mother-child reminiscing about children's emotions provides a context for children to understand past emotional events, and to integrate these events into a coherent autobiography or self-concept (e.g., Fivush, 1993; Nelson, 1993). Mother-child reminiscing that highlights shared positive emotion may foster positive aspects of the parent-child relationship (Nelson & Fivush, 2004, Wareham & Salmon, 2006). Moreover, supportive reminiscing of children's *negative* emotions including validation of children's feelings, identification and explanations of children's emotions and resolutions, appears to be uniquely associated with children's emotion regulation (Fivush et al., 2006) and self-esteem (Bohanek, Marin, & Fivush, 2008). When mothers are not able to engage in emotional discussion, however, and are dismissing or avoidant of children's negative emotions, then children may not develop adequate coping skills, thereby increasing risk for psychopathology (Koren-Karie, Oppenheim, & Getzler Yosef, 2004). Maltreating mothers engage in less emotion discussion (Shipman & Zeman, 1999), less emotion coaching, and more emotion invalidation than nonmaltreating mothers (Shipman et al., 2007), and these behaviors, in turn, have been shown to mediate the association between maltreatment and school-aged children's emotion regulation (Shipman et al., 2007). Given the importance of young children's emerging emotion knowledge for facilitating emotional competence and regulation, we set out to examine maternal elaborative reminiscing about children's emotions as an explanatory factor between maltreatment and children's emotion knowledge during the preschool years.

A primary mechanism by which early life stressors, such as maltreatment, may be translated into long term mental and physical well-being is through changes in HPA activity (Essex et al., 2011; McEwen, 1998). For the most part, the studies of the physiological ramifications of maltreatment have relied on retrospective reports of early maltreatment (e.g., Carpenter, et al., 2007; Gonzalez, Jenkins, Steiner, & Fleming, 2009; van der Vegt, van der Ende, Kirschbaum, Verhulst, & Teimerier, 2009), with relatively fewer studies examining prospective or concurrent processes (e.g., Bruce, Fisher, Pears, & Levine, 2009; Cicchetti & Rogosch, 2001). While retrospective studies are informative, they provide limited information on the specific features of parent-child interactions in maltreating dyads that exert regulatory control over the developing HPA axis. Thus, there is a gap in the current knowledge describing the process by which maltreatment “gets under the skin” influencing HPA development and producing enduring trait-like adrenocortical profiles.

Although there have not been any studies specifically examining associations between maternal elaborative reminiscing and child HPA functioning, supportive parental responses to children’s emotions predict children’s physiological regulation of the peripheral nervous system (Gottman, et al., 1996; Hooven, Gottman, & Katz, 1995). Moreover, severe disruptions in mother-infant communication have been associated with divergence between maternal and child cortisol levels (Crockett, Holmes, Granger, & Lyons-Ruth, 2013), and there are higher rates of child maltreatment among mothers who display higher levels of disrupted communication with her infant (Lyons-Ruth, Bronfman, & Parsons, 1999). As such children whose mothers engage in elaborative discussions of children’s past emotions may develop more adaptive physiological regulatory abilities to manage emotional arousal (Gottman et al., 1996). Conversely, in maltreated children, whose mothers may have difficulties with mother-child communication about emotions, we expected to find HPA axis dysregulation.

Hypotheses

The two main objectives of this project were to examine maternal elaborative reminiscing as a mediator between maltreatment and 1) child behavioral outcomes (receptive language and child emotion knowledge) and 2) indices of children’s physiological outcomes (cortisol levels and diurnal decline). As a critical step in assessing these mediation processes, we hypothesized that maltreating mothers would demonstrate less elaborative reminiscing with their preschool-aged children than would non-maltreating mothers. In this scenario, we expected that elaborative reminiscing would mediate associations between child maltreatment and child outcomes in behavioral and physiological domains. Behaviorally, we expected that elaborative reminiscing would be positively linked to child language and child emotion knowledge, and would account for significant variance in the association between child maltreatment and these outcomes. Physiologically, we expected that elaborative reminiscing would be positively linked to child cortisol levels and diurnal decline, and would account for significant variance in the association between child maltreatment and these outcomes. In summary, we expected that maltreatment would be related to multiple processes potentially related to developmental outcomes among maltreated children, including receptive language, emotion knowledge, and HPA axis functioning, reflected in

cortisol levels and responses, and that maternal elaborative reminiscing would mediate these associations.

Methods

Participants

The participants included 92 children between 3 and 6 years of age and their mothers. 43 families had substantiated child maltreatment with the mother named as a perpetrator. The other 49 families were demographically comparable to the maltreated families but had no history of prior involvement with the child welfare system. All children were residing with their biological mothers. Maltreated and nonmaltreated dyads did not differ on a number of important demographic characteristics (see Table 1). In addition, maltreating and nonmaltreating mothers did not differ in language abilities, with both groups performing approximately one standard deviation below the mean on standardized assessments of expressive and receptive language.

Maltreating families were recruited through the Department of Child Services (DCS). DCS Family Case Workers initially introduced our project to eligible participants with a verbal script and an informational flyer. They asked whether or not mothers would be interested in sharing their contact information with project staff, who then contacted interested families to discuss enrollment. Nonmaltreating families were recruited in the local community in locations that typically serve a similar demographic population such as the WIC office, the housing authority, and Head Start. All participating families provided informed consent and signed release forms granting access to their DCS records. The presence or absence of maltreatment was subsequently verified through extensive examinations of each family's case history and through maternal interview. Only families who have never received child protective services through DCS and indicated no maltreatment on the maternal interview were included in the nonmaltreating comparison sample.

Maltreatment Classifications

DCS records were coded using the Maltreatment Classification System (MCS; Barnett, Manly, & Cicchetti, 1993). The MCS utilizes operational criteria for determining the occurrence of subtypes of maltreatment. Subtype categories include sexual abuse, physical abuse, physical neglect, and emotional maltreatment. Sexual abuse is coded when any sexual contact or attempted sexual conduct occurred between the child and an adult. Physical abuse is determined by injuries that had been inflicted upon a child by nonaccidental means. Physical neglect is coded for failure of the primary caregiver to meet a child's needs for food, clothing, shelter, health care, education, hygiene, or safety. Emotional maltreatment is coded for chronic or extreme neglect or disregard of children's emotional needs (see Barnett et al., 1993). Additionally, the severity, chronicity, perpetrator, and the developmental timing of each maltreatment incident were assessed. The MCS allows for measurement of reliability of maltreatment classifications between coders. MCS ratings were supplemented by information obtained during the Maternal Maltreatment Classification Interview (MMCI; Cicchetti, Toth, & Manly, 2002), a structured interview based on the MCS. More than 50%

of the maltreated sample was double coded ($n = 32$) by two coders, and reliability was established ($\kappa = .84-1.0$)

Among the maltreated children, 8.3% experienced sexual abuse, 18.8% experienced physical abuse, 70.8% experienced physical neglect, and 50% experienced emotional maltreatment. Consistent with other samples of maltreated children, subtype comorbidity was high (Manly, Kim, Cicchetti & Rogosch, 2001), such that 54% of the maltreated children experienced more than one subtype. Given the limited sample size, subtype was not considered in subsequent analyses.

Procedure

Data for the current investigation was drawn from a longitudinal randomized clinical trial. Families in all conditions completed a baseline assessment consisting of one session in the home followed by one in the laboratory, typically scheduled within one week of each other. Data for the current investigation was drawn from the baseline assessment only. Research staff conducting the home- and lab-based baseline assessments were blind to families' maltreatment status. At the time of writing, participants were still being enrolled in this study; as such, this report provides data for only a subsample of the full sample. Likewise, cortisol assays had only been completed on 75 of the 94 families. As part of the home assessment, mothers collected three saliva samples (waking, midday, and bedtime) on their child for two consecutive days, and completed daily diaries each day of collection. The lab assessment included observations of mother-child reminiscing as well as individual assessments of maternal language, child language and child emotion knowledge, among other tasks.

Home Measures

Daily diary data—To control for variation in family routine, affect, and compliance, mothers completed daily diaries on each day cortisol was collected. On each morning of saliva collection, participants filled out a series of questions assessing morning routine (e.g., “did you wake your child up” “did your child wake before you”, “what time did you/your child wake”) and compliance (“did you/your child get out of bed before collecting your/your child’s first sample”, “was your/your child’s first sample collected immediately upon waking”). Participants also indicated their saliva collection times. Every evening, participants completed the Positive Affect-Negative Affect Scale, a 20-item scale that measures two distinct dimensions of positive (e.g., excited, proud, interested) and negative mood (e.g., nervous, afraid, irritable) on a 6-point likert scale, ranging from 0 = not at all to 5 = extremely (PANAS; Watson & Clark, 1994). Mothers also reported their child’s positive (e.g., happy, cheerful, joyful) and negative mood (e.g., sad, scared, mad) on a 6-point likert scale. Internal consistency was high, with alpha values higher than .63 on each day (lowest maternal positive = .85, and negative = .83; and lowest child positive = .77 and negative = .63).

Cortisol—Mothers were trained to collect saliva from their child via sponge (Salimetrics Child Swab, Salimetrics, State College, PA). Samples were collected on two consecutive weekend days when mother and child were home together. Collections were based off of

each individual's schedule, occurring at waking, before lunch (i.e., midday), and before bed (i.e., bedtime). Participants were instructed to drink water 10 minutes before collection (except at waking), and not to brush their teeth, eat, or drink within 20 minutes of providing salivary samples. Respondents were instructed to keep the saliva samples in the freezer and to bring samples to the laboratory on ice in provided portable coolers when they attended their lab assessment. Families without freezers were provided a cooler in which samples could be kept; these samples were picked up by staff and transported to the lab each morning following collection. Samples were transported on ice to the lab and then frozen at -80°C until assayed.

Samples were assayed for salivary cortisol using a highly sensitive enzyme immunoassay (Salimetrics, State College, PA). The test used 25 μl of saliva (for singlet determinations), had a range of sensitivity from 0.007 to 1.8 g/dl , and had average intra- and inter-assay coefficients of variation of less than 10% and 15%, respectively. All samples were assayed in duplicate and the average of the duplicates was used in all analyses. Cortisol values 3.0 $\mu\text{g/ml}$ and higher were trimmed, as were values with coefficients of variation exceeding 15% (unless the absolute value of the difference between repeats was less than .03 $\mu\text{g/ml}$).

Adherence to the sampling protocol: Numerous steps were taken to increase compliance and protocol adherence. Research staff trained mothers to collect saliva on their children, following which staff observed mothers collect a practice child sample and provided corrective feedback. To objectively record sample collection time, each sample was placed in a bottle sealed with a MEMS (Medication Electronic Monitoring System Aardex Ltd) cap and stored in the freezer. Caps recorded the date and time of each opening. Mothers were also given a cell phone to facilitate cortisol collection. Mothers estimated their child's waking, lunch and bedtimes for the following day; based on this schedule text message-based reminders were sent to the cell phone provided 20 minutes before each collection time and mothers were asked to respond. According to the MEMS, waking samples were collected 11 min after waking ($SD = 00:19$), midday samples were collected 4 hours 49 mins after waking ($SD = 1:18$), and bedtime samples were collected 12 hours and 55 mins after waking ($SD = 1:33$). Wake times ($r = .50, p < .001$) and waking collections ($r = .40, p < .001$) were highly consistent between the two collection days. However, midday and bedtime samples were not ($r = .18, ns$ and $r = .23, p = .08$, respectively). The nonmaltreatment and maltreatment groups did not differ base on wake time, waking collection time, before lunch collection time, or before bed collection time (p 's range .17 – .57).

Lab Measures

The Peabody Picture Vocabulary Test. (PPVT-4; Dunn & Dunn, 2007)—The PPVT-4 was utilized to assess parents' and children's receptive language. The PPVT-4 is an individually administered, multiple-choice test designed to assess receptive vocabulary skills in individuals aged 2–90 years old. The PPVT-4 was administered to all mothers to ensure there were no differences between maltreating and nonmaltreating groups. It was also administered to children as the primary language dependent variable. Standard scores were utilized in analyses.

Expressive Vocabulary Test, 2nd edition (EVT-2; K. Williams, 2007)—The EVT-2 is an individually administered vocabulary test designed to assess expressive vocabulary skills. This standardized measure was administered to all mothers to ensure that there were not differences in maternal expressive language ability between the maltreating and nonmaltreating groups.

Child emotional knowledge—The Affect Knowledge Task (AKT; Denham, 1986) was used to measure multiple aspects of child emotion knowledge. Children's understanding of emotion (nonverbal recognition and verbal labeling of emotional expressions) was assessed using felt faces that depict happy, sad, angry, and afraid expressions (Denham, 1986; Denham & Couchoud, 1990). First, children were asked to identify happy, sad, angry, and afraid facial expressions verbally, by naming them. This was followed by a second (receptive) task where children were asked to identify the four emotions non-verbally, by pointing (i.e., show me the sad face). Children received 2 points for a correct answer, and 1 point for correctly specifying only the emotion's positivity or negativity (e.g., choosing a sad rather than the correct angry face). Children's emotion knowledge was operationalized as the total score across the receptive and expressive emotion knowledge tasks.

Mother-child reminiscing—During the lab-based assessment mothers were asked to nominate on paper four past emotional events that were one-time occurrences and had been experienced by the parent and child together (see Fivush, et al., 2006; Salmon, Dadds, Allen, & Hawes, 2009). Consistent with the Autobiographical Emotional Events Dialogue (Koren-Karie Oppenheim, Haimovich, & Etzion-Carasso, 2003), the mother was instructed to think of a time her child felt happy, sad, scared, and angry, and wrote a brief reminder of each event on an index card. Cues elicited predominantly negative events as maternal reminiscing about negative emotions, in particular, tends to be more elaborative and coherent (i.e., Sales, Fivush, & Peterson, 2003), and more predictive of child well-being (Sales & Fivush, 2005) than does reminiscing about exclusively positive events. Mothers and children were then asked to sit together on a couch and mothers were instructed to discuss the events with their children, as they normally would at home, while the experimenter left the room. All mothers were instructed to discuss the happy event first, and the order of the remaining three events was counterbalanced across participants. Conversations typically lasted between 5 and 15 minutes in length. The conversations were videotaped.

Coding and Reliability

Reminiscing conversations were transcribed verbatim and transcripts were then coded for maternal elaborative quantity. Maternal elaborative quantity was coded with a frequency-based scheme where each utterance (subject-verb proposition) was coded (see Harley & Reese, 1999; Fivush & Sales, 2006; Fivush & Vasudeva, 2002; Reese & Newcombe, 2007; Valentino et al., 2014; Van Bergen, Salmon, Dadds, & Allen 2009 for similar schemes). Utterances were coded for the presence or absence of Wh-questions (open-ended elaborative questions), Yes/No questions (closed-ended elaborative questions), elaborative statements, and confirmations. Elaborative statements are utterances that provided the child with new information about the event (i.e., who was there, where it occurred, etc.). Confirmations included maternal positive affirmations of child contributions to the memory conversation

(i.e., “Yes, that’s right”). The total number of each type of elaborative utterance made by each mother (Wh- Questions, Y/N Questions, elaborative statements, confirmations) was counted and summed across event discussions. Inter-rater reliability was assessed with 20% of the transcripts. Intraclass correlation coefficients for the elaborative reminiscing variables ranged from .87–.98.

Analytic Strategy

The two main objectives of these analyses were to examine maternal elaborative reminiscing as a mediator between maltreatment and 1) child receptive language and child emotion knowledge and 2) indices of child diurnal cortisol. To test these objectives we created a composite maternal elaborative reminiscing variable. The four elaborative quantity reminiscing variables (Wh- questions, Y/N questions, elaborative statements, and confirmations) were square-root transformed to normalize their distributions and then averaged together; internal consistency was adequate, $\alpha = .76$. Although some prior research has classified confirmations as evaluations, and distinguished between evaluations and elaborations, we followed Fivush and Sales (2006), Fivush and Vasudeva (2002), and Valentino et al. (2014) and combined confirmations with the other elaborative quantity variables.

To determine the relationships among child maltreatment, maternal elaborative quantity, and child diurnal cortisol (objective 2), the six cortisol samples collected across the two days were combined into two composites, area under the curve with respect to ground (AUCg), and area under the curve with respect to increase (AUCi; see Pruessner, Kirschbaum, Meinlschmid, & Hellhammer, 2003 for equations). These composites measure the two key aspects of daily cortisol (Smyth, et al., 2013) namely, the overall cortisol secretion across the day and the diurnal cortisol slope, respectively. Because of the large range of time covered, time was coded in hours and minutes elapsed (as opposed to minutes elapsed). The AUCi measure was not skewed, thus no transformation was applied; however, AUCg was skewed (skew statistic = 2.22) and subjected to a natural log transformation. This transformation removed the skew (skew statistic = .04). Thus, logged AUCg and raw AUCi were used in all analyses with these variables. Only four children of the 75 had insufficient cortisol data to compute the composite measures.

All main analyses were conducted in Mplus (Mplus version 7.2; Muthen & Muthen, 2014) using full information maximum likelihood estimation to handle missing data. To test the main mediation objectives regarding the role of elaborative reminiscing in associations between maltreatment and 1) child behavioral and 2) physiological outcomes, we implemented the nonparametric, bias-corrected bootstrap method recommended by MacKinnon, Lockwood, and Williams (2004). Specifically we used 1,000 resamples to construct bias-corrected bootstrap 95% confidence intervals around the product coefficient of the indirect effects of maltreatment via maternal elaborative reminiscing on child receptive language and child emotion knowledge (objective 1). Similarly, we constructed bias-corrected bootstrap 95% confidence intervals around the product coefficient of the indirect effect of maltreatment via maternal elaborative reminiscing on child AUCg and AUCi (objective 2).

Results

Descriptive Analyses

Sample means and standard deviations for maternal elaborative reminiscing, child behavioral functioning and child physiological functioning by maltreatment group are presented in Table 2. Maltreating and nonmaltreating mothers differed significantly with respect to average maternal elaborative reminiscing, such that maltreating mothers were less elaborative ($M = 46.38$, $SD = 23.4$) than nonmaltreating mothers ($M = 59.17$, $SD = 29.1$). In addition, maltreated children's receptive language and emotion knowledge scores were both significantly lower than their nonmaltreated peers. No significant differences emerged with respect to child AUCg or child AUCi as a function of maltreatment group. Bivariate correlations among all primary study variables are presented in Table 3.

Indicative of a strong diurnal decline, child cortisol decreased from waking ($M = .41 \mu\text{g/ml}$, $SD = .44$) to midday ($M = .24 \mu\text{g/ml}$, $SD = .30$; $t(70) = 7.75$, $p < .0001$, and again from midday to bedtime ($M = .16 \mu\text{g/ml}$, $SD = .26$; $t(71) = 4.80$, $p < .0001$. Examining individual cortisol samples across the maltreatment groups revealed lower waking cortisol in maltreated children ($M = .32 \mu\text{g/ml}$, $SD = .32$), compared to nonmaltreated children ($M = .48 \mu\text{g/ml}$, $SD = .52$; $b = -.34$, $SE = .16$, $p = .04$). Cortisol levels at midday and bedtime were not different between the two groups ($b = .10$, $SE = .22$, *n.s.* and $b = -.35$, $SE = .29$, *n.s.*, respectively).

Mother and child person-specific (e.g., age, report of health status, household income, maternal marital status, medication use, average mood) and day-specific (e.g., number of hours slept, bed time, wake time, amount of time elapsed between awakening and the first sample) variables were examined as potential control variables for child cortisol AUCg and AUCi. Mother's positive mood as reported on the PANAS was related to child AUCi, $r = -.25$, $p = .04$, and there was a trend toward flatter rhythms in children who wake before their mothers $r = .17$, $p = .15$; $r = .22$, $p = .08$ on day one and two, respectively.

Main Analyses

Maternal reminiscing as a mediator of the association between maltreatment and child behavioral functioning—We first evaluated the role of maternal elaborative reminiscing in the associations between maltreatment with child behavioral outcomes (see Figure 1). Child age was included as a continuous covariate in the model. The model was saturated. Maltreatment was negatively associated with child receptive language ($b = -7.49$, $SE = 3.17$, $p < .05$) and child emotion knowledge ($b = -1.88$, $SE = .68$, $p < .01$). Child age was not associated with child receptive language scores ($b = .77$, $SE = 1.7$, *n.s.*), but was a significant predictor of child emotion knowledge ($b = 1.67$, $SE = 0.40$, $p < .001$) such that older children demonstrated greater knowledge of emotions. Child receptive language and emotion knowledge were significantly associated ($b = 21.9$, $SE = 4.56$, $p < .001$).

Maltreatment was significantly and negatively associated with maternal elaborative reminiscing ($b = -13.49$, $SE = 5.6$, $p < .05$). Elaborative reminiscing was significantly associated with child receptive language ($b = .13$, $SE = .05$, $p < .05$). The true indirect effect was estimated to lie between -4.27 and $-.30$ with 95% confidence. Because the 95%

confidence interval did not contain zero, we concluded that the indirect effect is significantly different from zero at $p < .05$; thus, the indirect pathway between maltreatment and child receptive language via maternal elaborative reminiscing was significant. Similarly, elaborative reminiscing was significantly associated with child emotion knowledge ($b = .02$, $SE = .01$, $p < .05$). The true indirect effect was estimated to lie between $-.87$ and $-.05$ with 95% confidence. Because the 95% confidence interval did not contain zero, we concluded that the indirect effect is significantly different from zero at $p < .05$; thus, the indirect pathway between maltreatment and child emotion knowledge via maternal elaborative reminiscing was significant.

Maternal reminiscing as a mediator of the association between maltreatment and child physiological functioning

—To address objective 2 regarding the role of maternal elaborative reminiscing in the association between child maltreatment and child physiological outcomes, a parallel model was tested with AUCg and AUCi as the dependent variables (see Figure 2). The model was saturated. The association between maltreatment and AUCg was a negative trend ($b = -.25$, $SE = .18$, $p = .15$), suggesting that maltreated children had lower levels of cortisol than did the nonmaltreated children. The direct association of maltreatment to AUCi was nonsignificant ($b = -.46$, $SE = .43$, $n.s.$). AUCg and AUCi were not significantly correlated ($b = .072$, $SE = .17$, $n.s.$).

Maltreatment was negatively associated with maternal elaborative reminiscing ($b = -12.79$, $SE = 5.65$, $p < .05$). Elaborative reminiscing was not associated with AUCg ($b = .004$, $SE = .004$, $n.s.$) and the true indirect effect was estimated to lie between $-.202$ and $.03$ with 95% confidence; therefore because the confidence interval includes zero, the indirect effect is not significantly different from zero. In contrast, elaborative reminiscing was marginally associated with AUCi ($b = -.017$, $SE = .009$, $p = .058$) such that more elaborative reminiscing predicted a greater diurnal decline. The true indirect effect was estimated to lie between $.037$ and $.69$ with 95% confidence. Because the confidence interval does not include zero, we concluded that the indirect effect is significantly different from zero at $p < .05$. This pattern of results remained after controlling for maternal positive mood, child waking before the mother, and child age.

Discussion

The current investigation underscores that the manner in which mothers communicate with their preschool-aged children about children's past emotional experiences has the potential to serve an important organizing and regulatory function for children across multiple developmental domains. When done well, highly elaborative mothers engage children in the co-construction of these past event dialogues by asking open-ended questions, providing details, and confirming children's contributions to the discussion (Fivush et al., 2006). In doing so, children not only practice and improve their language functioning, but they may also learn more about their feelings and how to regulate those feelings. In the absence of appropriate elaboration or scaffolding, however, as was the case with maltreating mothers, children are at risk for deviations in language, emotion knowledge and diurnal cortisol regulation.

A key finding in the current investigation was that maltreating mothers engaged in less elaborative reminiscing than did nonmaltreating mothers when discussing past emotional events with their preschool-aged children. This is consistent with prior work indicating that maltreating mothers engage in less verbal behavior with their infants and toddlers during free play (i.e., Alessandri, 1992, Eigsti & Cicchetti, 2004; Valentino et al., 2006), and engage in less discussion of children's emotions (Shipman & Zeaman, 1999) with their school-aged children. Importantly, the current study adds that maltreating mothers engage in less elaborative reminiscing as operationalized more broadly (not specific to emotion-language), and that elaborative reminiscing mediates associations between child maltreatment and child behavioral and physiological outcomes during early childhood.

There are several reasons why maternal elaborative reminiscing, particularly about children's emotions during past event discussion, may facilitate child cognitive, emotional and physiological functioning. First, elaborative reminiscing provides children with opportunities to actively engage in the mother-child dialogue through the use of open-ended questions and confirmations of children's contributions. Related to language development, prior research findings clearly demonstrate the influence of both the quantity and quality of parental verbal input on child language acquisition (i.e., Hart & Risley, 1995). When focused on children's past emotions, these mother-child dialogues provide an important context in which mothers can help children to identify their feelings, as well as to understand the causes, consequences and ways to cope with these emotions (Laible, 2004; Wareham & Salmon, 2006). Because these conversations revolve around past events, children may be in a better position to benefit from maternal elaborative discussion because they are not currently emotionally aroused by the situation and have greater reflective distance (Denham & Burton, 2003; Fivush, Brotman, Buckner, & Goodman, 2000). Furthermore, maternal elaborative reminiscing about children's past every-day emotional experiences may be another example of external parental regulatory behavior that scaffolds children's developing physiological capacity to handle everyday stressors. Importantly, while reminiscing was only related to blunted cortisol slopes and not overall production across the day, flattened rhythms (regardless of the levels) have been associated with a wide range of physical and mental disorders (e.g., Huber, Issa, Schik & Wolf, 2006; Smyth et al., 2009). Taken together, our findings suggest that maltreating mother's deficiencies in elaborative reminiscing might be a primary factor linking child maltreatment to long-term maladjustment.

Given that maltreating mothers are engaging in less elaborative reminiscing than are nonmaltreating mothers, and that poor reminiscing partially explains associations between child maltreatment and children's developmental outcomes across multiple domains, it is important to consider what, then, is contributing to why maltreating mothers are less elaborative in their reminiscing style. As noted above, there were not differences in receptive or expressive language between maltreating and nonmaltreating mothers, thus language ability does not account for these differences in elaborative reminiscing. One possibility may be mothers' own trauma history and/or her current strategies for emotion regulation. Maternal history of childhood maltreatment or trauma is one of the largest risk factors for child maltreatment (i.e., Dixon, Browne, & Hamilton-Giachritsis, 2005), as well as for adult psychopathology including post-traumatic stress symptoms, including traumatic avoidance (Anda et al., 2006). Avoidance is an emotion-regulation strategy characterized by attempts to

not think about prior traumatic experiences. When parents are currently experiencing traumatic avoidance symptoms, they may have difficulty separating their own emotions from those of their children, and may generalize this avoidant strategy to discussions of children's emotional experiences. In fact, there is some evidence that the extent to which maternal sexual abuse history has been resolved is predictive of the quality of maternal support and coherence during emotional discussions with her children (Koren-Karie, Oppenheim, & Getzler-Yosef, 2008).

Another possible explanation for maltreating mothers' reduced elaboration might stem from her sensitivity and responsivity to child cues. Sensitive caregivers both accurately perceive children's emotional signals and respond in an appropriate and contingent manner. High rates of parental insensitivity, atypical caregiving, and even frightened/frightening parental behavior, however, are characteristic of maltreating parents (Van Ijzendoorn, Schuengel, & Bakermans-Kranenburg, 1999). If maltreating mothers are not sensitive to recognizing children's emotional signals, then they may, too, be poor at identifying past shared instances where children felt different emotions, and subsequently engaging in a highly elaborative conversation. To date, no research has compared maternal sensitivity to either the quantity or quality of mother-child reminiscing about children's everyday past experiences.

Alternately maternal physiological regulation may have an important role in accounting for differences in reminiscing between maltreating and nonmaltreating mothers. Parenting a young child is challenging (Brame, Nagin, & Tremblay, 2001) causing daily frustrations and irritations (Crnic & Greenberg, 1990) that require emotional and physiological resources to cope with these demands. Recent work suggests that dysregulations in adrenocortical activity during mother-child interactions is associated with harsh parenting practices (Martorell & Bugental, 2006), reduced maternal sensitivity (Finegood, Blair, Granger, & Hibel, under review) and disruptions in mother-infant communication (Crockett et al., 2013). Given associations between child maltreatment and long-term alterations in stress physiology (e.g., van der Vegt, et al., 2009), and the high rates of intergenerational continuity in child maltreatment, it may be that mothers with less adaptive physiological control have the most difficulty engaging children in discussions of emotion.

Future research should focus on gaining a better understanding of what maltreating mothers are doing during emotion laden interactions and conversation (or, in this case, not doing), and why maltreating mothers are prone to these disrupted communication styles. A clear understanding of the mechanisms contributing to disrupted reminiscing among maltreating mothers will require examination of factors at multiple levels of analysis, and will inform interventions aimed at improving elaborative reminiscing among maltreating mothers. Successful interventions have the potential to disrupt the negative developmental cascades in children's functioning across multiple domains.

Limitations

The current study has limitations that should be acknowledged. First, recent work among at-risk mothers has indicated that the quality of reminiscing, in particular, is most relevant to children's developing self-representations (Valentino et al., 2014), and deviations in self-development are well-documented among maltreated children (Cicchetti & Valentino, 2006).

It will be important for future research to evaluate the quality of maternal reminiscing including the extent to which maltreating mothers facilitate the co-construction of emotionally supportive and coherent narratives of children's emotional experiences (Koren-Karie et al., 2004). Second, although child maltreatment occurred prior to the assessment of other study variables, maternal elaborative reminiscing and the child behavioral and physiological variables were obtained concurrently. As such, longitudinal research is necessary to confirm the direction of associations between maternal elaborative reminiscing and child language, emotion knowledge, and diurnal cortisol regulation. Additionally, the sample size limited our ability to test more complex structural equation models, and to evaluate differences in patterns as a function of maltreatment subtype. Future research with larger samples should consider maltreatment subtype both at the child-level and the maternal-level.

Clinical Implications

The identification of maternal elaborative reminiscing as an explanatory process that links maltreatment to preschool-aged children's cognitive, emotional, and physiological development has important clinical implications. Establishing maternal elaborative reminiscing as a mediating process between child maltreatment and children's functioning across multiple levels of analysis provides key support for the potential utility of a reminiscing-based intervention for maltreating mothers and their preschool-aged children (Valentino et al., 2013). Because maternal elaborative reminiscing partially accounts for associations between prior child maltreatment and current child outcomes, improving maternal elaborative reminiscing may be effective in facilitating children's cognitive, emotional, and physiological development. Importantly, maternal reminiscing style appears to be quite modifiable and responsive to treatment. Experimental research has demonstrated that elements of an elaborative reminiscing style can be taught to parents of middle and low socioeconomic statuses and of diverse backgrounds (Boland, Haden, & Ornstein, 2003; Peterson, et al., 1999; Reese & Newcombe, 2007; Salmon, et al., 2009, Van Bergen et al., 2009). In this work, mothers were able to be effectively trained in elaborative reminiscing following a relatively brief intervention of 4–6 sessions, and maternal improvements in elaborative reminiscing following training appear to endure over time (Boland et al. 2003, Peterson et al. 1999). Recent work in our lab with maltreated preschool-aged children and their mothers suggests that following a brief, in-home intervention that focuses on facilitating elaborative and emotionally supportive reminiscing, maltreating mothers significantly improve in their reminiscing. Moreover, in the context of these past event discussions, maltreated children improve in both memory and spontaneous emotion attributions (Valentino et al., 2013). Longitudinal evaluation of this brief intervention approach in a randomized clinical design is currently underway.

Acknowledgments

The authors wish to thank Jeanne Mattei, John Borkowski, Jennifer Lefever, and the H2H project staff for their invaluable assistance with this project. Additionally, we are grateful to the children and families that participated in this study and the Department of Child Services of St. Joseph County. This research was supported by grant 5 R01 HD071933-03 to K. Valentino.

References

- Alessandri S. Mother-child interactional correlates of maltreated and nonmaltreated children's play behavior. *Development and Psychopathology*. 1992; 4:257–270.
- Anda R, Felitti V, Bremner J, Walker J, Whitfield C, Perry B. The enduring effects of abuse and related adverse experiences in childhood - A convergence of evidence from neurobiology and epidemiology. *European Archives of Psychiatry and Clinical Neuroscience*. 2006; 256(3):174–186. [PubMed: 16311898]
- Barnett D, Ganiban J, Cicchetti D. Maltreatment, negative expressivity, and the development of Type D attachments from 12- to 24-months of age. *Monographs of the Society for Research in Child Development*. 1999; 64:97–118. [PubMed: 10597544]
- Barnett, D.; Manly, JT.; Cicchetti, D. Defining child maltreatment: The interference between policy and research. In: Cicchetti, D.; Toth, S., editors. *Child abuse, child development and social policy*. Norwood, NJ, US: Ablex; 1993. p. 7-73.
- Bohanek JG, Marin KA, Fivush R. Family narratives, self, and gender in early adolescence. *The Journal of Early Adolescence*. 2008; 28:153–176.
- Boland AM, Haden CA, Ornstein PA. Boosting children's memory by training mothers in the use of an elaborative conversational style as an event unfolds. *Journal of Cognition and Development*. 2003; 4:39–65.
- Bowlby, J. *Attachment and loss*. Vol. 1: Attachment. New York, NY, US: Basic Books; 1969.
- Brame B, Nagin DS, Tremblay RE. Developmental trajectories of physical aggression from school entry to late adolescence. *Journal of Child Psychology and Psychiatry*. 2001; 42:503–512. [PubMed: 11383966]
- Bruce J, Fisher PA, Pears KC, Levine S. Morning cortisol Levels in preschool-aged foster children: Differential effects of maltreatment type. *Developmental Psychobiology*. 2009; 51:14–23. [PubMed: 18720365]
- Bruce J, Fisher PA, Pears KC, Levine S. Morning cortisol Levels in preschool-aged foster children: Differential effects of maltreatment type. *Developmental Psychobiology*. 2009; 51:14–23. [PubMed: 18720365]
- Burns B, Phillips S, Wagner H, Barth R, Kolko D, Campbell Y. Mental health need and access to mental health services by youths involved with child welfare: A national survey. *Journal of the American Academy of Child Adolescent Psychiatry*. 2004; 43:960–970. [PubMed: 15266190]
- Calkins, SD. Regulatory competence and early disruptive behavior problems: The role of physiological regulation. In: Olson, SL.; Sameroff, A., editors. *Biopsychosocial regulatory processes in the development of childhood behavioral problems*. 2009. p. 86-115.
- Carpenter LL, Carvalho JP, Tyrka AR, Wier LM, Mello AF, Mello MF, Anderson GM, Wilkinson CW, Price LH. Decreased adrenocorticotropic hormone and cortisol responses to stress in healthy adults reporting significant child maltreatment. *Biological Psychiatry*. 2007; 62:1080–1087. [PubMed: 17662255]
- Cicchetti, D. The organization and coherence of socioemotional, cognitive, and representational development: Illustrations through a developmental psychopathology perspective on Down's syndrome and child maltreatment. In: Thompson, R., editor. *Socioemotional development: Nebraska Symposium on Motivation 1988*. Lincoln, NE: University of Nebraska Press; 1990. p. 259-367.
- Cicchetti D, Rogosch FA. Diverse patterns of neuroendocrine activity in maltreated children. *Development and Psychopathology*. 2001a; 13:677– 693. [PubMed: 11523854]
- Cicchetti, D.; Toth, SL. A multilevel perspective on child maltreatment. In: Lamb, M.; Garcia Coll, C., editors. *Handbook of child psychology and developmental science: Vol. 3. Socioemotional process*. 7. Hoboken, NJ: Wiley; 2015. p. 513-563.
- Cicchetti, D.; Toth, SL.; Manly, JT. *Maternal Maltreatment Classification Interview*. Rochester, NY, US: Mt. Hope Family Center; 2003. Unpublished measure
- Cicchetti, D.; Valentino, K. An ecological-transactional perspective on child maltreatment: Failure of the average expectable environment and its influence on child development. In: Cicchetti, D.;

- Cohen, DJ., editors. *Developmental psychopathology*. Vol. 3: Risk, disorder, and adaptation. 2. Hoboken, NJ, US: John Wiley & Sons, Inc; 2006. p. 129-201.
- Cicchetti, D.; Valentino, K. Toward the application of a multiple-levels-of-analysis perspective to research in development and psychopathology. In: Masten, A., editor. *Multilevel Dynamics in Developmental Psychology*, Minnesota Symposia on Child Development. Vol. 34. Mahwah, NJ: Lawrence Erlbaum Associates; 2007. p. 243-284.
- Coster WJ, Cicchetti D. Research on the communicative development of maltreated children: Clinical implications. *Topics in Language Disorders*. 1993; 13(4):25–38.
- Crnic KA, Greenberg MT. Minor parenting stresses with young children. *Child Development*. 1990; 61:1628–1637. [PubMed: 2245752]
- Crockett EE, Holmes BM, Granger DA, Lyons-Ruth K. Maternal disrupted communication during face-to-face interaction at 4 months: Relation to maternal and infant cortisol among at-risk families. *Infancy*. 2013; 18(6):1111–1134. [PubMed: 25506272]
- Denham, SA.; Burton, R. *Social and emotional prevention and intervention programming for preschoolers*. New York: Kluwer Academic/Plenum Publishers; 2003.
- Denham SA. Social cognition, prosocial behavior, and emotion in preschoolers: Contextual validation. *Child Development*. 1986; 57:194–201.
- Denham SA, Couchoud EA. Young preschooler's understanding of emotions. *Child Study Journal*. 1990; 20:171–192.
- Dixon L, Browne K, Hamilton-Giachritsis C. Risk factors of parents abused as children: A mediational analysis of the intergenerational continuity of child maltreatment (part I). *Journal of Child Psychology and Psychiatry and Allied Disciplines*. 2005; 46(1):47–57.
- Dunn, LM.; Dunn, DM. *Peabody Picture Vocabulary Test*. 4. Minneapolis, MN, US: NCS Pearson; 2007.
- Edwards A, Shipman K, Brown A. The socialization of emotional understanding: A comparison of neglectful and nonneglectful mothers and their children. *Child Maltreatment*. 2005; 10:293–304. [PubMed: 15983112]
- Egeland B, Sroufe LA. Attachment and early maltreatment. *Child Development*. 1981; 52:44–52. [PubMed: 7238152]
- Eigsti I, Cicchetti D. The impact of child maltreatment on expressive syntax at 60 months. *Developmental Science*. 2004; 7(1):88–102. [PubMed: 15323121]
- Essex MJ, Shirtcliff EA, Burka LR, Ruttle PL, Klein MH, Slattery MJ, Kalin NH, Armstrong JM. Influence of early life stress on later hypothalamic–pituitary–adrenal axis functioning and its covariation with mental health symptoms: A study of the allostatic process from childhood into adolescence. *Development and Psychopathology*. 2011; 23:1039–1058. [PubMed: 22018080]
- Finegood E, Blair D, Granger D, Hibel LC, Mills-Koonce WR. Investigators TFLP. Maternal salivary cortisol and cumulative risk predict parenting behavior in the postpartum period. *Developmental Psychology*. under review.
- Fivush, R. Emotional content of parent–child conversations about the past. In: Nelson, CA., editor. *Memory and affect in development: The Minnesota Symposia in Child Psychology*. Hillsdale, NJ: Erlbaum; 1993. p. 39-78.
- Fivush R, Brotman M, Buckner JP, Goodman SH. Gender differences in parent–child emotion narratives. *Sex Roles*. 2000; 42:233–253.
- Fivush R, Fromhoff F. Style and structure in mother child conversations about the past. *Discourse Processes*. 1988; 11(3):337–355.
- Fivush R, Haden C, Reese E. Elaborating on elaborations: Role of maternal reminiscing style in cognitive and socioemotional development. *Child Development*. 2006; 77:1568–1588. [PubMed: 17107447]
- Fivush R, Nelson K. Culture and language in the emergence of autobiographical memory. *Psychological Science*. 2004; 15(9):573–577. [PubMed: 15327626]
- Fivush R, Sales J. Coping, attachment, and mother-child narratives of stressful events. *Merrill - Palmer Quarterly*. 2006; 52(1):125–150.
- Fivush R, Vasudeva A. Remembering to relate: Socioemotional correlates of mother-child reminiscing. *Journal of Cognition and Development*. 2002; 3(1):73–90.

- Gottman JM, Katz LF, Hooven C. Parental meta-emotion philosophy and the emotional life of families: Theoretical models and preliminary data. *Journal of Family Psychology*. 1996; 10:243–268.
- Gonzalez A, Jenkins JM, Steiner M, Fleming AS. The relation between early life adversity, cortisol awakening response and diurnal salivary cortisol levels in postpartum women. *Psychoneuroendocrinology*. 2009; 34:76–86.10.1016/j.psyneuen.2008.08.012 [PubMed: 18835661]
- Gunnar M, Fisch R, Malone S. The effects of a pacifying stimulus on behavioral and adrenocortical responses to circumcision in the newborn. *Journal of the American Academy of Child Adolescent Psychiatry*. 1984; 23:34–38.
- Gunnar MR, Quevedo KM. Early care experiences and HPA axis regulation in children: a mechanism for later trauma vulnerability. *Progress in Brain Research*. 2007; 167:137–149. [PubMed: 18037012]
- Harley K, Reese E. Origins of autobiographical memory. *Developmental Psychology*. 1999; 35(5): 1338–1348. [PubMed: 10493658]
- Hart, TR.; Risley, B. Meaningful differences in the everyday experiences of young American Children. Paul H. Brookes Publishing; 1995.
- Hooven C, Gottman JM, Katz LF. Parental meta-emotion structure predicts marital quality and children's socioemotional outcomes. *Cognition and Emotion*. 1995; 9:229–264.
- Huber TJ, Issa K, Schik G, Wolf OT. The cortisol awakening response in blunted in psychotherapy inpatients suffering from depression. *Psychoneuroendocrinology*. 2006; 31:900–904. [PubMed: 16707227]
- Kopp CB. Regulation of distress and negative emotions: A developmental view. *Developmental Psychology*. 1989; 25(3):343.
- Koren-Karie N, Oppenheim D, Getzler-Yosef R. Shaping children's internal working models through mother-child dialogues: The importance of resolving past maternal trauma. *Attachment Human Development*. 2008; 10:465–483. [PubMed: 19016053]
- Koren-Karie N, Oppenheim D, Getzler Yosef R. Mothers who were severely abused during childhood and their children talk about emotions: Co-construction of narratives in light of maternal trauma. *Infant Mental Health Journal*. 2004; 25:300–317.
- Koren-Karie, N.; Oppenheim, D.; Haimovich, Z.; Etzion-Carasso, A. Dialogues of seven-year-olds with their mothers about emotional events: Development of typology. In: Emde, RN.; Wolf, DP.; Oppenheim, D., editors. *Revealing the inner worlds of young children: The MacArthur story stem batter and parent-child narratives*. Oxford: Oxford University Press; 2003. p. 338-354.
- Laible D. Mother-child discourse surrounding a child's past behavior at 30 months: Links to emotional understanding and early conscience development at 36 months. *Merrill-Palmer Quarterly*. 2004; 50:159–181.
- Lyons-Ruth, K.; Bronfman, E.; Parsons. Atypical maternal behavior and disorganized infant attachment strategies: Frightened, frightening, and atypical maternal behavior and disorganized infant attachment strategies. *Atypical patterns of infant attachment: Theory, research, and current directions*. In: Vondra, EJ.; Barnett, D., editors. *Monographs of the Society for Research in Child Development*. Vol. 64. 1999.
- MacKinnon DP, Lockwood CM, Williams J. Confidence limits for the indirect effect: Distribution of the product and resampling methods. *Multivariate Behavioral Research*. 2004; 39:99–128. [PubMed: 20157642]
- Manly JT, Kim JE, Rogosch FR, Cicchetti D. Dimensions of child maltreatment and children's adjustment: Contributions of developmental timing and subtype. *Development and Psychopathology*. 2001; 13:759–782. [PubMed: 11771907]
- Martorell GA, Bugental DB. Maternal variations in stress reactivity: Implications for harsh parenting practices with very young children. *Journal of Family Psychology*. 2006; 20(4):641. [PubMed: 17176199]
- McEwen BS. Stress, Adaptation, and Disease: Allostasis and Allostatic Load. *Annals New York Academy of Sciences*. 1998; 840:33–44.
- Muthen, LK.; Muthen, BO. *Mplus user's guide*. 7. Los Angeles, CA: Author; 1998–2014.

- Nelson K. The psychological and social origins of autobiographical memory. *Psychological Science*. 1993; 4:1–8.
- Nelson K, Fivush R. The emergence of autobiographical memory: A social cultural developmental theory. *Psychological Review*. 2004; 111:486. [PubMed: 15065919]
- Parke, RD.; Cassidy, J.; Burks, VM.; Carson, JL.; Boyum, L. Familial contributions to peer competence among young children: The role of interactive and affective processes. In: Parke, RD.; Ladd, GW., editors. *Family–peer relationships: Modes of linkage*. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc; 1992. p. 107-134.
- Pears K, Fisher PA. Developmental, cognitive, and neuropsychological functioning in preschool-aged foster children: Associations with prior maltreatment and placement history. *Developmental and Behavioral Pediatrics*. 2005; 26(2):112–122.
- Peterson C, Jesso B, McCabe A. Encouraging narratives in preschoolers: An intervention study. *Journal of Child Language*. 1999; 26(1):49. [PubMed: 10217889]
- Pruessner JC, Kirschbaum C, Meinlschmid G, Hellhammer DH. Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology*. 2003; 28:916–931. [PubMed: 12892658]
- Reese E. Predicting children’s literacy from mother-child conversations. *Cognitive Development*. 1995; 10:381–405.
- Reese E, Newcombe R. Training mothers in elaborative reminiscing enhances children’s autobiographical memory and narrative. *Child Development*. 2007; 78:1153–1170. [PubMed: 17650131]
- Rogosch F, Cicchetti D, Aber JL. The role of child maltreatment in early deviations in cognitive and affective processing abilities and later peer relationship problems. *Development and Psychopathology*. 1995; 7:591–609.
- Ruttle PL, Shirtcliff EA, Serbin LA, Ben-Dat Fisher D, Stack DM, Schwartzman AE. Disentangling psychobiological mechanisms underlying internalizing and externalizing behaviors in youth: Longitudinal and concurrent associations with cortisol. *Hormones and Behavior*. 2011; 59:123–132. [PubMed: 21056565]
- Sales J, Fivush R. Social and emotional functions of mother-child reminiscing about stressful events. *Social Cognition*. 2005; 23(1):70–90.
- Sales J, Fivush R, Peterson C. Parental Reminiscing About Positive and Negative Events. *Journal of Cognition and Development*. 2003; 4(2):185–209.
- Salmon K, Dadds M, Allen J, Hawes D. Can emotional language skills be taught during parent training for conduct problem children? *Child Psychiatry and Human Development*. 2009; 40:485–498. [PubMed: 19373551]
- Sameroff, AE. *The transactional model of development: How children and contexts shape each other*. Washington, DC, US: American Psychological Association; 2009.
- Shipman KL, Schneider R, Fitzgerald MM, Sims C, Swisher L, Edwards A. Maternal emotion socialization in maltreating and non-maltreating families: Implications for children’s emotion regulation. *Social Development*. 2007; 16:268–285.
- Shipman K, Zeman J. Emotional understanding: A comparison of physically maltreating and nonmaltreating mother-child dyads. *Journal of Clinical Child Psychology*. 1999; 28:407–417. [PubMed: 10446690]
- Smyth N, Hucklebridge F, Thorn L, Evan P, Clow A. Salivary Cortisol as a Biomarker in Social Science Research. *Social and Personality Psychology Compass*. 2013; 7(9):605–625.
- Sroufe LA. Early relationships and the development of children. *Infant Mental Health Journal*. 2000; 21(1–2):67–74.
- Sroufe L, Carlson E, Levy A, Egeland B. Implications of attachment theory for developmental psychopathology. *Development and Psychopathology*. 1999; 11:1–13. [PubMed: 10208353]
- Spangler G, Schieche M, Iig U, Maier U, Ackermann C. Maternal sensitivity as an external organizer for biobehavioral regulation in infancy. *Developmental Psychobiology*. 1994; 27:425–437. [PubMed: 7843497]
- Sparks A, Reese E. From reminiscing to reading: Home contributions to children’s developing language and literacy in low-income families. *First Language*. 2013; 33(1):89–109.

- Stacks A, Beeghly M, Partridge T, Dexter C. Effects of Placement Type on the Language Developmental Trajectories of Maltreated Children From Infancy to Early Childhood. *Child Maltreatment*. 2011; 16:287–299. [PubMed: 22146859]
- Stahmer AC, Hurlburt M, Horwitz SM, Landsverk J, Zhang J, Leslie LK. Associations between intensity child welfare involvement and child development among young children in child welfare. *Child Abuse & Neglect*. 2009; 33:598–611. [PubMed: 19818496]
- Stock CD, Fisher PA. Language delays among foster children: Implications for policy and practice. *Child Welfare*. 2006; 85(3):445–461. [PubMed: 16999381]
- Sullivan MW, Bennett DS, Carpenter K, Lewis M. Emotion knowledge in young neglected children. *Child Maltreatment*. 2008; 13:301–306. [PubMed: 18299632]
- Tamis-LeMonda CS, Shannon JD, Cabrera NJ, Lamb MJ. Fathers and mothers at play with their 2- and 3-year olds: contributions to language and cognitive development. *Child Development*. 2004; 75(6):1806–1820. [PubMed: 15566381]
- Tarullo AR, Gunnar MR. Child maltreatment and the developing HPA axis. *Hormones and Behavior*. 2006; 50(2006):632–639. [PubMed: 16876168]
- Thompson R. Conversation and developing understanding: Introduction to the special issue. *Merrill-Palmer Quarterly*. 2006; 52:1–16.
- Toth SL, Gravener-Davis JA, Guild DJ, Cicchetti D. Relational interventions for child maltreatment: Past, present, and future perspectives. *Development and Psychopathology*. 2013; 25:1601–1617. [PubMed: 24342858]
- U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families. *Child maltreatment 2013*. 2015. Available from <http://www.acf.hhs.gov/programs/cb/resource/child-maltreatment-2013>
- Valentino K, Cicchetti D, Toth S, Rogosch F. Mother-child play and emerging social behaviors among infants from maltreating families. *Developmental Psychology*. 2006; 42:474–485. [PubMed: 16756439]
- Valentino K, Comas M, Nuttall AK, Thomas T. Training maltreating parents in elaborative and emotion-rich reminiscing with their preschool-aged children. *Child Abuse & Neglect*. 2013; 37:585–595. [PubMed: 23548682]
- Valentino K, Nuttall AK, Comas M, McDonnell CG, Piper B, Thomas T, Fanuele S. Mother-child reminiscing and autobiographical memory specificity among preschool-aged children. *Developmental Psychology*. 2014; 50(4):1197–1207. [PubMed: 24219316]
- Van Bergen P, Salmon K, Dadds M, Allen J. Training mothers in emotion-rich elaborative reminiscing: Facilitating children's autobiographical memory and emotion knowledge. *Journal of Cognition and Development*. 2009; 10:162–187.
- van der Vegt EJM, van der Ende J, Kirschbaum C, Verhulst FC, Teimerier H. Early neglect and abuse predict diurnal cortisol patterns in adults A study of international adoptees. *Psychoneuroendocrinology*. 2009; 34:660–669. [PubMed: 19128884]
- Van Ijzendoorn MH, Schuengel C, Bakermans-Kranenburg MJ. Disorganized attachment in early childhood: Meta-analysis of precursors, concomitants, and sequelae. *Development & Psychopathology*. 1999; 11:225–249. [PubMed: 16506532]
- Wareham P, Salmon K. Mother-child reminiscing about everyday experiences: Implications for psychological interventions in the preschool years. *Clinical Psychology Review*. 2006; 26:535–554. [PubMed: 16815615]
- Watson, D.; Clark, LA. *The PANAS-X: Manual for the positive and negative affect schedule - expanded form*. Ames, IA, US: The University of Iowa; 1994.
- Williams, KT. *Expressive Vocabulary Test. 2*. Minneapolis, MN, US: Pearson Assessments; 2007.




Figure 1. Mediation model of child maltreatment on child behavioral functioning via maternal elaborative reminiscing.
Note: Path coefficients are unstandardized. Dashed lines indicate the influence of control variables. Error variances are omitted from the figure; * $p < .05$, ** $p < .01$, *** $p < .001$




Figure 2.

Mediation model of child maltreatment on child physiological functioning via maternal elaborative reminiscing.

Note: AUCg: Area under the curve with respect to ground; AUCi: Area under the curve with respect to increase. Path coefficients are unstandardized. Error variances are omitted from the figure; $^+ p < .10$, $^* p < .05$, $^{**} p < .01$.

Table 1

Sample characteristics by maltreatment group

	Nonmaltreated (n =43)	Maltreated (n = 49)
Variable	M (SD)/%	M(SD)/%
Maternal Age	30.36(6.5)	29.53(5.2)
Child Age	4.94(.98)	4.85(1.25)
Child Gender		
Male	46.5%	49%
Child Ethnicity		
African American	46.5%	34.7%
Caucasian	46.5%	34.7%
Other	7%	14.3%
Maternal Employment		
Employed	39.5%	37.5%
Family Income		
\$12,000/year	56%	59%
Maternal Language		
PPVT-4	85.12(10.06)	85.08(12.8)
EVT-2	87.62(14.1)	86.29(12.7)

Note: PPVT-4: Peabody Picture Vocabulary Test, 4th Edition; EVT-2: Expressive Vocabulary Test, 2nd Edition;

*
p<.05

Means and (standard deviations) of variables of interest, separated by maltreatment group. Asterisks indicate significant differences between groups

Table 2

	Non Maltreated			Maltreated		
	Day 1	Day 2	Average	Day 1	Day 2	Average
Maternal Reminiscing			59.17 (29.1)			46.38* (23.4)
Child PPVT-4 ^a			100.95 (14.9)			92.31*** (14.1)
Child Emotion Knowledge			13.67 (2.7)			11.5*** (4.4)
Child AUC _g ^b	2.67 (2.11)	3.07 (3.31)	2.97 (2.55)	2.29 (1.76)	2.11 (1.70)	2.18 (1.73)
Child AUC _i ^c	-2.40 (2.87)	-0.92 (2.86)	-1.40 (1.97)	-2.22 (2.13)	-1.13 (1.49)	-1.70 (1.48)

Note:

^a PPVT-4; Peabody Picture Vocabulary Test, 4th Edition;

^b AUC_g: Area under the curve with respect to ground;

^c AUC_i: Area under the curve with respect to increase. t-tests AUC_g were performed on natural log transformed variables. Negative AUC_i values reflect diurnal decline. Cortisol in units of µg/ml. p < .05,

** p < .01,

*** p < .001

Table 3

Bivariate correlations among primary study variables

Variable	1.	2.	3.	4.	5.
1. Maternal Elaborative Reminiscing	1.0	--	--	--	--
2. Child PPVT-4	.24*	1.0	--	--	--
3. Child Emotion Knowledge	.18 [†]	.48***	1.0	--	--
4. Child AUCg	.16	-.20	-.03	1.0	--
5. Child AUCi	-.24*	-.27*	-.15	.04	1.0
6. Child Age	-.10	-.05	.42**	-.24*	.12

Note: PPVT-4 = Peabody Picture Vocabulary Test, 4th Edition; AUCg: Area under the curve with respect to ground; AUCi: Area under the curve with respect to increase;

[†] p<.10,

* p < .05,

*** p < .01