

Functional polymorphisms of macrophage migration inhibitory factor as predictors of morbidity and mortality of pneumococcal meningitis

Athina Savva^{a,1,2}, Matthijs C. Brouwer^{b,1}, Thierry Roger^{a,1}, Mercedes Valls Serón^b, Didier Le Roy^a, Bart Ferwerda^b, Arie van der Ende^c, Pierre-Yves Bochud^a, Diederik van de Beek^{b,1}, and Thierry Calandra^{a,1,3}

^aInfectious Diseases Service, Centre Hospitalier Universitaire Vaudois and University of Lausanne, CH-10111 Lausanne, Switzerland; ^bDepartment of Neurology, Center of Infection and Immunity Amsterdam (CINIMA), Academic Medical Center, University of Amsterdam, 1105 AZ Amsterdam, The Netherlands; and ^cDepartment of Medical Microbiology, the Netherlands Reference Laboratory for Bacterial Meningitis, CINIMA, Academic Medical Center, University of Amsterdam, 1105 AZ Amsterdam, The Netherlands

Edited by Richard Bucala, Yale University School of Medicine, New Haven, CT, and accepted by the Editorial Board February 12, 2016 (received for review November 1, 2015)

Pneumococcal meningitis is the most frequent and critical type of bacterial meningitis. Because cytokines play an important role in the pathogenesis of bacterial meningitis, we examined whether functional polymorphisms of the proinflammatory cytokine macrophage migration inhibitory factor (MIF) were associated with morbidity and mortality of pneumococcal meningitis. Two functional MIF promoter polymorphisms, a microsatellite (−794 CATT₅₋₈; rs5844572) and a single-nucleotide polymorphism (−173 G/C; rs755622) were genotyped in a prospective, nationwide cohort of 405 patients with pneumococcal meningitis and in 329 controls matched for age, gender, and ethnicity. Carriages of the CATT₇ and −173 C high-expression MIF alleles were associated with unfavorable outcome ($P = 0.005$ and 0.003) and death ($P = 0.03$ and 0.01). In a multivariate logistic regression model, shock [odds ratio (OR) 26.0, $P = 0.02$] and carriage of the CATT₇ allele (OR 5.12, $P = 0.04$) were the main predictors of mortality. MIF levels in the cerebrospinal fluid were associated with systemic complications and death ($P = 0.0002$). *Streptococcus pneumoniae* strongly up-regulated MIF production in whole blood and transcription activity of high-expression MIF promoter *Luciferase* reporter constructs in THP-1 monocytes. Consistent with these findings, treatment with anti-MIF immunoglobulin G (IgG) antibodies reduced bacterial loads and improved survival in a mouse model of pneumococcal pneumonia and sepsis. The present study provides strong evidence that carriage of high-expression MIF alleles is a genetic marker of morbidity and mortality of pneumococcal meningitis and also suggests a potential role for MIF as a target of immune-modulating adjunctive therapy.

macrophage migration inhibitory factor | polymorphism | meningitis | sepsis | innate immunity

Acute community-acquired bacterial meningitis is a life-threatening disease associated with substantial morbidity and mortality and ranks among the top 10 infectious causes of death (1). *Streptococcus pneumoniae* is the most common cause of bacterial meningitis in adults of all age groups, accounting for 50–70% of cases in developed countries (2). Pneumococcal meningitis is associated with a mortality ranging from 19% to 37% (3, 4). Neurological sequelae such as hearing loss, focal deficits, and motor and cognitive impairments significantly affect the quality of life of survivors (5–7). Predisposing factors for pneumococcal meningitis include pneumonia, otitis, sinusitis, cerebrospinal fluid (CSF) leaks, splenectomy or asplenic states, debilitating conditions (i.e., alcoholism, cirrhosis, diabetes, and cancer), and primary or acquired immune deficiencies (i.e., multiple myeloma, hypogammaglobulinemia, sickle cell anemia, HIV/AIDS, and the use of immunosuppressive agents). Genetic studies of extreme phenotypes have revealed that patients with single-gene inborn errors in *MyD88*, *IRAK4*, and *NEMO* affecting the activation of the canonical TLR and IL-1R signaling pathways or in complement factor two are

prone to pneumococcal diseases (8, 9). In addition, case-control candidate gene studies identified polymorphisms of genes associated either with increased susceptibility to (*MBL2* and *PTPN22*) or with protection from (*TIRAP*, *NFKB1A*, and *NFKB1E*) pneumococcal disease (2, 8, 9). The increased susceptibility was related to reduced concentrations of the mannose binding lectin (variants of *MBL2*) or to an increased activity of the PTPN22 phosphatase (variants of *PTPN22*) (10, 11). The protection afforded by the polymorphic *TIRAP* variant is mediated by an attenuation of TLR2 signal transduction due to a defective recruitment of the *TIRAP* variant to TLR2 (12). The functional effects of the polymorphisms of the *NFKB1A* and *NFKB1E* genes coding for the inhibitors of NF- κ B (I κ B) are unknown (13).

Cytokines are critical effector molecules of the immune system and play a central role in the orchestration of host defenses against infection. Until now, no polymorphism of cytokine genes (including *TNF*, *IL6*, *IL10*, and *LTA*) has been associated with susceptibility to and outcome of invasive pneumococcal infection (2). Within this large family of mediators, macrophage migration inhibitory factor (MIF) occupies a special place (14, 15). Unrelated to classical cytokine families (tumor

Significance

Pneumococcal meningitis, the most frequent cause of bacterial meningitis in adults, is associated with substantial morbidity and mortality. In a prospective, nationwide cohort of patients with pneumococcal meningitis, macrophage migration inhibitory factor (MIF), a proinflammatory mediator, was identified as a previously unidentified genetic marker of patient's outcome. High-expression MIF alleles were associated with disease severity and death, a finding consistent with the harmful consequences of robust proinflammatory cytokine responses on brain edema and neuronal damage in the course of bacterial meningitis. These results provide strong evidence that functional MIF polymorphisms are genetic predictors of morbidity and mortality of pneumococcal meningitis and suggest that MIF is a potential target for immune-modulating adjunctive therapies.

Author contributions: A.S., M.C.B., T.R., D.v.d.B., and T.C. designed research; A.S., M.C.B., T.R., M.V.S., D.L.R., B.F., and A.v.d.E. performed research; A.S., M.C.B., T.R., P.-Y.B., D.v.d.B., and T.C. analyzed data; and A.S., M.C.B., T.R., P.-Y.B., D.v.d.B., and T.C. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. R.B. is a guest editor invited by the Editorial Board.

¹A.S., M.C.B., T.R., D.v.d.B., and T.C. contributed equally to this work.

²Present address: 4th Department of Internal Medicine, Attikon University Hospital, University of Athens Medical School, 12462 Athens, Greece.

³To whom correspondence should be addressed. Email: thierry.calandra@chuv.ch.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1520727113/-DCSupplemental.

necrosis factor, chemokines, interleukins, or interferons), MIF is a constitutively expressed proinflammatory cytokine acting at the interface of the immune and endocrine systems. Within the innate immune system, MIF positively regulates TLR4 expression, inhibits activation-induced and p53-dependent apoptosis of macrophages, and counter-regulates the anti-inflammatory and immunosuppressive effects of glucocorticoids in part by a down-regulation of mitogen-activated protein kinase phosphatase-1 (16–19). MIF is up-regulated in inflammatory, infectious, and autoimmune diseases functioning as a modulator of innate and adaptive immunity (20–22).

Functional polymorphisms of the *MIF* gene locus include a microsatellite repeat of five to eight CATT tetranucleotide (CATT_{5–8}) at position –794 (rs5844572) and a single-nucleotide polymorphism (SNP) of a G-to-C transition at position –173 (–173G/C; rs755622) (23, 24). Genetic studies have revealed a complex picture of the role of polymorphic *MIF* alleles in the pathogenesis of autoimmune diseases (20, 25). Few studies have been performed in patients with infectious diseases, especially in patients with bacterial sepsis (20, 25). We therefore examined the impact of the *MIF* gene locus on the susceptibility to, severity of, and outcome of pneumococcal meningitis in a large, nationwide cohort of patients. Functional studies of polymorphic *MIF* promoters were conducted in human monocytic cells stimulated with *S. pneumoniae* and analyzed by *Luciferase* reporter assays. Lastly, the effect of an anti-MIF treatment strategy was evaluated in a mouse model of *S. pneumoniae* pneumonia and sepsis.

Results

Pneumococcal Meningitis Cohort. A total of 461 patients with culture-proven, community-acquired *S. pneumoniae* meningitis and 343 controls matched for age (median, 59.4 y vs. 60.1 y), gender (female: 53% vs. 50.8%), and ethnicity (Caucasian: 94.2% vs. 96.0%) were enrolled in a prospective, nationwide cohort study. DNA samples were available from 434 patients and 329 controls, who were all Caucasians. The baseline characteristics of patients are shown in Table 1. Briefly, 80.4% of the patients were bacteremic, and 43.7% required intensive care unit (ICU) admission for shock or respiratory failure. During hospitalization, 79.6%

developed neurological complications, and 38.1% developed systemic complications. Outcome was unfavorable [defined as a Glasgow Outcome Score (GOS) 1–4] in 133 patients (32.8%). A total of 30 patients (7.5%) died (GOS of 1).

Association Between *MIF* Polymorphisms and Susceptibility to, Severity of, and Outcome of Pneumococcal Meningitis. Allelic frequencies and genotypes for the –173 G/C (rs755622) and CATT_{5–8} (rs5844572) polymorphisms of patients and controls are presented in Table S1. No deviation from the Hardy–Weinberg equilibrium (HWE) was observed for the –173 G/C SNP and CATT_{5–8} microsatellite. The allele frequencies, genotypes, and haplotypes of the –173 G/C and of the CATT_{5–8} polymorphisms were similar in patients and controls ($P > 0.5$), indicating that there was no association between *MIF* polymorphisms and susceptibility to pneumococcal meningitis. Carriage of the –173 C or of the CATT₇ high *MIF* expression alleles were associated with unfavorable outcome [–173 C: odds ratio (OR) 1.9, $P = 0.003$; CATT₇: OR 1.89, $P = 0.005$], respiratory failure (–173 C: OR 1.71, $P = 0.03$), and death (–173 C: OR: 2.6, $P = 0.01$; CATT₇: OR 2.27, $P = 0.03$) (Table 2). The association between the –173 C or CATT₇ high *MIF* expression allelic variants and death was also significant by using an additive mode of inheritance (Fig. 1). Carriage of the CATT₇ allele was associated with markers of inflammation such as C-reactive protein ($P = 0.02$) and erythrocyte sedimentation rate ($P = 0.02$) and with indicators of disease severity (i.e., a CSF leukocyte count of $<1,000$ cells per mm^3 ; $P = 0.05$) (22). The association between the CATT₇ and death remained significant in a multivariate logistic regression model (OR 5.12, $P = 0.04$) and tended to be associated with unfavorable outcome (OR 2.61, $P = 0.07$) after adjustment for relevant covariables (Table 3).

MIF Levels in the CSF and Morbidity and Mortality of Pneumococcal Meningitis. MIF CSF levels were measured in 242 patients (52% of the patients included in the study). The median concentration was 2.65 ng/mL (range: 0.025–323 ng/mL). The MIF CSF levels were similar in patients in whom antibiotic treatment was started

Table 1. Characteristics of controls and patients with pneumococcal meningitis

Characteristics	Controls ($n = 343$)	Patients ($n = 461$)	Caucasian patients with DNA and outcome ($n = 405$)
Age (years)	59.4 ± 18.1	60.1 ± 20.8	
Female gender	146 (49.2)	220 (47.0)	
Caucasian ethnicity	329 (96.0)	434 (94.2)	
Predisposing factors ($n = 405$)			
Otitis/sinusitis			191 (47.3)
Immunocompromised status			100 (24.7)
Body temperature in °C ($n = 401$)			39.0 ± 1.5
Glasgow coma scale <8 (indicating coma) ($n = 402$)			53 (13.2)
Bacteremia ($n = 357$)			287 (80.4)
Neurologic complications ($n = 404$)			257 (79.6)
Systemic complications ($n = 396$)			151 (38.1)
Complications ($n = 405$)			
Shock			25 (6.2)
Respiratory failure			93 (23.0)
ICU admission			177 (43.7)
Mechanical ventilation			140 (34.6)
Outcome			
GOS ($n = 405$)			
1: Death			30 (7.5)
2: Persistent vegetative state			1 (0.2)
3: Severe disability			18 (4.4)
4: Moderate disability			84 (20.7)
5: Good recovery			272 (67.2)
Hearing loss at discharge ($n = 368$)			41 (11.1)

Data are mean ± interquartile range or number (percent).

Table 2. Association between *MIF* gene polymorphisms and patient's outcome

Polymorphisms	Unfavorable outcome (GOS \leq 4; $n = 133$)		Respiratory failure ($n = 93$)		Death ($n = 30$)	
	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>
GC/CC vs. GG	1.9 (1.24–2.92)	0.003	1.71 (1.06–2.74)	0.03	2.6 (1.01–3.78)	0.01
55/5X vs. XX	0.88 (0.58–1.33)	0.54	0.87 (0.54–1.39)	0.56	0.66 (0.31–1.43)	0.30
66/6X vs. XX	0.98 (0.57–1.68)	0.93	1.05 (0.57–1.93)	0.89	1.48 (0.50–4.38)	0.48
77/7X vs. XX	1.89 (1.21–2.96)	0.005	1.41 (0.98–2.63)	0.06	2.27 (1.07–4.83)	0.03

The analyses were performed by using a univariate logistic regression model assuming a dominant mode of inheritance.

before or after the lumbar puncture (median [range]: 2.49 [0.02–323.22] vs. 3.49 [0.02–68.34] ng/mL, $P = 0.22$). MIF CSF levels were markedly higher in nonsurvivors than in survivors (10.80 [3.29–89.11] vs. 3.19 [0.02–323.22] ng/mL, $P = 0.0002$) and also higher in patients with than in patients without unfavorable outcome ($P = 0.0001$), respiratory failure ($P = 0.006$), or systemic complications ($P = 0.0004$) (Fig. S1).

***S. pneumoniae* Induces MIF Production in Whole Blood and MIF Promoter Activity in THP-1 Monocytes.** Given that *MIF* polymorphisms and MIF levels in the CSF were associated with morbidity and mortality of pneumococcal meningitis, we examined whether *S. pneumoniae* up-regulated blood levels of MIF and whether polymorphic *MIF* alleles affected the transcriptional activity of the MIF promoter. *S. pneumoniae* induced a strong and dose-dependent up-regulation of MIF production in human whole blood (Fig. 2A). After 24 h, the median concentrations of MIF in cell culture supernatants increased 2.4-, 3.8-, and 15.5-fold upon stimulation with 2×10^6 , 2×10^7 , and 2×10^8 CFU/mL *S. pneumoniae* ($P = 0.005$ and 0.001 , and $P < 10^{-8}$ compared with unstimulated control cells). We then evaluated the transcriptional activity of the four most common polymorphic MIF promoters (CATT₅/–173 G, CATT₆/–173 G, CATT₆/–173 C, and CATT₇/–173 C) in human THP-1 monocytes (Fig. 2B). The transcriptional activity was up-regulated after stimulation with *S. pneumoniae* and was highest with the CATT₆/–173 C and CATT₇/–173 C constructs ($P = 0.03$ and 0.003 compared with CATT₅/–173 G).

Anti-MIF Antibody Protects Against Lethal Pneumococcal Pneumonia and Sepsis. Given that high-expression *MIF* alleles and high MIF CSF levels were associated with poor patient outcome, we tested the impact of an anti-MIF treatment strategy in a mouse model of pneumococcal pneumonia and sepsis (Fig. 3). Compared with mice treated with control antibody, mice treated with anti-MIF antibody had lower bacterial counts in the lung (median [range] log CFU per gram of tissue: 3.8 [1–9.2] vs. 1 [1–7.9], $P = 0.008$) and higher survival (25% vs. 53.6%, $P = 0.04$), indicating that neutralization of MIF was associated with enhanced bacterial clearance and substantial survival benefit in this lethal model of pneumococcal sepsis.

Discussion

In this large, nationwide, and well-characterized cohort of patients with community-acquired pneumococcal meningitis, high-expression *MIF* alleles and elevated levels of MIF in the CSF were identified as predictors of morbidity and mortality. These results suggest that a vigorous MIF response is detrimental in patients with pneumococcal meningitis, a finding consistent with the harmful consequences of robust proinflammatory cytokine responses on brain edema and neuronal damage in the course of bacterial meningitis (26).

Up to now, studies of polymorphisms of cytokine genes have failed to demonstrate association between *TNF*, *IL6*, *IL10*, or *LTA* and the susceptibility to or the outcome of bacterial meningitis (2). MIF is therefore, to our knowledge, the first cytokine whose gene polymorphisms are associated with morbidity and mortality of bacterial meningitis. MIF may exert its effects by

promoting the migration, tissue seeding, and survival of myeloid cells (such as neutrophils and monocytes) that are important players in the pathogenesis of bacterial meningitis (26, 27). At a cellular level, MIF up-regulates proinflammatory responses of immune cells via the activation the MAPK and PI3K/Akt signaling pathways downstream of the CD74–CD44 and CXCR2–CXCR4 MIF receptor complex, which is likely to be detrimental in patients with bacterial meningitis (26, 27).

Few and fairly small studies have looked at MIF in patients with pneumococcal diseases (28, 29). In one study, elevated levels of MIF were detected in the CSF of 31 patients with bacterial meningitis (29). Albeit higher in 11 patients with pneumococcal meningitis than in 15 patients with meningococcal meningitis, MIF CSF levels were not associated with patient outcome. In a study that included 15 patients with pneumococcal meningitis, the frequency of the CATT₇ allele, but not that of the –173 C allele, was higher in patients with meningitis than in those with non-CNS pneumococcal infections (28). Associations between *MIF* genotypes and other clinical variables were not reported. Moreover, given the absence of uninfected control subjects, it was not possible to determine whether the carriage of a CATT₇ allele was a risk factor for the development of pneumococcal meningitis. The present study included a large number of patients and had sufficient power to examine these questions. Neither the CATT_{5–8} nor the –173 G/C *MIF* polymorphisms was a risk factor for susceptibility to pneumococcal meningitis. However, high-expression *MIF* alleles and high levels of MIF levels in the CSF were associated with markers of inflammation, indicators of disease severity, complications, and mortality.

Studies exploring the relationship between functional polymorphisms of the *MIF* gene locus and susceptibility to, severity of, and outcome of infectious and autoimmune diseases have yielded a complex global picture (20, 25). In clinical studies of infectious diseases, carriage of high-expression *MIF* alleles (CATT_{7/8}) or haplotypes (CATT_{6/7}/–173 C) was associated with severe complications of malaria (30), with mortality in patients




Fig. 1. Carriage of –173 C or CATT₇ high-expression *MIF* allelic variants is associated with death from pneumococcal meningitis. Proportions of survivors (open bars) and nonsurvivors (filled bars) for the –173 (G/G, G/C, and C/C) and CATT₇ (X/X, 7/X, and 7/7) genotypes. Logistic regression analysis was performed using dominant and additive modes of inheritance ($P = 0.01$ and 0.01 for –173 G/C and $P = 0.03$ and 0.01 for CATT₇, respectively).

Table 3. Logistic regression analysis of factors associated with death and unfavorable outcome

Prediction	OR (95% CI)	P
Death (n = 30/398)		
Shock	26.06 (5.36–48.09)	0.02
Carriage of CATT7 allele	5.12 (1.11–23.67)	0.04
Immunocompromised state	2.85 (1.16–6.98)	0.09
Male gender	2.48 (0.99–6.17)	0.05
Age (per year increase)	1.04 (1.01–1.08)	0.02
Unfavorable outcome (n = 133/405)		
Systemic complications on admission	15.8 (1.80–137.97)	0.01
Otitis or sinusitis	3.79 (2.21–6.49)	<0.001
Carriage of CATT7 allele	2.61 (0.91–7.50)	0.07
Male gender	1.67 (0.96–2.91)	0.07
Age (per year increase)	1.01 (0.99–1.03)	0.18
GCS > 12	0.54 (0.29–1.01)	0.06

The analyses were performed by using multivariate logistic regression models. GCS, Glasgow coma score.

with severe sepsis (31), and with survival in patients with community-acquired pneumonia (32). Carriage of the low-expression *MIF* allele (CATT₅) protected children from meningococemia (33), but it predisposed older adults to Gram-negative bacteremia (34). Notwithstanding the possibility that the impact of *MIF* polymorphisms may vary according to the age of the host, the site of infections, and the type of microorganism, it also is possible that confounding factors (such as selection biases, ambiguous phenotypes, patient and pathogen heterogeneity, and lack of power) account for these discrepant findings (2, 35, 36). To minimize the influence of potential confounders, we elected to conduct our study in a large and homogeneous cohort of patients with community-acquired bacterial meningitis caused by a single pathogen (*S. pneumoniae*). One of the strengths of the study was the use of a well-defined phenotype. Working with a nationwide cohort also protected against the potential risk of single-center biases. The choice of controls (patient's partners or proxies) limited the risk of socioeconomic and environmental mismatching. Power calculations indicated that the study had 80% power to detect ORs in the range of 1.5–3.0 for the main study endpoints.

The study also has some limitations. It included only Caucasians. Given that *MIF* polymorphisms are unequally distributed in different populations (20), our findings therefore might not be extrapolated to other races or ethnic groups. Blood samples were not available to measure MIF in the systemic circulation. However, one study looking at MIF levels in patients with CNS

infections did not find an association between MIF levels in the CSF and inflammation biomarkers in the systemic circulation (37–40). Attempts to correlate *MIF* polymorphisms with MIF CSF levels were unsuccessful. This result may be due to the relatively limited number of CSF and DNA specimen pairs. Additionally, the range of MIF CSF levels was rather broad, likely because lumbar punctures were performed in patients at different stages of the disease. It is also conceivable that carriage of a high-expression *MIF* allele plays a detrimental role at the very early stage of disease, promoting inflammation, blood–brain barrier disruption, and bacterial neuro-invasion, as shown both for MIF or IFN lambda in West Nile encephalitis (41, 42). MIF-driven early pathogenic effects may indeed not be correlated with CSF parameters measured just once at a later stage of the disease. In that scenario, both *MIF* polymorphisms and MIF expression levels might be independent markers of severity at different stages of the disease. Finally, given that MIF is constitutively expressed by innate immune cells and markedly up-regulated upon infection, the release of intracellular MIF pools from dying cells may have increased MIF levels in the CSF, compromising pairwise analyses of *MIF* gene polymorphisms and MIF protein levels.

A large body of experimental and clinical evidence indicates that MIF is an upstream mediator of the host antimicrobial defense response and a powerful immunomodulating cytokine. It has therefore been investigated as a target for therapeutic interventions in infectious, inflammatory, and autoimmune diseases (19, 32–34, 37–40, 43–45). The observation that *MIF* is a genetic marker of morbidity and mortality of pneumococcal meningitis is in line with the results obtained in animal models of sepsis and in clinical studies that have linked high levels of MIF with life-threatening bacterial infections (21, 22, 46–49). In a murine model of pneumococcal colonization, the clearance of pneumococci from the nasopharynx, a critical step in the pathogenesis of invasive pneumococcal diseases, was more rapid in wild-type mice than in MIF-deficient mice (45). This finding suggested that MIF might be a protective factor against pneumococcal diseases. However, in the present study, there was no relationship between *MIF* expression and susceptibility to pneumococcal meningitis. In a model of pneumococcal pneumonia, wild-type mice had higher bacterial load, higher innate immune cell counts in the lung parenchyma, more severe lung pathology, and a higher mortality rate than MIF-deficient mice (50), indicating that MIF played a detrimental role in severe pneumococcal infection. Consistent with this observation, high-expression *MIF* alleles and MIF CSF levels were predictors of disease severity and poor outcome in patients with pneumococcal meningitis. Furthermore, anti-MIF therapy using either neutralizing antibodies (present study) or a small-molecule inhibitor of MIF (50) promoted bacterial clearance and increased survival in a lethal pneumococcal




Fig. 2. *S. pneumoniae* induces MIF production in whole blood and polymorphic *MIF* alleles affect the transcriptional activity of the *MIF* promoter. (A) Whole blood from five healthy volunteers was stimulated for 24 h with *S. pneumoniae*. MIF concentrations were measured by ELISA. Bottom, median, and top lines of the box mark the 25th, 50th, and 75th percentiles, respectively. Vertical lines with whiskers show the range of values. $P = 0.005$ (2×10^6), $P = 0.001$ (2×10^7), and $P < 10^{-8}$ (2×10^8) compared with control. (B) THP-1 monocytes were transiently transfected with CATT₅/–173 G, CATT₆/–173 G, CATT₆/–173 C, and CATT₇/–173 C MIF promoter pGL3 and *Renilla* luciferase vectors. Cells were stimulated for 24 h with (filled bars) or without (open bars) *S. pneumoniae* (MOI 10). Data are means + SD of three independent experiments performed in triplicate. $P = 0.05$ (CATT₆/–173 G), $P = 0.04$ (CATT₆/–173 C), and $P = 0.007$ (CATT₇/–173 C) for *S. pneumoniae* vs. control. After *S. pneumoniae* stimulation, $P = 0.03$ (CATT₆/–173 C) and $P = 0.003$ (CATT₇/–173 C) compared with CATT₅/–173 G.




Fig. 3. Anti-MIF antibody protects against lethal pneumococcal pneumonia and sepsis. Mice were treated with anti-MIF or nonimmune (control) IgG (2 mg injected intraperitoneally) given 2 h before an intranasal inoculation of 10^4 CFU of *S. pneumoniae*. (A) Box plots of bacterial counts in the lung 48 h after infection. Data are from five independent experiments with a total of 34 mice per treatment group. Bottom, median, and top lines of the box mark the 25th, 50th, and 75th percentiles, respectively. Vertical lines with whiskers show the range of values. $P = 0.008$ (Mann–Whitney u test). (B) Kaplan–Meier survival plot. Data points are from three independent experiments with a total of 28 mice per treatment group. $P = 0.04$ (log-rank test).

sepsis. Likewise, humanized anti-MIF antibodies were shown to be protective in a lethal mouse *Escherichia coli* peritonitis model (51). A first-in-class anti-MIF monoclonal antibody (BAX69/imalumab) is currently under evaluation in phase 1/2a clinical trials in patients with metastatic colorectal cancer or with ovarian cancer and malignant ascites.

All together, the present study provides strong evidence that carriage of high-expression *MIF* alleles is a genetic marker of morbidity and mortality of pneumococcal meningitis and also suggests a potential role for MIF as a target of immune-modulating adjunctive therapies for bacterial meningitis and sepsis.

Materials and Methods

Bacteria Preparation and Whole-Blood Stimulation Assay. A serotype 19 *S. pneumoniae* strain (kind gift of Gilbert Greub, Institute of Microbiology, Centre Hospitalier Universitaire Vaudois and University of Lausanne) isolated from the CSF of a patient with acute community-acquired meningitis was used in all stimulation experiments. Bacteria were grown to log phase in brain–heart infusion medium, washed, and resuspended in PBS at an OD_{620} of 1.0. Whole blood from five healthy Caucasian subjects was drawn into S-monovette-tubes (Sarstedt) containing 16 U of heparin per mL, diluted fivefold in phenol red-free RPMI medium 1640 containing 2 mM L-glutamine (Life Technologies) (52). Blood was incubated at 37 °C with *S. pneumoniae* (53). Penicillin and streptomycin (100 μ g/mL) were added to the cultures 1 h after stimulation. Cells were pelleted ($500 \times g$ for 5 min at room temperature) 24 h after stimulation and cell-free supernatants were used to measure MIF concentrations by ELISA (R&D Systems). The lower limit of detection of the assay was 15.6 pg/mL lactate dehydrogenase (LDH) was measured by using the International Federation of Clinical Chemistry method to evaluate cell death. LDH concentration in samples (9.5–66 units/L) remained in the normal range (i.e., <250 units/L).

Patient Cohort and Controls. After identification by the Netherlands Reference Laboratory for Bacterial Meningitis (Academic Medical Center, Amsterdam) patients >16 y of age and with a positive CSF culture were enrolled between March 2006 and June 2011 in a nationwide prospective cohort study (54). The study was approved by the medical ethics committee of the University of Amsterdam. Treating physicians obtained informed written consent from either the patient or their legal representatives. Patients with culture-negative community-acquired or with hospital-acquired bacterial meningitis were excluded. Controls consisted of co-dwelling nonrelated patient proxies who shared environment and socioeconomic milieu and provided written informed consent (55). Demographic data including age, gender, and ethnicity for both controls and patients were collected. A detailed medical history and

extensive clinical and laboratory data were obtained from all patients. Outcome was evaluated at discharge according to the GOS, with the following scoring system: score of 1 for death; score of 2 for persistent vegetative state; score of 3 for severe disability, defined as a conscious patient who is dependent for daily activities; score of 4 for moderate disability, defined as a patient with some deficits, capable of living independently, but unable to return to work; and score of 5 for a good recovery (56). A favorable outcome was defined as a score of 5 and an unfavorable outcome as a score of 1–4. A CSF leukocyte count <1,000 per mm^3 was used as an indicator of disease’s severity (57).

Blood Collection and Genotyping. Blood from patients and controls was collected in sodium EDTA. DNA extraction was performed by the Gentra Puregene isolation kit (Qiagen) with quality control evaluation for DNA yield and purity. The –794 CATT_{5–8} microsatellite (rs5844572) and the –173 G/C SNP (rs755622) were genotyped as reported (33). Given the frequencies of the –173 C minor allele and of the CATT₇ allele in the Caucasian population, the study had 80% power to detect ORs of at least 1.5 for the susceptibility to infection, 1.9 for an unfavorable outcome, and 3.0 for mortality for these two alleles. These numbers were calculated by using the Quanto 1.2.3 software (biostats.usc.edu/software).

CSF Collection and MIF Measurement. CSF was obtained from the first lumbar puncture and centrifuged, and supernatant was aliquoted and stored at –80 °C until analysis. Human MIF levels in the CSF were measured by the Luminex technology using a Milliplex assay (Millipore). The lower limit of detection of the assay was 12 pg/mL.

MIF Promoter Activity. A fragment ranging from –1,073 to –129 bp of the *MIF* gene of the –794 CATT₅–173 G, –794 CATT₆–173 G, –794 CATT₆–173 C, or –794 CATT₇–173 C genotype was cloned in the pGL3-basic luciferase vector (Promega) (58). The human monocytic THP-1 (TIB-202; American Type Culture Collection) cell line was cultured in RPMI medium 1640 containing 2 mM L-glutamine, 50 μ M 2-mercaptoethanol, and 10% (vol/vol) heat-inactivated FCS (Sigma-Aldrich). A total of 40,000 THP-1 cells in 96-well plates were transfected with 350 ng of the luciferase reporter vector together with 50 ng of the *Renilla* pRL-TK vector (Promega) (59). After 7 h, cells were incubated with *S. pneumoniae* [multiplicity of infection (MOI) 10]. Penicillin and streptomycin (100 μ g/mL) were added to the cultures 1 h after stimulation, and incubation was continued for an additional 23 h. Luciferase and *Renilla* luciferase activities were measured by using the Dual-Luciferase Reporter Assay System (Promega). Results were expressed as the ratio of luciferase activity to *Renilla* luciferase activity. An empty vector served as a negative control (18).

Mouse Model of Pneumococcal Pneumonia and Sepsis. Animal experiments were approved by the Office Vétérinaire du Canton de Vaud (authorization no. 877.8) and performed according to the institution and Animal Research: Reporting of *In Vivo* Experiments guidelines (www.nc3rs.org.uk/arrive-guidelines). Female BALB/cByJ mice (10 wk old; Charles River Laboratories) were housed under specific pathogen-free conditions. Mice (8–10 per treatment group) were injected intraperitoneally with 2 mg of rabbit anti-MIF or nonimmune (control) IgG (obtained as described in ref. 37) administered 2 h before an intranasal inoculation of 10^8 CFU of *S. pneumoniae* in anesthetized mice. Survival was monitored at least twice daily until the end of the experiment. In selected experiments, mice were killed 48 h after infection to obtain blood and lung tissue for the measurements of bacterial counts.

Statistical Analyses. Graphs were generated with GraphPad Prism (Version 6.0; GraphPad Software). Data were analyzed by using STATA (Version 11.1; Stata-Corp LP) and IBM SPSS Statistics (Version 21.0; IBM). Genotype frequencies among patients and controls and HWE were calculated by using the χ^2 test. Tests for associations between *MIF* genotypes and susceptibility to and severity of pneumococcal meningitis were performed by using logistic regression models. Stepwise multivariate selection was performed by using a P value > 0.2 for removal of variables in the model. Sex, age, and polymorphism carriage were forced into the model. Polymorphisms were analyzed by using dominant (comparison of heterozygotes plus homozygotes vs. wild-type carriers) and additive modes of inheritance (assessment of the effect of each additional copy of the minor allele). Strength of relationships between CSF levels and continuous variables was evaluated by using Spearman’s correlation tests. Dichotomous variables were compared with the χ^2 test. Continuous variable for patient’s characteristics and disease parameters were assessed by using the Mann–Whitney U or Kruskal–Wallis tests. MIF promoter activity and concentrations of MIF released in whole blood after stimulation with *S. pneumoniae* were analyzed by using one-way ANOVA for multiple comparisons. The log-rank test was used to compare the Kaplan–Meier survival curves. All tests were two-tailed, and P values < 0.05 were considered to indicate statistical significance.

ACKNOWLEDGMENTS. We thank Agnieszka Wójtowicz for power calculations. This work was supported by Swiss National Science Foundation Grants 138488 (to T.C.), 145014 and 149511 (to T.R.), and 144054 (to P.-Y.B.); European Research Council Starting Grant 281156 (to D.v.d.B.); Netherlands

Organization for Health Research and Development Vidi Grant 016.116.358 (to D.v.d.B.); and Netherlands Organisation for Scientific Research Veni Grant 916.13.078 (to M.C.B.). A.S. was supported by the Porphyrogenis Foundation.

- Lozano R, et al. (2012) Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380(9859):2095–2128.
- Brouwer MC, et al. (2009) Host genetic susceptibility to pneumococcal and meningococcal disease: A systematic review and meta-analysis. *Lancet Infect Dis* 9(1):31–44.
- Brouwer MC, Tunkel AR, van de Beek D (2010) Epidemiology, diagnosis, and antimicrobial treatment of acute bacterial meningitis. *Clin Microbiol Rev* 23(3):467–492.
- Thigpen MC, et al.; Emerging Infections Programs Network (2011) Bacterial meningitis in the United States, 1998–2007. *N Engl J Med* 364(21):2016–2025.
- Edmond K, et al. (2010) Global and regional risk of disabling sequelae from bacterial meningitis: A systematic review and meta-analysis. *Lancet Infect Dis* 10(5):317–328.
- Heckenberg SG, Brouwer MC, van der Ende A, Hensen EF, van de Beek D (2012) Hearing loss in adults surviving pneumococcal meningitis is associated with otitis and pneumococcal serotype. *Clin Microbiol Infect* 18(9):849–855.
- Jit M (2010) The risk of sequelae due to pneumococcal meningitis in high-income countries: A systematic review and meta-analysis. *J Infect* 61(2):114–124.
- Casanova JL, Abel L, Quintana-Murci L (2011) Human TLRs and IL-1Rs in host defense: Natural insights from evolutionary, epidemiological, and clinical genetics. *Annu Rev Immunol* 29:447–491.
- Kasanmoentalib ES, Brouwer MC, van de Beek D (2013) Update on bacterial meningitis: Epidemiology, trials and genetic association studies. *Curr Opin Neurol* 26(3):282–288.
- Chapman SJ, et al. (2006) PTPN22 and invasive bacterial disease. *Nat Genet* 38(5):499–500.
- Roy S, et al.; Oxford Pneumococcal Surveillance Group (2002) MBL genotype and risk of invasive pneumococcal disease: A case-control study. *Lancet* 359(9317):1569–1573.
- Khor CC, et al. (2007) A Mal functional variant is associated with protection against invasive pneumococcal disease, bacteremia, malaria and tuberculosis. *Nat Genet* 39(4):523–528.
- Chapman SJ, et al. (2007) IkappaB genetic polymorphisms and invasive pneumococcal disease. *Am J Respir Crit Care Med* 176(2):181–187.
- Bloom BR, Bennett B (1966) Mechanism of a reaction in vitro associated with delayed-type hypersensitivity. *Science* 153(3731):80–82.
- David JR (1966) Delayed hypersensitivity in vitro: Its mediation by cell-free substances formed by lymphoid cell-antigen interaction. *Proc Natl Acad Sci USA* 56(1):72–77.
- Calandra T, et al. (1995) MIF as a glucocorticoid-induced modulator of cytokine production. *Nature* 377(6544):68–71.
- Mitchell RA, et al. (2002) Macrophage migration inhibitory factor (MIF) sustains macrophage proinflammatory function by inhibiting p53: Regulatory role in the innate immune response. *Proc Natl Acad Sci USA* 99(1):345–350.
- Roger T, Chanson AL, Knaup-Reymond M, Calandra T (2005) Macrophage migration inhibitory factor promotes innate immune responses by suppressing glucocorticoid-induced expression of mitogen-activated protein kinase phosphatase-1. *Eur J Immunol* 35(12):3405–3413.
- Roger T, David J, Glauser MP, Calandra T (2001) MIF regulates innate immune responses through modulation of Toll-like receptor 4. *Nature* 414(6866):920–924.
- Bucala R (2013) MIF, MIF alleles, and prospects for therapeutic intervention in autoimmunity. *J Clin Immunol* 33(Suppl 1):S72–S78.
- Calandra T, Roger T (2003) Macrophage migration inhibitory factor: A regulator of innate immunity. *Nat Rev Immunol* 3(10):791–800.
- Lue H, Kleemann R, Calandra T, Roger T, Bernhagen J (2002) Macrophage migration inhibitory factor (MIF): Mechanisms of action and role in disease. *Microbes Infect* 4(4):449–460.
- Baugh JA, et al. (2002) A functional promoter polymorphism in the macrophage migration inhibitory factor (MIF) gene associated with disease severity in rheumatoid arthritis. *Genes Immun* 3(3):170–176.
- Donn RP, Shelley E, Ollier WE, Thomson W; British Paediatric Rheumatology Study Group (2001) A novel 5'-flanking region polymorphism of macrophage migration inhibitory factor is associated with systemic-onset juvenile idiopathic arthritis. *Arthritis Rheum* 44(8):1782–1785.
- Renner P, Roger T, Calandra T (2005) Macrophage migration inhibitory factor: Gene polymorphisms and susceptibility to inflammatory diseases. *Clin Infect Dis* 41(Suppl 7):S513–S519.
- Mook-Kanamori BB, Geldhoff M, van der Poll T, van de Beek D (2011) Pathogenesis and pathophysiology of pneumococcal meningitis. *Clin Microbiol Rev* 24(3):557–591.
- Bernhagen J, et al. (2007) MIF is a noncognate ligand of CXCR2 chemokine receptors in inflammatory and atherogenic cell recruitment. *Nat Med* 13(5):587–596.
- Doernberg S, et al. (2011) Association of macrophage migration inhibitory factor (MIF) polymorphisms with risk of meningitis from *Streptococcus pneumoniae*. *Cytokine* 53(3):292–294.
- Østergaard C, Benfield T (2009) Macrophage migration inhibitory factor in cerebrospinal fluid from patients with central nervous system infection. *Crit Care* 13(3):R101.
- Awandare GA, et al. (2009) MIF (macrophage migration inhibitory factor) promoter polymorphisms and susceptibility to severe malarial anemia. *J Infect Dis* 200(4):629–637.
- Lehmann LE, et al. (2009) A MIF haplotype is associated with the outcome of patients with severe sepsis: A case control study. *J Transl Med* 7:100.
- Yende S, et al. (2009) The influence of macrophage migration inhibitory factor gene polymorphisms on outcome from community-acquired pneumonia. *FASEB J* 23(8):2403–2411.
- Renner P, et al. (2012) A functional microsatellite of the macrophage migration inhibitory factor gene associated with meningococcal disease. *FASEB J* 26(2):907–916.
- Das R, et al. (2014) Functional polymorphisms in the gene encoding macrophage migration inhibitory factor are associated with Gram-negative bacteremia in older adults. *J Infect Dis* 209(5):764–768.
- Bochud PY, Bochud M, Telenti A, Calandra T (2007) Innate immunogenetics: A tool for exploring new frontiers of host defence. *Lancet Infect Dis* 7(8):531–542.
- Chapman SJ, Hill AV (2012) Human genetic susceptibility to infectious disease. *Nat Rev Genet* 13(3):175–188.
- Calandra T, et al. (2000) Protection from septic shock by neutralization of macrophage migration inhibitory factor. *Nat Med* 6(2):164–170.
- Beishuizen A, Thijs LG, Haanen C, Vermes I (2001) Macrophage migration inhibitory factor and hypothalamo-pituitary-adrenal function during critical illness. *J Clin Endocrinol Metab* 86(6):2811–2816.
- Emonts M, et al. (2007) Association between high levels of blood macrophage migration inhibitory factor, inappropriate adrenal response, and early death in patients with severe sepsis. *Clin Infect Dis* 44(10):1321–1328.
- Sprong T, et al. (2007) Macrophage migration inhibitory factor (MIF) in meningococcal septic shock and experimental human endotoxemia. *Shock* 27(5):482–487.
- Arjona A, et al. (2007) Abrogation of macrophage migration inhibitory factor decreases West Nile virus lethality by limiting viral neuroinvasion. *J Clin Invest* 117(10):3059–3066.
- Lazear HM, et al. (2015) Interferon- λ restricts West Nile virus neuroinvasion by tightening the blood-brain barrier. *Sci Transl Med* 7(284):284ra59.
- Calandra T, Spiegel LA, Metz CN, Bucala R (1998) Macrophage migration inhibitory factor is a critical mediator of the activation of immune cells by exotoxins of Gram-positive bacteria. *Proc Natl Acad Sci USA* 95(19):11383–11388.
- Roger T, et al. (2013) Macrophage migration inhibitory factor deficiency is associated with impaired killing of gram-negative bacteria by macrophages and increased susceptibility to *Klebsiella pneumoniae* sepsis. *J Infect Dis* 207(2):331–339.
- Das R, et al. (2014) Macrophage migration inhibitory factor promotes clearance of pneumococcal colonization. *J Immunol* 193(2):764–772.
- Bernhagen J, et al. (1993) MIF is a pituitary-derived cytokine that potentiates lethal endotoxaemia. *Nature* 365(6448):756–759.
- Lehmann LE, et al. (2001) Plasma levels of macrophage migration inhibitory factor are elevated in patients with severe sepsis. *Intensive Care Med* 27(8):1412–1415.
- Roger T, Glauser MP, Calandra T (2001) Macrophage migration inhibitory factor (MIF) modulates innate immune responses induced by endotoxin and Gram-negative bacteria. *J Endotoxin Res* 7(6):456–460.
- Calandra T, Froidevaux C, Martin C, Roger T (2003) Macrophage migration inhibitory factor and host innate immune defenses against bacterial sepsis. *J Infect Dis* 187(Suppl 2):S385–S390.
- Weiser JN, et al. (2015) Macrophage migration Inhibitory factor is detrimental in pneumococcal pneumonia and a target for therapeutic immunomodulation. *J Infect Dis* 212(10):1677–1682.
- Kerschbaumer RJ, et al. (2012) Neutralization of macrophage migration inhibitory factor (MIF) by fully human antibodies correlates with their specificity for the β -sheet structure of MIF. *J Biol Chem* 287(10):7446–7455.
- Roger T, et al. (2011) Histone deacetylase inhibitors impair innate immune responses to Toll-like receptor agonists and to infection. *Blood* 117(4):1205–1217.
- Lugrin J, et al. (2009) Histone deacetylase inhibitors repress macrophage migration inhibitory factor (MIF) expression by targeting MIF gene transcription through a local chromatin deacetylation. *Biochim Biophys Acta* 1793(11):1749–1758.
- Woehrl B, et al. (2011) Complement component 5 contributes to poor disease outcome in humans and mice with pneumococcal meningitis. *J Clin Invest* 121(10):3943–3953.
- Little J, et al. (2002) Reporting, appraising, and integrating data on genotype prevalence and gene-disease associations. *Am J Epidemiol* 156(4):300–310.
- Jennett B, Bond M (1975) Assessment of outcome after severe brain damage. *Lancet* 1(7905):480–484.
- van de Beek D, et al. (2004) Clinical features and prognostic factors in adults with bacterial meningitis. *N Engl J Med* 351(18):1849–1859.
- Roger T, Ding X, Chanson AL, Renner P, Calandra T (2007) Regulation of constitutive and microbial pathogen-induced human macrophage migration inhibitory factor (MIF) gene expression. *Eur J Immunol* 37(12):3509–3521.
- Delaloye J, et al. (2009) Innate immune sensing of modified vaccinia virus Ankara (MVA) is mediated by TLR2-TLR6, MDA-5 and the NALP3 inflammasome. *PLoS Pathog* 5(6):e1000480.