

# Complete Genome Sequence of Methanogenic Archaeon ISO4-G1, a Member of the *Methanomassiliicoccales*, Isolated from a Sheep Rumen

William J. Kelly, Dong Li, Suzanne C. Lambie, Jeyamalar Jeyanathan, Faith Cox, Yang Li, Graeme T. Attwood, Eric Altermann, Sinead C. Leahy

AgResearch Limited, Grasslands Research Centre, Palmerston North, New Zealand

**Methanogenic archaeon ISO4-G1 is a methylotrophic methanogen belonging to the order *Methanomassiliicoccales* that was isolated from a sheep rumen. Its genome has been sequenced to provide information on the genetic diversity of rumen methanogens in order to develop technologies for ruminant methane mitigation.**

Received 16 February 2016 Accepted 24 February 2016 Published 7 April 2016

**Citation** Kelly WJ, Li D, Lambie SC, Jeyanathan J, Cox F, Li Y, Attwood GT, Altermann E, Leahy SC. 2016. Complete genome sequence of methanogenic archaeon ISO4-G1, a member of the *Methanomassiliicoccales*, isolated from a sheep rumen. *Genome Announc* 4(2):e00221-16. doi:10.1128/genomeA.00221-16.

**Copyright** © 2016 Kelly et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sinead C. Leahy, [sinead.leahy@agresearch.co.nz](mailto:sinead.leahy@agresearch.co.nz).

Members of the order *Methanomassiliicoccales* are methylotrophic methanogens first detected in the rumen (1, 2) and subsequently in a variety of other anaerobic environments (3–5). Currently, the only complete genome sequence available for a rumen isolate is from *Thermoplasmatales* archaeon BRNA1 (NCBI reference sequence: NC\_020892.1). Here, we report the genome sequence of an isolate from a sheep rumen, which is designated as methanogenic archaeon ISO4-G1 and belongs to a different group within the order *Methanomassiliicoccales* (6, 7).

The complete genome sequence of ISO4-G1 was determined by pyrosequencing 3-kb mate-paired libraries on a 454 GS FLX platform with titanium chemistry combined with reads from a MiSeq 2 × 300-bp sequencing run (Macrogen, South Korea). Pyrosequencing reads were assembled using the Newbler assembler version 2.7 (Roche 454 Life Sciences, USA) and combined with the MiSeq data using the SPAdes assembler version 3.0 (8), resulting in 4 contigs in a single scaffold. Gap closure was managed using the Staden package (9), and gaps were closed using standard PCR techniques with Sanger sequencing. Protein-encoding genes were identified by Glimmer (10), and a GAMOLA/ARTEMIS (11, 12) software suite was used to manage genome annotation. Assignment of protein function to open reading frames was performed manually using results from BLASTp and the COG (Clusters of Orthologous Groups), Pfam, and TIGRFAM databases (13–15).

The genome sequence of methanogenic archaeon ISO4-G1 consists of a single 1,593,503-bp circular chromosome, with a GC content of 55.5%, and 1,501 predicted protein-coding genes representing 92.0% of the genome. The ISO4-G1 genome does not contain plasmid, prophage, or CRISPR sequences. Analysis of the genome suggests that ISO4-G1 relies on hydrogen-dependent methylotrophic methanogenesis to produce energy, with methanol and methylamines as substrates. The complement of methane formation genes is very similar to that reported for “*Candidatus Methanoplasma termitum*” (16). Genes for the production of coenzyme M and tryptophan biosynthesis are missing from the genome. The genome encodes a large number of transporters, in-

cluding 15 ABC transporters predicted to be involved in Fe<sup>3+</sup> or siderophore uptake. Like the other members of the order *Methanomassiliicoccales*, ISO4-G1 has a complete operon (AUP07\_0651–654) predicted to encode pyrrolysine biosynthesis together with a specific aminoacyl-tRNA synthetase that enables read-through of the amber stop codon UAG (5, 17). A total of 25 genes encoding pyrrolysine-containing proteins were identified in the ISO4-G1 genome, 9 of which were mono-/di-/tri-methylamine: corrinoid methyltransferases, as reported for other members of the *Methanomassiliicoccales* (5). ISO4-G1 also has a gene (AUP07\_0971) predicted to encode a unique pyrrolysine-containing nonribosomal peptide synthase (5,216 amino acid residues) whose function is unknown. Genomic information from this group of organisms will complement genome sequences from other rumen methanogens and will be used to design strategies aimed at reducing methane emissions from ruminant livestock (18).

**Nucleotide sequence accession number.** This whole-genome sequencing project has been deposited at DDBJ/EMBL/GenBank under the accession number CP013703.

## ACKNOWLEDGMENTS

The ISO4-G1 genome sequencing project was supported by the New Zealand Pastoral Greenhouse Gas Research Consortium (PGgRc) and the New Zealand Agricultural Greenhouse Gas Research Centre (NZAGRC).

## FUNDING INFORMATION

This work, including the efforts of William J Kelly, Dong Li, Suzanne C Lambie, Jeyamalar Jeyanathan, Faith Cox, Yang Li, Graeme Trevor Attwood, Eric Altermann, and Sinead Christine Leahy, was funded by the New Zealand Pastoral Greenhouse Gas Research Consortium and the New Zealand Agricultural Greenhouse Gas Research Centre.

The funders had no role in study design or data collection or interpretation.

## REFERENCES

1. Tajima K, Nagamine T, Matsui H, Nakamura M, Aminov RI. 2001. Phylogenetic analysis of archaeal 16S rRNA libraries from the rumen sug-

- gests the existence of a novel group of archaea not associated with known methanogens. *FEMS Microbiol Lett* 200:67–72. <http://dx.doi.org/10.1111/j.1574-6968.2001.tb10694.x>.
2. Janssen PH, Kirs M. 2008. Structure of the archaeal community of the rumen. *Appl Environ Microbiol* 74:3619–3625. <http://dx.doi.org/10.1128/AEM.02812-07>.
  3. Paul K, Nonoh JO, Mikulski L, Brune A. 2012. “*Methanoplasmatales*,” *Thermoplasmatales*-related archaea in termite guts and other environments, are the seventh order of methanogens. *Appl Environ Microbiol* 78:8245–8253. <http://dx.doi.org/10.1128/AEM.02193-12>.
  4. Iino T, Tamaki H, Tamazawa S, Ueno Y, Ohkuma M, Suzuki K, Igarashi Y, Haruta S. 2013. *Candidatus* Methanogranum caenicola: a novel methanogen from the anaerobic digested sludge, and proposal of *Methanomassiliicoccaceae* fam. nov. and *Methanomassiliicoccales* ord. nov., for a methanogenic lineage of the class *Thermoplasmata*. *Microbes Environ* 28:244–250. <http://dx.doi.org/10.1264/j sme2.ME12189>.
  5. Borrel G, Parisot N, Harris HM, Peyretailade E, Gaci N, Tottey W, Bardot O, Raymann K, Gribaldo S, Peyret P, O’Toole PW, Brugère JF. 2014. Comparative genomics highlights the unique biology of *Methanomassiliicoccales*, a *Thermoplasmatales*-related seventh order of methanogenic archaea that encodes pyrrolysine. *BMC Genomics* 15:679. <http://dx.doi.org/10.1186/1471-2164-15-679>.
  6. Jeyanathan J. 2010. Investigation of rumen methanogens in New Zealand livestock. Ph.D. thesis. Massey University, Palmerston North, New Zealand.
  7. Seedorf H, Kittelmann S, Henderson G, Janssen PH. 2014. Rim-DB: a taxonomic framework for community structure analysis of methanogenic archaea from the rumen and other intestinal environments. *PeerJ* 2:e494. <http://dx.doi.org/10.7717/peerj.494>.
  8. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
  9. Staden R, Beal KF, Bonfield JK. 2000. The Staden package, 1998. *Methods Mol Biol* 132:115–130.
  10. Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res* 27:4636–4641. <http://dx.doi.org/10.1093/nar/27.23.4636>.
  11. Altermann E, Klaenhammer TR. 2003. GAMOLA: a new local solution for sequence annotation and analyzing draft and finished prokaryotic genomes. *Omics* 7:161–169. <http://dx.doi.org/10.1089/15362310332246557>.
  12. Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945. <http://dx.doi.org/10.1093/bioinformatics/16.10.944>.
  13. Tatusov RL, Galperin MY, Natale DA, Koonin EV. 2000. The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res* 28:33–36. <http://dx.doi.org/10.1093/nar/28.1.33>.
  14. Punta M, Coghill PC, Eberhardt RY, Mistry J, Tate J, Boursnell C, Pang N, Forslund K, Ceric G, Clements J, Heger A, Holm L, Sonnhammer EL, Eddy SR, Bateman A, Finn RD. 2012. The Pfam protein families database. *Nucleic Acids Res* 40:D290–D301. <http://dx.doi.org/10.1093/nar/gkr1065>.
  15. Haft DH, Selengut JD, Richter RA, Harkins D, Basu MK, Beck E. 2013. TIGRFAMs and genome properties in 2013. *Nucleic Acids Res* 41:D387–D395. <http://dx.doi.org/10.1093/nar/gks1234>.
  16. Lang K, Schuldes J, Klingl A, Poehlein A, Daniel R, Brune A. 2015. New mode of energy metabolism in the seventh order of methanogens as revealed by comparative genome analysis of “*Candidatus* Methanoplasma termitum.” *Appl Environ Microbiol* 81:1338–1352. <http://dx.doi.org/10.1128/AEM.03389-14>.
  17. Borrel G, Gaci N, Peyret P, O’Toole PW, Gribaldo S, Brugere JF. 2014. Unique characteristics of the pyrrolysine system in the 7th order of methanogens: implications for the evolution of a genetic code expansion cassette. *Archaea* 2014:374146. <http://dx.doi.org/10.1155/2014/374146>.
  18. Leahy SC, Kelly WJ, Ronimus RS, Wedlock N, Altermann E, Attwood GT. 2013. Genome sequencing of rumen bacteria and archaea and its application to methane mitigation strategies. *Animal* 7(suppl 2):235–243. <http://dx.doi.org/10.1017/S1751731113000700>.