



Published in final edited form as:

*Lung*. 2016 February ; 194(1): 25–29. doi:10.1007/s00408-015-9832-5.

## Airway Vagal Neuroplasticity Associated with Respiratory Viral Infections

Eric J. Zaccone<sup>1</sup> and Bradley J. Udem<sup>1</sup>

<sup>1</sup>Division of Allergy & Clinical Immunology, Department of Medicine, Johns Hopkins University School of Medicine, Johns Hopkins Asthma & Allergy Center, 5501 Hopkins Bayview Circle, Rm. 3A.44, Baltimore, MD 21224-6801, USA

### Abstract

Respiratory virus infections leads to coughing, sneezing, and increases in reflex parasympathetic bronchoconstriction and secretions. These responses to viral infection are exclusively or largely secondary to changes in the function of the nervous system. For many with underlying airway pathologies such as asthma and COPD, this neuroplasticity can lead to disease exacerbations and hospitalization. Relatively little is understood about the cellular and molecular mechanisms that underlie the changes in neuronal control of the respiratory tract during viral infection, but the evidence supports the idea that changes occur in the physiology of both the sensory and autonomic innervation. Virus infection can lead to acute increases in the activity of sensory nerves as well as to genetic changes causing alterations in sensory nerve phenotype. In addition, respiratory viral infections are associated with changes in the control of neurotransmitter release from cholinergic nerve endings terminating at the level of the airway smooth muscle.

### Keywords

Cough; Respiratory virus; TRPV1; Nodose; Airway sensory nerves; Brain-derived neurotrophic growth factor; Vagal afferent

### Introduction

The immune system is effective in clearing most respiratory viral infections within a period of a few weeks. In order to avoid the immune system, viruses have evolved mechanisms to escape the respiratory tract of the host where it can then go on to infect another person. This has been accomplished by causing profound alterations in airway neuronal control, leading to coughing, sneezing, and increases in reflex parasympathetic mucus secretions. The increase in parasympathetic drive can also lead to reflex bronchoconstriction. In most healthy subjects, these symptoms can be trying but are self-limiting and often little more than an inconvenience. In some cases, the symptoms such as urge to cough can persist into a chronic or sub acute state [1, 2]. For those suffering with underlying airway pathologies such

---

<sup>©</sup>Bradley J. Udem, bundem@jhmi.edu.

**Compliance with Ethical Standards**

**Conflict of Interest** None.

as asthma or COPD, the changes in neural control may contribute to acute exacerbations of the disease [1, 3, 4].

Experimental viral infection in laboratory animals [5–11] and in human volunteers [12–15] leads to increases in cough and also increases in bronchial responsiveness to stimuli such as methacholine and histamine. The increase in cough associated with virus infection is consistent with alterations in the sensory nerve function in the airways. The increase in bronchial responsiveness involves multiple mechanisms but changes in both afferent (sensory) and efferent (parasympathetic) nerve function are likely involved.

The hyperresponsiveness associated with airway inflammation in laboratory animals typically requires an intact vagal nervous system. The classical allergen-induced airway hyperresponsiveness to methacholine in mice is entirely prevented in animals in which either the vagus nerves are cut or if only the C-fiber neurons are depleted from the vagal sensory ganglia [16, 17]. Likewise, virus-induced airway hyperresponsiveness of guinea pigs to histamine challenge also requires intact vagus nerves [6, 18]. This indicates that the hyperresponsiveness to histamine and methacholine in these studies is actually a bronchial hyper-*neuronal* responsiveness (Fig. 1).

In human volunteers, experimental viral infections consistently lead to an increase in the bronchial responsiveness to histamine [12, 14]. Histamine effectively sensitizes vagal afferent C-fibers in a manner that would likely lead to increased reflex parasympathetic drive [19]. Indeed, the increase in response to histamine following respiratory viral infection was abolished by cholinergic muscarinic receptor antagonism [12]. Virus infection of rats and guinea pigs can also augment the cholinergic nerve evoked bronchoconstriction, even when the sensory input to the brainstem is bypassed [6, 7, 9]. Considered together, the evidence for increases in urge to cough and bronchial vagal responsiveness associated with viral infections likely involves neural plasticity within both the sensory and autonomic systems. The sensory and autonomic innervation of the respiratory tract has recently been extensively reviewed [20, 21].

## Sensory Nerves

Sensory nerves innervating the airways are responsible for initiating respiratory sensations and reflexes, including coughing and sneezing. In addition by regulating parasympathetic reflex tone, sensory nerves play a major role in controlling mucus secretion and airway caliber. The majority of sensory nerve fibers that project to airways are derived from cell bodies located in the vagal jugular or nodose ganglia. The vagal C-fibers are relatively insensitive to the physiological milieu but respond vigorously to noxious stimuli and to mediators associated with inflammation [22, 23]. As Sherrington said of nociceptors in the somatosensory system they are equipped to provide the organ, “with a so to say sense of its own potential injury” [24]. In addition to nociceptive C-fibers, low-threshold mechanosensory sensory nerves innervate the lower respiratory tract that can be activated in a rapidly or slowly adapting fashion by the mechanical forces of respiration. These nerves are generally less susceptible to inflammatory- associated stimuli [22, 23].

Respiratory viruses commonly infect and replicate in airway epithelial cells leading to local inflammatory reactions. The inflammation is associated with the production of a host of inflammatory mediators, many of which can lead to the acute activation and sensitization of nociceptive terminals that reside within the epithelial layer [20]. This acute activation (action potential discharge) of sensory nerves can potentially lead to coughing, sneezing and parasympathetic driven bronchial smooth muscle contraction and secretions. These responses to viral infection would be expected to terminate as the infection and inflammation subside. Longer lasting effects of a viral infection may involve phenotypic changes to the nerves.

Airway inflammation not only activates airway C-fibers but also can phenotypically switch the low-threshold mechanosensitive A-fibers such that they take on characteristics of nociceptive C-fibers [25–29]. Induction of genes that are normally not expressed by a cell has been defined as “phenotypic switching.” In the somatosensory system, inflammation is associated with a phenotypic switch of low-threshold mechanosensitive A $\beta$ -fibers that respond to light touch such that they begin to express the tachykinin substance P. Light touch of the inflamed skin can then lead to tachykinergic transmission in the spinal cord, altering the communication between the skin and central nervous system [30]. Large A-fiber neurons in the trachea and lungs typically do not express the genes for the synthesis of neuropeptides like substance P and calcitonin gene-related peptide (CGRP). In the presence of allergic inflammation of guinea pig airways, these neurons begin to express both substance P and CGRP [26, 28]. Likewise, four days after a respiratory viral infection substance P is synthesized by large A $\delta$  cough receptors that terminate in the guinea pig trachea [31]. In addition to changing the neurotransmitters that A-fiber neurons produce, allergic inflammation also induces the synthesis of key ion channels involved in nerve activation. For example, during allergic airway inflammation transient receptor potential cation channel subfamily V member 1 (TRPV1) is turned on in rapidly and slowly adapting mechanosensitive A-fibers in rat lungs such that they become responsive to capsaicin [29]. Allergic inflammation also leads to de novo production of functional TRPV1 channels in tracheal A $\delta$  cough receptors in the guinea pig trachea [32]. This is relevant because TRPV1 is not only the receptor for capsaicin but also a pivotal channel in the activation mechanisms of many disparate stimuli including heat, acid, and certain inflammatory mediators [33, 34]. We have now found that viral infection with intranasal parainfluenza 3 (PIV3) leads to TRPV1 induction in trachea A $\delta$  cough receptors in guinea pig trachea. In vehicle treated animals, ~16 % of the tracheal-specific nodose neurons expressed TRPV1, whereas 4 days after PIV3 infection, ~50 % of the neurons expressed TRPV1. The most likely mechanism by which a local infection in the airway epithelium leads to changes in gene expression in the cell bodies involves the production of neurotrophic factors.

Neurotrophic factors can be produced during inflammatory reactions, including those associated with viral infections [35]. The neurotrophin family of neurotrophic factors comprise nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), neurotrophin 3 (NT 3), and neurotrophin 4 (NT 4) [36]. Unlike other inflammatory mediators, these factors can interact with high affinity receptors on nerve terminals in a manner that leads to changes in gene transcription in cell bodies situated within the distant sensory ganglia [37, 38]. Which neurotrophin is most likely to interact with airway sensory



the level of the smooth muscle [7]. However, there is good evidence that viral infections can increase the amount of acetylcholine released from the postganglionic nerve terminals. A key mechanism that contributes to the effect is a reduction in the activity of prejunctional muscarinic M2 receptors [7, 9, 11, 45]. Muscarinic M2 receptors are expressed in the postganglionic nerve terminal membrane where they serve as feedback inhibitory autoreceptors. When the acetylcholine release is increased, it acts on these terminal M2 receptors leading to a reduction in the release of acetylcholine per impulse [46]. Following viral infections, this inhibitory feedback mechanism is inhibited in guinea pig and rat airways [9, 10]. The virally mediated decrease in the cholinergic inhibitory feedback loop is associated with cytokine mediated decreases in M2 receptor gene expression in neurons within the airway parasympathetic ganglia [45].

## Conclusions


In summary, it is likely that respiratory viral infections can lead to airway inflammation and the production of inflammatory mediators that would be expected to stimulate nociceptive C-fibers. In addition, viral infection may lead to changes in gene expression within the sensory cell bodies causing phenotypic changes in a manner that is relevant to their neurotransmission (neuropeptide production) and activation profile (e.g., TRP channel expression). The phenotypic changes in sensory nerves following viral infection is likely due to increases in the production of neurotrophic factors by the epithelial cells and may persist beyond the period of active infection. Viral infection may also alter the function of parasympathetic nerves to further increase the reflex bronchoconstriction and secretions. The net effect of this sensory and autonomic neuroplasticity is an increase in the classical neuronal symptoms such as sneezing, coughing, reflex secretions, and bronchoconstriction. For those with underlying airway disorders, the viral-associated neuromodulation likely contributes to disease exacerbations. A better understanding of the molecular mechanisms underlying this neuroplasticity may lead to novel therapeutic approaches aimed at decrease viral-associated airways disorders.

## References

1. Footitt J, Johnston SL. Cough and viruses in airways disease: mechanisms. *Pulm Pharmacol Ther.* 2009; 22(2):108–113. [PubMed: 19480062]
2. McGarvey LP, Nishino T. Acute and chronic cough. *Pulm Pharmacol Ther.* 2004; 17(6):351–354. [PubMed: 15564075]
3. Dulek DE, Peebles RS Jr. Viruses and asthma. *Biochim Biophys Acta.* 1810; 11:1080–1090.
4. Sethi S. Molecular diagnosis of respiratory tract infection in acute exacerbations of chronic obstructive pulmonary disease. *Clin Infect Dis.* 2011; 52(Suppl 4):S290–S295. [PubMed: 21460287]
5. Broadley KJ, Blair AE, Kidd EJ, Bugert JJ, Ford WR. Bradykinin-induced lung inflammation and bronchoconstriction: role in parainfluenza-3 virus-induced inflammation and airway hyperreactivity. *J Pharmacol Exp Ther.* 2010; 335(3):681–692. [PubMed: 20847038]
6. Buckner CK, Songsiridej V, Dick EC, Busse WW. In vivo and in vitro studies on the use of the guinea pig as a model for virus-provoked airway hyperreactivity. *Am Rev Respir Dis.* 1985; 132(2):305–310. [PubMed: 2992331]
7. Fryer AD, Jacoby DB. Parainfluenza virus infection damages inhibitory M2 muscarinic receptors on pulmonary parasympathetic nerves in the guinea-pig. *Br J Pharmacol.* 1991; 102(1):267–271. [PubMed: 1646059]

8. Elwood W, Lotvall JO, Barnes PJ, Chung KF. Airway hyperresponsiveness to acetylcholine and to tachykinins after respiratory virus infection in the guinea pig. *Ann Allergy*. 1993; 70(3):231–236. [PubMed: 7680845]
9. Sorkness R, Clough JJ, Castleman WL, Lemanske RF Jr. Virus-induced airway obstruction and parasympathetic hyperresponsiveness in adult rats. *Am J Respir Crit Care Med*. 1994; 150(1):28–34. [PubMed: 8025764]
10. Lee AM, Fryer AD, van Rooijen N, Jacoby DB. Role of macrophages in virus-induced airway hyperresponsiveness and neuronal M2 muscarinic receptor dysfunction. *Am J Physiol Lung Cell Mol Physiol*. 2004; 286(6):L1255–L1259. [PubMed: 15136297]
11. Nie Z, Scott GD, Weis PD, Itakura A, Fryer AD, Jacoby DB. Role of TNF-alpha in virus-induced airway hyperresponsiveness and neuronal M(2) muscarinic receptor dysfunction. *Br J Pharmacol*. 2011; 164(2b):444–452. [PubMed: 21457223]
12. Laitinen LA, Elkin RB, Empey DW, Jacobs L, Mills J, Nadel JA. Bronchial hyperresponsiveness in normal subjects during attenuated influenza virus infection. *Am Rev Respir Dis*. 1991; 143(2): 358–361. [PubMed: 1990953]
13. Grunberg K, Kuijpers EA, de Klerk EP, de Gouw HW, Kroes AC, Dick EC, Sterk PJ. Effects of experimental rhinovirus 16 infection on airway hyperresponsiveness to bradykinin in asthmatic subjects in vivo. *Am J Respir Crit Care Med*. 1997; 155(3):833–838. [PubMed: 9117013]
14. Grunberg K, Timmers MC, Smits HH, de Klerk EP, Dick EC, Spaan WJ, Hiemstra PS, Sterk PJ. Effect of experimental rhinovirus 16 colds on airway hyperresponsiveness to histamine and interleukin-8 in nasal lavage in asthmatic subjects in vivo. *Clin Exp Allergy*. 1997; 27(1):36–45. [PubMed: 9117878]
15. Curley FJ, Irwin RS, Pratter MR, Stivers DH, Doern GV, Vernaglia PA, Larkin AB, Baker SP. Cough and the common cold. *Am Rev Respir Dis*. 1988; 138(2):305–311. [PubMed: 3057962]
16. McAlexander MA, Gavett SH, Kollarik M, Udem BJ. Vagotomy reverses established allergen-induced airway hyperreactivity to methacholine in the mouse. *Respir Physiol Neurobiol*. 2015; 212–214:20–24.
17. Trankner D, Hahne N, Sugino K, Hoon MA, Zuker C. Population of sensory neurons essential for asthmatic hyperreactivity of inflamed airways. *Proc Natl Acad Sci USA*. 2014; 111(31):11515–11520. [PubMed: 25049382]
18. Costello RW, Evans CM, Yost BL, Belmonte KE, Gleich GJ, Jacoby DB, Fryer AD. Antigen-induced hyperreactivity to histamine: role of the vagus nerves and eosinophils. *Am J Physiol*. 1999; 276(5 Pt 1):L709–L714. [PubMed: 10330026]
19. Lee LYMR. Histamine enhances vagal pulmonary C-fiber responses to capsaicin and lung inflation. *Respir Physiol*. 1993; 93(1):83–96. [PubMed: 8367619]
20. Lee LY, Yu J. Sensory nerves in lung and airways. *Compr Physiol*. 2014; 4(1):287–324. [PubMed: 24692141]
21. Udem BJ, Potenzieri C. Autonomic neural control of intrathoracic airways. *Compr Physiol*. 2012; 2(2):1241–1267. [PubMed: 23798300]
22. Lee, L-Y.; Udem, BJ. *Advances in vagal afferent neurobiology*. Boca Raton: CRC Taylor and Francis; 2005. *Bronchopulmonary vagal afferent nerves*.
23. Coleridge HMCJ. Impulse activity in afferent vagal C-fibres with endings in the intrapulmonary airways of dogs. *Respir Physiol*. 1977; 29:125–142. [PubMed: 866808]
24. Sherrington, C. *The integrated action of the nervous system*. Vol. 1. New Haven: Yale University Press; 1906.
25. Fischer AMG, Saria A, Philippin B, Kummer W. Induction of tachykinin gene and peptide expression in guinea pig nodose primary afferent neurons by allergic airway inflammation. *J Clin Invest*. 1996; 98:2284–2291. [PubMed: 8941645]
26. Chuaychoo BHD, Myers AC, Kollarik M, Udem BJ. Allergen-induced substance P synthesis in large-diameter sensory neurons innervating the lungs. *J Allergy Clin Immunol*. 2005; 116(2):325–331. [PubMed: 16083787]
27. Lieu TUB. Allergen and BDNF-induced TRP channel expression in nodose cough-fiber neurons. *FASEB J*. 2011; 25(864):814.

28. Myers ACKR, Udem BJ. Allergic inflammation-induced neuropeptide production in rapidly adapting afferent nerves in guinea pig airways. *Am J Physiol Lung Cell Mol Physiol.* 2002; 282:L775–L781. [PubMed: 11880304]
29. Zhang GLR, Wiggers M, Snow DM, Lee LY. Altered expression of TRPV1 and sensitivity to capsaicin in pulmonary myelinated afferents following chronic airway inflammation in the rat. *J Physiol.* 2008; 586:5771–5786. [PubMed: 18832423]
30. Neumann S, Doubell TP, Leslie T, Woolf CJ. Inflammatory pain hypersensitivity mediated by phenotypic switch in myelinated primary sensory neurons. *Nature.* 1996; 384(6607):360–364. [PubMed: 8934522]
31. Carr MJ, Hunter DD, Jacoby DB, Udem BJ. Expression of tachykinins in nonnociceptive vagal afferent neurons during respiratory viral infection in guinea pigs. *Am J Respir Crit Care Med.* 2002; 165(8):1071–1075. [PubMed: 11956047]
32. Lieu TKM, Myers AC, Udem BJ. Neurotrophin and GDNF family ligand receptor expression in vagal sensory nerve subtypes innervating the adult guinea pig respiratory tract. *Am J Physiol Lung Cell Mol Physiol.* 2011; 300(5):L790–L798. [PubMed: 21335521]
33. Caterina MJJD. The vanilloid receptor: a molecular gateway to the pain pathway. *Annu Rev Neurosci.* 2001; 24:487–517. [PubMed: 11283319]
34. Veldhuis NA, Poole DP, Grace M, McIntyre P, Bunnnett NW. The G protein-coupled receptor-transient receptor potential channel axis: molecular insights for targeting disorders of sensation and inflammation. *Pharmacol Rev.* 2015; 67(1):36–73. [PubMed: 25361914]
35. Tortorolo L, Langer A, Polidori G, Vento G, Stampachiachere B, Aloe L, Piedimonte G. Neurotrophin overexpression in lower airways of infants with respiratory syncytial virus infection. *Am J Respir Crit Care Med.* 2005; 172(2):233–237. [PubMed: 15879412]
36. Huang EJ, Reichardt LF. Neurotrophins: roles in neuronal development and function. *Annu Rev Neurosci.* 2001; 24:677–736. [PubMed: 11520916]
37. Chowdary PD, Che DL, Cui B. Neurotrophin signaling via long-distance axonal transport. *Annu Rev Phys Chem.* 2012; 63:571–594. [PubMed: 22404590]
38. Zweifel LSKR, Ginty DD. Functions and mechanisms of retrograde neurotrophin signalling. *Nat Rev Neurosci.* 2005; 22(3):146–152.
39. Ye XM, Zhong NS, Liu CL, Chen RC. Cough reflex sensitivity is increased in guinea pigs with parainfluenza virus infection. *Exp Lung Res.* 2011; 37(3):186–194. [PubMed: 21417816]
40. Carr MJKM, Meeker SN, Udem BJ. A role for TRPV1 in bradykinin-induced excitation of vagal airway afferent nerve terminals. *J Pharmacol Exp Ther.* 2003; 304:1275–1279. [PubMed: 12604706]
41. Kollarik MUB. Mechanisms of acid-induced activation of airway afferent nerve fibres in guinea-pig. *J Physiol.* 2002; 543:591. [PubMed: 12205192]
42. Grace M, Birrell MA, Dubuis E, Maher SA, Belvisi MG. Transient receptor potential channels mediate the tussive response to prostaglandin E2 and bradykinin. *Thorax.* 2012; 67(10):891–900. [PubMed: 22693178]
43. Tan Y, Yang T, Liu S, Liu H, Xiang Y, Qu F, Li H, Qin X. Infection with respiratory syncytial virus alters peptidergic innervation in the lower airways of guinea-pigs. *Exp Physiol.* 2008; 93(12): 1284–1291. [PubMed: 18603600]
44. Auais A, Adkins B, Napchan G, Piedimonte G. Immunomodulatory effects of sensory nerves during respiratory syncytial virus infection in rats. *Am J Physiol Lung Cell Mol Physiol.* 2003; 285(1):L105–L113. [PubMed: 12639840]
45. Rynko AE, Fryer AD, Jacoby DB. Interleukin-1beta mediates virus-induced m2 muscarinic receptor dysfunction and airway hyperreactivity. *Am J Respir Cell Mol Biol.* 2014; 51(4):494–501. [PubMed: 24735073]
46. Fryer AD, Maclagan J. Muscarinic inhibitory receptors in pulmonary parasympathetic nerves in the guinea-pig. *Br J Pharmacol.* 1984; 83(4):973–978. [PubMed: 6518347]



**Fig. 1.**

A schematic highlighting the potential effects of viral infection on vagus nerve activity. (I) Viral infection of epithelial cells can lead to release of mediators that stimulate action potential discharge in afferent nerves, thereby alerting the central nervous system. Viral infection can also lead to the production and release of neurotrophic factors that can (II) influence gene expression in the vagal jugular and nodose ganglia in a manner that can lead to relevant phenotypic changes in the airway sensory nerves. (III) Action potentials arriving in the brainstem are integrated, ultimately leading to sensations and to (IV) increases in preganglionic parasympathetic drive. (V) This will increase neurotransmission at the neuro-effector junctions, causing bronchoconstriction and mucus secretion; effects further amplified by inhibition of inhibitory muscarinic autoreceptors. See text for references