

Commentary

Open Access

PPAR γ , neuroinflammation, and disease

Robert E Mrak*¹ and Gary E Landreth²

Address: ¹Department of Pathology, University of Arkansas for Medical Sciences, Little Rock, Arkansas, USA and ²Alzheimer Research Laboratory, Department of Neurosciences, Case Western Reserve University School of Medicine, Cleveland, Ohio, USA

Email: Robert E Mrak* - mrakroberte@uams.edu; Gary E Landreth - gel2@po.cwru.edu

* Corresponding author

Published: 14 May 2004

Received: 02 April 2004

Journal of Neuroinflammation 2004, 1:5

Accepted: 14 May 2004

This article is available from: <http://www.jneuroinflammation.com/content/1/1/5>

© 2004 Mrak and Landreth; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Peroxisome proliferator-activated receptors (PPARs) are a class of nuclear transcription factors that are activated by fatty acids and their derivatives. One of these, PPAR γ , regulates responsiveness to insulin in adipose cells, and PPAR γ -activating drugs such as pioglitazone are used in the treatment of type 2 diabetes. PPAR γ acts in myeloid-lineage cells, including T-cells and macrophages, to suppress their activation and their elaboration of inflammatory molecules. PPAR γ activation also suppresses the activated phenotype in microglia, suggesting that PPAR γ -activating drugs may be of benefit in chronic neuroinflammatory diseases. Some, but not all, nonsteroidal anti-inflammatory agents (indomethacin and ibuprofen in particular) also have activating effects on PPAR γ .

Discussion and conclusions: These observations suggest on the one hand a role for PPAR γ -activating drugs in the treatment of chronic neuroinflammatory diseases-as shown for a patient with secondary progressive multiple sclerosis by Pershadsingh et al. in this issue of the *Journal of Neuroinflammation*-and suggest on the other hand a possible explanation for confusing and contradictory results in trials of nonsteroidal anti-inflammatory agents in Alzheimer's disease.

Introduction

There are still times in modern medicine when a single patient can enlighten our understanding of a disease or disease process, and can serve as an impetus for further discovery. In this issue of *Journal of Neuroinflammation*, Harrihar Pershadsingh and his colleagues [1] describe stabilization and, indeed, clinical improvement in a patient with progressive secondary multiple sclerosis who was treated with pioglitazone over a three-year period. These observations suggest that larger, controlled trials of such treatment may be warranted.

The possible connection between pioglitazone and multiple sclerosis is a fascinating story in itself, and one that not only provides interesting parallels between chronic CNS inflammatory diseases and chronic peripheral diseases,

but also illuminates an area of current interest for diseases such as Alzheimer's disease as well.

Discussion

Pioglitazone is currently used in the treatment of type 2 diabetes. The mechanism of action involves activation of a nuclear transcription factor known as the peroxisome proliferator-activated receptor gamma, or PPAR γ , that controls lipid metabolism in adipocytes, and sensitizes these cells to insulin. PPAR γ agonists also suppress T-cell activation suggesting that they may be useful in treating inflammatory diseases. Moreover, activation of PPAR γ in microglia (as well as in macrophages) results in decreased activation of these cells, with decreased expression of pro-inflammatory cytokines and related molecules. This suggests that PPAR γ agonists might be useful in a number of

central nervous system diseases with inflammatory components.

Peroxisomes, or microbodies as they were originally known, were discovered by early electron microscopists in the 1950s [2]. Christian de Duve, in Brussels, Belgium, subsequently isolated these structures, demonstrated hydrogen peroxide generation, and renamed them peroxisomes [3]. The discovery of PPARs grew out of the War on Cancer in the 1970s. A class of drugs was identified that promoted cancer-like growths in animals, but that did not cause DNA damage [4]. What these drugs did do was to stimulate proliferation of peroxisomes in target cells. At the time, this suggested a specific genetic trigger for tumorigenesis, and there ensued two decades of attempts to identify the receptor for these peroxisome proliferation-activating drugs.

By the 1990s, when PPARs were identified and shown to be transcription factors [5], interest had waned in cancer circles. PPARs are a class of transcription factors that are activated by fatty acids and their derivatives. They were found to control a number of genes, most of which have little or nothing to do with peroxisomes. PPAR γ is important both in fat cell metabolism and in modulating cellular responsiveness to insulin - hence the connection with diabetes [6]. PPAR γ -activating drugs were subsequently found to regulate T-cell responsiveness [7,8] and to suppress macrophage and microglia activation [9-11]. Both of these actions are relevant to multiple sclerosis, and may have implications for other chronic neurodegenerative diseases as well. In addition to pioglitazone, some (but not all) nonsteroidal anti-inflammatory drugs (in particular indomethacin and ibuprofen) have activating effects on PPAR γ in addition to their effects on cyclooxygenase [12]. NSAID use has been linked with decreased risk of Alzheimer's disease in epidemiological studies [13-15], but prospective trials of NSAIDs in Alzheimer patients have yielded contradictory and inconclusive results [16-18]. The NSAID-PPAR γ connection might explain some of these contradictions, as the only one of these clinical trials that reported a benefit was also the only one that used a PPAR γ -activating drug [16]. There are currently two clinical trials in progress testing the efficacy of PPAR γ agonists in AD patients.

Conclusions

Pioglitazone and related drugs activate PPAR γ , and activation of PPAR γ suppresses T-cell, macrophage, and microglial immune responses. If suppression of these immune responses is of potential benefit for inflammatory diseases of the brain, then pioglitazone should provide therapeutic benefit in multiple sclerosis. Multiple sclerosis, of course, is notoriously variable in its course, but the secondary progressive variant is an exception to this. Pershadsingh et

al. show clinical stabilization in such a patient, treated for three years with the PPAR γ -activating drug pioglitazone. This single clinical case thus provides support for a link between PPAR γ activation and suppression of neuroinflammation, and suggests avenues of research for the further treatment of multiple sclerosis as well as other chronic neuroinflammatory diseases.

List of abbreviation used

PPAR – peroxisome proliferator-activated receptor

NSAID – nonsteroidal anti-inflammatory drug

Competing interests

None declared.

Acknowledgments

Supported in part by NIH PO1 AG 12411, NIH P30 AG19606, and NIH ROI AG 37989

References

- Pershadsingh HA, Heneka MT, Saini R, Amin NM, Broeske DJ, Feinstein DL: **Effect of pioglitazone treatment in a patient with secondary multiple sclerosis.** *J Neuroinflamm* 2004, **1**:3.
- Rhodin J: **Further studies on the nephron ultrastructure in mouse. Terminal part of proximal convolution.** In: *Electron Microscopy: Proceedings of the Stockholm Conference* Edited by: Sjostrand FS, Rhodin J. New York: Academic Press; 1957.
- De Duve C, Baudhuin P: **Peroxisomes (microbodies and related particles).** *Physiol Rev* 1966, **46**:323-357.
- Lake BG: **Mechanisms of hepatocarcinogenicity of peroxisome-proliferating drugs and chemicals.** *Annu Rev Pharmacol Toxicol* 1995, **35**:483-507.
- Issemann I, Green S: **Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators.** *Nature* 1990, **347**:645-650.
- Willson TM, Lambert MH, Kliewer SA: **Peroxisome proliferator activated receptor γ and metabolic disease.** *Annu Rev Biochem* 2001, **70**:341-367.
- Clark RB, Bishop-Bailey D, Estrada-Hernandez T, Hla T, Puddington L, Padula SJ: **The nuclear receptor PPAR gamma and immunoregulation: PPAR gamma mediates inhibition of helper T cell responses.** *J Immunol* 2000, **164**:1364-71.
- Schmidt S, Moric E, Schmidt M, Sastre M, Feinstein DL, Heneka MT: **Anti-inflammatory and antiproliferative actions of PPAR-gamma agonists on T lymphocytes derived from MS patients.** *J Leukoc Biol* 2004, **75**:478-485.
- Jiang C, Ting AT, Seed B: **PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines.** *Nature* 1998, **391**:82-86.
- Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK: **The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation.** *Nature* 1998, **391**:79-82.
- Combs CK, Bates P, Karlo JC, Landreth GE: **Regulation of beta-amyloid stimulated proinflammatory responses by peroxisome proliferator-activated receptor alpha.** *Neurochem Int* 2001, **39**:449-457.
- Lehmann JM, Lenhard JM, Oliver BB, Ringold GM, Kliewer SA: **Peroxisome proliferator-activated receptors alpha and gamma are activated by indomethacin and other non-steroidal anti-inflammatory drugs.** *J Biol Chem* 1997, **272**:3406-3410.
- Breitner JCS, Gau BA, Welsh KA, Plassman BL, McDonald WM, Helms MJ, Anthony JC: **Inverse association of anti-inflammatory treatments and Alzheimer's disease: initial results of a co-twin control study.** *Neurology* 1994, **44**:227-232.
- Andersen K, Launer LJ, Ott A, Hoes A, Breteler MMB, Hofman A: **Do nonsteroidal anti-inflammatory drugs decrease the risk for Alzheimer's disease?: The Rotterdam Study.** *Neurology* 1995, **45**:1441-1445.

15. Stewart WF, Kawas C, Corrada M, Metter EJ: **Risk of Alzheimer's disease and duration of NSAID use.** *Neurology* 1997, **48**:626-632.
16. Rogers J, Kirby LC, Hempelman SR, Berry DL, McGeer PL, Kaszniak AW, Zilinski J, Cofield M, Mansukhani L, Willson P, Kogan F: **Clinical trial of indomethacin in Alzheimer's disease.** *Neurology* 1993, **43**:1609-1611.
17. Scharf S, Mander A, Ugoni A, Vajda F, Christophidis N: **A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer's disease.** *Neurology* 1999, **53**:197-201.
18. Van Gool WA, Weinstein HC, Scheltens P, Walstra GJ, Scheltens PK: **Effect of hydroxychloroquine on progression of dementia in early Alzheimer's disease: an 18-month randomised, double-blind, placebo-controlled study.** *Lancet* 2001, **358**:455-460.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

