



Published in final edited form as:

Cell Rep. 2016 April 12; 15(2): 398–410. doi:10.1016/j.celrep.2016.03.029.

## Enhanced Transcriptional Activity and Mitochondrial Localization of STAT3 Co-induce Axon Regrowth in Adult Central Nervous System

Xueting Luo<sup>1,#</sup>, Marcio Ribeiro<sup>1</sup>, Eric R. Bray<sup>1</sup>, Do-Hun Lee<sup>1</sup>, Benjamin J. Yungler<sup>1</sup>, Saloni T. Mehta<sup>1</sup>, Kinjal A. Thakor<sup>1</sup>, Francisca Diaz<sup>2</sup>, Jae K. Lee<sup>1</sup>, Carlos T. Moraes<sup>2</sup>, John L. Bixby<sup>1</sup>, Vance P. Lemmon<sup>1</sup>, and Kevin K. Park<sup>1,\*</sup>

<sup>1</sup>Department of Neurological Surgery, Miami Project to Cure Paralysis; University of Miami Miller School of Medicine, Miami, FL 33136, USA

<sup>2</sup>Department of Neurology; University of Miami Miller School of Medicine, Miami, FL 33136, USA

### SUMMARY

STAT3 is a transcription factor central to axon regrowth with an enigmatic ability to act in different subcellular regions independently of its transcriptional roles. However, its roles in mature central nervous system (CNS) neurons remain unclear. Here we show that along with nuclear translocation, STAT3 translocates to mitochondria in mature CNS neurons upon cytokine stimulation. Loss- and gain-of-function studies using knockout mice and viral expression of various STAT3 mutants demonstrate that STAT3's transcriptional function is indispensable for CNS axon regrowth whereas mitochondrial STAT3 enhances bioenergetics and further potentiates regrowth. STAT3's localization, functions, and growth-promoting effects are regulated by MEK, an effect further enhanced by *Pten* deletion leading to extensive axon regrowth in the mouse optic pathway and spinal cord. These results highlight CNS neuronal dependence on STAT3 transcriptional activity with mitochondrial STAT3 providing ancillary roles, and illustrate a critical contribution for MEK in enhancing diverse STAT3 functions and axon regrowth.


### Abstract

\*Corresponding author: Kevin K. Park, kpark@miami.edu.

#Current Address, Department of Ophthalmology, Shanghai First People's Hospital, Shanghai Jiao Tong University School of Medicine; Shanghai Key Laboratory of Fundus Disease, 100 Hai Ning Road, Shanghai 200080, Shanghai, China

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Author Contributions:** K.K.P supervised the project. X.L. and K.K.P planned and performed experiments, analyzed data and wrote the manuscript. C.T.M. and F.D. provided the transgenic mice. M.R., D.H.L., B.J.Y., E.R.B., S.M. and K.A.T. performed the experiments. J.L.B., V.P.L and J.K.L advised and analyzed the data.



## INTRODUCTION

The CNS is composed of distinct neuronal types, most of which fail to regenerate after injury. While significant progress has been made in identifying genes involved in regulating axon regrowth, the mechanisms underlying regeneration failure remain elusive (Case and Tessier-Lavigne, 2005; Cregg et al., 2014; Huebner and Strittmatter, 2009; Sun and He, 2010). The interleukin-6 (IL-6) family of cytokines promotes axon regeneration (Cafferty et al., 2004; Cao et al., 2006; Leaver et al., 2006; Leibinger et al., 2009; Muller et al., 2009; Pernet et al., 2012; Zigmond, 2011), and identification of downstream molecules that mediate cytokines' actions in neurons has provided some insights into signaling pathways and neuron-intrinsic molecular mechanisms that regulate axon regeneration process. The IL6 family of cytokines activates many signaling molecules, triggered by dimerization of gp130 and phosphorylation of Janus kinases (JAKs). These proteins serve as docking sites for several adaptors that link the receptor to signal transducer and activator of transcription (STAT), MAP kinases and PI3K/Akt (Ernst and Jenkins, 2004). In addition, they activate other central regulators of cell growth including mammalian target of rapamycin (mTOR),  $\text{NF}\kappa\beta$  and Yes-associated protein (YAP) (Gallagher et al., 2007; Saleh et al., 2013; Smith et al., 2009; Taniguchi et al., 2015). However, activating these target modules individually in neurons results in only limited regeneration (Pernet et al., 2005; Pernet et al., 2013), indicative of multifaceted mechanisms that mediate axon regeneration.

STAT3 is a transcription factor involved in many biological processes. Genetic and pharmacological inhibition of STAT3 reduces axon regeneration both in central and peripheral nervous systems (Bareyre et al., 2011; Leibinger et al., 2013), demonstrating its role in promoting axon regeneration. Most of STAT3's functions are attributed to transcriptional regulation. However, recent studies have demonstrated that STAT3 can also function outside the nucleus; STAT3 proteins are found in the mitochondria where they

locally regulate metabolic functions in non-neuronal cells (Gough et al., 2009; Wegrzyn et al., 2009). In vitro cytoplasmic STAT3 is required for ciliary neurotrophic factor (CNTF)-induced axon growth in embryonic neurons (Selvaraj et al., 2012), and mitochondrially localized STAT3 promotes neurite outgrowth in PC12 cells (Zhou and Too, 2011). Intriguingly, these studies indicated that the transcriptional activity of STAT3 is not required for these axon growth promoting-effects. Conversely, both nuclear and cytoplasmic STAT3 are required to induce axon growth in peripheral neurons, (Pellegrino and Habecker, 2013), indicating that the mechanisms by which STAT3 promotes axon growth might be different among distinct cell types and under different conditions. It is unclear whether STAT3's transcription-independent functions are broadly involved in different systems, and whether they play a role in axon regeneration in the mature CNS. Here we examine STAT3's transcription independent roles in adult CNS neurons. Our study shows that STAT3 localizes to both nucleus and mitochondria in adult retinal ganglion cells (RGCs) in response to cytokine, and enhancing STAT3's transcription activity and localization to mitochondria together improves optic nerve regeneration. Further, STAT3's transcription activity, mitochondrial localization and growth promoting effects are enhanced by MEK, an effect further enhanced by *Pten* deletion with extensive axon regrowth in mouse optic pathway and spinal cord. These results establish STAT3's transcription dependent- and independent-contributions to promoting axon regrowth in the mature CNS, and suggest that MEK enhancement of STAT3's versatile functions is integral in this function.

## RESULTS

### CNTF induces phosphorylation and translocation of STAT3 to distinct subcellular regions in CNS neurons: MEK-dependent translocation of STAT3 in mitochondria

To gain insights into STAT3's mode of action and its potential regulators in promoting axon regeneration in mature CNS neurons, we examined activation status of STAT3 and several other key factors linked to cytokine-induced cell growth (Ernst and Jenkins, 2004; Gallagher et al., 2007; Taniguchi et al., 2015) in postmitotic postnatal (P) 3 cortical neurons following CNTF treatment. Elevating CNTF levels promotes axon regeneration *in vivo* and increases neurite outgrowth in vitro (Gallagher et al., 2007; Leaver et al., 2006; Pernet et al., 2012; Selvaraj et al., 2012; Smith et al., 2009; Sun et al., 2011). As expected, addition of recombinant CNTF (rCNTF) led to rapid activation of STAT3 (Figure S1); the levels of STAT3 phosphorylation at serine 727 (S727) and tyrosine 705 (Y705) remained elevated for at least 20 hours after adding rCNTF. Similarly, the level of ERK phosphorylation rapidly increased before returning to the basal level. The level of phosphorylated S6 (pS6), an indicator of mTOR activity moderately increased in the absence of Akt activation. No obvious change was detected for I $\kappa$ B $\alpha$ , YAP and  $\beta$ -catenin (Figure S1). To further examine STAT3's mode of action in CNS neurons, we investigated STAT3 expression in adult RGCs. In responses to growth factors STAT3 is phosphorylated at Y705 and S727, binds to DNA and regulates transcription. Consistently, immunohistochemistry shows that intravitreal injection of AAV expressing the secretable form of CNTF (Leaver et al., 2006) (AAV-CNTF), known to promote RGC axon regeneration (Leaver et al., 2006; Pernet et al., 2012), elevates the levels of phosphorylated STAT3 (pSTAT3) at both Y705 and S727 in RGCs (Figure 1A). Notably however, we observed that the cellular localization patterns of

pSTAT3(Y705) and pSTAT3(S727) are distinct; pSTAT3(Y705) seems to localize exclusively in the nucleus whereas pSTAT3(S727) is also found in the cytoplasm and RGC processes (Figure 1A). Are there functional roles for STAT3 proteins located outside RGC nucleus? Since past studies have shown mitochondrial STAT3's ability to promote neurite growth and that translocation of STAT3 to mitochondria requires phosphorylation at S727 (Gough et al., 2009; Tammineni et al., 2013; Wegrzyn et al., 2009), we examined whether STAT3 translocates to mitochondria in RGCs after CNTF treatment. First, we used confocal microscopy and quantified the levels of pSTAT3(S727) co-localizing with the mitochondrial protein ATP synthase  $\beta$  (ATPsyn $\beta$ ). Compared to controls, there was a significant increase in the pSTAT3(S727)/ATPsyn $\beta$  co-localization in RGCs in CNTF-treated retina (Figures 1B and 1C), indicative of STAT3's presence in RGC mitochondria. Additionally, we examined STAT3's subcellular localization using ascorbate peroxidase (APEX) protein tagging technique. APEX is a monomeric peroxidase that endures strong electron microscope fixation, preserving ultrastructural details (Martell et al., 2012). At EM level, STAT3 is detected in inner mitochondrial membrane in CNTF-treated cortical neurons (Figure 1D). Since phosphorylation at the S727 residue is critical for STAT3 mitochondrial localization (Gough et al., 2009; Tammineni et al., 2013), and that CNTF induces activation of ERK (Figure S1), a protein kinase shown to cause S727 phosphorylation (Zhou and Too, 2011), we examined whether MEK (the upstream activator of ERK) induces STAT3 phosphorylation and translocation to mitochondria. Pharmacological MEK inhibitors almost completely blocked ERK phosphorylation, and CNTF-mediated STAT3 S727 phosphorylation in cortical neurons. In contrast, pSTAT3 (Y705) level was not affected by these inhibitors (Figure 1E). Additionally, subcellular fractionation reveals that total STAT3 and pSTAT3(S727) levels in the mitochondria increased about 2 and 2.5 folds, respectively after CNTF treatment (Figures 1F and 1G). Increases in the level of total STAT3 (but not pSTAT3(S727)) were also detected in the nuclear fraction but only after exposing the film for a prolonged period (Figure 1F). Refamitinib (MEK inhibitor) significantly reduced the amounts of pSTAT3(S727) and total STAT3 in the mitochondria (Figures 1F and 1G). Together, these data demonstrate that CNTF promotes phosphorylation and translocation of STAT3 to both the nuclei and mitochondria in post-mitotic CNS neurons, and suggests that mitochondrial presence of STAT3 is MEK-dependent.

### **DNA binding domain of STAT3 is essential for CNTF induced-RGC axon regeneration**

Translocation of STAT3 to different subcellular locations in RGCs raises a question of the extent to which the transcriptional activity-dependent (nuclear) and -independent functions of STAT3 contribute to promoting CNS axon regeneration. To examine this question, we designed optic nerve regeneration experiment using *Stat3<sup>fl/fl</sup>* mice crossed to *R26-tdTomato* reporter (*Stat3<sup>fl/fl</sup>;R26-tdTomato*) in which *Stat3*-null RGCs can be created, and STAT3 expression reconstituted with AAV2 expressing various STAT3 mutants (Figure 2A). Intravitreal injection of AAV-Cre in this mouse line results in concomitant *Stat3* deletion and tdTomato labeling in transduced cells. We injected AAV-Cre together with AAV expressing either wild-type STAT3 (STAT3wt) or DNA-binding mutant version of STAT3 (STAT3dbm) (Table S1)(Bromberg et al., 1998; Bromberg et al., 1999; Horvath et al., 1995) in *Stat3<sup>fl/fl</sup>;R26-tdTomato* mice. Five days later animals received optic nerve crush. This injection scheme (Figure 2A) allows the onset of *Stat3* ablation in adult RGCs without the

tdTomato presence in the optic nerve before the injury (Sun et al., 2011), thus allowing tdTomato labeling only in regenerating, and not in degenerated axons. Immediately after injury, animals also received injection of either AAV-GFP or AAV-CNTF. At 19 days post-injury, we examined the extent of tdTomato-labeled (i.e. *Stat3* KO) regenerating axons.

Consistent with the previous studies (Park et al., 2008; Sun et al., 2011), AAV-Cre injection infected the majority of RGCs (~90%) as evident by Cre and tdTomato immunoreactivity in the *R26-tdTomato* mice (Figure 2B). In mice without *Stat3* deletion, AAV-CNTF resulted in significant RGC axon regeneration (Figures 2C and 2D). *Stat3* deletion completely eliminated CNTF-induced regeneration. Reconstituting *Stat3* KO-RGCs with STAT3wt (i.e. with AAV-STAT3wt) rescued CNTF induced- regeneration. In contrast, AAV-STAT3dbm failed to rescue regeneration (Figures 2C and 2D). Using quantitative RT-PCR, we validated that STAT3dbm is transcriptionally inactive as it does not increase the expression of *Socs3* (data not shown), a prominent STAT3 target gene. RGC transduction efficiency was approximately 60% for both AAV-STAT3wt and AAV-STAT3dbm (Figure 2E). Additionally, we assessed RGC survival. The increase in RGC survival induced by CNTF is reduced by *Stat3* deletion. Reconstitution with STAT3wt, but not with STAT3dbm, rescued CNTF's RGC survival effect (Figures S2A and S2B). Together, these data establish that the DNA binding domain of STAT3 is essential for CNTF induced-RGC axon regeneration, strongly suggesting a requirement for transcriptional activity.

### Mitochondrial STAT3's involvement in CNS axon regeneration

While our data point to a requirement for STAT3's transcriptional function, distribution of phosphorylated STAT3 proteins in the RGC cytoplasm suggests that non-nuclear STAT3 may additionally facilitate regeneration. To directly evaluate the role of mitochondrial STAT3 in RGC axon regeneration, we generated AAV expressing the DNA binding mutant form of STAT3, that also contains a mitochondrial targeting sequence (STAT3mts) taken from the cytochrome c oxidase subunit VIII gene (Szczepanek et al., 2011). AAV-STAT3mts alone failed to increase regeneration significantly compared to the GFP controls (Figures 3A–3C). It was previously shown that the constitutively active form of STAT3 proteins (STAT3ca) dimerizes and translocates to the nucleus, and overexpression of STAT3ca promotes axon regeneration (Pernet et al., 2013). Consistently, we observed that AAV-STAT3ca increases regeneration (Figures 3B and 3C). AAV-STAT3wt alone led to some increase in regeneration, but this was statistically insignificant (Figure 3C). Interestingly, co-injection of AAV-STAT3mts with AAV-STAT3ca (but not with AAV-STAT3wt; data not shown) resulted in further enhancement in regeneration compared to the AAV-STAT3ca group (Figure 3C). RGC transduction efficiency was similar for these AAV viruses carrying different forms of STAT3, ~60% (Figure 3D). The improved regeneration seen after STAT3ca and STAT3mts combination could be due to separate pools of virally expressed STAT3 in the nucleus and mitochondria, acting cooperatively. Alternatively, it could occur if the concentration of virally expressed STAT3 proteins was somewhat higher in the double compared to each single AAV group. This is unlikely as the various animal groups received equivalent final titers of each AAV (see Experimental Procedures). Moreover, we observed that overexpressed STAT3ca is detected predominantly in the nuclear and cytosolic fractions, and not in the mitochondrial fraction. Conversely, overexpressed STAT3mts is localized

mostly in the mitochondrial and not in the nuclear fraction (Figure S2C), indicating that the enhanced regeneration in the double AAV group is due to virally expressed STAT3 acting in different cellular compartments. Furthermore, we examined RGC survival and found no difference among the animal groups indicating that the enhanced regeneration is not due to differences in cell survival (Figures 3E and 3F). Together, these data indicate that differentially localized STAT3 proteins in the nucleus and mitochondria cooperate to promote RGC axon regeneration.

### **Mitochondrial STAT3 increases cellular ATP levels, and complex IV activity: contribution of mitochondrial electron transport chain (ETC) to regeneration**

Our data indicate that mitochondrial STAT3 has facilitative roles in enhancing RGC axon regeneration, but the underlying mechanism(s) is not known. In non-neuronal cells, mitochondrial STAT3 has been shown to enhance electron transport chain (ETC) complex functions and oxidative phosphorylation (OXPHOS) efficiency (Szczepanek et al., 2011; Wegrzyn et al., 2009). Given that mitochondrial ATP production is significantly compromised in injured neurons, and that axon extension requires substantial energy (Mattson, 2007), it is plausible that the improved regeneration seen after expression of STAT3<sub>mts</sub> and STAT3<sub>ca</sub> might be due to its ability to enhance mitochondrial bioenergetics in injured neurons. While ATP is generated primarily in the mitochondria via ETC, under certain proliferative states (i.e. during retinal development or tumors), cells can rely mostly on glycolysis to produce ATP (Agathocleous et al., 2012). To first examine the extent to which the ETC contributes to axon regeneration, we assessed RGC axon regeneration in cytochrome *c* oxidase (COX) deficient-mice (Fukui et al., 2007). The *cox10* gene encodes haem A farnesyl transferase, necessary for the assembly of COX (Mogi et al., 1994), the terminal complex of the ETC. In *Cox10* ablated-cells, COX is unstable and rapidly degraded, leading to defects in mitochondrial ATP production (Fukui et al., 2007). *Cox10*<sup>fl/fl</sup> mice received intravitreal AAV-Cre injection (Figure S3A) followed by optic nerve crush/AAV-CNTF injection 8 weeks later. Wild-type animals receiving AAV-CNTF showed significant axon regeneration. In contrast, *Cox10* KO animals showed markedly reduced regeneration (Figures S3B and S3C). Interestingly, RGC survival after crush in the *Cox10* KO-mice was similar to the WT mice (Figure S3D), indicating that the ETC requirements for promoting RGC survival and axon regeneration are different. To directly assess the effects of STAT3 expression on mitochondrial functions, we measured the levels of intracellular ATP in WT cortical neurons, and observed that expression of STAT3<sub>mts</sub> (Figures S4A and S4B) leads to significantly higher levels of ATP compared to the control (Figure S4C). We also observed that STAT3<sub>mts</sub> expression in cortical neurons significantly increases mitochondrial membrane potential (i.e. an indicator of mitochondrial function) (Figure S4D). Additionally, we measured the levels of ETC complex activity, and observed that complex IV (but not complex I) activity was significantly higher in STAT3<sub>mts</sub> group compared to the control (Figures S4E and S4F). These data indicate that mitochondrial ETC is critical for axon regeneration and STAT3 targeted to mitochondria enhances mitochondrial energetics in CNS neurons.

### MEK promotes STAT3 phosphorylation at S727 and enhances expression of STAT3-induced genes in CNS neurons

In addition to STAT3, CNTF also induces activation of ERK, the primary downstream target of MEK (Figure S1). We reasoned that MEK/ERK may enhance transcriptional and mitochondrial STAT3 functions and support regeneration. To test this, we injected AAV expressing the constitutively active form of MEK (MEKca) (Table S1)(Boehm et al., 2007), and observed that this alone results in an increase in the level of pSTAT3(S727), but not pSTAT3(Y705) in the retina (Figure S5A). In retinas, we observed a marked increase in the levels of pERK and pSTAT3(S727) caused by AAV-MEKca in RGCs (Figures S5B–S5D). Next, we examined the effects of MEKca overexpression on the level of STAT3-target gene transcripts. Among the major STAT3 target genes examined (Figure S5F), STAT3ca expression resulted in moderate upregulation of *Ccckn1a*, *Socs1*, *Socs3*, *Irf1* and *Atf3*. STAT3wt did not significantly change gene expression levels compared to GFP control, except for *Socs3* (data not shown). Co-expression of MEKca with STAT3ca significantly enhanced expression of each of these genes except for ATF3, compared to STAT3ca alone. These data demonstrate that MEKca promotes phosphorylation of STAT3 (S727) and further enhances expression of STAT3-induced genes in CNS neurons.

### MEK enhances STAT3-mediated axon regeneration

Given its ability to modulate STAT3 at multiple levels, we next tested the effect of MEKca on RGC axon regeneration in the presence of STAT3ca expression. Consistent with the previous study (Pernet et al., 2005), AAV-MEKca alone enhanced RGC survival, but did not significantly enhance regeneration (Figures 4A, 4B, 4D and 4E). However, co-injection of AAV-MEKca with AAV-STAT3ca markedly increases regeneration compared to AAV-STAT3ca alone (Figures 4A and 4B). Co-injection of AAV-MEKca with AAV-STAT3wt led to a slight increase in regeneration, albeit much lower than the MEKca+STAT3ca group (Figure 4B). Four weeks post-injury, MEKca+STAT3ca treatment led to a 6-fold increase in axon numbers at 1 mm from the lesion site compared to the STAT3ca alone group (Figure 4B). Injection of AAV-MEKca led to transduction of about 70% of RGCs (Figure 4C). In the combined group, nearly all of the AAV-STAT3ca transduced RGCs (i.e. ~60% of RGCs) are also transduced by AAV-MEKca (data not shown). Whole-mount retinal immunohistochemistry shows similar number of surviving RGCs in the MEKca alone group compared to the MEKca/STAT3ca combined group (Figures 4D and 4E).

### STAT3/MEK with *Pten* deletion- effects on regenerating RGC axon number and length

Deletion of *Pten* promotes RGC axon regeneration (Park et al., 2008). This regeneration does not rely on STAT3 or B-raf (a major upstream activator of MEK)(O'Donovan et al., 2014; Sun et al., 2011), but depends on Akt/mTOR activation (Park et al., 2008; Sun et al., 2011). Since regeneration by *Pten* deletion involves distinct molecules, we next examined the extent of regeneration after co-overexpressing STAT3ca and MEKca in PTEN KO-neurons. *Pten*<sup>fl/fl</sup> mice received an AAV cocktail containing AAV-Cre, AAV-STAT3ca and AAV-MEKca. Two weeks after AAV injection, animals received optic nerve crush. *Pten* deletion alone promoted axon regeneration. Co-treatment of *Pten* deletion with either MEKca or STAT3ca modestly increased regeneration. Mice receiving the triple AAV

treatment (i.e. AAV-Cre, AAV-STAT3ca and AAV-MEKca) showed massive axon regeneration (Figures 5A and 5B). Single or double AAV-treated animals had only a few regenerating axons near the optic chiasm (i.e. 4–5 mm from the optic disc). In comparison, a large number of axons reached the chiasm in the triple AAV-mice (Figures 5A and 5B). Consistent with the single AAV injections, this triple AAV caused Cre expression in the majority of RGCs (approximately 90%) whereas about 70% and 60% of RGCs expressed MEKca and STAT3ca, respectively. We also observed that Cre expressing-RGCs also expressed both STAT3ca and HA (i.e. tag for MEKca) (Figures 5C and 5D). We also used tissue clearing and whole nerve imaging for analysis of axonal trajectories (Luo et al., 2014; Yungher et al., 2015). Three-dimensional examination in the cleared optic nerve shows that the degree of axon turning (i.e. U-turns) is similar among the animal groups, indicating that the higher number of axons distally near the chiasm in the triple AAV-mice is unlikely to be due to differences in axon navigation (Figures S6A and S6B). Analysis of RGC survival showed similar RGC numbers among all groups (Figure S6C). These data demonstrate that simultaneously modulating PTEN, STAT3 and MEK in neurons increase both the number of regenerating axons and the speed at which they extend when compared to single or double treatments.

To further assess the extent of RGC axon regeneration, we examined regeneration in the cleared, whole brains (Figures 6A and 6B) (Video 1). Four to 5 weeks post-injury, regenerating axons exit the optic chiasm and continue to extend extensively into the brain. Various growth pattern are seen with axons making turns and arbors within the brain (Figures S6D–S6G). No regenerating axons are seen in the brain of control animals receiving AAV-GFP (n=8; Figures S6H and S6I). To assess the progression of regeneration, animals were analyzed 10 weeks after injury. Extensive axon regeneration was seen in the brain as evident in the 3D projection and individual brain slices from the whole brain imaging (Figures 6C–6E) (Videos 2, 3). Consistent with previous studies demonstrating axon misguidance, many axons grew aberrantly throughout the brain (Luo et al., 2013; Pernet et al., 2012) (Figures 6C–6E) (Videos 2, 3).

### **STAT3/MEK with *Pten* deletion- effects on axon sprouting in the corticospinal tract (CST)**

In addition to axon regeneration, which typically refers to re-extension of injured axons, another form of axon remodeling that also contributes significantly to functional recovery after injury, is collateral sprouting of spared, uninjured axons (Tuszynski and Steward, 2012). It has been previously shown that *Pten* deletion or STAT3wt overexpression each promotes axonal sprouting to some degree after spinal cord injury (SCI) (Lang et al., 2013; Lee et al., 2014; Liu et al., 2010). Thus we examined whether the triple AAV treatment synergistically promotes extensive sprouting of uninjured axons after SCI. To this end, we assessed the degree of sprouting of CST neurons after unilateral pyramidotomy (Figure 7A). *Pten<sup>fl/fl</sup>;R26-tdTomato* mice received intracortical injection of AAV-Cre alone or AAV-Cre together with AAV-STAT3ca and AAV-MEKca at 1 week prior to injury. Two additional groups included were; i) *R26-tdTomato* receiving AAV-Cre injection (WT group) and ii) *R26-tdTomato* receiving AAV-Cre, AAV-STAT3ca and AAV-MEKca. The completeness of lesion was assessed by the degree of loss of PKC $\gamma$  immunostaining in the lesioned CST tract (Lee et al., 2014). The lesion resulted in essentially complete loss of PKC $\gamma$ , confirming the



severity of pyramidotomy (Figure S7A). At 4 weeks after injury, deletion of *Pten* alone elicited a significant increase in contralateral sprouting of CST axons (i.e. tdTomato<sup>+</sup> axons) from the intact side (Figures 7B and 7C). Co-treatment with AAV-STAT3ca and AAV-MEKca also enhanced axonal sprouting to some degree. Strikingly, we observed extensive axonal sprouting in animals subjected to the triple AAV (Figures 7B and 7C). Immunohistochemical analysis shows that many CST neurons were transduced by all three AAV types as evidenced by the concomitant tdTomato, STAT3 and HA (i.e. tag for MEKca) immunoreactivity in the same neurons (Figure S7B). The total numbers of tdTomato-labeled axons in the brainstem at the level of the medullary pyramid were similar among the different animal groups (Figures S7C and S7D). These data demonstrate markedly improved axonal sprouting in injured spinal cord after targeting STAT3 and MEK in *Pten*-deleted animals.

## DISCUSSION

Although significant progress has been made in identifying genetic factors involved in axon regeneration failure (Benowitz and Yin, 2007; Cho and Cavalli, 2014; Di Giovanni and Rathore, 2012; Gobrecht et al., 2014; Hammarlund and Jin, 2014; Jiao et al., 2005; Moore and Goldberg, 2011; Omura et al., 2015; Parikh et al., 2011; Planchamp et al., 2008; Saijilafu et al., 2013; Watkins et al., 2013), our understanding of this process remains incomplete. Even with some “fruitful” strategies so far, the number of regrowing axons beyond the lesion site and the extent to which these axons reach their targets remain generally low in adult mammals. One gene consistently implicated in axonal regeneration, in a variety of neuronal types, is Stat3. The mechanisms underlying the ability of STAT3 to promote regeneration, however, remain incompletely understood. This is reflected by the fact that merely overexpressing STAT3 in neurons yields limited axon regeneration *in vivo*. Our study not only extends the previous findings demonstrating mitochondrial presence of STAT3, but also provides direct *in vivo* evidence of site-specific STAT3 functions in coordinating CNS axon regeneration. Moreover, our results show that MEK is a critical factor that modulates STAT3's phosphorylation status, localization and functions in RGCs and cortical neurons. The strong regeneration phenotypes that we see, particularly in the STAT3ca and MEKca combined animals point to the importance of appropriately locating and activating STAT3 through MEK.

Previous studies using combinatorial approaches including *Pten* KO mice with CNTF<sup>12, 42</sup> or b-raf (O'Donovan et al., 2014)(an upstream regulator of MEK) also promoted long distance RGC axon regeneration (Luo et al., 2013; Sun et al., 2011), but the number of regenerating axons in these past studies dropped significantly with distances from the retina, with only a few RGC axons reaching the brain. In contrast, in the triple AAV mice used in this study, hundreds of axons extended into the brain and continued to elongate long distances, densely populating the brain. The same manipulations in CST neurons led to extensive collateral sprouting after SCI.

Recent studies have demonstrated STAT3's versatile ability to induce cell proliferation and axon growth by acting outside the nucleus, a function that is regulated by phosphorylation. However, whether such a mechanism exists in post-mitotic CNS neurons remained

undetermined. We show that STAT3 proteins are phosphorylated at both the tyrosine and serine residues in adult RGCS in response to CNTF. Further, we show that the level of pSTAT3(S727) is increased in mitochondria after CNTF stimulation in a MEK dependent-manner. Thus, our study is in agreement with previous studies demonstrating mitochondrial accumulation of STAT3 in different cell lines (Gough et al., 2009; Wegrzyn et al., 2009; Zhou and Too, 2011). Specifically, others have demonstrated that nerve growth factor (NGF) causes STAT3 phosphorylation exclusively on S727 and promotes mitochondrial translocation, and it is the mitochondrial (not the nuclear) STAT3 that mediates NGF's growth-promoting effects (Zhou and Too, 2011). It has also been demonstrated that cytoplasmic STAT3 is responsible for CNTF's axon growth promoting effects in cultured embryonic neurons (Selvaraj et al., 2012). Thus, while our study is in line with these previous studies describing STAT3's non-nuclear functions contributing to axon growth, our results also demonstrate the indispensable role of the transcriptional functions of STAT3 in adult CNS axon regeneration. Our results demonstrating a requirement for STAT3's transcriptional functions, in contrast to being nonessential in embryonic neurons (Selvaraj et al., 2012), suggest that neurons may acquire a reliance on STAT3's transcription activity to induce axon growth over the course of development. On the other hand, embryonic neurons are perhaps equipped with more versatile and flexible sets of transcription factors that can be activated to induce axon growth in the absence of STAT3's transcriptional regulation.

Regeneration of RGC axons in zebrafish, a regeneration-competent species, also involves JAK/STAT, but SOCS3 which normally inhibits STAT3's transcriptional activity did not block regeneration (Elsaedi et al., 2014). Whether any particular or all of STAT3's versatile actions contribute in the regeneration-competent species remains unknown. Another notable finding in our study is that although mitochondrial STAT3 alone is insufficient to promote regeneration, it further improves regeneration when the levels of nuclear active STAT3 is increased. Overall, our data indicate that distinctly distributed STAT3 in the nucleus and mitochondria modulate synergistic nuclear transcription and yet undefined mitochondrial mechanisms, and this bi-functionality of STAT3 induces optimal CNS axon regeneration. Interestingly, mitochondria STAT3 was also shown to facilitate the production of ROS, and this allowed neurite extension (Zhou and Too, 2011), suggesting that STAT3 may modulate different aspects of mitochondrial functions which coordinately improve regeneration.

Several studies have demonstrated that CNTF and STAT3 modulate mitochondrial bioenergetics; CNTF enhances respiratory capacity in sensory neurons (Chowdhury et al., 2014). STAT3 prevents a decrease in complex I activity under ischemia (Szczepanek et al., 2011). Furthermore, STAT3 ablated-cells exhibit defective ATP production (Sarafian et al., 2010; Wegrzyn et al., 2009). However, the precise mechanisms by which STAT3 locally modulates mitochondrial functions are currently unknown. Our study shows that STAT3 localizes to mitochondrial inner membrane in response to CNTF, and STAT3<sub>mts</sub> enhances cellular ATP levels, membrane potential and complex IV activity. Since *Cox10* ablated-RGCs have significantly reduced axon regeneration, improved regeneration seen after expression of STAT3<sub>mts</sub> could be due, at least in part, to its ability to maintain or enhance mitochondrial ATP production following injury. We also observed that *Cox* deletion reduces axon regeneration in PTEN ablated cells (data now shown), underlining the idea that the effects of ETC might be a general phenomenon for CNS axon regeneration.

It is notable that MEKca leads to considerable regeneration when it is combined with STAT3ca. MEK/ERK pathway is known to activate or silence many different proteins downstream, and thus controls many distinct cellular processes including gene transcription, translation as well as protein transport. Our data indicate that MEK-dependent localization of STAT3 to mitochondria could be one mechanism by which MEK facilitates STAT3-induced axon regeneration. Recently, it was demonstrated that retrogradely transported ERK activates histone acetyltransferases including p300/CBP-associated factor in the cell body, leading to enhanced acetylation and expression of several regeneration-associated genes (Puttagunta et al., 2014). Similarly, axon injury causes retrograde transport of axonal STAT3 to the cell body (Ben-Yaakov et al., 2012). Thus, in addition to acting downstream of growth factors at the soma, STAT3 and ERK may represent a complementary “dual” injury signal that collectively links the axon and soma response. In sum, our study extends STAT3's diverse abilities to function in distinct cellular regions in mature CNS neurons, but also indicates that in mature neurons, STAT3's transcription regulation is at the “center” of growth promotion with mitochondrial STAT3 likely playing facilitative roles, overall providing a more clear idea about how STAT3 could be manipulated to induce CNS axon regeneration. Delineating STAT3's divergent functions and understanding how these processes are regulated may lead to more targeted therapy for promoting functional axon regrowth.

## EXPERIMENTAL PROCEDURES

Further details of methods and materials are provided in Supplemental Experimental Procedures.

### Animals

All experimental procedures were performed in compliance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) at University of Miami. Animals used are; C57BL/6J, *Pten<sup>fl/fl</sup>* (stock number, 006440; Jackson Laboratory, Bar Harbor, ME, USA), *Stat3<sup>fl/fl</sup>* mice (stock number, 016923; Jackson Laboratory), *R26 loxP-stop-loxP-tdTomato* (*R26-tdTomato*; a generous gift from Dr. Fan Wang, Duke University), *Cox10<sup>fl/fl</sup>* (Fukui et al., 2007) and *Pten<sup>fl/fl</sup>;R26-tdTomato*, *Stat3<sup>fl/fl</sup>;R26-tdTomato*. For all surgical procedures, mice were anaesthetized with ketamine and xylazine, and buprenorphine (0.05mg/kg, Bedford Lab, Bedford, OH, USA) was administered as post-operative analgesic. At least 5 animals were used for each groups in the axon regeneration and RGC survival experiment unless indicated otherwise. The precise animal numbers are also indicated in the Figure Legends.

### Optic nerve crush

For optic nerve crush injury, the optic nerve was exposed intraorbitally and crushed with jeweler's forceps (Dumont #5, Roboz, Gaithersburg, MD, USA) for 10 seconds approximately 1 mm behind the optic disc.

## Intravitreal injection

Female or male mice (5–7 weeks old) received unilateral AAV injection. For each intravitreal injection, a glass micropipette was inserted into the peripheral retina, just behind the ora serrata, and was deliberately angled to avoid damage to the lens. To anterogradely label regenerating RGC axons, animals received intravitreal injection of 2  $\mu$ l of cholera toxin  $\beta$  subunit (CTB)-Alexa 555 or CTB-Alexa 488 (2  $\mu$ g/ $\mu$ l, Invitrogen, Waltham, MA, USA) with a Hamilton syringe (Hamilton, Reno, NE, USA) at 2–5 days before euthanasia.

## Statistics

GraphPad Prism (GraphPad Software, La Jolla, CA, USA) was used for statistical analyses. Data were analyzed using ANOVA and the Bonferroni within-groups comparison with additional testing using Dunnett's test or Student's t-test. Values of  $P < 0.05$  were considered significant. All error bars indicate standard error of mean (SEM) unless indicated otherwise in the Figure Legends.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

This work was supported by grants from NEI 1R01EY022961-01 (KKP), NINDS R01NS009923-43 (KKP), NICHD R01HD057632 (VPL and JLB), 5R01EY010804 (CTM), 5R01NS079965 (CTM), U.S Army W81XWH-05-1-0061 (KKP), W81XWH-12-1-0319 (KKP), Ziegler Foundation (KKP), Pew Charitable Trust (KKP), Craig H. Neilsen Foundation (KKP), and the Buoniconti Fund (KKP, VPL, JKL, JLB). We thank Aline-Ohashi Hida for her help with *Cox10<sup>fl/fl</sup>* mice, Melissa Carballosa-Gautam at the Miami Project to Cure Paralysis Imaging Core, Margaret Rebecca Bates and Vania Wolff Almeida at the University of Miami Electron Microscopy Core, Yadira Salgueiro in Park Lab and Pingping Jia at the University of Miami Viral Vector Core.


## References

- Agathocleous M, Love NK, Randlett O, Harris JJ, Liu J, Murray AJ, Harris WA. Metabolic differentiation in the embryonic retina. *Nat Cell Biol.* 2012; 14:859–864. [PubMed: 22750943]
- Bareyre FM, Garzorz N, Lang C, Misgeld T, Buning H, Kerschensteiner M. In vivo imaging reveals a phase-specific role of STAT3 during central and peripheral nervous system axon regeneration. *Proc Natl Acad Sci U S A.* 2011; 108:6282–6287. [PubMed: 21447717]
- Ben-Yaakov K, Dagan SY, Segal-Ruder Y, Shalem O, Vuppalanchi D, Willis DE, Yudin D, Rishal I, Rother F, Bader M, et al. Axonal transcription factors signal retrogradely in lesioned peripheral nerve. *EMBO J.* 2012; 31:1350–1363. [PubMed: 22246183]
- Benowitz LI, Yin Y. Combinatorial treatments for promoting axon regeneration in the CNS: strategies for overcoming inhibitory signals and activating neurons' intrinsic growth state. *Dev Neurobiol.* 2007; 67:1148–1165. [PubMed: 17514713]
- Boehm JS, Zhao JJ, Yao J, Kim SY, Firestein R, Dunn IF, Sjöström SK, Garraway LA, Weremowicz S, Richardson AL, et al. Integrative genomic approaches identify IKBKE as a breast cancer oncogene. *Cell.* 2007; 129:1065–1079. [PubMed: 17574021]
- Bromberg JF, Horvath CM, Besser D, Lathem WW, Darnell JE Jr. Stat3 activation is required for cellular transformation by v-src. *Mol Cell Biol.* 1998; 18:2553–2558. [PubMed: 9566875]
- Bromberg JF, Wrzeszczynska MH, Devgan G, Zhao Y, Pestell RG, Albanese C, Darnell JE Jr. Stat3 as an oncogene. *Cell.* 1999; 98:295–303. [PubMed: 10458605]

- Cafferty WB, Gardiner NJ, Das P, Qiu J, McMahon SB, Thompson SW. Conditioning injury-induced spinal axon regeneration fails in interleukin-6 knock-out mice. *J Neurosci*. 2004; 24:4432–4443. [PubMed: 15128857]
- Cao Z, Gao Y, Bryson JB, Hou J, Chaudhry N, Siddiq M, Martinez J, Spencer T, Carmel J, Hart RB, et al. The cytokine interleukin-6 is sufficient but not necessary to mimic the peripheral conditioning lesion effect on axonal growth. *J Neurosci*. 2006; 26:5565–5573. [PubMed: 16707807]
- Case LC, Tessier-Lavigne M. Regeneration of the adult central nervous system. *Curr Biol*. 2005; 15:R749–753. [PubMed: 16169471]
- Cho Y, Cavalli V. HDAC signaling in neuronal development and axon regeneration. *Curr Opin Neurobiol*. 2014; 27:118–126. [PubMed: 24727244]
- Chowdhury SR, Saleh A, Akude E, Smith DR, Morrow D, Tessler L, Calcutt NA, Fernyhough P. Ciliary neurotrophic factor reverses aberrant mitochondrial bioenergetics through the JAK/STAT pathway in cultured sensory neurons derived from streptozotocin-induced diabetic rodents. *Cell Mol Neurobiol*. 2014; 34:643–649. [PubMed: 24682898]
- Cregg JM, DePaul MA, Filous AR, Lang BT, Tran A, Silver J. Functional regeneration beyond the glial scar. *Exp Neurol*. 2014; 253:197–207. [PubMed: 24424280]
- Di Giovanni S, Rathore K. p53-Dependent pathways in neurite outgrowth and axonal regeneration. *Cell Tissue Res*. 2012; 349:87–95. [PubMed: 22271139]
- Elsaiedi F, Bemben MA, Zhao XF, Goldman D. Jak/Stat signaling stimulates zebrafish optic nerve regeneration and overcomes the inhibitory actions of Socs3 and Sfpq. *J Neurosci*. 2014; 34:2632–2644. [PubMed: 24523552]
- Ernst M, Jenkins BJ. Acquiring signalling specificity from the cytokine receptor gp130. *Trends Genet*. 2004; 20:23–32. [PubMed: 14698616]
- Fukui H, Diaz F, Garcia S, Moraes CT. Cytochrome c oxidase deficiency in neurons decreases both oxidative stress and amyloid formation in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A*. 2007; 104:14163–14168. [PubMed: 17715058]
- Gallagher D, Gutierrez H, Gavalda N, O'Keefe G, Hay R, Davies AM. Nuclear factor-kappaB activation via tyrosine phosphorylation of inhibitor kappaB-alpha is crucial for ciliary neurotrophic factor-promoted neurite growth from developing neurons. *J Neurosci*. 2007; 27:9664–9669. [PubMed: 17804627]
- Gobrecht P, Leibinger M, Andreadaki A, Fischer D. Sustained GSK3 activity markedly facilitates nerve regeneration. *Nat Commun*. 2014; 5:4561. [PubMed: 25078444]
- Gough DJ, Corlett A, Schlessinger K, Wegrzyn J, Larner AC, Levy DE. Mitochondrial STAT3 supports Ras-dependent oncogenic transformation. *Science*. 2009; 324:1713–1716. [PubMed: 19556508]
- Hammarlund M, Jin Y. Axon regeneration in *C. elegans*. *Curr Opin Neurobiol*. 2014; 27:199–207. [PubMed: 24794753]
- Horvath CM, Wen Z, Darnell JE Jr. A STAT protein domain that determines DNA sequence recognition suggests a novel DNA-binding domain. *Genes Dev*. 1995; 9:984–994. [PubMed: 7774815]
- Huebner EA, Strittmatter SM. Axon regeneration in the peripheral and central nervous systems. *Results Probl Cell Differ*. 2009; 48:339–351. [PubMed: 19582408]
- Jiao J, Huang X, Feit-Leithman RA, Neve RL, Snider W, Dartt DA, Chen DF. Bcl-2 enhances Ca(2+) signaling to support the intrinsic regenerative capacity of CNS axons. *EMBO J*. 2005; 24:1068–1078. [PubMed: 15719013]
- Lang C, Bradley PM, Jacobi A, Kerschensteiner M, Bareyre FM. STAT3 promotes corticospinal remodelling and functional recovery after spinal cord injury. *EMBO Rep*. 2013; 14:931–937. [PubMed: 23928811]
- Leaver SG, Cui Q, Bernard O, Harvey AR. Cooperative effects of bcl-2 and AAV-mediated expression of CNTF on retinal ganglion cell survival and axonal regeneration in adult transgenic mice. *Eur J Neurosci*. 2006; 24:3323–3332. [PubMed: 17229081]
- Lee DH, Luo X, Yungher BJ, Bray E, Lee JK, Park KK. Mammalian target of rapamycin's distinct roles and effectiveness in promoting compensatory axonal sprouting in the injured CNS. *J Neurosci*. 2014; 34:15347–15355. [PubMed: 25392502]

- Leibinger M, Andreadaki A, Diekmann H, Fischer D. Neuronal STAT3 activation is essential for CNTF- and inflammatory stimulation-induced CNS axon regeneration. *Cell Death Dis.* 2013; 4:e805. [PubMed: 24052073]
- Leibinger M, Muller A, Andreadaki A, Hauk TG, Kirsch M, Fischer D. Neuroprotective and axon growth-promoting effects following inflammatory stimulation on mature retinal ganglion cells in mice depend on ciliary neurotrophic factor and leukemia inhibitory factor. *J Neurosci.* 2009; 29:14334–14341. [PubMed: 19906980]
- Liu K, Lu Y, Lee JK, Samara R, Willenberg R, Sears-Kraxberger I, Tedeschi A, Park KK, Jin D, Cai B, et al. PTEN deletion enhances the regenerative ability of adult corticospinal neurons. *Nat Neurosci.* 2010; 13:1075–1081. [PubMed: 20694004]
- Luo X, Salgueiro Y, Beckerman SR, Lemmon VP, Tsoulfas P, Park KK. Three-dimensional evaluation of retinal ganglion cell axon regeneration and pathfinding in whole mouse tissue after injury. *Exp Neurol.* 2013; 247:653–662. [PubMed: 23510761]
- Luo X, Yungker B, Park KK. Application of tissue clearing and light sheet fluorescence microscopy to assess optic nerve regeneration in unsectioned tissues. *Methods Mol Biol.* 2014; 1162:209–217. [PubMed: 24838970]
- Martell JD, Deerinck TJ, Sancak Y, Poulos TL, Mootha VK, Sosinsky GE, Ellisman MH, Ting AY. Engineered ascorbate peroxidase as a genetically encoded reporter for electron microscopy. *Nat Biotechnol.* 2012; 30:1143–1148. [PubMed: 23086203]
- Mattson MP. Mitochondrial regulation of neuronal plasticity. *Neurochem Res.* 2007; 32:707–715. [PubMed: 17024568]
- Mogi T, Saiki K, Anraku Y. Biosynthesis and functional role of haem O and haem A. *Mol Microbiol.* 1994; 14:391–398. [PubMed: 7885224]
- Moore DL, Goldberg JL. Multiple transcription factor families regulate axon growth and regeneration. *Dev Neurobiol.* 2011; 71:1186–1211. [PubMed: 21674813]
- Muller A, Hauk TG, Leibinger M, Marienfeld R, Fischer D. Exogenous CNTF stimulates axon regeneration of retinal ganglion cells partially via endogenous CNTF. *Mol Cell Neurosci.* 2009; 41:233–246. [PubMed: 19332123]
- O'Donovan KJ, Ma K, Guo H, Wang C, Sun F, Han SB, Kim H, Wong JK, Charron J, Zou H, et al. B-RAF kinase drives developmental axon growth and promotes axon regeneration in the injured mature CNS. *J Exp Med.* 2014; 211:801–814. [PubMed: 24733831]
- Omura T, Omura K, Tedeschi A, Riva P, Painter MW, Rojas L, Martin J, Lisi V, Huebner EA, Latremoliere A, et al. Robust Axonal Regeneration Occurs in the Injured CAST/Ei Mouse CNS. *Neuron.* 2015; 86:1215–1227. [PubMed: 26004914]
- Parikh P, Hao Y, Hosseinkhani M, Patil SB, Huntley GW, Tessier-Lavigne M, Zou H. Regeneration of axons in injured spinal cord by activation of bone morphogenetic protein/Smad1 signaling pathway in adult neurons. *Proc Natl Acad Sci U S A.* 2011; 108:E99–107. [PubMed: 21518886]
- Park KK, Liu K, Hu Y, Smith PD, Wang C, Cai B, Xu B, Connolly L, Kramvis I, Sahin M, et al. Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway. *Science.* 2008; 322:963–966. [PubMed: 18988856]
- Pellegrino MJ, Habecker BA. STAT3 integrates cytokine and neurotrophin signals to promote sympathetic axon regeneration. *Mol Cell Neurosci.* 2013; 56:272–282. [PubMed: 23831387]
- Pernet V, Hauswirth WW, Di Polo A. Extracellular signal-regulated kinase 1/2 mediates survival, but not axon regeneration, of adult injured central nervous system neurons in vivo. *J Neurochem.* 2005; 93:72–83. [PubMed: 15773907]
- Pernet V, Joly S, Dalkara D, Jordi N, Schwarz O, Christ F, Schaffer DV, Flannery JG, Schwab ME. Long-distance axonal regeneration induced by CNTF gene transfer is impaired by axonal misguidance in the injured adult optic nerve. *Neurobiol Dis.* 2012
- Pernet V, Joly S, Jordi N, Dalkara D, Guzik-Kornacka A, Flannery JG, Schwab ME. Misguidance and modulation of axonal regeneration by Stat3 and Rho/ROCK signaling in the transparent optic nerve. *Cell Death Dis.* 2013; 4:e734. [PubMed: 23868067]
- Planchamp V, Bermel C, Tonges L, Ostendorf T, Kugler S, Reed JC, Kermer P, Bahr M, Lingor P. BAG1 promotes axonal outgrowth and regeneration in vivo via Raf-1 and reduction of ROCK activity. *Brain.* 2008; 131:2606–2619. [PubMed: 18757464]

- Puttagunta R, Tedeschi A, Soria MG, Hervera A, Lindner R, Rathore KI, Gaub P, Joshi Y, Nguyen T, Schmandke A, et al. PCAF-dependent epigenetic changes promote axonal regeneration in the central nervous system. *Nat Commun.* 2014; 5:3527. [PubMed: 24686445]
- Sajilafu, Zhang BY, Zhou FQ. Signaling pathways that regulate axon regeneration. *Neurosci Bull.* 2013; 29:411–420. [PubMed: 23846598]
- Saleh A, Roy Chowdhury SK, Smith DR, Balakrishnan S, Tessler L, Martens C, Morrow D, Schartner E, Frizzi KE, Calcutt NA, et al. Ciliary neurotrophic factor activates NF-kappaB to enhance mitochondrial bioenergetics and prevent neuropathy in sensory neurons of streptozotocin-induced diabetic rodents. *Neuropharmacology.* 2013; 65:65–73. [PubMed: 23022047]
- Sarafian TA, Montes C, Imura T, Qi J, Coppola G, Geschwind DH, Sofroniew MV. Disruption of astrocyte STAT3 signaling decreases mitochondrial function and increases oxidative stress in vitro. *PLoS One.* 2010; 5:e9532. [PubMed: 20224768]
- Selvaraj BT, Frank N, Bender FL, Asan E, Sendtner M. Local axonal function of STAT3 rescues axon degeneration in the pmn model of motoneuron disease. *J Cell Biol.* 2012; 199:437–451. [PubMed: 23109669]
- Smith PD, Sun F, Park KK, Cai B, Wang C, Kuwako K, Martinez-Carrasco I, Connolly L, He Z. SOCS3 deletion promotes optic nerve regeneration in vivo. *Neuron.* 2009; 64:617–623. [PubMed: 20005819]
- Sun F, He Z. Neuronal intrinsic barriers for axon regeneration in the adult CNS. *Curr Opin Neurobiol.* 2010; 20:510–518. [PubMed: 20418094]
- Sun F, Park KK, Belin S, Wang D, Lu T, Chen G, Zhang K, Yeung C, Feng G, Yankner BA, et al. Sustained axon regeneration induced by co-deletion of PTEN and SOCS3. *Nature.* 2011; 480:372–375. [PubMed: 22056987]
- Szczepanek K, Chen Q, Derecka M, Salloum FN, Zhang Q, Szelag M, Cichy J, Kukreja RC, Dulak J, Lesnefsky EJ, et al. Mitochondrial-targeted Signal transducer and activator of transcription 3 (STAT3) protects against ischemia-induced changes in the electron transport chain and the generation of reactive oxygen species. *J Biol Chem.* 2011; 286:29610–29620. [PubMed: 21715323]
- Tammineni P, Anugula C, Mohammed F, Anjaneyulu M, Larner AC, Sepuri NB. The import of the transcription factor STAT3 into mitochondria depends on GRIM-19, a component of the electron transport chain. *J Biol Chem.* 2013; 288:4723–4732. [PubMed: 23271731]
- Taniguchi K, Wu LW, Grivennikov SI, de Jong PR, Lian I, Yu FX, Wang K, Ho SB, Boland BS, Chang JT, et al. A gp130-Src-YAP module links inflammation to epithelial regeneration. *Nature.* 2015; 519:57–62. [PubMed: 25731159]
- Tuszynski MH, Steward O. Concepts and methods for the study of axonal regeneration in the CNS. *Neuron.* 2012; 74:777–791. [PubMed: 22681683]
- Watkins TA, Wang B, Huntwork-Rodriguez S, Yang J, Jiang Z, Eastham-Anderson J, Modrusan Z, Kaminker JS, Tessier-Lavigne M, Lewcock JW. DLK initiates a transcriptional program that couples apoptotic and regenerative responses to axonal injury. *Proc Natl Acad Sci U S A.* 2013; 110:4039–4044. [PubMed: 23431164]
- Wegrzyn J, Potla R, Chwae YJ, Sepuri NB, Zhang Q, Koeck T, Derecka M, Szczepanek K, Szelag M, Gornicka A, et al. Function of mitochondrial Stat3 in cellular respiration. *Science.* 2009; 323:793–797. [PubMed: 19131594]
- Yungher BJ, Luo X, Salgueiro Y, Blackmore MG, Park KK. Viral vector-based improvement of optic nerve regeneration: characterization of individual axons' growth patterns and synaptogenesis in a visual target. *Gene Ther.* 2015; 22:811–821. [PubMed: 26005861]
- Zhou L, Too HP. Mitochondrial localized STAT3 is involved in NGF induced neurite outgrowth. *PLoS One.* 2011; 6:e21680. [PubMed: 21738764]
- Zigmond RE. gp130 cytokines are positive signals triggering changes in gene expression and axon outgrowth in peripheral neurons following injury. *Front Mol Neurosci.* 2011; 4:62. [PubMed: 22319466]



**Figure 1. CNTF induces phosphorylation and translocation of STAT3 proteins to distinct subcellular regions: MEK-dependent mitochondrial translocation**

(A) Retinal sections showing the levels of pSTAT3(Y705) and pSTAT3(S727) immunoreactivity (red) in RGCs (i.e. labeled with the TUJ1 antibody, green) 2 weeks following intravitreal AAV-GFP (“control”) or AAV-CNTF injection (“CNTF”). DAPI in blue. Arrow, RGC process.

(B) Single plane confocal image showing co-labeling of mitochondrial protein ATPsynβ and pSTAT3(S727) in RGCs in AAV-CNTF-treated retina.

(C) Quantification of pSTAT3(S727) and ATPsynβ co-localization. For each retina, 15–20 individual RGCs were measured, and 3 retinas were analyzed per group. Value of the control group (AAV-GFP) is represented as 100%. \*\* $P < 0.01$  by unpaired Student's t-test.


(D) Upper left image, brightfield image showing a P3 rat cortical neuron transfected with APEX-STAT3, (green arrow head) and an untransfected cell (red arrow head). Bottom left image, electron micrographs of a neuron transfected with APEX-STAT3 treated with recombinant CNTF (rCNTF; 50 ng/ml); the inset is a magnification of the white boxed area, showing mitochondria. Upper right image, APEX-STAT3 without rCNTF. Bottom right image, HA-APEX with rCNTF. DAB signal is detected in the inner membrane (red arrow) of mitochondria in APEX-STAT3 cells. On the other hand, while DAB signal is found throughout the cytoplasm, it is not detected in the mitochondria of HA-APEX cells. Scale bars: low magnification 1  $\mu$ m, inset 100 nm.

(E) Western blot of P3 rat cortical neurons under various conditions; no treatment, treated with rCNTF (50 ng/mL) alone or with MEK inhibitors U0126 (5  $\mu$ M) or refamitinib (Refm, 5  $\mu$ M) or vehicle (Veh, DMSO) for 1 hour. The control (“cntrl”) group received neither rCNTF nor MEK inhibitors.



(F) Representative western blot of fractionation-derived nuclear, cytosolic and mitochondrial proteins from rat P3 cortical neurons showing the levels of STAT3 protein expression 4 hours after rCNTF (50 ng/mL). Total STAT3 and pSTAT3(S727) are increased in the mitochondria after rCNTF. Increases in the level of total STAT3 (but not pSTAT3(S727)) was also detected; data shown are result of 15 second exposure for the cytosol and mitochondrial fractionations, and 1 minutes exposure for the nuclear fraction. Voltage-dependent anion channel (VDAC) mitochondrial marker, nucleolin (NCL) nuclear marker and vinculin (VCL), cytosolic marker were used to verify the relative purity of fractionated proteins.

(G) Densitometric quantification of mitochondrial STAT3 levels in western blot normalized against VDAC. Both the total and pSTAT3(S727) protein levels are increased in mitochondria after rCNTF treatment, an effect that was reduced by refamitinib. Value for the control group is represented as 100%. Values are means obtained from at least 6 biological replicates. \* $P < 0.05$ , \*\* $P < 0.01$  by one-way ANOVA followed by Dunnett's test. Scale bars, 20  $\mu\text{m}$  in (A) and (B).



**Figure 2. DNA binding domain of STAT3 is essential for CNTF induced-RGC axon regeneration** (A) Time line of STAT3 *in vivo* rescue experiment. AAV-Cre together with either AAV-GFP or AAV vectors carrying different versions of STAT3 were injected intravitreally in *R26-tdTomato* or *Stat3<sup>fl/fl</sup>;R26-tdTomato* mice. Five days later, animals received optic nerve crush plus AAV-GFP or AAV-CNTF injection.


(B) Whole-mount retinal staining showing Cre, tdTomato and TUJ1 immunoreactivity 2 weeks after AAV-Cre injection in a *R26-tdTomato* mouse. (B') Higher magnification of the boxed area in (B).

(C) Representative optic nerve sections showing the degree of axon regeneration following various treatments. Axons are labeled with tdTomato via Cre recombination in *Stat3<sup>fl/fl</sup>;R26-tdTomato* or *R26-tdTomato* mice. STAT3wt, but not STAT3dbm rescues AAV-CNTF induced regeneration. Red asterisks, lesion site.

(D) Quantification of tdTomato-labeled regenerating axons in various animal groups; “GFP”, AAV-Cre/AAV-GFP co- injected *R26-tdTomato* mice. “CNTF + GFP”, AAV-Cre/AAV-GFP co-injected *R26-tdTomato* mice with AAV-CNTF injection at the time of injury. “STAT3 KO + CNTF + GFP”, AAV-Cre/AAV-GFP co-injected *Stat3<sup>fl/fl</sup>;R26-tdTomato* mice with AAV-CNTF injection at the time of injury. “STAT3 KO + CNTF + STAT3wt”, AAV-Cre and AAV-STAT3wt co-injected *Stat3<sup>fl/fl</sup>;R26-tdTomato* mice with AAV-CNTF injection at the time of injury. “STAT3 KO + CNTF + STAT3dbm”, AAV-Cre and AAV-STAT3dbm co-injected *Stat3<sup>fl/fl</sup>;R26-tdTomato* mice with AAV-CNTF injection at the time of injury. n=5 animals/group. \*\*\**P*<0.001 compared to the “GFP group” using two-way repeated measures ANOVA followed by Bonferroni post-test.

(E) Whole-mount retinal immunohistochemistry showing transduction efficacy of AAV-STAT3wt and AAV-STAT3dbm in RGCs. C57BL/6 mice received intravitreal AAV injection

and the tissues were analyzed 2 week later. Approximately 60% of RGCs (TUJ1<sup>+</sup>, green) are transduced as evident by STAT3 immunoreactivity (red) compared to undetectable STAT3 seen in the AAV-GFP injected retina. Scale bars: 100  $\mu\text{m}$  in (B) and (C), 20  $\mu\text{m}$  in (B') and (E).



**Figure 3. Site-specific STAT3 effects - mitochondrial STAT3 facilitates axon regeneration in the presence of STAT3ca**

(A) Time line of the experiment.


(B) Representative optic nerve sections showing the degree of axon regeneration in animals treated with either AAV-GFP, AAV-STAT3ca or AAV-STAT3mts alone, or AAV-STAT3ca/AAV-STAT3mts together 2 weeks prior to injury. Axons are anterogradely labeled by CTB. Red asterisks, lesion site.

(C) Quantification of CTB-labeled regenerating axons in various animal groups (AAV-GFP, AAV-STAT3wt, AAV-STAT3ca or AAV-STAT3mts alone, or AAV-STAT3ca/AAV-STAT3mts).  $n=4-7$  animals/group.  $*P<0.05$ ,  $***P<0.001$ ,  $****P<0.0001$  by two-way repeated measures ANOVA followed by Bonferroni post-test.

(D) whole-mount retinal immunohistochemistry showing transduction efficacy of AAV-STAT3ca and AAV-STAT3mts. AAV was injected in adult WT mice, and retinas were removed 2 weeks post-injection. Approximately 60% of RGCs (TUJ1<sup>+</sup>, green) are transduced as evident by STAT3 immunoreactivity (red) compared to undetectable STAT3 seen in the AAV-GFP injected retina.

(E) Representative retinal whole-mount images with TUJ1 staining at 2 weeks post-injury in animals subjected to either AAV-GFP, AAV-STAT3ca or AAV-STAT3mts alone, or AAV-STAT3ca/AAV-STAT3mts.

(F) Quantification of RGC survival. Values are represented as the percentage of the contralateral uninjured retina.  $n=6-7$  animals/group. Scale bars, 100 μm in (B), 20 μm in (D) and (E).



#### Figure 4. MEK enhances STAT3-mediated axon regeneration


(A) Representative optic nerve sections showing the degree of axon regeneration 4 weeks after injury in animals treated with either AAV-GFP, AAV-STAT3ca or AAV-MEKca alone, or AAV-STAT3ca/AAV-MEKca. Axons are labeled anterogradely with CTB. Red asterisks, lesion site.

(B) Quantification of CTB-labeled regenerating axons in the various animal groups.  $n=4-6$  animals/group.  $*P<0.05$ ,  $**P<0.01$ ,  $***P<0.001$ ,  $****P<0.0001$  compared to the “GFP” control group using two-way repeated measures ANOVA followed by Bonferroni post-test.

(C) Whole-mount retinal immunohistochemistry showing transduction efficacy of the AAV-MEKca. Adult mice received AAV injection, and retinas removed 2 weeks post-injection. Approximately 70% of RGCs are transduced by the AAV-MEKca as evident by the expression of HA (HA-MEKca) (red) in TUJ1<sup>+</sup> RGCs (green).

(D) Representative retinal whole-mount images with TUJ1 staining at 4 weeks post-injury in animals subjected to either AAV-GFP, AAV-STAT3ca or AAV-MEKca alone, or combined.

(E) Quantification of RGC survival. Values are represented as the percentage of the contralateral uninjured retina.  $n=6$  animals/group.  $***P<0.001$ ,  $****P<0.0001$  compared to the “GFP” control group using Bonferroni post-test. Scale bars; 100 μm in (A), 20 μm in (C) and (D).




**Figure 5. Simultaneously modulating PTEN, STAT3 and MEK in RGCs results in extensive axon regeneration**

(A) Representative optic nerve sections showing the degree of axon regeneration 4 weeks after injury in *Pten<sup>fl/fl</sup>* animals treated with either AAV-Cre alone or AAV-Cre/AAV-MEKca or AAV-Cre/AAV-STAT3ca or all three AAV together (i.e. AAV-Cre/AAV-STAT3ca/AAV-MEKca). Axons are labeled anterogradely with CTB. Red asterisks, lesion site.

(B) Quantification of CTB-labeled regenerating axons in the various animal groups. n=6-7 animals/group. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\*\* $P < 0.0001$  compared to the “PTEN KO” group using two-way repeated measures ANOVA followed by Bonferroni post-test.

(C) Whole-mount retinal staining showing AAV transduction efficacy after injection of the triple AAV mix (AAV-Cre, AAV-MEKca and AAV-STAT3ca) shown by the expression of Cre (green), STAT3 (red), HA (i.e. tag for MEKca) (white) in TUJ1<sup>+</sup> RGCs (blue). Adult mice received AAV injection, and retinas removed 2 weeks post-injection.

(D) Euler diagram showing the percentages of RGC transduction by each AAV and the co-transduction levels. Scale bars, 100  $\mu\text{m}$  in (A), 20  $\mu\text{m}$  in (C).



**Figure 6. Whole brain imaging reveals extensive RGC axons regeneration following the triple AAV treatment**


(A) Mouse brains with or without the tissue clearing procedure.

(B) Dorsal view of a cleared brain of uninjured mouse injected with CTB555 (pseudo-colored in white) and CTB488 in the left and right eyes, respectively, showing the entire trajectories of RGC axons from each eye. Tissues were processed 3 days after CTB injection.

(C) Image from axonal projection in the whole, cleared brain of a triple AAV-treated animal 10 weeks after injury.

(D) Higher magnification of the yellow boxed area in (C).

(E) Higher magnification of the blue boxed area in (C). CTB<sup>+</sup> axons grow beyond the optic chiasm and continue to extend to the intact optic nerve or into the brain. OX, optic chiasm; SC, superior colliculus. Scale bars, 2 mm in (A), 500  $\mu$ m in (B) and (C), 50  $\mu$ m in (E).



**Figure 7. Modulation of PTEN, STAT3 and MEK results in extensive CST sprouting after SCI** (A) Schematic of pyramidotomy model. Corticospinal tract axons (green) originating from the motor cortex are unilaterally transected. Intracortical injection of AAVs into the contralateral brain side allows the examination of treatment effects on axonal sprouting of intact nerves (red).

(B) Quantifications of tdTomato<sup>+</sup> crossing axons counted at < 50, 250 and 500 μm lateral to the midline in the wild-type control (“WT”), single (“PTEN KO”), double (“STAT3ca + MEKca”) and triple AAV (“PTEN KO + STAT3ca + MEKca”) animal groups normalized against the numbers of total labeled CST axons in the medulla. n=5–6 animals/group. \**P*<0.05, \*\**P*<0.01, \*\*\**P*<0.001, \*\*\*\**P*<0.0001 compared to the WT group using two-way repeated measures ANOVA followed by Bonferroni post-test.

(C) Representative of C5 coronal sections in low magnification. High magnification of the boxed regions in (C) are shown in (C') (midline region) and (C'') (lateral region) of the single, double and triple AAV animal groups. Scale bar, 500 μm.