

# Risk of Infectious Complications in Hemato-Oncological Patients Treated with Kinase Inhibitors

Mark Reinwald, Tobias Boch, Wolf-Karsten Hofmann and Dieter Buchheidt

Department of Hematology and Oncology, Mannheim University Hospital, University of Heidelberg, Mannheim, Germany.

## Supplementary Issue: Tyrosine Kinases in Cancer

**ABSTRACT:** Infectious complications are a major cause of morbidity and mortality in patients with hemato-oncological diseases. Although disease-related immunosuppression represents one factor, aggressive treatment regimens, such as chemotherapy, stem cell transplantation, or antibody treatment, account for a large proportion of infectious side effects. With the advent of targeted therapies affecting specific kinases in malignant diseases, the outcome of patients has further improved. Nonetheless, dependent on the specific pathway targeted or off-target activity of the kinase inhibitor, therapy-associated infectious complications may occur. We review the most common and approved kinase inhibitors targeting a variety of hemato-oncological malignancies for their immunosuppressive potential and evaluate their risk of infectious side effects based on preclinical evidence and clinical data in order to raise awareness of the potential risks involved.

**KEYWORDS:** TKI, kinase inhibitors, infections, infectious complications, targeted therapy, EGFR, BCR-ABL, BRAF, VEGF, BCR signaling, mTOR, JAK

**SUPPLEMENT:** Tyrosine Kinases in Cancer

**CITATION:** Reinwald et al. Risk of Infectious Complications in Hemato-Oncological Patients Treated with Kinase Inhibitors. *Biomarker Insights* 2015:10(S3) 55–68  
doi: 10.4137/BMI.S22430.

**TYPE:** Review

**RECEIVED:** May 06, 2015. **RESUBMITTED:** September 13, 2015. **ACCEPTED FOR PUBLICATION:** September 15, 2015.

**ACADEMIC EDITOR:** Karen Pulford, Editor in Chief

**PEER REVIEW:** Three peer reviewers contributed to the peer review report. Reviewers' reports totaled 804 words, excluding any confidential comments to the academic editor.

**FUNDING:** Authors disclose no funding sources.

**COMPETING INTERESTS:** MR received personal fees (lecture honoraria) from Roche and Pfizer, and grants and fees from Gilead. Other authors disclose no potential conflicts of interest.

**CORRESPONDENCE:** mark.reinwald@medma.uni-heidelberg.de

**COPYRIGHT:** © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

Paper subject to independent expert blind peer review. All editorial decisions made by independent academic editor. Upon submission manuscript was subject to anti-plagiarism scanning. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties. This journal is a member of the Committee on Publication Ethics (COPE).

Published by Libertas Academica. Learn more about this journal.

## Introduction

Over the last decade, the introduction of novel, innovative targeted therapies in hemato-oncological diseases has made a significant impact on the current therapeutic strategies and is associated with an outstanding clinical benefit for hematological diseases such as chronic myeloid leukemia (CML),<sup>1</sup> myelofibrosis (MF), and polycythemia vera (PV),<sup>2</sup> with additional improvement in solid tumors such as renal cell cancer,<sup>3</sup> lung cancer,<sup>4</sup> and melanoma.<sup>5</sup> Besides antibody therapy targeting specific epitopes, the mainstay of targeted therapies is the inhibition of specific tyrosine or serine/threonine kinases that have been implicated in disease pathogenesis and prognosis for a variety of diseases. Thus, kinase inhibitors (KIs) have undoubtedly become one of the major advances in the treatment of malignant diseases in recent years.

Yet, unpredictable and unexpected adverse effects have been reported with the use of targeted therapies, with several of these effects being recognized after long-term clinical use such as reversible posterior leukoencephalopathy by anti-vascular endothelial growth factor (VEGF) agents<sup>6</sup> or osteonecrosis of the jaw, which is associated with bisphosphonate treatment.

As several of these targeted agents affect critical components of the immune system, they are associated with infectious

complications with potentially life-threatening consequences. This is of importance as infections in cancer patients represent a typical, often therapy-associated complication,<sup>7</sup> and it has been noted that for several hematologic malignancies, infections are the major contributor to nonrelapse mortality.<sup>8</sup>

Therefore, in this review, we give an overview on the infectious complications of different tyrosine kinase inhibitor (TKI) classes affecting different hemato-oncological diseases based on preclinical and clinical evidence. We review KIs such as epidermal growth factor receptor (EGFR) inhibitors (eg, erlotinib), multikinase inhibitors such as sorafenib, the fusion gene resulting from a fusion of the breakpoint cluster region-gene and the Abelson murine leukemia viral oncogene homolog (BCR-ABL) inhibitors (eg, imatinib), inhibitors of mammalian target of rapamycin (mTOR), inhibitors of the Janus kinase (JAK), BRAF inhibitors such as vemurafenib, and most recently, inhibitors of the B-cell receptor (BCR) signaling pathway such as ibrutinib and idelalisib. The focus of this review is on the infections associated with these classes of drugs in order to raise awareness of these complications in the treating physician. The specific KI, its mode of action and associated frequency of infectious complications are summarized in table 1 while typical infections associated with this KI and possible prophylactic measures are summarized in Table 2.



## Inhibitors of BCR-ABL Tyrosine Kinase

Based on the results from the pivotal International Randomized Study of Interferon and STI571 (IRIS) trial in 2002, imatinib, as the first approved inhibitor of BCR-ABL tyrosine kinase, heralded the age of KI therapy and revolutionized the treatment of CML<sup>9</sup> and later on also gastrointestinal stromal tumors (GISTs) due to its additional activity in targeting c-Kit.<sup>10</sup> BCR-ABL, a fusion protein that results from the translocation (9;22) is the major hallmark that drives the malignant phenotype of CML, its inhibition suppressing the growth advantage of the transformed cells and potentially inducing even molecular remissions. In addition, for the subgroup of patients with acute lymphoblastic leukemia (ALL) and 9;22 translocation, inhibition of BCR-ABL added to conventional chemotherapy is the standard of care. The spectrum of KIs inhibiting BCR-ABL has grown with dasatinib, nilotinib, bosutinib, and most recently, ponatinib, further broadening the therapeutic armamentarium with the capacity of targeting mutations conferring resistance to imatinib.<sup>11–14</sup> However, all BCR-ABL-targeting KIs also potentially affect other targets such as SRC-family kinases as well as c-Kit, platelet-derived growth factor receptor (PDGFR)-a and -b, and ephrin receptor kinase<sup>11</sup>, thus carrying the potential for infectious complications.

Besides inducing neutropenia and therefore increasing the likelihood of infections, preclinical studies have shown that imatinib also inhibits CD4+ and CD8+ T-cell proliferation.<sup>15,16</sup> In addition, the inhibitory effect on T-cell activity and proliferation has also been demonstrated for nilotinib<sup>17</sup> and dasatinib.<sup>18</sup> Furthermore, differential immunosuppressive effects between these KIs have been observed probably due to individual off-target kinase activity of these different agents.<sup>19,20</sup> Besides its effect on T cells, recent data have shown that TKIs impair B-cell immune responses in CML through off-target inhibition of kinases important for B-cell signaling.<sup>21</sup> Taken together, there is evidence of a potential immunosuppressive effect of TKIs affecting BCR-ABL, most likely due to their off-target activity.

There are suggestions that the observed immunosuppressive effect translates into an increased risk of infections clinically; nonetheless, specific data on these complications are rare in the literature: data from the initial clinical trials showed a rate of 15% upper respiratory infections in patients treated with imatinib compared to 8% in those treated with interferon/cytarabine; however, the rate of grade 3/4 reactions was similar.<sup>9</sup> Reactivations of hepatitis B under imatinib treatment have been repeatedly reported,<sup>22–26</sup> and one trial evaluated varicella zoster virus (VZV) infections,<sup>27</sup> occurring in only 2% of CML patients treated with imatinib. Similarly, another group found a low infection rate for CML patients under imatinib treatment.<sup>28</sup> For nilotinib, data from the initial trials are rather scarce; infections of any kind are not listed as nonhematological adverse effects in the Evaluating Nilotinib Efficacy and Safety in clinical Trials (ENEST trial)<sup>29</sup> and its

three-year follow-up.<sup>30</sup> A retrospective multicenter analysis of imatinib-resistant or imatinib-intolerant CML patients who had been treated with nilotinib revealed infections occurring in 9% of patients, yet only 1% of them represented grade 3/4 infections.<sup>31</sup> Similar to imatinib, there is one case of hepatitis B reactivation in a nilotinib-treated patient.<sup>32</sup>

Dasatinib has been reported to show the highest off-target activity of KIs targeting BCR-ABL, and *in vitro* data hint at the strongest immunosuppressive effect for this TKI.<sup>20</sup> In the clinical trials, infections of any grade occurred in 27 (11%) of dasatinib-treated patients and 18 (7%) imatinib-treated patients. In the dasatinib arm, five patients died due to infection, whereas one patient died in the imatinib arm; however, the investigators deemed these infections not drug related.<sup>33</sup> Interestingly, the majority of infections did not occur in neutropenia. In a safety analysis of two major clinical trials for dasatinib evaluating 1150 patients for infectious complications, serious infections were rare and only one grade 3–4 opportunistic infection was observed for dasatinib.<sup>34</sup> In contrast to imatinib and nilotinib, however there seems to be a potential impact of dasatinib on infectious side effects: In a retrospective review of CML and ALL patients treated with dasatinib, three or more cycles of dasatinib significantly increased the risk of infection with predominantly bacterial infections,<sup>35</sup> and even opportunistic infections such as *Pneumocystis jirovecii* have been recently described.<sup>36</sup> The majority of infections occurred during neutropenia, thus confounding the effects. In another publication for patients with Ph+–ALL, the infection rate was 18% for all-grade infections and 8% for grade 3/4 infections.<sup>37</sup> In addition, reactivation of latent viral infections such as cytomegalovirus (CMV)<sup>38</sup> and even hepatitis B reactivation has been described.

For the newest TKIs targeting BCR-ABL bosutinib and ponatinib, data on infectious complications can only be derived by evaluating available safety data from the initial clinical trials. In the randomized comparison of imatinib and bosutinib (Bosutinib Efficacy and Safety in Newly Diagnosed Chronic Myeloid Leukemia [BELA] trial), the rate of upper respiratory infections has been reported to be 12% for bosutinib and 8% for imatinib, yet all these consisted of grade 1/2 infections.<sup>39</sup> For ponatinib, the rate of febrile neutropenia has been reported to be 1–6%, depending on the type of disease, with the highest rates observed for ALL, probably based on intensive neutropenia.<sup>40</sup>

Taken together, there is some, albeit minor, evidence of a slightly increased rate of infections in BCR-ABL-positive patients treated with TKIs, probably reflecting not inhibition of BCR-ABL itself but off-target kinase inhibition involved in immune system function. The broader the spectrum of the TKI, the higher the potential for immunosuppressive side effects, thus showing a differential effect, with dasatinib having the highest potential for infectious complications.



**Table 1.** Assessment of clinical evidence of risk of infections and corresponding references.

KINASE INHIBITOR TREATMENT REGIMEN	PATHWAY	UNDERLYING MALIGNANCY	SIGNIFICANT EVIDENCE OF KI-ASSOCIATED INFECTIOUS COMPLICATIONS <sup>Y</sup>	ALL GRADE INFECTIONS/ GRADE 3/4 INFECTION IN CLINICAL TRIALS [%]	REFERENCE(S) <sup>€</sup>
Imatinib	BCR-ABL, c-KIT	CML	Minor	15/0,2	[9]
		GIST	None	11/0	[167]
		Ph+-ALL	Minor	N/A. /38–52 (combined w. chemotherapy)	[168] [169]
Dasatinib	BCR-ABL, c-KIT	CML	Minor	10.5/ 2	[33]
		Ph+-ALL	Minor	18/8	[37]
Nilotinib	BCR-ABL	CML	None	9/1	[31]
		Ph+-ALL	None	N/A.	N/A.
Bosutinib	BCR-ABL	CML	None	12/0	[39]
Ponatinib	BCR-ABL	CML	None	6/6	[40]
		Ph+-ALL	None	6/6	[40]
Erlotinib	EGFR	NSCLC	None	5/1	[53]
Gefitinib	EGFR	NSCLC	None	N/A./3	[54]
Afatinib	EGFR	NSCLC	None	0/0	[42]
Crizotinib	ALK-4-EML	NSCLC	None	32/0	[47]
		ALK-pos. NHL	None	N/A/0	[48]
Ceritinib	ALK4-EML	NSCLC	None	N/A./0	[57]
Vemurafenib	BRAF	Melanoma	None	0/0*	[64]
Dabrafenib	BRAF	Melanoma	None	0/0*	[170]
Trametinib <sup>§</sup>	MEK	Melanoma	None	0/0*	[60]
Temsirolimus	mTOR	MCL	Major	25/7	[102]
		RCC	Major	27/5	[103]
Everolimus	mTOR	RCC	Major	37/13	[105]
		pNET	Major	N/A./5	[171]
		TB	Major	22/14	[172]
		Hs BRCA <sup>§</sup>	Major	9–15/N/A.	[110]
Sunitinib	Multikinase	RCC	None	0/0*	[173]
		pNET	None	0/0*	[174]
		GIST	None	0/0*	[175]
Sorafenib	Multikinase	RCC	None	0/0*	[128]
		HCC	None	0/0*	[176]
		FLT3-pos. AML <sup>¥</sup>	Moderate	N/A./55	[130]
Regorafenib	Multikinase	CRC	None	10/1	[132]
		GIST	None	0/0*	[133]
Pazopanib	Multikinase	STS	None	0/0*	[177]
		RCC	None	0/0*	[138]
Axitinib	Multikinase	RCC	None	0/0*	[178]
Ruxolitinib	JAK2	MF	Major	50/N/A.	[179]
		PV	Major	42/4	[73]
Tofacitinib	JAK3	RA	Major	23/5	[180]
Ibrutinib	BTK	CLL	Major	33/12	[157]
		MCL	Major	23/6	[145]
Idelalisib	PIK3delta	CLL <sup>#</sup>	Major	28/19	[164]
		NHL	Major	25/7	[146]

**Notes:** Kinase inhibitors are shown according to target/indication. <sup>Y</sup>Graded to none, minor, moderate, and major. \*Infectious complications were not mentioned in the safety data or the supplements. <sup>§</sup>In combination with dabrafenib. <sup>#</sup>In combination with rituximab. <sup>€</sup>If possible randomized trial. <sup>¥</sup>Combined with chemotherapy. <sup>§</sup>In combination with exemestan.

**Abbreviations:** CML, chronic myeloid leukemia; GIST, gastrointestinal stromal tumor; Ph+-ALL, Philadelphia-chromosome-positive acute lymphoblastic leukemia; NSCLC, non-small cell lung cancer; ALK-pos. NHL, ALK-positive non-Hodgkin's lymphoma; MCL, mantle cell lymphoma; RCC, renal cell cancer; pNET, pancreatic neuroendocrine tumor; TB, tuberculoma; Hs BRCA, hormone-sensitive breast cancer; HCC, hepatocellular carcinoma; AML, acute myeloid leukemia; CRC, colorectal carcinoma; STS, soft-tissue sarcoma; MF, myelofibrosis; PV, polycythemia vera; RA, rheumatoid arthritis; CLL, chronic lymphocytic leukemia.



## Inhibitors of EGFR-Activating Mutations and ALK-4-EML Rearrangement

In non-small-cell lung cancer (NSCLC), activating mutations of the EGFR and rearrangement of anaplastic lymphoma kinase (ALK) have been identified as drivers of cancer progression and initiation and inhibition of these pathways in mutated disease have been associated with significant improvement in patients' outcome (reviewed by Minuti et al.<sup>4</sup>). The first available TKIs for metastatic NSCLC were gefitinib and erlotinib targeting EGFR, and EGFR overexpression has been observed in lung cancer and was associated with impaired survival.<sup>41</sup> Recently, with afatinib, a second-generation EGFR KI has been introduced in the clinical treatment routine and has shown promising activity against the EGFR T790 mutation, which confers resistance to erlotinib and gefitinib,<sup>42</sup> but it is also active in first-line treatment.<sup>43,44</sup>

Besides targeting EGFR, it was discovered that a characteristic gene rearrangement involving the ALK gene, including its intracellular tyrosine kinase domain, can be observed in 3–6% of NSCLC patients, with the most frequent rearrangement partner being the echinoderm microtubule-associated protein-like 4 (EML4) gene.<sup>45</sup> The resulting kinase activity can be successfully inhibited by the TKI crizotinib. Initially developed for the treatment of ALK-positive anaplastic large cell lymphoma, crizotinib has shown significant activity with prolonged survival not only in second-line therapy<sup>46</sup> but also most recently in first-line therapy in NSCLC,<sup>47</sup> and in small series, it has shown promising activity in relapsed ALK-positive non-Hodgkin's lymphoma.<sup>48</sup> A second TKI targeting ALK-positive NSCLC, ceritinib, has recently been approved for AL4-EML4-rearranged NSCLC patients intolerant or resistant to crizotinib.<sup>49</sup>

From a clinician's point of view, EGFR-TKIs have a rather mild side-effect profile usually consisting of hepatotoxicity, rash, diarrhea, and interstitial lung disease, with supposedly ethnical differences in incidence<sup>50</sup>; infections seem to be a rather rare event as the EGFR pathway is not involved in immunoprocessing. However, preclinically, it has been discovered that airway epithelial surface signaling mediated by EGFR is one way of activating innate immune responses to a variety of infectious and noninfectious stimuli in the respiratory system,<sup>51</sup> which in theory might translate into a higher rate of respiratory infections in patients receiving EGFR-TKI therapy. Yet, when analyzing the data from the randomized trials comparing erlotinib with chemotherapy, the rate of febrile neutropenia was 0% in contrast to the chemotherapy-based therapy arm where the rate of pneumonitis was similar.<sup>52</sup> Data from the maintenance trial for erlotinib show a rate of grade 3/4 infections of 1% vs. 0% in the placebo arm.<sup>53</sup>

In the NCIC CTG BR19 trial, which tests gefitinib after surgical resection of the tumor in an adjuvant setting, Common Toxicity Criteria (CTC) grade 3/4 infections are reported in 3% of the patients and 1% of placebo-treated patients<sup>54</sup> and

in the trials for gefitinib maintenance again tested vs. placebo infectious complications are not mentioned.<sup>55</sup> Afatinib, which is an irreversible Erb blocker and has supposedly the highest activity, although reported to have a higher rate of adverse events in general, shows similar rates of infectious side effects as other EGFR-TKIs<sup>56</sup> even when compared to placebo. Infectious side effects did not seem to be frequent with afatinib treatment,<sup>42</sup> suggesting that even potent inhibition of EGFR or Erb does not increase the rate of infectious side effects.

Taken together, the data from the trials suggest that EGFR-TKI treatment seems safe in terms of infectious complications and does not have clinically relevant immunosuppressive properties.

For ALK-rearranged NSCLC and its primary TKI crizotinib, data on potential immunosuppressive properties are even more scarce: in the initial trial comparing crizotinib to chemotherapy, a higher rate of grade 1/2 upper respiratory infections compared to chemotherapy was observed but grade 3/4 infections were only present in the chemotherapy arm,<sup>46</sup> a finding that was later also seen in the first-line trial.<sup>47</sup> For ceritinib, available data concerning this topic are also lacking. The only clinical trial published does not describe infections as typically occurring Adverse Events (AEs)<sup>57</sup>; however, the limited data surely forbid drawing a definite conclusion. Interestingly, both drugs differ in their off-target activity as crizotinib targets c-met, whereas insulin-like growth factor 1 receptor (IGF-1R) and insulin receptor (InsR) are additionally inhibited by ceritinib. Both TKIs affect the proto-oncogene 1 receptor kinase (ROS1) at clinically relevant concentrations.

In summary, the inhibition of EGFR via TKI treatment has little potential detrimental immunosuppressive effects and does not seem to increase the risk of infections. For ALK-rearranged malignancies, the (limited) data also suggest a rare occurrence of infectious complications. Nonetheless, based on the limited number of patients who have yet been exposed to crizotinib or ceritinib, the final conclusion concerning this matter can probably not be drawn at this moment.

## Inhibitors of BRAF/MEK

Activating mutations of BRAF, which induce constitutive activation of the MAPK signaling pathway, have been implicated in induction and in maintaining the malignant phenotype in a variety of cancers. Vemurafenib and dabrafenib, both inhibiting BRAF, have been approved for the treatment of melanoma and have also been successfully used in other BRAF-mutated cancer entities<sup>58,59</sup>; yet most patients suffer a relapse later in the course of their disease. More recently, the introduction of MEK-inhibitor trametinib combined with dabrafenib has led to further increase in overall survival in melanoma patients,<sup>60</sup> underlining the potential of inhibition of the MAPK pathways by TKI therapy.

As the MAPK pathway has been implicated in immune system functions, especially in the processing of pattern recognition receptors such as toll-like receptors,<sup>61</sup> pharmacological



inhibition might, at least in theory, cause immunosuppressive properties of these drugs. Clinical data on potential infectious complications in patients treated with BRAF or MEK inhibitors are however scarce. It has been recently shown that melanoma patients treated with vemurafenib had a significant decrease in lymphocyte count, especially the CD4+ T-cell population and significantly reduced secretion of interferon- $\gamma$  and interleukin 9; the effect was not observed for dabrafenib.<sup>62</sup> An extended analysis could find infections in 9/102 patients, especially if the patients had been additionally treated with steroids.<sup>63</sup> However, neither in the pivotal trial comparing vemurafenib with dacarbazine in melanoma patients<sup>64,65</sup> nor in the combination trials of vemurafenib with MEK inhibitors cometinib<sup>66</sup> or trametinib,<sup>60</sup> a significant rate of infectious side effects is mentioned in the safety data.

In summary, from the available data, a strong clinical evidence of severe immunosuppression with consecutive increased risk of infection cannot yet be demonstrated, although a heightened awareness should be applied if patients receive concomitant steroids.

### Inhibitors of JAK

The family of JAK consists of four kinases – JAK1, JAK2, JAK3, and TYK – and plays a major role in hematopoiesis as knockout studies with JAK2-deficient mice showed an impaired development of their hematopoiesis leading to death at day 13 of gestation.<sup>67</sup> However, there is evidence that

different JAKs induce varied transcriptional changes, typically via the signal transducer and activator of transcription (STAT) family pathway, and are involved in several diseases. The activating mutation V617F of JAK2 has been identified as one of the hallmarks in the pathogenesis of myeloproliferative neoplasms and has been detected in 95% of patients with PV and to a lesser extent in 50–60% of patients with MF and essential thrombocythemia.<sup>68</sup> In addition, it has also been found in a significant proportion of patients with myeloid malignancies, such as myelodysplastic syndrome, acute myeloid leukemia (AML), and CML.<sup>69</sup> The activation of the JAK2 V617F kinase domain causes the constitutive activation of proteins STAT5 and STAT3, which consecutively induce malignant cell transformation.<sup>70</sup> Interestingly, STAT3, targeted via JAK3, has also been implicated in a variety of different autoimmune diseases such as rheumatoid arthritis (RA) and psoriasis and JAK3 inhibitors have been introduced as another class of disease-modifying drugs in rheumatoid arthritis (DMARD).<sup>71</sup>

Based on the CML success story, where TKIs dramatically affected outcome and clinical course of the disease, inhibitors of JAK were developed; these inhibitors, however, do not specifically inhibit the mutated kinase but JAKs in general. Although several other compounds are in development, the two US Food and Drug Administration (FDA)-approved inhibitors are ruxolitinib and tofacitinib. Ruxolitinib targets JAK1 and JAK2, and after being initially approved for MF,<sup>72</sup> it has also quite recently been approved for PV.<sup>73</sup> In addition,

**Table 2.** Assessment of specific kinase inhibitors infections and prophylaxis recommendations based on available evidence.

PATHWAY TARGETED KINASE	AVAILABLE DRUGS	TYPICAL INFECTIONS REPORTED	PROPHYLAXIS RECOMMENDED*
BCR-ABL	Imatinib, Dasatinib, Nilotinib, Ponatinib, Bosutinib	HSV reactivation CMV reactivation Hepatitis Reactivation Febrile neutropenia URTI	May be considered
EGFR/ALK	Erlotinib, Gefitinib, Afatinib, Crizotinib, Ceritinib	URTI	None
BRAF/MEK	Vemurafenib, Dabrafenib, Trametinib	No specific	None
mTOR	Temsirolimus, Everolimus	VZV Reactivation HSV Reactivation Invasive Aspergillosis PcP	Aciclovir/Cotrimoxazole <i>should</i> be considered
Multikinase (esp. VEGF)	Sorafenib, Sunitinib, Regorafenib, Axitinib, Pazopanib	No specific	None
JAK	Ruxolitinib, Tofacitinib	VZV Reactivation HSV Reactivation Invasive Aspergillosis PcP	Aciclovir/Cotrimoxazole <i>should</i> be considered
BCR-Pathway-Inhibitory	Ibrutinib, Idelalisib	Pneumonia, URTI	May be considered

**Note:** \*Recommendation of anti-infective prophylaxis based on the data on frequency, type, and severity of infections in the current available literature, prone to change in the future.

**Abbreviations:** HSV, herpes simplex virus; CMV, cytomegalovirus; URTI, upper respiratory tract infection; PcP, *P. jirovecii* pneumonia.



tofacitinib, an inhibitor of JAK1 and JAK3, has been approved for refractory RA based on a placebo-controlled trial, showing its potential as a DMARD.<sup>74</sup>

The indication that a drug targeting JAK does have a significant use in autoimmune diseases clearly hints at immunosuppressive properties and thus warrants further investigation. Indeed, preclinical data clearly show an influence of ruxolitinib and tofacitinib on components of the immune system. It has been demonstrated that tofacitinib not only suppresses cytokine production of CD4+ T lymphocytes in RA patients<sup>75,76</sup> but also inhibits proliferation of these cells.<sup>77</sup> For ruxolitinib, this profound effect has also been confirmed: Schönberg et al could clearly show a decrease in natural killer cells of patients treated with ruxolitinib, and this effect was clearly linked to an increase in infections in their study.<sup>78</sup> The same group also found that the drug impairs T-cell function by decreasing their potential of producing proinflammatory cytokines, and thus, Th1 and Th17 cells were reduced *in vivo* and *in vitro*.<sup>79</sup> Lastly, even dendritic cells' function and migration is hampered by ruxolitinib, further aggravating immune system dysfunction.<sup>80</sup>

Clinical trials and increased clinical exposure clearly underline the potential of infectious complications of JAK inhibitors. Indeed, for tofacitinib, pooled data of all patients in the randomized trials covering approximately 4800 patients suggest a significant incidence of infection and infection-related mortality, which is however similar to what is observed in treatment with other biological agents for RA.<sup>81</sup> Similar to these other biological agents, frequent herpes zoster reactivation and even *Mycobacterium tuberculosis* reactivation has been recognized. Quite interesting is the efficacy and immunosuppressive property of tofacitinib in inflammatory bowel disease<sup>82</sup> and also its potential role as an immunosuppressive agent in kidney transplantation.<sup>83</sup> Interestingly, in both of these trials, infections were the major problem not only compared to placebo but also compared to cyclosporine A in kidney transplanted patients. This is also supplanted by the prospective placebo-controlled trial of tofacitinib in RA patients, where infections were associated with tofacitinib treatment and even serious infections occurred in this treatment arm.<sup>84</sup> Typical infections represented herpes zoster, CMV, and even Epstein-Barr virus reactivation associated with posttransplant lymphoproliferative disease.

For ruxolitinib, clinical data also indicate an increased risk of infection: in the recent trials for PV, ruxolitinib treatment was associated with an increased rate of infection (42% vs. 37%) underscored by the increase in herpes zoster reactivation (6% vs. 0%).<sup>73</sup> In addition, in a recent phase 2 trial for patients with AML relapse treated with ruxolitinib, the most frequent grade 3 or 4 nonhematologic event was infection (most frequently, pneumonia; 15 of 26; 58%).<sup>85</sup> Furthermore, reports about severe opportunistic infections have been published for patients under ruxolitinib treatment, such as *Cryptococcus neoformans* pneumonia,<sup>86</sup> hepatitis B reactivation,<sup>87</sup> toxoplasmosis chorioretinitis,<sup>88</sup> disseminated

tuberculosis,<sup>89</sup> and even JC virus-associated progressive multifocal encephalopathy.<sup>90</sup> On the other hand, recent data on its potential in treating steroid-refractory graft-versus-host disease (GvHD) clearly emphasize the immunosuppressive properties of ruxolitinib: in a murine model, treatment with ruxolitinib led to improvement in GvHD while maintaining a graft-versus-leukemia effect.<sup>91</sup> This preclinical effect was also observed by Spoerl et al, who could clearly demonstrate the activity of ruxolitinib in the treatment of GvHD and also confirmed that clinically by successfully treating six patients harboring steroid-refractory GvHD with ruxolitinib.<sup>92</sup> This interesting finding was confirmed by observing a complete response rate of 84% in 52 patients with steroid-refractory GvHD in a multicenter setting.<sup>93</sup>

Thus, taken together, there is sufficient preclinical and clinical evidence for an increased risk of infectious complications in treatment with JAK inhibitors, and physicians using these drugs should be alert. Heine et al therefore recently proposed a risk stratification and recommendation in which they propagate an acyclovir and cotrimoxazole prophylaxis as well as a basic screening program prior to beginning ruxolitinib treatment,<sup>94</sup> which from our point of view should be considered, giving the clear evidence mentioned above.

### TKIs as Inhibitors of mTOR

The mTOR is a serine/threonine kinase that affects cell growth, proliferation, survival, autophagy, metabolism, and cytoskeletal organization.<sup>95</sup> The mTOR pathway located at a central hub for different signaling cascades plays a pivotal role in the pathogenesis of many malignancies and has thus become a target of interest for therapeutic inhibition. The first drug affecting the mTOR pathway was rapamycin, from which the enzyme draws its name. Rapamycin was originally approved as an immunosuppressant in the United States, yet soon after its approval, its antineoplastic effect was described for a variety of malignancies<sup>96</sup>; however, the clinical benefit of rapamycin has been found to be disappointing, further leading to the development of two drugs targeting mTOR, temsirolimus and everolimus.

Temsirolimus is available in both intravenous and oral formulations and is approved for the treatment of advanced-stage metastatic renal cell carcinoma (RCC) and relapsed or refractory mantle cell lymphoma.<sup>97</sup> Everolimus, which is available as an oral formulation, has been approved by the FDA and the European Medical Agency (EMA) for the treatment of pancreatic neuroendocrine tumors (pNET), advanced RCC, subependymal giant cell astrocytoma associated with tuberous sclerosis, and, in combination with exemestane, for advanced hormone-receptor-positive, HER2-negative breast cancer. In addition, everolimus has also received approval as an immunosuppressant for liver and kidney organ transplantation,<sup>98</sup> and has shown efficacy in treating refractory GvHD.<sup>99</sup>

It is prudent to assume that drugs that are preventing organ rejection predispose to induce infectious adverse effects,



and physicians should be aware of this risk while treating patients with mTOR inhibitors. From a transplant physician's perspective, mTOR inhibitors might be promising. Considering the different options for immunosuppression for solid organ rejection, there is evidence that infectious complications for mTOR inhibitors are less compared to those for classic calcineurin inhibitors such as cyclosporine or mycophenolate mofetil, at least for specific infections such as CMV.<sup>100,101</sup>

The immunosuppressive effects and the risk of opportunistic infections are thus well recognized in solid organ transplantation recipients, but potentially less in patients suffering from solid tumors. As the exposure of patients to these drugs is continuously increasing, more patients are at risk for infectious complications. For temsirolimus, data from the randomized controlled trials clearly suggest a risk of infections for mantle cell lymphoma, where all-grade infections were present in 25% of patients compared to only 9% of patients in the control arm; also the severity of infections was higher.<sup>102</sup> In the pivotal clinical trial, which established temsirolimus as the standard of care for advanced renal cell cancer, the risk of infection for temsirolimus was nearly doubled compared to interferon and the addition of temsirolimus to interferon further significantly increased the infection rate.<sup>103</sup> Combination of temsirolimus and conventional chemotherapy such as temozolomide was also associated with further increased infectious potential.<sup>104</sup>

In the Renal Cell cancer treatment with Oral RAD001 given Daily (RECORD-1) trial, which established everolimus as the drug of choice for patients progressing under anti-VEGF therapy, the number of adverse events as well as infections was significantly higher compared to placebo (37% vs. 18%) with 13% severe infections and several infection-related deaths due to opportunistic infections such as invasive aspergillosis.<sup>105</sup> Indeed, an expert panel recommended to be cautious and aware of infectious complications in patients treated with everolimus, especially if the CD4 cell count is  $<200/\mu\text{L}$  and recommended a basal screening program to evaluate the antibody immune status of the patient prior to treatment with everolimus.<sup>106</sup> Infectious complications were also observed for pediatric and adolescent patients with tuberous sclerosis treated with everolimus with even fatal course.<sup>107</sup> For patients with hormone-sensitive, advanced breast cancer, two clinical trials analyzed the combination of everolimus with exemestan (Breast Cancer Trials of Oral Everolimus-2 [BOLERO-2] trial)<sup>108</sup> or temsirolimus with letrozole (Randomized phase III placebo-controlled trial of letrozole plus oral temsirolimus as first-line endocrine therapy in postmenopausal women with locally advanced or metastatic breast cancer2 [HORIZON] trial).<sup>109</sup> The addition of the mTOR inhibitor lead to increased infectious adverse events<sup>110</sup> compared to the arm with placebo and aromatase inhibitor. In a large meta-analysis on infectious episodes from 1924 renal cell cancer patients treated with temsirolimus and everolimus in clinical trials, it was shown that infections are twice as frequent and even 2.6 times more frequent for grade 3/4

infections.<sup>111</sup> However, there was no significant difference between temsirolimus and everolimus, suggesting that the observed immunosuppression is class specific and not dependent on the type of mTOR inhibitor. Another recent interesting publication evaluating the occurrence and the grading of infections in patients treated with mTOR inhibitors in clinical trials and comparing these with control patients not receiving mTOR inhibition showed unequivocally that incidence of grade 1–5 infections was significantly higher with single-agent mTOR inhibitors compared with the controls (27% vs. 8%; odds ratio, 4.26;  $P < 0.0001$ ) and also that the clinically relevant grade 3/4 infections were much more frequent. Also, the combination of mTOR inhibitors and chemotherapy further aggravated the incidence and severity of infections compared to single-agent mTOR inhibition.<sup>112</sup>

Indeed, classic opportunistic infections such as *P. jirovecii* pneumonia, which represent a rarity in patients with solid tumors, have been reported in oncologic patients exposed to mTOR inhibitors, underlining the immunosuppressive potential<sup>113,114</sup> of these class inhibitors.

An aggressive diagnostic approach, such as bronchoalveolar lavage for patients with mTOR treatment and pulmonary infiltrates, is recommended as the main differential diagnosis is the frequent noninfectious pneumonitis<sup>115</sup> and the potential microorganisms causing opportunistic infections, such as fungi or *Pneumocystis*, can thus be much more easily diagnosed.

In conclusion, treatment with mTOR inhibitors everolimus and temsirolimus is clearly associated with an increased risk of all-grade and high-grade infections. Indeed, based on the frequency, type, and severity of infections encountered, an antiviral (eg, aciclovir) and even a *P. jirovecii* prophylaxis may be instituted, based on the individual patients' immune status. Although these agents have influenced the therapeutic landscape and armamentarium in various malignancies and thus patients' survival, a heightened awareness even for atypical infections and a stringent diagnostic approach concerning infectious complications is warranted.

### Multikinase Inhibitors

In this paragraph, we summarize the potential immunosuppressive properties of multikinase inhibitors, such as sorafenib, sunitinib, regorafenib, axitinib, and pazopanib, as these KIs mediate their effect by targeting several kinases such as vascular endothelial growth factor receptor (VEGFR), PDGFR, FLT-3, c-Kit, and RET. Most studies suggest that the antiangiogenic properties of these multikinase inhibitors are the major contributor to their clinical efficacy, at least in solid tumors. Treatment with sunitinib and sorafenib is associated with a significant increase in neutropenia,<sup>116</sup> and if these cytopenias have an impact on the incidence of infectious complications is not yet clear.

Sunitinib is approved for the treatment of renal cell cancer, GIST, and pNET based on several trials demonstrating its clinical efficacy. Sunitinib's mode of action is believed to be



primarily mediated by antiangiogenic effects.<sup>117</sup> Although it has been recognized that sunitinib induces leukopenia and a potent lymphopenia,<sup>118</sup> the placebo-controlled trials for several malignancies suggest that this finding does not translate into an increase in opportunistic infections<sup>119–122</sup> as infections are rarely reported in the safety data. Sunitinib has been approved since 2006, and thus, a great number of patients have been exposed to this drug until now; reports on infections because of sunitinib treatment are sporadic, suggesting a presumably negligible immunosuppressive potential.

Sorafenib was introduced into the clinical setting in 2005 and is currently approved for palliative treatment of hepatocellular carcinoma (HCC), renal cell cancer, and metastatic thyroid carcinoma<sup>123</sup>; in addition, it has activity in FLT3-mutated leukemias.<sup>124</sup> Compared to sunitinib, myelosuppression and therefore a possible neutropenia is less frequent. Sorafenib treatment affects signal transduction pathways during T-cell activation and even impairs production of interferon gamma independent of the MAPK pathway.<sup>125</sup> It also inhibits the activation and induces apoptosis of peripheral T-cells.<sup>126</sup>

Apparently, this preclinical effect does not seem to translate into a higher rate of infections in solid tumor patients treated with sorafenib. In a recent meta-analysis evaluating postapproval safety data of more than 2000 patients, infections are not reported as significant AEs.<sup>127</sup> Safety data from the large placebo-controlled phase 3 trials suggest a higher noninfection-related AE rate for sorafenib for renal cell cancer<sup>128</sup> in general, but infectious complications were not among them. For HCC, this is equally true, and there are even data that sorafenib has some activity in inhibiting hepatitis C virus, one major and frequent driver in oncogenic transformation to HCC.<sup>129</sup>

On the other hand, a recent multicenter placebo-controlled trial evaluating the effect of adding sorafenib to conventional chemotherapy in patients with de novo AML older than 60 years found a significant increase in infections and infection-related mortality within 60 days after the start of therapy with 15 infection-related deaths in the sorafenib arm versus only 4 in the placebo arm ( $P < 0.015$ ).<sup>130</sup>

These rather conflicting results in myeloid malignancies compared to the data on sorafenib in solid tumor treatment might, in theory, reflect that the (supposedly lymphocyte based) immunosuppressive effect of sorafenib itself is negligible in oncologic patients without prevalent granulocytopenia. In AML patients with long-lasting and profound myelosuppression, the immunosuppression induced by sorafenib affecting the lymphocyte compartment might become more apparent and cause these observed complications.

Regorafenib is a multikinase inhibitor with a very broad spectrum that blocks the activity of several protein kinases, such as VEGFR1, VEGFR2, VEGFR3, TIE2, KIT, RET, RAF1, BRAF, PDGFR, and FGFR.<sup>131</sup> It has been approved for the treatment of advanced colorectal cancer and refractory GIST in 2013 as it has displayed an improvement of survival

in these patients in several trials.<sup>132,133</sup> Typical adverse events consist of hand-foot reactions, stomatitis, (noninfectious) diarrhea, and hypertension, but infections are rarely reported.

Although regorafenib targets a myriad of kinases, the literature does not report about an increase in infections, and the number of patients exposed to regorafenib is much less compared to those exposed to sorafenib or sunitinib. The safety data from the placebo-controlled clinical trials do not indicate a heightened prevalence of infection in the study arms with regorafenib, which presumably indicates a marginal effect on the occurrence of infectious side effects.

Pazopanib is a second-generation small-molecule TKI especially targeting VEGFR-1/2/3 while showing a lower affinity against Platelet-derived Growth-Factor receptor (PDGFR)-, FGFR-1/2, and c-KitR. It has shown excellent activity in a variety of malignant diseases such as renal cell cancer<sup>134</sup> and soft-tissue sarcoma,<sup>135</sup> both for which it has been approved, but also in other malignant diseases such as ovarian carcinoma<sup>136</sup> or NSCLC.<sup>137</sup> Similar to regorafenib, data in the literature on potential immunosuppressive side effects leading to infections are almost nonexistent, and the final phase 3 safety data do not imply infectious complications as frequent adverse events.<sup>138</sup>

Lastly, axitinib is a TKI of VEGF, PDGFR- $\alpha$ , and c-Kit and has recently been approved for patients with renal cell cancer refractory to sunitinib or cytokine treatment based on the AXIS trial, which demonstrated an increase in progression-free survival compared to sorafenib<sup>139</sup>; *in vitro* data suggest that axitinib is 40–50 times more potent in inhibiting VEGF, its major target, compared to first-generation multikinase inhibitors.<sup>140</sup> Preclinical data suggest that axitinib has a much less suppressive effect on lymphocytes compared to sunitinib or sorafenib<sup>141</sup>; the (scarce) clinical data on potential infectious complications suggest that this drug seems safe in terms of immunosuppression.

Taken together, the multikinase inhibitors mentioned above, whose main target is the VEGF pathway, are not associated with infectious complications; a finding that can also be observed for bevacizumab, an established antibody targeting VEGF, for whom a potential immunosuppressive effect leading to infectious complications has not yet been observed and for whom a large amount of clinical data is available due to its broad indication and long time past approval.

### Inhibitors of the BCR Signaling Pathway

B cells are an essential component of the immune defense because they present antigens, produce neutralizing antibodies, and maintain the lymphoid architecture. For the treatment of B-cell malignancies such as non-Hodgkin's lymphoma (NHL) or even autoimmune diseases such as RA, using pharmacologic agents to target the B cells has shown promising results leading to approval of rituximab, a chimeric anti-CD20 antibody. It has been recognized that infectious complications could be a side effect of prolonged B-cell-depleting treatment,





eg, with rituximab, especially under maintenance therapy,<sup>142</sup> although the exact influence of rituximab on infectious complications is still controversially discussed.<sup>143</sup>

Quite recently, two drugs inhibiting the Bruton's tyrosine kinase (BTK) or the phosphatidylinositol 3-kinase delta (PIK3delta) and thus inhibiting the critical components of the activation of the BCR pathway have emerged and have made a profound impact on the therapeutic landscape of indolent NHL and chronic lymphatic leukemia (CLL).

Ibrutinib, an irreversible inhibitor of the BTK, has shown excellent activity even in high-risk relapsed CLL<sup>144</sup> and mantle cell lymphoma<sup>145</sup> leading to a rapid approval. The second TKI, idelalisib, is targeting another component of the BCR signaling pathway, the PIK3delta; it has also demonstrated significant clinical activity as a single agent in follicular lymphoma<sup>146</sup> and for relapsed CLL in combination with rituximab.<sup>147</sup>

Interestingly, for both of these enzymes, it has been shown that hereditary mutations are associated with an increase in infections. Inactivating mutations of the BTK are the cause of X-linked agammaglobulinemia and lead to deficient development of B lymphocytes, thus causing hypogammaglobulinemia, profoundly reduced levels of serum antibodies, and reduced levels of circulating B cells.<sup>148</sup> BTK deficiency impairs B-cell and also monocytic and dendritic cell functions.<sup>149,150</sup> This leads to a markedly increased incidence and severity of infections often causing even lethal complications.<sup>148</sup> The importance of BTK in the defense against a variety of organisms such as bacteria, virus, and even fungi has been demonstrated preclinically.<sup>151-153</sup> Recently, the so-called activated PI3K- $\delta$  syndrome (APDS) was described, which although leading to a gain-of-function mutation of that enzyme; patient-derived lymphocytes had increased levels of phosphatidylinositol 3,4,5-trisphosphate; and phosphorylated AKT protein was prone to activation-induced cell death, thus leading, in fact, to immunosuppression and an increase in, especially, upper airway infections.<sup>154</sup> In addition, patients with this hereditary mutation had a substantial deficiency in naive T cells but an overrepresentation of senescent effector T cells, leading to sinupulmonary infections and viremia due to CMV and/or Epstein-Barr virus.<sup>155</sup> Inhibition of this pathway, however, has also detrimental effects on the immune system, suggesting that there is a strict balance that has to be kept to avoid infections and ensure immune system function.<sup>156</sup>

Based on these preclinical data, it is prudent to assume that there is a substantial potential for infectious complications when affecting these two critical enzymes, and albeit limited, the clinical data support these observations. In the initial phase, 1/2 trial for treatment of relapsed CLL with ibrutinib pneumonia grade 3/4 was observed in 12% of patients and 33% of all adverse events were upper respiratory infections<sup>157</sup>; these occurred typically in the beginning of the treatment, and with continuous neutrophil recovery, the incidence of these events subsided. For patients with high-risk CLL, ibrutinib

in combination with rituximab showed grade 3 infections in 13% of patients, with pneumonia the most typical infectious complication.<sup>158</sup> Interestingly, in that study, the investigators observed a trend toward decrease in serum IgM levels and a continuous decrease in CD4+ lymphocytes, which were more than halved after 12 months of treatment, which the authors interpreted as treatment response, as the coevolution and interdependence between T cells and leukemia cells was reduced. In the randomized comparison of ibrutinib versus ofatumumab (the Ibrutinib versus Ofatumumab in Patients with Relapsed or Refractory Chronic Lymphocytic Leukemia [RESONATE] trial) for relapsed CLL patients, the incidence of infections of any grade was significantly higher in the ibrutinib arm (70% vs. 54%), whereas all-grade infections were similar.<sup>144</sup> These findings suggest that BTK inhibition might be more immunosuppressive than targeting CD20. Lastly, in the trial that leads to the approval of ibrutinib for mantle cell lymphoma, the incidence of upper respiratory tract infections was 23%; however, these were all graded 1 and 2, which, in contrast to treatment in CLL, might reflect the inherently lesser disease-related immunosuppression of mantle cell lymphoma compared to CLL.<sup>145</sup> In a recent trial for patients with Waldenstrom's macroglobulinemia, the rate of infections was slightly less compared to the previous studies (8%), again indicating that the amount of infectious complications is dependent on the underlying disease and its inherent lesser immune system activity.<sup>159</sup> In a recently published analysis of a three-year follow-up of 132 patients with CLL and SLL (small lymphocytic lymphoma) treated with ibrutinib, the rate of overall Infectious side effects and rate of  $\geq$  grade 3 infections was up to 51% in relapsed/refractory patients; due to occurring VZV reactivation, an antiviral prophylaxis was instituted.<sup>160</sup>

For idelalisib, targeting the PIK3delta pathway, the clinical data also suggest an immunosuppressive potential causing infectious side effects. In the initial phase 1 study for CLL patients, serious adverse events were frequently present with  $\geq$  grade 3 pneumonia occurring in 20% and additionally febrile neutropenia and bacteremia occurring in 5.6% and 9.3% of patients with even fatal clinical course.<sup>161,162</sup> Even more striking, two of the identified organisms were *P. jirovecii*, and even fungal pneumonia was identified in two further patients suggesting that there is a profound inducible immunosuppression associated with idelalisib treatment. Yet, it has to be kept in mind that patients had received intensive treatment regimens such as purine analogues or alemtuzumab prior to idelalisib and CLL itself carries a high risk of infection.<sup>163</sup> In the first-line trial for combination of idelalisib and rituximab in newly diagnosed CLL patients, adverse events occurred frequently, especially pneumonia in 28% of patients, with 19% classified as serious adverse events and also infection-related fatalities.<sup>164</sup> In the large double-blind phase 3 trial analyzing treatment of idelalisib and rituximab compared to rituximab alone for patients with relapsed CLL, the most frequent serious adverse events in the two groups were pneumonia, pyrexia, and febrile



neutropenia.<sup>147</sup> Idelalisib for the treatment of NHL, which by itself probably carries a lower disease-related immunosuppression, was however also associated with infectious complications, especially pneumonia, occurring in 17% of patients in the phase 1 trial<sup>165</sup> and even three infection-related deaths, a finding that was also confirmed in the large placebo-controlled phase 3 trial for follicular lymphoma<sup>146</sup> and, more recently, the phase 1 trial for mantle cell lymphoma.<sup>162</sup>

The potential immunosuppressive properties of inhibitors of the BCR pathway are probably, similar to JAK inhibitors, hinted at by a recent (preclinical) publication regarding its use in treating GvHD. In a murine model of sclerodermatous chronic GvHD, ibrutinib treatment delayed progression, improved survival, and ameliorated clinical and pathological manifestations of mice with GvHD, and the authors found that animals lacking BTK and IL-2 inducible T cell kinase, which is also inhibited by ibrutinib, did not develop cGvHD, indicating that these molecules are critical for its pathogenesis.<sup>166</sup>

So, in summary, the treatment of inhibitors of the BCR signal cascade ibrutinib and idelalisib is associated with infectious complications, the typical manifestation consisting of respiratory infections such as pneumonia. The incidence and severity of these infections is probably dependent on concomitant disease and other immunosuppressants such as steroid treatment in CLL patients, probably representing the group with the highest risk. Individual prophylaxis regimens, such as PcP prophylaxis with cotrimoxazole or antiviral prophylaxis, should be considered in individual patients based on their immune status.

## Summary

Infectious complications are a typical treatment-related complication of modern therapies in hemato-oncological treatment regimens. Although often considered as representing just a *pill treatment*, KIs bear the potential of causing severe and even life-threatening infections depending on the pathway involved and the associated off-target kinase activity.

In case of infectious complications, the preclinical and clinical evidence of KIs targeting the angiogenesis or the EGF pathway suggests that these KIs do not aggravate the incidence or the intensity of infections. However, treatment with KIs interfering with critical immune system components such as MTOR inhibitors, JAK inhibitors, or the new KIs affecting the BCR pathways is associated with an increased occurrence of infections and even with the risk of fatal complications. Based on the frequency, type, and severity of infections in patients treated with these drugs, a prophylaxis approach based on individual patient's immune status, concomitant medication (eg, steroids), and comorbidities should be considered. A heightened awareness and a stringent diagnostic approach should be instituted in these patients as life-threatening infections may occur.

## Author Contributions

Conceived and designed the experiments: MR, DB. Analyzed the data: MR, DB, TB. Wrote the first draft of the manuscript: MR. Contributed to the writing of the manuscript: MR, DB, TB, WKH. Agree with manuscript results and conclusions: MR, DB, TB, WKH. Jointly developed the structure and arguments for the paper: MR, DB, WKH. All authors reviewed and approved of the final manuscript.

## REFERENCES

- Jabbour E, Cortes J, Ravandi F, O'Brien S, Kantarjian H. Targeted therapies in hematology and their impact on patient care: chronic and acute myeloid leukemia. *Semin Hematol*. 2013;50:271–83.
- Plosker GL. Ruxolitinib: a review of its use in patients with myelofibrosis. *Drugs*. 2015;75:297–308.
- Daste A, Grellety T, Gross-Goupil M, Ravaud A. Protein kinase inhibitors in renal cell carcinoma. *Expert Opin Pharmacother*. 2014;15:337–51.
- Minuti G, D'Incecco A, Landi L, Cappuzzo F. Protein kinase inhibitors to treat non-small-cell lung cancer. *Expert Opin Pharmacother*. 2014;15:1203–13.
- Dossett LA, Kudchadkar RR, Zager JS. BRAF and MEK inhibition in melanoma. *Expert Opin Drug Saf*. 2015;14:559–70.
- Tlemsani C, Mir O, Boudou-Rouquette P, et al. Posterior reversible encephalopathy syndrome induced by anti-VEGF agents. *Target Oncol*. 2011;6:253–8.
- Zembower TR. Epidemiology of infections in cancer patients. *Cancer Treat Res*. 2014;161:43–89.
- Augustson BM, Begum G, Dunn JA, et al. Early mortality after diagnosis of multiple myeloma: analysis of patients entered onto the United Kingdom Medical Research Council trials between 1980 and 2002 – Medical Research Council Adult Leukaemia Working Party. *J Clin Oncol*. 2005;23:9219–26.
- O'Brien SG, Guilhot F, Larson RA, et al; IRIS Investigators. Imatinib compared with interferon and low-dose cytarabine for newly diagnosed chronic-phase chronic myeloid leukemia. *N Engl J Med*. 2003;348:994–1004.
- ESMO/European Sarcoma Network Working Group. Gastrointestinal stromal tumours: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2014;25(suppl 3):iii21–6.
- McCormack PL, Keam SJ. Dasatinib: a review of its use in the treatment of chronic myeloid leukaemia and Philadelphia chromosome-positive acute lymphoblastic leukaemia. *Drugs*. 2011;71:1771–95.
- Syed YY, McCormack PL, Plosker GL. Bosutinib: a review of its use in patients with Philadelphia chromosome-positive chronic myelogenous leukemia. *Bio Drugs*. 2014;28:107–20.
- Jabbour E, Cortes J, Giles F, O'Brien S, Kantarjian H. Drug evaluation: nilotinib – a novel Bcr-Abl tyrosine kinase inhibitor for the treatment of chronic myelocytic leukemia and beyond. *IDrugs*. 2007;10:468–79.
- Hoy SM. Ponatinib: a review of its use in adults with chronic myeloid leukaemia or Philadelphia chromosome-positive acute lymphoblastic leukaemia. *Drugs*. 2014;74:793–806.
- Seggewiss R, Loré K, Greiner E, et al. Imatinib inhibits T-cell receptor-mediated T-cell proliferation and activation in a dose-dependent manner. *Blood*. 2005;105:2473–9.
- Chen J, Schmitt A, Giannopoulos K, et al. Imatinib impairs the proliferation and function of CD4+CD25+ regulatory T cells in a dose-dependent manner. *Int J Oncol*. 2007;31:1133–9.
- Chen J, Schmitt A, Chen B, et al. Nilotinib hampers the proliferation and function of CD8+ T lymphocytes through inhibition of T cell receptor signalling. *J Cell Mol Med*. 2008;12:2107–18.
- Fei F, Yu Y, Schmitt A, et al. Dasatinib exerts an immunosuppressive effect on CD8+ T cells specific for viral and leukemia antigens. *Exp Hematol*. 2008;36:1297–308.
- Rohon P, Porkka K, Mustjoki S. Immunoprofiling of patients with chronic myeloid leukemia at diagnosis and during tyrosine kinase inhibitor therapy. *Eur J Haematol*. 2010;85:387–98.
- Hayashi Y, Nakamae H, Katayama T, et al. Different immunoprofiles in patients with chronic myeloid leukemia treated with imatinib, nilotinib or dasatinib. *Leuk Lymphoma*. 2012;53:1084–9.
- de Lavallade H, Khoder A, Hart M, et al. Tyrosine kinase inhibitors impair B-cell immune responses in CML through off-target inhibition of kinases important for cell signaling. *Blood*. 2013;122:227–38.
- Kang BW, Lee SJ, Moon JH, et al. Chronic myeloid leukemia patient manifesting fatal hepatitis B virus reactivation during treatment with imatinib rescued by liver transplantation: case report and literature review. *Int J Hematol*. 2009;90:383–7.



23. Ikeda K, Shiga Y, Takahashi A, et al. Fatal hepatitis B virus reactivation in a chronic myeloid leukemia patient during imatinib mesylate treatment. *Leuk Lymphoma*. 2006;47:155–7.
24. Thia TJ, Tan HH, Chuah TH, Chow WC, Lui HF. Imatinib mesylate-related fatal acute hepatic failure in a patient with chronic myeloid leukaemia and chronic hepatitis B infection. *Singapore Med J*. 2008;49:e86–9.
25. Walker EJ, Simko JP, Ko AH. Hepatitis B viral reactivation secondary to imatinib treatment in a patient with gastrointestinal stromal tumor. *Anticancer Res*. 2014;34:3629–34.
26. Lakhani S, Davidson L, Priebe DA, Sherker AH. Reactivation of chronic hepatitis B infection related to imatinib mesylate therapy. *Hepatol Int*. 2008;2:498–9.
27. Mattiuzzi GN, Cortes JE, Talpaz M, et al. Development of Varicella-Zoster virus infection in patients with chronic myelogenous leukemia treated with imatinib mesylate. *Clin Cancer Res*. 2003;9:976–80.
28. Breccia M, Girmenia C, Latagliata R, et al. Low incidence rate of opportunistic and viral infections during imatinib treatment in chronic myeloid leukemia patients in early and late chronic phase. *Mediterr J Hematol Infect Dis*. 2011;3:e2011021.
29. Saglio G, Kim DW, Issaragrisil S, et al. Nilotinib versus imatinib for newly diagnosed chronic myeloid leukemia. *N Engl J Med*. 2010;362:2251–9.
30. Larson RA, Hochhaus A, Hughes TP, et al. Nilotinib vs imatinib in patients with newly diagnosed Philadelphia chromosome-positive chronic myeloid leukemia in chronic phase: ENESTnd 3-year follow-up. *Leukemia*. 2012;26:2197–203.
31. Koren-Michowitz M, le Coutre P, Duyster J, et al. Activity and tolerability of nilotinib: a retrospective multicenter analysis of chronic myeloid leukemia patients who are imatinib resistant or intolerant. *Cancer*. 2010;116:4564–72.
32. Lai GM, Yan SL, Chang CS, Tsai CY. Hepatitis B reactivation in chronic myeloid leukemia patients receiving tyrosine kinase inhibitor. *World J Gastroenterol*. 2013;19:1318–21.
33. Kantarjian HM, Shah NP, Cortes JE, et al. Dasatinib or imatinib in newly diagnosed chronic-phase chronic myeloid leukemia: 2-year follow-up from a randomized phase 3 trial (DASISION). *Blood*. 2012;119:1123–9.
34. Al-Ameri A, Kantarjian H, Burton E, et al. Low risk of infectious events in patients (pts) with chronic myeloid leukemia (CML) in chronic phase (CP) treated with dasatinib. ASH Annual Meeting 2009, December 7, 2009, Abstract 3291, New Orleans, USA. 2015.
35. Rodriguez GH, Ahmed SI, Al Akhrass F, Rallapalli V, Safdar A. Characteristics of, and risk factors for, infections in patients with cancer treated with dasatinib and a brief review of other complications. *Leuk Lymphoma*. 2012;53:1530–5.
36. Chang H, Hung YS, Chou WC. *Pneumocystis jirovecii* pneumonia in patients receiving dasatinib treatment. *Int J Infect Dis*. 2014;25:165–7.
37. Lilly MB, Ottmann OG, Shah NP, et al. Dasatinib 140 mg once daily versus 70 mg twice daily in patients with Ph-positive acute lymphoblastic leukemia who failed imatinib: results from a phase 3 study. *Am J Hematol*. 2010;85:164–70.
38. Garcia-Munoz R, Galar A, Moreno C, et al. Parvovirus B19 acute infection and a reactivation of cytomegalovirus and herpesvirus 6 in a chronic myeloid leukemia patient during treatment with dasatinib (BMS-354825). *Leuk Lymphoma*. 2007;48:2461–4.
39. Gambacorti-Passerini C, Cortes JE, Lipton JH, et al. Safety of bosutinib versus imatinib in the phase 3 BELA trial in newly diagnosed chronic phase chronic myeloid leukemia. *Am J Hematol*. 2014;89:947–53.
40. Cortes JE, Talpaz M, Kantarjian H. Ponatinib in Philadelphia chromosome-positive leukemias. *N Engl J Med*. 2014;370:577.
41. Brabender J, Danenberg KD, Metzger R, et al. Epidermal growth factor receptor and HER2-neu mRNA expression in non-small cell lung cancer is correlated with survival. *Clin Cancer Res*. 2001;7:1850–5.
42. Miller VA, Hirsh V, Cadranel J, et al. Afatinib versus placebo for patients with advanced, metastatic non-small-cell lung cancer after failure of erlotinib, gefitinib, or both, and one or two lines of chemotherapy (LUX-Lung 1): a phase 2b/3 randomised trial. *Lancet Oncol*. 2012;13:528–38.
43. Sequist LV, Yang JC, Yamamoto N, et al. Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. *J Clin Oncol*. 2013;31:3327–34.
44. Yang JC, Wu YL, Schuler M, et al. Afatinib versus cisplatin-based chemotherapy for EGFR mutation-positive lung adenocarcinoma (LUX-Lung 3 and LUX-Lung 6): analysis of overall survival data from two randomised, phase 3 trials. *Lancet Oncol*. 2015;16:141–51.
45. Koivunen JP, Mermel C, Zejnullahu K, et al. EML4-ALK fusion gene and efficacy of an ALK kinase inhibitor in lung cancer. *Clin Cancer Res*. 2008;14:4275–83.
46. Shaw AT, Kim DW, Nakagawa K, et al. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med*. 2013;368:2385–94.
47. Solomon BJ, Mok T, Kim DW, et al; PROFILE 1014 Investigators. First-line crizotinib versus chemotherapy in ALK-positive lung cancer. *N Engl J Med*. 2014;371:2167–77.
48. Gambacorti Passerini C, Farina F, Stasia A, et al. Crizotinib in advanced, chemoresistant anaplastic lymphoma kinase-positive lymphoma patients. *J Natl Cancer Inst*. 2014;106:djt378.
49. Khozin S, Blumenthal GM, Zhang L, et al. FDA approval: ceritinib for the treatment of metastatic anaplastic lymphoma kinase-positive non-small cell lung cancer. *Clin Cancer Res*. 2015;21(11):2436–9.
50. Takeda M, Okamoto I, Nakagawa K. Pooled safety analysis of EGFR-TKI treatment for EGFR mutation-positive non-small cell lung cancer. *Lung Cancer*. 2015;88:74–9.
51. Burgel PR, Nadel JA. Epidermal growth factor receptor-mediated innate immune responses and their roles in airway diseases. *Eur Respir J*. 2008;32:1068–81.
52. Rosell R, Carcereny E, Gervais R, et al; Spanish Lung Cancer Group in collaboration with Groupe Français de Pneumo-Cancérologie and Associazione Italiana Oncologia Toracica. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol*. 2012;13:239–46.
53. Cappuzzo F, Ciuleanu T, Stelmakh L, et al; SATURN Investigators. Erlotinib as maintenance treatment in advanced non-small-cell lung cancer: a multicentre, randomised, placebo-controlled phase 3 study. *Lancet Oncol*. 2010;11:521–29.
54. Goss GD, O'Callaghan C, Lorimer I, et al. Gefitinib versus placebo in completely resected non-small-cell lung cancer: results of the NCIC CTG BR19 study. *J Clin Oncol*. 2013;31:3320–6.
55. Gaafar RM, Surmont VF, Scagliotti GV, et al; EORTC Lung Cancer Group and the Italian Lung Cancer Project. A double-blind, randomised, placebo-controlled phase III intergroup study of gefitinib in patients with advanced NSCLC, non-progressing after first line platinum-based chemotherapy (EORTC 08021/ILCP 01/03). *Eur J Cancer*. 2011;47:2331–40.
56. Wu YL, Zhou C, Hu CP, et al. Afatinib versus cisplatin plus gemcitabine for first-line treatment of Asian patients with advanced non-small-cell lung cancer harbouring EGFR mutations (LUX-Lung 6): an open-label, randomised phase 3 trial. *Lancet Oncol*. 2014;15:213–22.
57. Shaw AT, Kim DW, Mehra R, et al. Ceritinib in ALK-rearranged non-small-cell lung cancer. *N Engl J Med*. 2014;370:1189–97.
58. Munoz J, Schlette E, Kurzrock R. Rapid response to vemurafenib in a heavily pretreated patient with hairy cell leukemia and a BRAF mutation. *J Clin Oncol*. 2013;31:e351–2.
59. Kim KB, Cabanillas ME, Lazar AJ, et al. Clinical responses to vemurafenib in patients with metastatic papillary thyroid cancer harboring BRAF(V600E) mutation. *Thyroid*. 2013;23:1277–83.
60. Robert C, Karaszewska B, Schachter J, et al. Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med*. 2015;372:30–9.
61. Li H, Hu D, Fan H, et al. beta-Arrestin 2 negatively regulates Toll-like receptor 4 (TLR4)-triggered inflammatory signaling via targeting p38 MAPK and interleukin 10. *J Biol Chem*. 2014;289:23075–85.
62. Schilling B, Sondermann W, Zhao F, et al. Differential influence of vemurafenib and dabrafenib on patients' lymphocytes despite similar clinical efficacy in melanoma. *Ann Oncol*. 2014;25:747–53.
63. Sondermann W, Griewank KG, Schilling B, et al. Corticosteroids augment BRAF inhibitor vemurafenib induced lymphopenia and risk of infection. *PLoS One*. 2015;10:e0124590.
64. Chapman PB, Hauschild A, Robert C, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med*. 2011;364:2507–16.
65. McArthur GA, Chapman PB, Robert C, et al. Safety and efficacy of vemurafenib in BRAF(V600E) and BRAF(V600K) mutation-positive melanoma (BRIM-3): extended follow-up of a phase 3, randomised, open-label study. *Lancet Oncol*. 2014;15:323–32.
66. Larkin J, Ascierto PA, Dréno B, et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med*. 2014;371:1867–76.
67. Neubauer H, Cumano A, Muller M, Wu H, Huffstadt U, Pfefferer K. Jak2 deficiency defines an essential developmental checkpoint in definitive hematopoiesis. *Cell*. 1998;93:397–409.
68. Jones AV, Kreil S, Zoi K, et al. Widespread occurrence of the JAK2 V617F mutation in chronic myeloproliferative disorders. *Blood*. 2005;106:2162–8.
69. Levine RL, Loriaux M, Huntly BJ, et al. The JAK2V617F activating mutation occurs in chronic myelomonocytic leukemia and acute myeloid leukemia, but not in acute lymphoblastic leukemia or chronic lymphocytic leukemia. *Blood*. 2005;106:3377–9.
70. Staerk J, Constantinescu SN. The JAK-STAT pathway and hematopoietic stem cells from the JAK2 V617F perspective. *JAKSTAT*. 2012;1:184–90.
71. Vyas D, O'Dell KM, Bandy JL, Boyce EG. Tofacitinib: the first Janus kinase (JAK) inhibitor for the treatment of rheumatoid arthritis. *Ann Pharmacother*. 2013;47:1524–31.
72. Harrison C, Kiladjian JJ, Al-Ali HK, et al. JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med*. 2012;366:787–98.
73. Vannucchi AM, Kiladjian JJ, Griesshammer M, et al. Ruxolitinib versus standard therapy for the treatment of polycythemia vera. *N Engl J Med*. 2015;372:426–35.
74. Burmester GR, Blanco R, Charles-Schoeman C, et al; ORAL Step investigators. Tofacitinib (CP-690,550) in combination with methotrexate in patients with active rheumatoid arthritis with an inadequate response to tumour necrosis factor inhibitors: a randomised phase 3 trial. *Lancet*. 2013;381:451–60.



75. Maeshima K, Yamaoka K, Kubo S, et al. The JAK inhibitor tofacitinib regulates synovitis through inhibition of interferon-gamma and interleukin-17 production by human CD4+ T cells. *Arthritis Rheum*. 2012;64:1790–8.
76. Migita K, Miyashita T, Izumi Y, et al. Inhibitory effects of the JAK inhibitor CP690,550 on human CD4(+) T lymphocyte cytokine production. *BMC Immunol*. 2011;12:51.
77. Sonomoto K, Yamaoka K, Kubo S, et al. Effects of tofacitinib on lymphocytes in rheumatoid arthritis: relation to efficacy and infectious adverse events. *Rheumatology (Oxford)*. 2014;53:914–8.
78. Schönberg K, Rudolph J, Vonnahme M, et al. JAK inhibition impairs NK cell function in myeloproliferative neoplasms. *Cancer Res*. 2015;75(11):2187–99.
79. Parampalli YS, Stubig T, Cornez I, et al. JAK1/2 inhibition impairs T cell function in vitro and in patients with myeloproliferative neoplasms. *Br J Haematol*. 2015;169(6):824–33.
80. Heine A, Held SA, Daecke SN, et al. The JAK-inhibitor ruxolitinib impairs dendritic cell function in vitro and in vivo. *Blood*. 2013;122:1192–202.
81. Cohen S, Radominski SC, Gomez-Reino JJ, et al. Analysis of infections and all-cause mortality in phase II, phase III, and long-term extension studies of tofacitinib in patients with rheumatoid arthritis. *Arthritis Rheumatol*. 2014;66:2924–37.
82. Sandborn WJ, Ghosh S, Panes J, et al; Study A3921063 Investigators. Tofacitinib, an oral Janus kinase inhibitor, in active ulcerative colitis. *N Engl J Med*. 2012;367:616–24.
83. Vincenti F, Tesesco Silva H, Busque S, et al. Randomized phase 2b trial of tofacitinib (CP-690,550) in de novo kidney transplant patients: efficacy, renal function and safety at 1 year. *Am J Transplant*. 2012;12:2446–56.
84. Fleischmann R, Kremer J, Cush J, et al; ORAL Solo Investigators. Placebo-controlled trial of tofacitinib monotherapy in rheumatoid arthritis. *N Engl J Med*. 2012;367:495–507.
85. Pemmaraju N, Kantarjian H, Kadia T, et al. A phase I/II study of the Janus kinase (JAK)1 and 2 inhibitor ruxolitinib in patients with relapsed or refractory acute myeloid leukemia. *Clin Lymphoma Myeloma Leuk*. 2015;15:171–6.
86. Wysham NG, Sullivan DR, Allada G. An opportunistic infection associated with ruxolitinib, a novel janus kinase 1,2 inhibitor. *Chest*. 2013;143:1478–9.
87. Caocci G, Murgia F, Podda L, Solinas A, Atzeni S, La Nasa G. Reactivation of hepatitis B virus infection following ruxolitinib treatment in a patient with myelofibrosis. *Leukemia*. 2014;28:225–7.
88. Goldberg RA, Reichel E, Oshry LJ. Bilateral toxoplasmosis retinitis associated with ruxolitinib. *N Engl J Med*. 2013;369:681–3.
89. Colomba C, Rubino R, Siracusa L, et al. Disseminated tuberculosis in a patient treated with a JAK2 selective inhibitor: a case report. *BMC Res Notes*. 2012;5:552.
90. Wathes R, Moule S, Milojkovic D. Progressive multifocal leukoencephalopathy associated with ruxolitinib. *N Engl J Med*. 2013;369:197–8.
91. Choi J, Cooper ML, Alahmari B, et al. Pharmacologic blockade of JAK1/JAK2 reduces GvHD and preserves the graft-versus-leukemia effect. *PLoS One*. 2014;9:e109799.
92. Spoerl S, Mathew NR, Bscheider M, et al. Activity of therapeutic JAK 1/2 blockade in graft-versus-host disease. *Blood*. 2014;123:3832–42.
93. Zeiser R, Lengerke C, Spoerl S, et al. 41st Annual Meeting of the European Society for Blood and Marrow Transplantation 2015, March 23, 2015, oral presentation No. 36, Istanbul, Turkey. High response rates in patients treated with ruxolitinib for corticosteroid-refractory Graft-versus-host disease: analysis including 13 Transplant Centers. 2015.
94. Heine A, Brossart P, Wolf D. Ruxolitinib is a potent immunosuppressive compound: is it time for anti-infective prophylaxis? *Blood*. 2013;122:3843–4.
95. Chiarini F, Evangelisti C, McCubrey JA, Martelli AM. Current treatment strategies for inhibiting mTOR in cancer. *Trends Pharmacol Sci*. 2015;36:124–35.
96. Guba M, von Breitenbuch P, Steinbauer M, et al. Rapamycin inhibits primary and metastatic tumor growth by antiangiogenesis: involvement of vascular endothelial growth factor. *Nat Med*. 2002;8:128–35.
97. Schulze M, Stock C, Zaccagnini M, Teber D, Rassweiler JJ. Temsirolimus. *Recent Results Cancer Res*. 2014;201:393–403.
98. Gurk-Turner C, Maniatisitkul W, Cooper M. A comprehensive review of everolimus clinical reports: a new mammalian target of rapamycin inhibitor. *Transplantation*. 2012;94:659–68.
99. Mielke S, Lutz M, Schmidhuber J, et al. Salvage therapy with everolimus reduces the severity of treatment-refractory chronic GVHD without impairing disease control: a dual center retrospective analysis. *Bone Marrow Transplant*. 2014;49:1412–8.
100. Guba M, Pratschke J, Hugo C, et al; SMART-Study Group. Renal function, efficacy, and safety of sirolimus and mycophenolate mofetil after short-term calcineurin inhibitor-based quadruple therapy in de novo renal transplant patients: one-year analysis of a randomized multicenter trial. *Transplantation*. 2010;90:175–83.
101. Rittà M, Costa C, Solidoro P, et al. Everolimus-based immunosuppressive regimens in lung transplant recipients: impact on CMV infection. *Antiviral Res*. 2015;113:19–26.
102. Hess G, Herbrecht R, Romaguera J, et al. Phase III study to evaluate temsirolimus compared with investigator's choice therapy for the treatment of relapsed or refractory mantle cell lymphoma. *J Clin Oncol*. 2009;27:3822–9.
103. Hudes G, Carducci M, Tomczak P, et al. Temsirolimus, interferon alfa, or both for advanced renal-cell carcinoma. *N Engl J Med*. 2007;356:2271–81.
104. Sarkaria JN, Galanis E, Wu W, et al. Combination of temsirolimus (CCI-779) with chemoradiation in newly diagnosed glioblastoma multiforme (GBM) (NCCTG trial N027D) is associated with increased infectious risks. *Clin Cancer Res*. 2010;16:5573–80.
105. Motzer RJ, Escudier B, Oudard S, et al; RECORD-1 Study Group. Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial. *Lancet*. 2008;372:449–56.
106. Porta C, Osanto S, Ravaud A, et al. Management of adverse events associated with the use of everolimus in patients with advanced renal cell carcinoma. *Eur J Cancer*. 2011;47:1287–98.
107. Trelinska J, Dachowska I, Kotulska K, Fendler W, Jozwiak S, Mlynarski W. Complications of mammalian target of rapamycin inhibitor anticancer treatment among patients with tuberous sclerosis complex are common and occasionally life-threatening. *Anticancer Drugs*. 2015;26:437–42.
108. Baselga J, Campone M, Piccart M, et al. Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer. *N Engl J Med*. 2012;366:520–9.
109. Wolff AC, Lazar AA, Bondarenko I, et al. Randomized phase III placebo-controlled trial of letrozole plus oral temsirolimus as first-line endocrine therapy in postmenopausal women with locally advanced or metastatic breast cancer. *J Clin Oncol*. 2013;31:195–202.
110. Pritchard KI, Burris HA 3rd, Ito Y, et al. Safety and efficacy of everolimus with exemestane vs. exemestane alone in elderly patients with HER2-negative, hormone receptor-positive breast cancer in BOLERO-2. *Clin Breast Cancer*. 2013;13:421–32.
111. Kaymakcalan MD, Je Y, Sonpavde G, et al. Risk of infections in renal cell carcinoma (RCC) and non-RCC patients treated with mammalian target of rapamycin inhibitors. *Br J Cancer*. 2013;108:2478–84.
112. Rafii S, Roda D, Geuna E, et al. Higher risk of infections with PI3 K-AKT-mTOR pathway inhibitors in patients with advanced solid tumors on phase I clinical trials. *Clin Cancer Res*. 2015;21:1869–76.
113. Carbonnaud M, Molin Y, Souquet PJ, Tantin A, Lombard-Bohas C, Walter T. *Pneumocystis jirovecii* pneumonia under everolimus in two patients with metastatic pancreatic neuroendocrine tumors. *Invest New Drugs*. 2014;32:1308–10.
114. Fillatre P, Decaux O, Jouneau S, et al. Incidence of *Pneumocystis jirovecii* pneumonia among groups at risk in HIV-negative patients. *Am J Med*. 2014;127:1242–7.
115. White DA, Camus P, Endo M, et al. Noninfectious pneumonitis after everolimus therapy for advanced renal cell carcinoma. *Am J Respir Crit Care Med*. 2010;182:396–403.
116. Funakoshi T, Latif A, Galsky MD. Safety and efficacy of addition of VEGFR and EGFR-family oral small-molecule tyrosine kinase inhibitors to cytotoxic chemotherapy in solid cancers: a systematic review and meta-analysis of randomized controlled trials. *Cancer Treat Rev*. 2014;40:636–7.
117. Chung EK, Stadler WM. Vascular endothelial growth factor pathway-targeted therapy as initial systemic treatment of patients with renal cancer. *Clin Genitourin Cancer*. 2008;6(suppl 1):S22–8.
118. Powles T, Chowdhury S, Bower M, et al. The effect of sunitinib on immune subsets in metastatic clear cell renal cancer. *Urol Int*. 2011;86:53–9.
119. Michaelson MD, Oudard S, Ou YC, et al. Randomized, placebo-controlled, phase III trial of sunitinib plus prednisone versus prednisone alone in progressive, metastatic, castration-resistant prostate cancer. *J Clin Oncol*. 2014;32:76–82.
120. Crown JP, Diéras V, Staroslawska E, et al. Phase III trial of sunitinib in combination with capecitabine versus capecitabine monotherapy for the treatment of patients with pretreated metastatic breast cancer. *J Clin Oncol*. 2013;31:2870–8.
121. Carrato A, Swieboda-Sadlej A, Staszewska-Skurczynska M, et al. Fluorouracil, leucovorin, and irinotecan plus either sunitinib or placebo in metastatic colorectal cancer: a randomized, phase III trial. *J Clin Oncol*. 2013;31:1341–7.
122. Ready NE, Pang HH, Gu L, et al. Chemotherapy with or without maintenance sunitinib for untreated extensive-stage small-cell lung cancer: a randomized, double-blind, placebo-controlled phase II study-CALGB 30504 (alliance). *J Clin Oncol*. 2015;33(15):1660–5.
123. Hasskarl J. Sorafenib: targeting multiple tyrosine kinases in cancer. *Recent Results Cancer Res*. 2014;201:145–64.
124. Man CH, Fung TK, Ho C, et al. Sorafenib treatment of FLT3-ITD(+) acute myeloid leukemia: favorable initial outcome and mechanisms of subsequent nonresponsiveness associated with the emergence of a D835 mutation. *Blood*. 2012;119:5133–43.
125. Houben R, Voigt H, Noelke C, Hofmeister V, Becker JC, Schrama D. MAPK-independent impairment of T-cell responses by the multikinase inhibitor sorafenib. *Mol Cancer Ther*. 2009;8:433–40.



126. Zhao W, Gu YH, Song R, Qu BQ, Xu Q. Sorafenib inhibits activation of human peripheral blood T cells by targeting LCK phosphorylation. *Leukemia*. 2008;22:1226–33.
127. Fishman MN, Tomshine J, Fulp WJ, Foreman PK. A systematic review of the efficacy and safety experience reported for Sorafenib in advanced renal cell carcinoma (RCC) in the post-approval setting. *PLoS One*. 2015;10:e0120877.
128. Escudier B, Eisen T, Stadler WM, et al. Sorafenib for treatment of renal cell carcinoma: final efficacy and safety results of the phase III treatment approaches in renal cancer global evaluation trial. *J Clin Oncol*. 2009;27:3312–8.
129. Descamps V, Helle F, Louandre C, et al. The kinase-inhibitor sorafenib inhibits multiple steps of the Hepatitis C Virus infectious cycle in vitro. *Antiviral Res*. 2015;118:93–102.
130. Serve H, Krug U, Wagner R, et al. Sorafenib in combination with intensive chemotherapy in elderly patients with acute myeloid leukemia: results from a randomized, placebo-controlled trial. *J Clin Oncol*. 2013;31:3110–8.
131. Crona DJ, Keisler MD, Walko CM. Regorafenib: a novel multitargeted tyrosine kinase inhibitor for colorectal cancer and gastrointestinal stromal tumors. *Ann Pharmacother*. 2013;47:1685–96.
132. Grothey A, Van Cutsem E, Sobrero A, et al; CORRECT Study Group. Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet*. 2013;381:303–12.
133. Demetri GD, Reichardt P, Kang YK, et al; GRID Study Investigators. Efficacy and safety of regorafenib for advanced gastrointestinal stromal tumours after failure of imatinib and sunitinib (GRID): an international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet*. 2013;381:295–302.
134. Motzer RJ, Hutson TE, Cella D, et al. Pazopanib versus sunitinib in metastatic renal-cell carcinoma. *N Engl J Med*. 2013;369:722–31.
135. van der Graaf WT, Blay JY, Chawla SP, et al; EORTC Soft Tissue and Bone Sarcoma Group; PALETTE Study Group. Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet*. 2012;379:1879–86.
136. Pignata S, Lorusso D, Scambia G, et al; MITO 11 Investigators. Pazopanib plus weekly paclitaxel versus weekly paclitaxel alone for platinum-resistant or platinum-refractory advanced ovarian cancer (MITO 11): a randomised, open-label, phase 2 trial. *Lancet Oncol*. 2015;16(5):561–8.
137. Scagliotti GV, Felip E, Besse B, et al. An open-label, multicenter, randomized, phase II study of pazopanib in combination with pemetrexed in first-line treatment of patients with advanced-stage non-small-cell lung cancer. *J Thorac Oncol*. 2013;8:1529–37.
138. Sternberg CN, Hawkins RE, Wagstaff J, et al. A randomised, double-blind phase III study of pazopanib in patients with advanced and/or metastatic renal cell carcinoma: final overall survival results and safety update. *Eur J Cancer*. 2013;49:1287–96.
139. Rini BI, Escudier B, Tomczak P, et al. Comparative effectiveness of axitinib versus sorafenib in advanced renal cell carcinoma (AXIS): a randomised phase 3 trial. *Lancet*. 2011;378:1931–9.
140. Sonpavde G, Hutson TE, Rini BI. Axitinib for renal cell carcinoma. *Expert Opin Investig Drugs*. 2008;17:741–8.
141. Stehle F, Schulz K, Fahldieck C, et al. Reduced immunosuppressive properties of axitinib in comparison with other tyrosine kinase inhibitors. *J Biol Chem*. 2013;288:16334–47.
142. van Oers MH, Van Glabbeke M, Giurgea L, et al. Rituximab maintenance treatment of relapsed/resistant follicular non-Hodgkin's lymphoma: long-term outcome of the EORTC 20981 phase III randomized intergroup study. *J Clin Oncol*. 2010;28:2853–8.
143. Nissen JC, Hummel M, Brade J, et al. The risk of infections in hematologic patients treated with rituximab is not influenced by cumulative rituximab dosage – a single center experience. *BMC Infect Dis*. 2014;14:364.
144. Byrd JC, Brown JR, O'Brien S, et al; RESONATE Investigators. Ibrutinib versus ofatumumab in previously treated chronic lymphoid leukemia. *N Engl J Med*. 2014;371:213–23.
145. Wang ML, Rule S, Martin P, et al. Targeting BTK with ibrutinib in relapsed or refractory mantle-cell lymphoma. *N Engl J Med*. 2013;369:507–16.
146. Gopal AK, Kahl BS, de Vos S, et al. PI3Kdelta inhibition by idelalisib in patients with relapsed indolent lymphoma. *N Engl J Med*. 2014;370:1008–18.
147. Furman RR, Sharman JP, Coutre SE, et al. Idelalisib and rituximab in relapsed chronic lymphocytic leukemia. *N Engl J Med*. 2014;370:997–1007.
148. Winkelstein JA, Marino MC, Lederman HM, et al. X-linked agammaglobulinemia: report on a United States registry of 201 patients. *Medicine (Baltimore)*. 2006;85:193–202.
149. Taneichi H, Kanegane H, Sira MM, et al. Toll-like receptor signaling is impaired in dendritic cells from patients with X-linked agammaglobulinemia. *Clin Immunol*. 2008;126:148–54.
150. Ormsby T, Schlecker E, Ferdin J, et al. Btk is a positive regulator in the TREM-1/DAP12 signaling pathway. *Blood*. 2011;118:936–45.
151. Köprülü AD, Kastner R, Wienerroither S, et al. The tyrosine kinase Btk regulates the macrophage response to *Listeria monocytogenes* infection. *PLoS One*. 2013;8:e60476.
152. Lee KG, Xu S, Kang ZH, et al. Bruton's tyrosine kinase phosphorylates Toll-like receptor 3 to initiate antiviral response. *Proc Natl Acad Sci U S A*. 2012;109:5791–6.
153. Strijbis K, Tafesse FG, Fairn GD, et al. Bruton's Tyrosine Kinase (BTK) and Vav1 contribute to Dectin1-dependent phagocytosis of *Candida albicans* in macrophages. *PLoS Pathog*. 2013;9:e1003446.
154. Angulo I, Vadas O, Garcon F, et al. Phosphoinositide 3-kinase delta gene mutation predisposes to respiratory infection and airway damage. *Science*. 2013;342:866–71.
155. Lucas CL, Kuehn HS, Zhao F, et al. Dominant-activating germline mutations in the gene encoding the PI(3)K catalytic subunit p110delta result in T cell senescence and human immunodeficiency. *Nat Immunol*. 2014;15:88–97.
156. Xie S, Chen M, Yan B, He X, Chen X, Li D. Identification of a role for the PI3K/AKT/mTOR signaling pathway in innate immune cells. *PLoS One*. 2014;9:e94496.
157. Byrd JC, Furman RR, Coutre SE, et al. Targeting BTK with ibrutinib in relapsed chronic lymphocytic leukemia. *N Engl J Med*. 2013;369:32–42.
158. Burger JA, Keating MJ, Wierda WG, et al. Safety and activity of ibrutinib plus rituximab for patients with high-risk chronic lymphocytic leukaemia: a single-arm, phase 2 study. *Lancet Oncol*. 2014;15:1090–9.
159. Treon SP, Tripsas CK, Meid K, et al. Ibrutinib in previously treated Waldenström's macroglobulinemia. *N Engl J Med*. 2015;372:1430–40.
160. Byrd JC, Furman RR, Coutre SE, et al. Three-year follow-up of treatment-naïve and previously treated patients with CLL and SLL receiving single-agent ibrutinib. *Blood*. 2015;125:2497–506.
161. Brown JR, Byrd JC, Coutre SE, et al. Idelalisib, an inhibitor of phosphatidylinositol 3-kinase p110delta, for relapsed/refractory chronic lymphocytic leukemia. *Blood*. 2014;123:3390–7.
162. Kahl BS, Spurgeon SE, Furman RR, et al. A phase 1 study of the PI3Kdelta inhibitor idelalisib in patients with relapsed/refractory mantle cell lymphoma (MCL). *Blood*. 2014;123:3398–405.
163. Young JA. Epidemiology and management of infectious complications of contemporary management of chronic leukemias. *Infect Disord Drug Targets*. 2011;11:3–10.
164. O'Brien S. Update on a phase 2 study of idelalisib in combination with rituximab in treatment-naïve patients = 65 years with chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL). In: Meeting of the American Society of Hematology 2014. ASH Annual Meeting 2014, December 6, 2014, Abstract 1994, San Francisco, USA. 2014.
165. Flinn IW, Kahl BS, Leonard JP, et al. Idelalisib, a selective inhibitor of phosphatidylinositol 3-kinase-delta, as therapy for previously treated indolent non-Hodgkin lymphoma. *Blood*. 2014;123:3406–13.
166. Dubovsky JA, Flynn R, Du J, et al. Ibrutinib treatment ameliorates murine chronic graft-versus-host disease. *J Clin Invest*. 2014;124:4867–76.
167. Dagher R, Cohen M, Williams G, et al. Approval summary: imatinib mesylate in the treatment of metastatic and/or unresectable malignant gastrointestinal stromal tumors. *Clin Cancer Res*. 2002;8:3034–8.
168. Daver N, Thomas D, Ravandi F, et al. Final report of a phase II study of imatinib mesylate with hyper-CVAD for the frontline treatment of adult patients with Philadelphia chromosome positive (Ph+) acute lymphoblastic leukemia. *Haematologica*. 2015;100(5):653–61.
169. Yanada M, Takeuchi J, Sugiura I, et al; Japan Adult Leukemia Study Group. High complete remission rate and promising outcome by combination of imatinib and chemotherapy for newly diagnosed BCR-ABL-positive acute lymphoblastic leukemia: a phase II study by the Japan Adult Leukemia Study Group. *J Clin Oncol*. 2006;24:460–6.
170. Hauschild A, Grob JJ, Demidov LV, et al. Dabrafenib in BRAF-mutated metastatic melanoma: a multicentre, open-label, phase 3 randomised controlled trial. *Lancet*. 2012;380:358–65.
171. Pavel ME, Hainsworth JD, Baudin E, et al; RADIANT-2 Study Group. Everolimus plus octreotide long-acting repeatable for the treatment of advanced neuroendocrine tumours associated with carcinoid syndrome (RADIANT-2): a randomised, placebo-controlled, phase 3 study. *Lancet*. 2011;378:2005–12.
172. Franz DN, Belousova E, Sparagana S, et al. Everolimus for subependymal giant cell astrocytoma in patients with tuberous sclerosis complex: 2-year open-label extension of the randomised EXIST-1 study. *Lancet Oncol*. 2014;15:1513–20.
173. Motzer RJ, Hutson TE, Tomczak P, et al. Sunitinib versus interferon alfa in metastatic renal-cell carcinoma. *N Engl J Med*. 2007;356:115–24.
174. Raymond E, Dahan L, Raoul JL, et al. Sunitinib malate for the treatment of pancreatic neuroendocrine tumors. *N Engl J Med*. 2011;364:501–13.
175. Shirao K, Nishida T, Doi T, et al. Phase I/II study of sunitinib malate in Japanese patients with gastrointestinal stromal tumor after failure of prior treatment with imatinib mesylate. *Invest New Drugs*. 2010;28:866–75.



176. Kudo M, Imanaka K, Chida N, et al. Phase III study of sorafenib after transarterial chemoembolisation in Japanese and Korean patients with unresectable hepatocellular carcinoma. *Eur J Cancer*. 2011;47:2117–27.
177. Sleijfer S, Ray-Coquard I, Papai Z, et al. Pazopanib, a multikinase angiogenesis inhibitor, in patients with relapsed or refractory advanced soft tissue sarcoma: a phase II study from the European organisation for research and treatment of cancer-soft tissue and bone sarcoma group (EORTC study 62043). *J Clin Oncol*. 2009;27:3126–3132.
178. Hutson TE, Lesovoy V, Al-Shukri S, et al. Axitinib versus sorafenib as first-line therapy in patients with metastatic renal-cell carcinoma: a randomised open-label phase 3 trial. *Lancet Oncol*. 2013;14:1287–94.
179. Cervantes F, Vannucchi AM, Kiladjian JJ, et al; COMFORT-II Investigators. Three-year efficacy, safety, and survival findings from COMFORT-II, a phase 3 study comparing ruxolitinib with best available therapy for myelofibrosis. *Blood*. 2013;122:4047–53.
180. Wollenhaupt J, Silverfield J, Lee EB, et al. Safety and efficacy of tofacitinib, an oral Janus kinase inhibitor, for the treatment of rheumatoid arthritis in open-label, longterm extension studies. *J Rheumatol*. 2014;41:837–52.