



HHS Public Access

Author manuscript

Nat Rev Mol Cell Biol. Author manuscript; available in PMC 2016 May 02.

Published in final edited form as:

Nat Rev Mol Cell Biol. 2013 November ; 14(11): 699–712. doi:10.1038/nrm3679.

RNA in unexpected places: long non-coding RNA functions in diverse cellular contexts

Sarah Geisler^{1,2} and Jeff Coller¹

Sarah Geisler: sarah.geisler@bsse.ethz.ch; Jeff Coller: jmc71@case.edu

¹Center for RNA Molecular Biology, Case Western Reserve University, Cleveland, Ohio 44106, USA

Abstract

The increased application of transcriptome-wide profiling approaches has led to an explosion in the number of documented long non-coding RNAs (lncRNAs). While these new and enigmatic players in the complex transcriptional milieu are encoded by a significant proportion of the genome, their functions are mostly unknown. Early discoveries support a paradigm in which lncRNAs regulate transcription via chromatin modulation, but new functions are steadily emerging. Given the biochemical versatility of RNA, lncRNAs may be used for various tasks, including post-transcriptional regulation, organization of protein complexes, cell-cell signalling and allosteric regulation of proteins.

Dedicated consortiums, such as the ENCODE (Encyclopedia of DNA Elements) project, have markedly expanded our knowledge of what lies in the dark recesses of the genome through their extensive annotation efforts¹. These findings in conjunction with previous studies looking specifically at transcriptional outputs have underscored the pervasiveness with which genomes are transcribed^{2,3}. An important implication of these findings is that whereas only a minuscule fraction of the human genome encodes proteins, nearly 60% is represented in processed transcripts that seem to lack protein-coding capacity⁴. Together with observations that more sophisticated organisms tend to have more non-coding DNA, this raises the possibility that the barren regions between genes are actually 'elysian fields' rich with information⁵. The implications of this are undeniably intriguing, but we are still far from ascribing biological functions to the vast array of non-coding RNA (ncRNA) transcripts. With thousands of documented ncRNAs, pervasive transcription has been described in virtually all eukaryotic organisms^{6,7}.

For the better part of the past decade, particular attention has focused on the exploding class of transcripts referred to as long non-coding RNAs (lncRNAs), arbitrarily defined as being longer than 200 nucleotides^{7,8}. Given the prevalence of lncRNA expression, it has been posited that lncRNAs might constitute a significant fraction of the functional output of mammalian genomes⁷⁻⁹. Such notions have been met with considerable, and quite possibly

²Present address: Department of Biosystems Science and Engineering, Eidgenössische Technische Hochschule (ETH) Zürich, 4058 Basel, Switzerland.

Competing interests statement

The authors declare no competing financial interests.

legitimate, scepticism¹⁰. Indeed, the documentation of pervasive transcription has far outpaced the molecular characterization of the transcripts produced. Although some lncRNA transcripts may represent transcriptional noise, a small but steadily growing list has authentic biological roles^{6,11–13}. For example, lncRNAs have been implicated in regulating imprinting, dosage compensation, cell cycle regulation, pluripotency, retro-transposon silencing, meiotic entry and telomere length, to name just a few^{12,13}. Despite these advances, most lncRNAs remain partially uncharacterized. Additionally, there has been a heavy focus so far on the ways that lncRNAs regulate chromatin states, and this emphasis probably underrepresents the full repertoire of lncRNA function. Nonetheless, the rapidly growing lncRNA field is already changing not just our perspective of genomic content, but also the way we think about genes.

In this Review, we focus on the functional attributes of RNA and highlight the unconventional, and perhaps underappreciated, biological contributions of lncRNAs, including the diverse mechanisms through which lncRNAs participate in transcriptional regulation. We touch briefly on the roles of lncRNAs in regulating chromatin states, as this has been explored in several recent reviews (see REFS 8,9,13–15). In addition, we highlight roles beyond transcription whereby lncRNAs function in various cellular contexts, including post-transcriptional regulation, post-translational regulation of protein activity, organization of protein complexes, cell-cell signalling, as well as recombination.

A biochemically versatile polymer

RNA is a versatile molecule making it well suited for a myriad of functions. It is this feature that inspired the ‘RNA world hypothesis’ in which it was postulated that billions of years ago, RNA provided the precursors of all life¹⁶. The multifunctionality of RNA stems from several unique physiochemical properties. First, and perhaps most obvious, is its ability to base pair with other nucleic acids (FIG. 1a). RNA is, therefore, particularly adept at recognizing both RNA and DNA targets through simple one-to-one base pairing interactions. By comparison, proteins such as transcription activator-like effectors (TALEs) and PUF proteins require 100 times more genomic sequence space than an RNA to achieve sequence-specific binding¹⁷. Moreover, because two RNA transcripts can base pair at any point during the life cycle of the target mRNA, regulatory RNAs can influence transcription, processing, editing, translation or degradation of target mRNAs. Second, RNA molecules can fold into intricate three-dimensional structures that provide complex recognition surfaces (FIG. 1b). This structure expands the large variety of molecular targets that RNA can bind with high affinity and specificity. RNA structures can even be selected for *in vitro* to bind to anything from small molecules to proteins¹⁸. Third, in terms of both expression and structure, RNA is dynamic. More explicitly, because RNA can be rapidly transcribed and degraded, it is well suited for dynamic, transient expression (FIG. 1c). Moreover, without the need to be translated, a regulatory RNA gene could transition faster from being transcriptionally inactive to fully functional. In addition, as conformational changes can be triggered by ligand binding, RNA structures themselves can be very dynamic¹⁹. Fourth, RNA is malleable and therefore provides an excellent platform for evolutionary innovation (FIG. 1d). Specifically, unencumbered by amino acid-coding potential, regulatory RNAs are less restricted in terms of their conservation. As such, RNAs

are more tolerant of mutations, which could allow for the rapid evolution of diverse cellular activities. Last, RNA-dependent events can have the capacity to be heritable. This idea is supported by the demonstration of RNA-templated modifications to the genome (FIG. 1e). For example, retroviral genomic integrations as well as the presence of thousands of processed pseudogenes suggest that information housed within mature RNA transcripts can be integrated back into the genome^{20,21}. These instances of RNA-mediated events that have manifested in genomic change suggest it is possible for other RNA-dependent events to become heritable. Importantly, these defining properties of RNAs raise exciting possibilities as to what roles lncRNAs could have in the cell. Although various functional roles have now been attributed to lncRNAs, it is likely that as we dig deeper into the molecular biology of lncRNAs more functions will emerge.

lncRNAs as regulators of transcription

The number of lncRNAs with described functions is steadily increasing, and many of these reports revolve around their regulatory capacity. For example, lncRNAs often function as important *cis*- and *trans*-acting modulators of protein-coding gene expression⁸. A common theme has emerged in which lncRNAs regulate transcription via chromatin modulation (for reviews, see REFS 8,13,15). lncRNAs across a broad range of eukaryotes affect chromatin context, suggesting that this is a conserved function despite the fact that the transcripts themselves are often not conserved¹². Numerous lncRNAs physically associate with, and potentially target, histone-modifying activities to specific loci^{22,23} (TABLE 1). lncRNAs such as *HOTAIR* (HOX transcript antisense RNA), *ANRIL* (also known as CDKN2B anti-sense RNA 1) and *KCNQ1OT1* (KCNQ1 opposite strand or antisense transcript 1) have even been shown to bind more than one histone-modifying complex. As such, a paradigm in which lncRNAs can act as scaffolds that organize the concerted actions of chromatin-modifying complexes spatially and temporally is emerging^{15,24–28} (FIG. 2a; TABLE 1). For example, *HOTAIR* physically associates not only with Polycomb repressive complex 2 (PRC2) but also with LSD1 (Lys-specific demethylase 1)²⁴. PRC2 and LSD1 are responsible for the deposition of the repressive histone mark trimethylated Lys27 of histone H3 (H3K27me3) and removal of active H3K4me2 marks, respectively. Moreover, global analyses suggest that a large number of other lncRNAs can also bind PRC2 and LSD1 (REF. 22). In addition, other lncRNAs have been shown to bind overlapping but distinct combinations of histone-modifying complexes. For example, *KCNQ1OT1* binds PRC2 and the methyltransferase G9A (also known as EHMT2), whereas *ANRIL* binds PRC1 and PRC2 (REFS 26–28) (TABLE 1). *HOTAIR* and other lncRNAs have, therefore, been proposed to function as scaffolds that coordinate the targeting of distinct repressive histone-modifying complexes to target loci²⁵. However, within this framework, the detailed mechanism of how lncRNAs target specific DNA regions remains unclear.

Additionally, at least in some cases, lncRNA expression may influence epigenetic events through transcription-dependent mechanisms²⁹. The mammalian lncRNA *Airm* (antisense of *Igf2r* non-coding RNA) has been suggested to interfere with transcription during its regulation of *Igf2r* (insulin-like growth factor 2 receptor) because *Airm* transcription rather than the lncRNA product itself is required for silencing³⁰ (TABLE 1). Similarly, an antisense RNA has also been postulated to repress mRNA expression at the yeast *IME4* locus through

transcriptional interference³¹ (TABLE 1). In some instances (for example, the *GAL10-1*, *IME1* and *PHO84* loci in yeast), movement of the polymerase along the DNA locus can result in the deposition of histone modifications, which in turn repress expression from nearby promoters. This may be one mechanism of transcription-dependent lncRNA regulation^{32–35} (FIG. 2b; TABLE 1). Conversely, in flies non-coding transcription through Polycomb response elements is thought to counteract silencing during the switch from repressed to active states³⁶. Moreover, lncRNA transcription in various organisms can modulate the binding of regulatory factors^{37–40} (FIG. 2c; TABLE 1). An interesting example is the pair of *cis*-acting lncRNAs, *ICR1* and *PWR1*, which dictate the variegated expression of *FLO11* mRNA in yeast. Specifically, transcription of *ICR1* is thought to reset the *FLO11* locus by inhibiting recruitment of the Flo8 or Sfl1 transcription factors, which promote *FLO11* mRNA repression and activation, respectively. After this reset, if Flo8 binds it drives expression of *PWR1*, which in turn interferes with *ICR1* expression in *cis*. *ICR1* and *PWR1* lncRNAs therefore represent a ‘toggle switch’, resulting in *FLO11* mRNA expression when *PWR1* is expressed and *FLO11* mRNA repression when *ICR1* is expressed^{37,41}. By contrast, the lncRNA *SRG1* exerts chromatin regulation by directing a high level of nucleosomes to the region of the phosphoglycerate dehydrogenase *SER3* mRNA promoter³⁸. In these particular *cis*-acting instances, it is often unclear whether the phenotype associated with the locus arises from the lncRNA itself or rather from changes in DNA-protein interactions that arise from polymerase movement.

lncRNAs have now also been implicated in transcriptional upregulation by enhancers^{42,43}. A specific type of lncRNA, termed enhancer RNA (eRNA), displays enhancer-like activity and upregulates expression via the Mediator complex^{43,44} (TABLE 1). As studies suggest that classic enhancer elements are widely expressed, eRNAs may frequently be important for enhancer function at chromatin^{42,45,46}.

Aside from modulating chromatin, lncRNAs can regulate transcription through additional mechanisms. For example, lncRNAs can influence the transcription machinery directly. During heat shock, lncRNAs generated from Alu SINE elements mediate transcriptional repression through direct contact with RNA polymerase II (Pol II) (FIG. 2d; TABLE 1). This interaction inhibits transcription of specific mRNAs during heat shock⁴⁷. Furthermore, several lncRNAs can regulate the binding and/or activity of transcription factors. At the *DHFR* (dihydrofolate reductase) locus, expression of an upstream lncRNA impairs the assembly of the transcription pre-initiation complex in *trans* through the formation of an RNA-DNA triplex structure⁴⁸ (FIG. 2e; TABLE 1).

Moreover, several lncRNAs act directly on specific transcription factors. For instance, during the heat shock response, heat shock factor 1 (HSF1) is activated through the combined actions of a lncRNA, *HSR1* (heat shock RNA 1), and a surprising protein interaction partner and co-activator, translation elongation factor eEF1A⁴⁹ (TABLE 1). In another example, the *GAS5* (growth arrest specific 5) lncRNA folds into a structure that mimics the DNA-binding site of the glucocorticoid receptor, and the resulting interaction represses GR-mediated transcription⁵⁰ (FIG. 2f; TABLE 1). By contrast, the lncRNA *Evf2* (also known as *Dlx6os1*) can act either as a co-activator or co-repressor, depending on whether it recruits the transcriptional activator DLX2 or the transcriptional repressor MeCP2 (methyl-CpG

binding-protein 2) to specific DNA regulatory elements^{51,52} (FIG. 2f; TABLE 1). Furthermore, binding of lncRNAs generated from the *CCND1* (cyclin D1) promoter allosterically promotes a conformational switch in the TLS (translocated in liposarcoma) protein factor from an inactive to active form. Active TLS inhibits histone acetyltransferases, ultimately leading to repression of *CCND1* transcription⁵³ (FIG. 2f; TABLE 1). The lncRNA transcript thus indirectly promotes a repressive chromatin environment.

By contrast, the *NRON* (non-coding repressor of NFAT) lncRNA indirectly represses transcription by inhibiting nucleocytoplasmic shuttling of the transcription factor NFAT (nuclear factor of activated T cells)⁵⁴. The transport of NFAT, which is imported from the cytoplasm into the nucleus in response to calcium-dependent signals, is inhibited by *NRON*. *NRON* binds the transport receptor importin- β , and knockdown of *NRON* results in nuclear accumulation of NFAT, suggesting that *NRON* competes with NFAT for importin- β interaction⁵⁴ (FIG. 2g; TABLE 1).

Interestingly, lncRNAs have been indirectly linked to both gene activation and repression through the organization of nuclear subdomains. For instance, the lncRNAs *TUG1* (taurine upregulated 1) and *MALAT1* (metastasis associated lung adenocarcinoma transcript 1; also known as *NEAT2*) have been linked to repressive Polycomb group bodies and more active interchromatin granules, respectively (TABLE 1). Both lncRNAs bind Polycomb 2, but *TUG1* binds methylated Polycomb 2 and *MALAT1* binds the unmethylated protein⁵⁵. The methylation status of Polycomb 2 therefore dictates a switch in both its lncRNA-binding specificity and nuclear subcompartment localization. Importantly, this switch is accompanied by movement of Polycomb 2 target genes between active and repressive nuclear domains and ultimately influences downstream gene expression⁵⁵.

lncRNAs, therefore, can regulate transcription through several mechanisms (FIG. 2). Given the decades of research focused on transcriptional control from a transcription factor-centric point of view, it is interesting to speculate about the purpose of this additional layer of RNA-based regulation. Even at the yeast *GAL* locus, arguably one of the most extensively studied DNA loci during the past 50 years, a hidden layer of lncRNA-based regulation has now been described^{35,56}. Indeed there has been a growing interest in such RNA-based control during the past decade⁵⁷, and we and others have speculated that this extra layer of regulation reinforces the control that is imposed by protein factors at a locus. Notably, the impressive diversity of transcriptional regulatory mechanisms discussed here might just be the tip of the iceberg, with additional means of lncRNA-mediated transcriptional regulation to be uncovered in the future.

Regulators of mRNA processing

mRNA transcripts often have a complicated post-transcriptional existence⁵⁸. Immediately in the wake of transcription, nascent pre-mRNAs are spliced and processed into one of potentially many isoforms. Importantly, alternative splicing and editing contribute to increasing gene isoform diversity.

In some cases, lncRNA genes that have an anti-sense orientation to known protein-coding genes, also known as natural antisense transcripts (NATs), can influence how an mRNA arising from the sense strand is processed. For example, NATs influence splicing patterns of mRNAs at the neuroblastoma *MYC*, *c-ErbAalpha* (also known as Thra) and *ZEB2* (zinc-finger E-box binding homeobox 2) loci in mammalian cells^{59–62} (FIG. 3a; TABLE 1). In the case of neuroblastoma *MYC* and *c-ErbAalpha*, the NAT and pre-mRNA were suggested to form RNA-RNA duplexes, which then inhibit splicing^{61,62}. At the *ZEB2* locus, NAT expression inhibits splicing of an internal ribosome entry site (IRES)-containing intron. Translation of *ZEB2* relies on this IRES, and therefore expression of the NAT indirectly facilitates expression of ZEB2 protein⁵⁹. The mechanism by which NATs influence splicing is unclear, but it has been postulated to involve splice-site masking and a subsequent block in spliceosome recruitment⁶³.

The *MALAT1* lncRNA also affects splicing, but through a more indirect mechanism. This lncRNA, which is retained in the nucleus and associates with interchromatin granules, has been implicated in alternative splicing through the modulation of active Ser/Arg splicing factors, named after characteristic Ser- and Arg- rich domains. Ser/Arg proteins are important regulators of alternative splicing, and *MALAT1* interacts with, and influences the nuclear distribution and levels of, phosphorylated Ser/Arg proteins. Importantly, depletion of *MALAT1* changes the alternative splicing patterns of the pre-mRNAs that they target⁶⁴.

In addition to modulating splicing, overlapping antisense lncRNAs have in principle the potential to direct mRNA editing (FIG. 3b). During editing, ADAR (adenosine deaminase acting on RNA) enzymes catalyze adenosine to inosine conversion in double-stranded RNA, and this conversion can influence RNA structure, splicing patterns, coding potential and targeting by microRNAs (miRNAs)⁶⁵. In *Drosophila melanogaster*, editing of *Rnp4F* (RNA-binding protein 4F) mRNA depends on developmentally restricted expression of a long isoform of the partially overlapping *Sas10* transcript (TABLE 1). Although, in this case, an mRNA isoform with an extended 3' untranslated region (UTR) provides the source of an antisense RNA, lncRNAs could act in a similar manner to direct editing⁶⁶. Given that many, if not most, mammalian genomic loci produce multiple RNA transcripts from both strands with at least partial overlap, the potential for double-stranded RNA editing substrates is extensive⁶⁷. With many of these pervasive transcripts anticipated to be lncRNAs, lncRNAs are likely to help diversify the transcriptome and proteome through control of RNA editing.

Modulators of post-transcriptional control

Following processing and nuclear export, mRNAs are subjected to various post-transcriptional regulatory pathways that modulate gene expression levels. For example, the overall level of protein produced from an mRNA depends on translation efficiency, mRNA turnover kinetics and small RNA-mediated translational repression. A growing number of reports implicate lncRNAs in control of these post-transcriptional events.

Translation control

The mouse *Uchl1AS* lncRNA produced from the *Uchl1* (ubiquitin carboxyl-terminal esterase L1) locus was shown to upregulate translation of *Uchl1* mRNA through a repeat

element (FIG. 3c; TABLE 1). In this instance, sense and antisense transcripts are oriented in a 5' head-to-head fashion such that the mature lncRNA contains a 73-nucleotide motif complementary to the 5' end of the *Uchl1* mRNA. This sequence-specific interaction serves to position the effector domain, which is contained in the non-overlapping 3' region of *Uchl1AS* and consists of a SINEB2 repeat element that upregulates protein expression without changing *Uchl1* mRNA levels. Bioinformatic analysis has identified 59 other cDNAs with similar antisense orientations and SINEB2 elements, suggesting that this regulatory mechanism might be used at other loci⁶⁸. lncRNA-mediated translational regulation has also been documented in yeast, in which an antisense *KCSI* lncRNA was suggested to regulate translation of the inositol pyrophosphate synthase *KCSI* mRNA expressed from the same locus. Through an unknown mechanism, which is thought to involve base pairing interactions between the antisense and sense RNAs, expression of *KCSI* antisense RNA results in the production of truncated KCS1 protein⁶⁹ (TABLE 1).

mRNA stability control

lncRNAs have also been implicated in both positive and negative regulation of mRNA stability. For instance, Alu repeat-containing lncRNAs are involved in targeting mRNA transcripts for Staufen-mediated decay (SMD)⁷⁰. SMD is induced by Staufen 1 (STAU1) binding to a double-stranded structure in mRNA 3' UTRs. Through imperfect base pairing interactions with Alu elements in the 3' UTR, Alu repeat-containing lncRNAs create STAU1-binding sites that *trans*-activate SMD and destabilize the target mRNA (FIG. 3d; TABLE 1).

By contrast, *BACE1AS*, an antisense lncRNA that arises from the *BACE1* (beta-site APP-cleaving enzyme 1) locus, increases stability of *BACE1* mRNA⁷¹ (TABLE 1). *BACE1AS* and *BACE1* mRNA form an RNA-RNA duplex, which has been suggested to stabilize the mRNA by abrogation of miRNA-induced repression. More specifically, the antisense transcript and miR-485-5p compete for binding to the same region in the *BACE1* mRNA^{71,72} (FIG. 3e; TABLE 1). *BACE1* mRNA encodes β -secretase, the rate-limiting enzyme in amyloid- β synthesis. Regulation of *BACE1* expression, therefore, has important implications in Alzheimer's disease. Intriguingly, *BACE1AS* levels are increased in the brains of patients with Alzheimer's disease, which perhaps suggests that the regulation of this lncRNA might be relevant in this condition⁷¹.

miRNA sponges

Aside from competing with small RNAs for binding sites on target mRNAs, lncRNAs also can act as decoys to attenuate small RNA regulation, for example through sequestration of proteins or RNA-dependent effectors. The competing endogenous RNA (ceRNA) hypothesis is based on this idea. It postulates that a widespread network of crosstalk exists between coding and non-coding RNAs that manifests through competition for miRNA binding⁷³. Examples of potential ceRNAs include *LINCMD1*, *HULC* (highly upregulated in liver cancer), *PTENP1* (PTEN pseudogene 1), *IPS1* (*INDUCED BY PHOSPHATE STARVATION 1*) and *CDR1as* (CDR1 antisense; also known as *ciRS-7*)⁷⁴⁻⁷⁸ (TABLE 1). Specifically, the muscle-specific lncRNA *LINCMD1* regulates muscle differentiation by binding and sequestering miR-133 and miR-135 (REF. 74). Normally, these miRNAs

negatively regulate expression of the *MAML1* (mastermind-like 1) and *MEF2C* (myocyte enhancer factor 2C) transcription factors, which drive muscle-specific gene expression. So, by sequestering these miRNAs, *LINCMD1* indirectly activates *MAML1* and *MEF2C*⁷⁴. Similarly, the *HULC* lncRNA has been suggested to act as a ‘sponge’ that inhibits miR-372 by sequestering it away from potential mRNA targets⁷⁵. This regulatory principle is shared with pseudogenes, which can also act as miRNA decoys to upregulate expression of their cognate genes. This has been shown, for example, in the case of the pseudogene *PTENP1* (REF. 79).

The *Arabidopsis thaliana* lncRNA *IPS1* also sequesters miR-399 away from its target mRNAs⁷⁶. Whereas most miRNAs in plants have perfect complementarity to their targets, which results in mRNA cleavage, *IPS1* contains an imperfect binding site for miR-399. Thus, miR-399 binding to *IPS1* does not result in its cleavage but instead limits the levels of miR-399 available for other targets. This ability to evade cleavage is an important aspect of *IPS1* regulation, because mutant *IPS1* with perfect complementarity to miR-399 no longer regulates miR-399 (REF. 76).

More recently, another example of lncRNA-based miRNA sponges has been described, but these RNAs are unique in that they have a circular structure^{77,78}. In humans, the highly stable circular RNA (circRNA) *CDR1as* has numerous miR-7-binding sites^{77,78} (FIG. 3f; TABLE 1). Importantly, a similar *CDR1as* genomic locus can be found across eutherian mammals, suggesting that, unlike many other lncRNAs, this RNA might be conserved⁷⁷. Moreover, bioinformatic analyses indicate that there may be thousands of expressed circRNAs across a broad range of multicellular eukaryotes⁷⁸.

lncRNAs can, therefore, modulate gene expression by diverse post-transcriptional regulatory pathways (FIG. 3c–f; TABLE 1). Whereas some lncRNAs seem to influence translation, others operate at the RNA level. As more and more lncRNAs are functionally characterized, we will probably see additional examples of post-transcriptional regulation by lncRNAs.

Regulators of protein activity

In addition to lncRNA-mediated modulation of gene expression events through effects on mRNAs, lncRNAs can also act at the protein level. Indeed, some of the same lncRNAs that affect mRNAs, such as *GAS5*, *EVF2* and *CCND1*, alter the activity of transcription factors (TABLE 2). However, the ability of lncRNAs to bind and modulate protein activity extends beyond factors involved in transcription.

For example, a new class of lncRNAs flanked by small nucleolar RNA (snoRNA) sequences, termed sno-lncRNAs, influence splicing patterns via physical interactions with an alternative splicing regulator in human cell lines⁸⁰. These sno-lncRNAs are derived from introns and are nuclear-enriched. A particularly abundant member of the sno-lncRNA family, generated from the 15q11-q13 chromosomal region, directly associates with the FOX2 alternative splicing factor (FIG. 4a; TABLE 2). Importantly, sno-lncRNA knockdown results in changes in FOX2-regulated splicing, and it has been speculated that the sno-lncRNA might inhibit FOX2 function via a sequestration mechanism⁸⁰. Similarly, the *Caenorhabditis elegans*

lncRNA *mcs-1* has been suggested to influence the processing of small RNAs via Dicer inhibition⁸¹. The *mcs-1* lncRNA forms an extensive double-stranded helix, but is not cleaved by Dicer due to inhibitory secondary structures flanking this helix (FIG. 4b; TABLE 2). It has been suggested that *mcs-1* competitively binds either Dicer or accessory double-stranded RNA-binding proteins to preclude processing of small RNAs from double-stranded RNA precursors⁸¹.

Flaviviruses, such as West Nile virus, also produce a highly structured lncRNA termed subgenomic flavivirus RNA (*sfRNA*), which is resistant to destruction by host nucleases. *sfRNA* is essential for pathogenicity and is thought to stall the host 5' to 3' exonuclease, XRN1, during viral RNA genome degradation⁸². The inhibition of XRN1 induced by *sfRNA* is even strong enough to stabilize host cellular mRNAs⁸³ (TABLE 2). Although this is an example of a viral lncRNA that inhibits a host cellular enzyme, it illustrates that structured lncRNAs have the capacity to inhibit wide-ranging enzymatic activities.

A ultraviolet (UV) light-induced lncRNA, *gadd7*, has also been shown to influence cellular mRNA stability⁸⁴. This lncRNA, however, does so by modulating the activity of the RNA-binding protein TDP43 (TAR DNA-binding protein 43). TDP43 has been implicated in pre-mRNA splicing as well as mRNA transport, translation and stability^{85–88}. It binds 3' UTR elements in a large number of genes, and this binding can result in either the stabilization or destabilization of mRNA targets^{84,86–88}. The association of *gadd7* with TDP43 impairs TDP43 binding to several of its targets (FIG. 4c; TABLE 2). For example, by preventing TDP43 association with cyclin-dependent kinase 6 (*CDK6*) mRNA, *gadd7* alters the role of TDP43 in modulating mRNA stability⁸⁴. Interestingly *gadd7* is not the only lncRNA that TDP43 binds. TDP43 also associates the *MALAT1* and *NEAT1* (also known as *Men ε/β*) lncRNAs⁸⁹. As both *MALAT1* and TDP43 are implicated in control of alternative splicing, it will be interesting to further explore this interaction in future studies.

Scaffolds for higher-order complexes

RNA transcripts associate with proteins to form ribonucleoprotein particles (RNPs). Compared with other RNAs such as snRNAs and rRNAs, we know very little about the composition of RNPs formed by lncRNAs. Some specific lncRNA-protein interactions have been characterized, but the lncRNA interaction network in cells is likely to be more complicated than single lncRNAs interacting with single proteins. Indeed there are indications that lncRNAs can act as scaffolds to organize higher-order complexes.

Some of the lncRNAs involved in chromatin-dependent events (such as *HOTAIR*, *KCNQ1OT1* and *ANRIL*) have been suggested to act as scaffolds that coordinate the activities of histone-modifying complexes^{15,25} (FIG. 2; TABLE 2). There are also notable examples of classic ncRNAs such as the RNA component of telomerase (*TERC*) and signal recognition particle (*SRP*) RNA that can act as scaffolds at telomeres and on translating ribosomes during protein targeting to the endoplasmic reticulum (ER), respectively^{90,91} (TABLE 2). Although the *SRP* and *TERC* ncRNAs are not generally considered to be lncRNAs, they demonstrate that RNA is particularly adept as a scaffold and that many lncRNAs could function as scaffolds in diverse contexts.

The telomerase RNP complex is responsible for adding telomeric repeats to chromosomal ends and thereby maintains their length in replicating cells. The RNA component of telomerase is not only responsible for templating the addition of telomeric repeats but also provides a scaffold that organizes telomeric regulatory proteins⁹⁰ (FIG. 4d; TABLE 2). Interestingly, other lncRNAs generated from telomeric repeats, termed *TERRA*, have a distinct role in telomere biology. Rather than extending telomere ends, these lncRNAs promote telomere shortening via exonuclease 1-dependent resection of chromosome ends⁹². lncRNA-mediated events thus serve critical functions in telomere homeostasis.

The SRP is a highly conserved RNP complex, consisting of the *SRPRNA* transcript and six proteins, which directs proteins to the ER. SRP co-translationally binds the signal sequence in nascent peptides, stalls translational elongation and then targets the ribosome-nascent chain complex to the SRP receptor on the ER. Whereas specific protein domains in SRP mediate peptide recognition and arrest of translational elongation, *SRPRNA* provides a scaffold to organize and coordinate distally occurring events at the sites of peptide exit and elongation factor binding on the ribosome⁹¹ (TABLE 2).

In addition to serving as scaffolds for specific multi-protein complexes, lncRNAs have been implicated in nuclear organization through the scaffolding of sub-nuclear domains⁹³. Indeed, RNA, both coding and non-coding, has been implicated in the nucleation of histone locus bodies, interchromatin granules, paraspeckles and nuclear stress bodies⁹⁴. Perhaps the best-studied lncRNA of this type is *NEATI*, which is important for the *de novo* assembly of paraspeckles (subnuclear domains that may mediate retention of hyperedited mRNAs in the nucleus)^{95,96}. Interestingly, the nascent lncRNA is important for this because ongoing *NEATI* lncRNA transcription is required for paraspeckle maintenance⁹⁷.

It is enticing to speculate that other uncharacterized lncRNAs may serve as scaffolds to organize and hold together other higher-order complexes. Imagine what might have been missed through the routine treatment of protein preparations with nuclease to remove RNA contaminants before purification and identification of interacting partners. Perhaps lncRNAs could even hold together enzymes involved in fundamental metabolic processes such as glycolysis or the Krebs cycle. Indeed, the orchestration of electron transport factors on the inner lumen of the mitochondria illustrates that spatial arrangements of enzymes can partly facilitate the catalysis of reactions by overcoming the limits imposed by diffusion. Similarly to cell membranes, lncRNA might also help facilitate this purpose by bringing enzymes closer together. Perhaps this is not such a far stretch, as metabolic enzymes such as aconitase and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) are known to have RNA-binding activity^{98–100}.

Signalling molecules

RNA can be transferred between cells in small vesicles known as exosomes^{101–103}. Not to be confused with the molecular machine with the same name that mediates RNA degradation, these exosomes are membrane-bound vesicles of endosomal origin that are released from various cell types in mammals. Upon fusion with another cell, both their RNA and protein cargo can be transferred¹⁰². The RNAs that have been found in exosomes,

termed exosomal shuttle RNAs (exRNAs), do not simply reflect the RNA composition of the cell of origin, suggesting that there may be selective loading of RNAs into exosomes¹⁰⁴. Because transmitted RNAs can function in the recipient cell, it has been suggested that exRNAs might be used as a signal to change gene expression patterns in the recipient cell^{101–104}. Although exosomes contain large amounts of exRNAs, so far miRNAs and mRNAs have been a primary focus of study^{103,105,106} (FIG. 4e). However, a recent report characterizing the full complement of human plasma-derived exRNA indicates that lncRNAs are indeed present in exosomes¹⁰⁶. The presence of lncRNAs certainly raises the exciting possibility that they might provide signals that impinge on various gene expression events.

Vehicles for increasing genetic diversity

Genetic diversity is crucial for the survival of a species and, within individuals, genetic innovation is of paramount importance to the adaptive immune system. Diversity in developing lymphocytes is achieved through genomic rearrangements in the form of class switch recombination (CSR; also known as isotype switching) and V(D)J recombination events. Interestingly, non-coding transcription has been implicated in both forms of recombination^{107,108}. Through CSR, the constant regions of antibodies are exchanged. As such, this process increases the range of effectors that a particular antibody can interface with and, therefore, increases its versatility^{109,110}. During CSR, the non-coding switch region (S region) is transcribed, and the lncRNAs generated from this S region are likely to be important guides in dictating the locations of recombination. The nascent lncRNA forms an RNA-DNA hybrid or R-loop structure, which displaces one strand of DNA and this, in turn, is thought to facilitate targeting of activation-induced deaminase (AID), the enzyme that initiates CSR^{107,111,112} (FIG. 4f). Transcription through non-coding regions also has a role in V(D)J recombination, the genomic rearrangement that generates diversity in antigen receptor-binding pockets in antibodies^{113,114}. During V(D)J recombination, chromatin accessibility has been suggested to affect recombinase targeting¹¹⁴, and production of non-coding transcripts from the mouse *Tcra* (T cell receptor alpha chain) locus can trigger changes in chromatin structure that then influence recombination¹⁰⁸.

It is tempting to postulate that non-coding transcription might also increase genetic diversity outside the immune system. During meiosis, sites of recombination are not distributed randomly but tend to occur in discrete locations¹¹⁵. Intriguingly, in fission yeast these hot spots correlate with lncRNA-expressing loci¹¹⁶. How exactly lncRNAs contribute to recombination-site selection is currently unclear, but one possibility is that this could involve similar mechanisms to those used during recombination in lymphocytes.

Conclusions and perspectives

Amidst the exciting discoveries being made during this time of genome exploration, RNA is taking centre stage. The burgeoning lncRNA field has a strong part in this, and lncRNAs have now been demonstrated to regulate all aspects of gene expression, including transcription (FIG. 2), processing and post-transcriptional control pathways (FIG. 3). Likewise, lncRNAs have also been shown to regulate protein function and organize multiprotein complex assembly. Now with hints that lncRNAs might participate in cell-cell

communication and recombination, the possible reach of lncRNA functions seems endless (FIG. 4). With most biologists trained to dissect function based on a protein-centric view of the cell, the task of functionally characterizing this new RNA world seems daunting. It is important, therefore, as we move forward, to utilize and develop more functional characterization methods that play to the strengths of RNA. Indeed technical advances are already underway that have the promise of greatly improving the *in vivo* functional characterization of lncRNAs. For instance, techniques to probe RNA chemical structure have often been limited to *in vitro* studies, but recently developed chemical probes that can be used in living cells have the promise of greatly improving our ability to determine *in vivo* RNA structures¹¹⁷. Additionally, the application of high-throughput microfluidics-based screening technologies towards the functional analysis of pre-programmed RNA libraries has the potential to streamline the process of discovering functional motifs within lncRNAs¹¹⁸. Last, recently developed RNA aptamers such as Spinach have adapted GFP tagging for RNA transcripts to allow RNA fusions to be imaged in living cells¹¹⁹.

Much like the multifunctional nature of a Swiss army knife, RNA has the biochemical diversity to function in diverse contexts. It may, however, take some time to determine in which contexts the cell uses some of the more exotic RNA tools. With eyes open to new possibilities, undoubtedly we will be surprised by what we find.

Acknowledgments

The authors are most grateful to T. Nilsen and K. Baker for insights and suggestions. The authors regret that not all contributions of their colleagues could be discussed due to space constraints. Work in the authors' laboratory is funded by the National Institute of General Medical Sciences (NIGMS) (GM080465).

Glossary

Chromatin	Condensed DNA structure that is associated with histone proteins and other DNA-binding proteins
Transcription activator-like effectors	TALEs). Naturally found in some bacteria, TALEs are proteins that bind DNA through repeat domains, and their code for sequence specificity has been elucidated allowing sequence specific TALEs to be engineered
PUF proteins	A family of sequence-specific RNA-binding proteins, which bind 3' untranslated regions within mRNAs to repress target mRNA translation
Pseudogenes	Dysfunctional relatives of normal genes thought to arise from duplication or retrotransposition
Chromatin-modifying complexes	Protein complexes that catalyse the covalent chemical modification of chromatin

Enhancers	Short regions of DNA that enhance the expression of genes at varying distances. Effects can be mediated by transcription factor binding to these sites
Alu SINE elements	Highly abundant retrotransposons of the short interspersed nuclear elements (SINE) family
Nuclear subdomains	Non-membrane bound subcompartments of eukaryotic nuclei where factors with similar functions colocalize
<i>GAL</i> locus	An inducible locus in yeast comprising the <i>GAL1</i> and <i>GAL10</i> genes, which are required for galactose metabolism
Alternative splicing	An mRNA processing step whereby exons can be alternatively used to generate different isoforms of the same gene
Internal ribosome entry sites	IRESs). Nucleotide sequence that allows cap-independent translation initiation within the middle of an mRNA transcript
Spliceosome	The macromolecular machinery (composed of both RNA and protein) responsible for pre-mRNA splicing
miRNAs	(miRNAs). A class of short (~ 23 nucleotides) endogenous non-coding RNAs that control gene expression post-transcriptionally through either translational repression or mRNA degradation
Competing endogenous RNA	(ceRNA). RNA transcripts (both coding and non-coding), which share microRNA-targeting sites and thus regulate each other via direct competition for microRNA binding
Circular RNA	(circRNA). As opposed to conventional linear RNA transcripts, the 5' and 3' ends of circular RNAs are covalently linked together.
Small nucleolar RNA	(snoRNA). A class of small RNA molecules that guide the chemical modification of other RNA transcripts
sno-lncRNAs	(small nucleolar long non-coding RNAs). Class of intron-derived long non-coding RNA flanked by snoRNA ends
Dicer	An RNase III family endoribonuclease responsible for the processing of pre-miRNAs into short double-stranded RNAs to be loaded into the RNA-induced silencing (RISC) complex
Adaptive immune system	A system of specialized cells that create immunological memory via specific antibodies after an initial response to a pathogen

References

1. The ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human genome. *Nature*. 2012; 489:57–74. [PubMed: 22955616] Provides an overview of a series of papers release as part of the ENCODE project in which landmarks of biochemical function (regions of

transcription, transcription factor association and histone modifications, among others) were attributed to 80% of the genome

2. Carninci P, et al. The transcriptional landscape of the mammalian genome. *Science*. 2005; 309:1559–1563. [PubMed: 16141072]
3. Koch E, Jourquin E, Ferrier P, Andrau J-C. Genome-wide RNA polymerase II: not genes only! *Trends Biochem. Sci.* 2008; 33:265–273. [PubMed: 18467100]
4. Djebali S, et al. Landscape of transcription in human cells. *Nature*. 2012; 489:101–108. [PubMed: 22955620]
5. Taft RJ, Pheasant M, Mattick JS. The relationship between non-protein-coding DNA and eukaryotic complexity. *Bioessays*. 2007; 29:288–299. [PubMed: 17295292]
6. Berretta J, Morillon A. Pervasive transcription constitutes a new level of eukaryotic genome regulation. *EMBO Rep.* 2009; 10:973–982. [PubMed: 19680288]
7. Mercer TR, Dinger ME, Mattick JS. Long non-coding RNAs: insights into functions. *Nature Rev. Genet.* 2009; 10:155–159. [PubMed: 19188922]
8. Rinn JL, Chang HY. Genome regulation by long noncoding RNAs. *Annu. Rev. Biochem.* 2012; 81:145–166. [PubMed: 22663078]
9. Mercer TR, Mattick JS. Structure and function of long noncoding RNAs in epigenetic regulation. *Nature Struct. Mol. Biol.* 2013; 20:300–307. [PubMed: 23463315]
10. Ponting CP, Belgard TG. Transcribed dark matter: meaning or myth? *Hum. Mol. Genet.* 2010; 19:R162–R168. [PubMed: 20798109]
11. Ponting CP, Oliver PL, Reik W. Evolution and functions of long noncoding RNAs. *Cell*. 2009; 136:629–641. [PubMed: 19239885]
12. Wilusz JE, Sunwoo H, Spector DL. Long noncoding RNAs: functional surprises from the RNA world. *Genes Dev.* 2009; 23:1494–1504. [PubMed: 19571179]
13. Nagano T, Fraser P. No-nonsense functions for long noncoding RNAs. *Cell*. 2011; 145:178–181. [PubMed: 21496640]
14. Lee J. T Epigenetic regulation by long noncoding RNAs. *Science*. 2012; 338:1435–1439. [PubMed: 23239728]
15. Kugel JF, Goodrich JA. Non-coding RNAs: key regulators of mammalian transcription. *Trends Biochem. Sci.* 2012; 37:144–151. [PubMed: 22300815]
16. Robertson MP, Joyce G. F The origins of the RNA world. *Cold Spring Harb. Perspect. Biol.* 2012; 4 pii:a003608.
17. Filipovska A, Rackham O. Modular recognition of nucleic acids by PUF, TALE and PPR proteins. *Mol. Biosyst.* 2012; 8:699–708. [PubMed: 22234420]
18. Stoltenburg R, Reinemann C, Strehlitz B. SELEX — a (r)evolutionary method to generate high-affinity nucleic acid ligands. *Biomol. Eng.* 2007; 24:381–403. [PubMed: 17627883]
19. Serganov A, Patel DJ. Molecular recognition and function of riboswitches. *Curr Opin. Struct. Biol.* 2012; 22:279–286. [PubMed: 22579413]
20. Pei B, et al. The GENCODE pseudogene resource. *Genome Biol.* 2012; 13:R51. [PubMed: 22951037]
21. Stoye JP. Studies of endogenous retroviruses reveal a continuing evolutionary saga. *Nature Rev. Microbiol.* 2012; 10:395–406. [PubMed: 22565131]
22. Khalil AM, et al. Many human large intergenic noncoding RNAs associate with chromatin-modifying complexes and affect gene expression. *Proc. Natl Acad. Sci. USA.* 2009; 106:11667–11672. [PubMed: 19571010]
23. Zhao J, et al. Genome-wide identification of polycomb-associated RNAs by RIP-seq. *Mol. Cell.* 2010; 40:939–953. [PubMed: 21172659]
24. Tsai M-C, et al. Long noncoding RNA as modular scaffold of histone modification complexes. *Science*. 2010; 329:689–693. [PubMed: 20616235] Illustrates an elegant example of a dominant theme in the lncRNA field whereby lncRNAs physically associate with histone-modifying

complexes to regulate chromatin states. Importantly, shows that *HOTAIR* can also act as a scaffold to organize the concerted actions of two enzymatic activities

25. Spitale RC, Tsai M-C, Chang HY. RNA templating the epigenome: long noncoding RNAs as molecular scaffolds. *Epigenetics*. 2011; 6:539–543. [PubMed: 21393997]
26. Yap KL, et al. Molecular interplay of the noncoding RNA *ANRIL* and methylated histone H3 lysine 27 by polycomb CBX7 in transcriptional silencing of *INK4a*. *Mol. Cell*. 2010; 38:662–674. [PubMed: 20541999]
27. Pandey RR, et al. *Kcnq1ot1* antisense noncoding RNA mediates lineage-specific transcriptional silencing through chromatin-level regulation. *Mol. Cell*. 2008; 32:232–246. [PubMed: 18951091]
28. Kotake Y, et al. Long non-coding RNA *ANRIL* is required for the PRC2 recruitment to and silencing of *p15^{INK4B}* tumor suppressor gene. *Oncogene*. 2011; 30:1956–1962. [PubMed: 21151178]
29. Kornienko AE, Guenzl PM, Barlow DP, Pauler FM. Gene regulation by the act of long non-coding RNA transcription. *BMC Biol*. 2013; 11:59. [PubMed: 23721193]
30. Latos PA, et al. *Airn* transcriptional overlap, but not its lncRNA products, induces imprinted *Igf2r* silencing. *Science*. 2012; 338:1469–1472. [PubMed: 23239737] Provides a particularly compelling example of a mammalian lncRNA *AIRN* which represses target expression by transcriptional interference, as transcriptional overlap of the lncRNA with the target promoter rather than the lncRNA transcript itself is sufficient to interfere with Pol II recruitment
31. Hongay CF, Grisafi PL, Galitski T, Fink GR. Antisense transcription controls cell fate in *Saccharomyces cerevisiae*. *Cell*. 2006; 127:735–745. [PubMed: 17110333]
32. van Werven FJ, et al. Transcription of two long noncoding RNAs mediates mating-type control of gametogenesis in budding yeast. *Cell*. 2012; 150:1170–1181. [PubMed: 22959267]
33. Camblong J, Iglesias N, Fickentscher C, Dieppo G, Stutz F. Antisense RNA stabilization induces transcriptional gene silencing via histone deacetylation in *S. cerevisiae*. *Cell*. 2007; 131:706–717. [PubMed: 18022365]
34. Carrozza MJ, et al. Histone H3 methylation by Set2 directs deacetylation of coding regions by Rpd3S to suppress spurious intragenic transcription. *Cell*. 2005; 123:581–592. [PubMed: 16286007]
35. Houseley J, Rubbi L, Grunstein M, Tollervey D, Vogelauer M. A ncRNA modulates histone modification and mRNA induction in the yeast GAL gene cluster. *Mol. Cell*. 2008; 32:685–695. [PubMed: 19061643]
36. Schmitt S, Prestel M, Paro R. Intergenic transcription through a polycomb group response element counteracts silencing. *Genes Dev*. 2005; 19:697–708. [PubMed: 15741315]
37. Bumgarner SL, et al. Single-cell analysis reveals that noncoding RNAs contribute to clonal heterogeneity by modulating transcription factor recruitment. *Mol. Cell*. 2012; 45:470–482. [PubMed: 22264825]
38. Hainer SJ, Pruneski JA, Mitchell RD, Monteverde RM, Martens JA. Intergenic transcription causes repression by directing nucleosome assembly. *Genes Dev*. 2011; 25:29–40. [PubMed: 21156811] Demonstrates a novel mode of *SER3* gene repression by the yeast lncRNA *SRG1* that involves the direction of nucleosome occupancy at the *SER3* promoter
39. Hirota K, et al. Stepwise chromatin remodelling by a cascade of transcription initiation of non-coding RNAs. *Nature*. 2008; 456:130–134. [PubMed: 18820678]
40. Lefevre P, Witham J, Lacroix CE, Cockerill PN, Bonifer C. The LPS-induced transcriptional upregulation of the chicken lysozyme locus involves CTCF eviction and noncoding RNA transcription. *Mol. Cell*. 2008; 32:129–139. [PubMed: 18851839]
41. Bumgarner SL, Dowell RD, Grisafi P, Gifford DK, Fink GR. Toggle involving *cis*-interfering noncoding RNAs controls variegated gene expression in yeast. *Proc. Natl Acad. Sci. USA*. 2009;

- 106:18321–18326. [PubMed: 19805129] Provides the first description of the pair of *cis* interfering lncRNAs at the *FL011* locus in yeast, where a regulatory circuit toggles between two states of expression depending on the identity of the lncRNA expressed
42. Flynn RA, Chang HY. Active chromatin and noncoding RNAs: an intimate relationship. *Curr Opin. Genet. Dev.* 2012; 22:172–178. [PubMed: 22154525]
43. Ørom UA, et al. Long noncoding RNAs with enhancer-like function in human cells. *Cell.* 2010; 143:46–58. [PubMed: 20887892] Describes a novel class of lncRNAs that, instead of repressing, activates target genes to function as RNA-dependent enhancers of gene expression
44. Lai F, et al. Activating RNAs associate with Mediator to enhance chromatin architecture and transcription. *Nature.* 2013; 494:497–501. [PubMed: 23417068]
45. Kim T-K, et al. Widespread transcription at neuronal activity-regulated enhancers. *Nature.* 2010; 465:182–187. [PubMed: 20393465]
46. Wang D, et al. Reprogramming transcription by distinct classes of enhancers functionally defined by eRNA. *Nature.* 2011; 474:390–394. [PubMed: 21572438]
47. Mariner PD, et al. Human Alu RNA is a modular transacting repressor of mRNA transcription during heat shock. *Mol. Cell.* 2008; 29:499–509. [PubMed: 18313387]
48. Martianov I, Ramadass A, Serra Barros A, Chow N, Akoulitchev A. Repression of the human dihydrofolate reductase gene by a non-coding interfering transcript. *Nature.* 2007; 445:666–670. [PubMed: 17237763]
49. Shamovsky I, Ivannikov M, Kandel ES, Gershon D, Nudler E. RNA-mediated response to heat shock in mammalian cells. *Nature.* 2006; 440:556–560. [PubMed: 16554823]
50. Kino I, Hurt DE, Ichijo T, Nader N, Chrousos GP. Noncoding RNA gas5 is a growth arrest- and starvation-associated repressor of the glucocorticoid receptor. *Sci Signal.* 2010; 3 ra8.
51. Feng J, et al. The *Evf-2* noncoding RNA is transcribed from the *Dlx-5/6* ultraconserved region and functions as a *Dlx-2* transcriptional coactivator. *Genes Dev.* 2006; 20:1470–1484. [PubMed: 16705037]
52. Bond AM, et al. Balanced gene regulation by an embryonic brain ncRNA is critical for adult hippocampal GABA circuitry. *Nature Neurosci.* 2009; 12:1020–1027. [PubMed: 19620975]
53. Wang X, et al. Induced ncRNAs allosterically modify RNA-binding proteins in *cis* to inhibit transcription. *Nature.* 2008; 454:126–130. [PubMed: 18509338]
54. Willingham AT, et al. A strategy for probing the function of noncoding RNAs finds a repressor of NFAT. *Science.* 2005; 309:1570–1573. [PubMed: 16141075] The first study to functionally screen 512 evolutionarily conserved putative lncRNAs. Identified the *NRON* lncRNA as a repressor of NFAT nuclear trafficking
55. Yang L, et al. ncRNA- and Pc2 methylation-dependent gene relocation between nuclear structures mediates gene activation programs. *Cell.* 2011; 147:773–788. [PubMed: 22078878] Describes a particularly interesting paradigm whereby the post-translational modification status of a protein effector constitutes a switch in lncRNA-binding specificity and consequently determines the nuclear subdomain localization of target genes
56. Geisler S, Lojek L, Khalil AM, Baker KE, Coller J. Decapping of long noncoding RNAs regulates inducible genes. *Mol. Cell.* 2012; 45:279–291. [PubMed: 22226051]
57. Mattick JS. Non-coding RNAs: the architects of eukaryotic complexity. *EMBO Rep.* 2001; 2:986–991. [PubMed: 11713189]
58. Moore MJ. From birth to death: the complex lives of eukaryotic mRNAs. *Science.* 2005; 309:1514–1518. [PubMed: 16141059]
- 59.

- Beltran M, et al. A natural antisense transcript regulates *Zeb2/Sip1* gene expression during *Snail1* - induced epithelial-mesenchymal transition. *Genes Dev.* 2008; 22:756–769. [PubMed: 18347095] Provides an example of a NAT lncRNA that regulates splicing of the sense encoded mRNA, but with a twist in that the NAT increases protein levels of its target by preventing the splicing of a 5' UTR IRES-containing intron
60. Hastings ML, Milcarek C, Martincic K, Peterson ML, Munroe SH. Expression of the thyroid hormone receptor gene, *erbA α* , in B lymphocytes: alternative mRNA processing is independent of differentiation but correlates with antisense RNA levels. *Nucleic Acids Res.* 1997; 25:4296–4300. [PubMed: 9336460]
61. Krystal GW, Armstrong BC, Battey J. F N-myc mRNA forms an RNA-RNA duplex with endogenous antisense transcripts. *Mol. Cell. Biol.* 1990; 10:4180–4191. [PubMed: 1695323]
62. Munroe SH, Lazar MA. Inhibition of c-erbA mRNA splicing by a naturally occurring antisense RNA. *J. Biol. Chem.* 1991; 266:22083–22086. [PubMed: 1657988]
63. Faghihi MA, Wahlestedt C. Regulatory roles of natural antisense transcripts. *Nature Rev. Mol. Cell. Biol.* 2009; 10:637–643. [PubMed: 19638999]
64. Tripathi V, et al. The nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by modulating SR splicing factor phosphorylation. *Mol. Cell.* 2010; 39:925–938. [PubMed: 20797886]
65. Hundley HA, Bass BL. ADAR editing in double-stranded UTRs and other noncoding RNA sequences. *Trends Biochem. Sci.* 2010; 35:377–383. [PubMed: 20382028]
66. Peters NT, Rohrbach JA, Zalewski BA, Byrkett CM, Vaughn JC. RNA editing and regulation of *Drosophila 4f-rnp* expression by *sas-10* antisense readthrough mRNA transcripts. *RNA.* 2003; 9:698–710. [PubMed: 12756328]
67. Katayama S, et al. Antisense transcription in the mammalian transcriptome. *Science.* 2005; 309:1564–1566. [PubMed: 16141073]
68. Carrieri C, et al. Long non-coding antisense RNA controls *Uchl1* translation through an embedded SINEB2 repeat. *Nature.* 2012; 491:454–457. [PubMed: 23064229] Discovers and characterizes the first lncRNA of a potentially new class of partially antisense SINE2B repeat-containing lncRNAs which upregulates translation of targets
69. Nishizawa M, et al. Nutrient-regulated antisense and intragenic RNAs modulate a signal transduction pathway in yeast. *PLoS Biol.* 2008; 6:2817–2830. [PubMed: 19108609]
70. Gong C, Maquat LE. lncRNAs transactivate STAU1 -mediated mRNA decay by duplexing with 3' UTRs via Alu elements. *Nature.* 2011; 470:284–288. [PubMed: 21307942] Provides the first evidence that Alu element-containing lncRNAs can transactivate SMD by imperfectly base pairing with 3' UTR Alu elements in target mRNAs
71. Faghihi MA, et al. Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feedforward regulation of β -secretase. *Nature Med.* 2008; 14:723–730. [PubMed: 18587408]
72. Faghihi MA, et al. Evidence for natural antisense transcript-mediated inhibition of microRNA function. *Genome Biol.* 2010; 11:R56. [PubMed: 20507594]
73. Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP. A ceRNA hypothesis: the Rosetta Stone of a hidden RNA language? *Cell.* 2011; 146:353–358. [PubMed: 21802130]
74. Cesana M, et al. A long noncoding RNA controls muscle differentiation by functioning as a competing endogenous RNA. *Cell.* 2011; 147:358–369. [PubMed: 22000014]
75. Wang J, et al. CREB up-regulates long non-coding RNA, HULC expression through interaction with microRNA-372 in liver cancer. *Nucleic Acids Res.* 2010; 38:5366–5383. [PubMed: 20423907]
76. Franco-Zorrilla JM, et al. Target mimicry provides a new mechanism for regulation of microRNA activity. *Nature Genet.* 2007; 39:1033–1037. [PubMed: 17643101]
- 77.

- Hansen TB, et al. Natural RNA circles function as efficient microRNA sponges. *Nature*. 2013 <http://dx.doi.org/10.1038/nature11993> References 77 and 78 provide powerful evidence that circRNAs, covalently linked by the head-to-tail splicing of exons, can function as miRNA sponges to suppress miRNA activity
78. Memczak, S., et al. Circular RNAs are a large class of animal RNAs with regulatory potency. *Nature*. 2013. <http://dx.doi.org/10.1038/nature11928>
79. Poliseno L, et al. A coding-independent function of gene and pseudogene mRNAs regulates tumour biology. *Nature*. 2010; 465:1033–1038. [PubMed: 20577206]
- 80.
- Yin Q-F, et al. Long noncoding RNAs with snoRNA ends. *Mol. Cell*. 2012; 48:219–230. [PubMed: 22959273] Describes the discovery of a new class of intron-derived lncRNAs flanked by snoRNAs and shows that one in particular associates with splicing regulators to alter splicing patterns
81. Hellwig S, Bass BL. A starvation-induced noncoding RNA modulates expression of Dicer-regulated genes. *Proc. Natl Acad. Sci. USA*. 2008; 105:12897–12902. [PubMed: 18723671]
82. Pijlman GP, et al. A highly structured, nuclease-resistant, noncoding RNA produced by flaviviruses is required for pathogenicity. *Cell Host Microbe*. 2008; 4:579–591. [PubMed: 19064258]
83. Moon SL, et al. A noncoding RNA produced by arthropod-borne flaviviruses inhibits the cellular exoribonuclease XRN1 and alters host mRNA stability. *RNA*. 2012; 18:2029–2040. [PubMed: 23006624]
84. Liu X, Li D, Zhang W, Guo M, Zhan Q. Long non-coding RNA *gaddl* interacts with TDP-43 and regulates *Cdk6* mRNA decay. *EMBO J*. 2012; 31:4415–4427. [PubMed: 23103768]
85. Buratti E, Baralle FE. The multiple roles of TDP-43 in pre-mRNA processing and gene expression regulation. *RNA Biol*. 2010; 7:420–429. [PubMed: 20639693]
86. Colombrita C, et al. TDP-43 and FUS RNA-binding proteins bind distinct sets of cytoplasmic messenger RNAs and differently regulate their post-transcriptional fate in motoneuron-like cells. *J. Biol. Chem*. 2012; 287:15635–15647. [PubMed: 22427648]
87. Strong MJ, et al. TDP43 is a human low molecular weight neurofilament (hNFL) mRNA-binding protein. *Mol. Cell. Neurosci*. 2007; 35:320–327. [PubMed: 17481916]
88. Volkening K, Leystra-Lantz C, Yang W, Jaffee H, Strong MJ. Tar DNA binding protein of 43 kDa (TDP-43), 14-3-3 proteins and copper/zinc superoxide dismutase (SOD1) interact to modulate NFL mRNA stability. Implications for altered RNA processing in amyotrophic lateral sclerosis (ALS). *Brain Res*. 2009; 1305:168–182. [PubMed: 19815002]
89. Tollervy JR, et al. Characterizing the RNA targets and position-dependent splicing regulation by TDP-43. *Nature Neurosci*. 2011; 14:452–458. [PubMed: 21358640]
90. Zappulla DC, Cech TR. Yeast telomerase RNA: a flexible scaffold for protein subunits. *Proc. Natl Acad. Sci. USA*. 2004; 101:10024–10029. [PubMed: 15226497]
91. Halic M, et al. Structure of the signal recognition particle interacting with the elongation-arrested ribosome. *Nature*. 2004; 427:808–814. [PubMed: 14985753]
92. Pfeiffer V, Lingner J. TERRA promotes telomere shortening through exonuclease 1-mediated resection of chromosome ends. *PLoS Genet*. 2012; 8:e1002747. [PubMed: 22719262]
93. Carmo-Fonseca M, Rino J. RNA seeds nuclear bodies. *Nature Cell Biol*. 2011; 13:110–112. [PubMed: 21283118]
94. Shevtsov SP, Dundr M. Nucleation of nuclear bodies by RNA. *Nature Cell Biol*. 2011; 13:167–173. [PubMed: 21240286]
95. Sunwoo H, et al. MEN ϵ/β nuclear-retained non-coding RNAs are up-regulated upon muscle differentiation and are essential components of paraspeckles. *Genome Res*. 2009; 19:347–359. [PubMed: 19106332]
96. Sasaki YTF, Ideue I, Sano M, Mituyama T, Hirose T. MEN ϵ/β noncoding RNAs are essential for structural integrity of nuclear paraspeckles. *Proc. Natl Acad. Sci. USA*. 2009; 106:2525–2530. [PubMed: 19188602]
- 97.

- Mao YS, Sunwoo H, Zhang B, Spector DL. Direct visualization of the co-transcriptional assembly of a nuclear body by noncoding RNAs. *Nature Cell Biol.* 2011; 13:95–101. [PubMed: 21170033] Uses a live-cell imaging system to directly visualize paraspeckle protein recruitment and shows that *NEAT1* lncRNA transcription regulates paraspeckle maintenance with the lncRNA potentially acting as an assembly platform
98. Ciesla J. Metabolic enzymes that bind RNA: yet another level of cellular regulatory network? *Acta Biochim. Pol.* 2006; 53:11–32. [PubMed: 16410835]
99. Hentze MW, Argos P. Homology between IRE-BP, a regulatory RNA-binding protein, aconitase, and isopropylmalate isomerase. *Nucleic Acids Res.* 1991; 19:1739–1740. [PubMed: 1903202]
100. Mukhopadhyay R, Jia J, Arif A, Ray PS, Fox PL. The GAIT system: a gatekeeper of inflammatory gene expression. *Trends Biochem. Sci.* 2009; 34:324–331.
101. Lotvall J, Valadi H. Cell to cell signalling via exosomes through esRNA. *Cell Adh. Migr.* 2007; 1:156–158. [PubMed: 19262134]
102. Ramachandran S, Palanisamy V. Horizontal transfer of RNAs: exosomes as mediators of intercellular communication. *Wiley Interdiscip. Rev. RNA.* 2012; 3:286–293. [PubMed: 22012863]
103. Valadi H, et al. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nature Cell Biol.* 2007; 9:654–659. [PubMed: 17486113]
104. Skog J, et al. Glioblastoma microvesicles transport RNA and proteins that promote tumour growth and provide diagnostic biomarkers. *Nature Cell Biol.* 2008; 10:1470–1476. [PubMed: 19011622]
105. Mathivanan S, Fahner CJ, Reid GE, Simpson RJ. ExoCarta 2012: database of exosomal proteins, RNA and lipids. *Nucleic Acids Res.* 2012; 40:D1241–D1244. [PubMed: 21989406]
106. Huang X, et al. Characterization of human plasma-derived exosomal RNAs by deep sequencing. *BMC Genomics.* 2013; 14:319. [PubMed: 23663360] Characterizes, for the first time, the RNA content of exosomes by RNA sequencing and reveals that lncRNAs are indeed present in these membrane-bound vesicles
107. Yu K, Chedin F, Hsieh C-L, Wilson TE, Lieber MR. R-loops at immunoglobulin class switch regions in the chromosomes of stimulated B cells. *Nature Immunol.* 2003; 4:442–451. [PubMed: 12679812]
108. Abarrategui I, Krangel MS. Noncoding transcription controls downstream promoters to regulate T-cell receptor- α recombination. *EMBO J.* 2007; 26:4380–4390. [PubMed: 17882258]
109. Pone EJ, Xu Z, White CA, Zan H, Casali P. B cell TLRs and induction of immunoglobulin class-switch DNA recombination. *Front. Biosci.* 2012; 17:2594–2615.
110. Stavnezer J, Amemiya CT. Evolution of isotype switching. *Semin. Immunol.* 2004; 16:257–275. [PubMed: 15522624]
111. Hackney JA, et al. DNA targets of AID evolutionary link between antibody somatic hypermutation and class switch recombination. *Adv. Immunol.* 2009; 101:163–189. [PubMed: 19231595]
112. Seising E. Ig class switching: targeting the recombinational mechanism. *Curr. Opin. Immunol.* 2006; 18:249–254. [PubMed: 16616473]
113. Abarrategui I, Krangel MS. Regulation of T cell receptor- α gene recombination by transcription. *Nature Immunol.* 2006; 7:1109–1115. [PubMed: 16936730]
114. Cobb RM, Oestreich KJ, Osipovich OA, Oltz EM. Accessibility control of V(D)J recombination. *Adv. Immunol.* 2006; 91:45–109. [PubMed: 16938538]
115. Petes TD. Meiotic recombination hot spots and cold spots. *Nature Rev. Genet.* 2001; 2:360–369. [PubMed: 11331902]
116. Wahls WP, Siegel ER, Davidson MK. Meiotic recombination hotspots of fission yeast are directed to loci that express non-coding RNA. *PLoS ONE.* 2008; 3:e2887. [PubMed: 18682829]
117. Spitale RC, et al. RNA SHAPE analysis in living cells. *Nature Chem. Biol.* 2013; 9:18–20. [PubMed: 23178934]

118. Martin L, et al. Systematic reconstruction of RNA functional motifs with high-throughput microfluidics. *Nature Methods*. 2012; 9:1192–1194. [PubMed: 23142872]
119. Paige JS, Wu KY, Jaffrey SR. RNA mimics of green fluorescent protein. *Science*. 2011; 333:642–646. [PubMed: 21798953]
120. Tsai M-C, Spitale RC, Chang H. Y Long intergenic noncoding RNAs: new links in cancer progression. *Cancer Res*. 2011; 71:3–7. [PubMed: 21199792]
121. Zhao J, Sun BK, Erwin JA, Song JJ, Lee J. I Polycomb proteins targeted by a short repeat RNA to the mouse X chromosome. *Science*. 2008; 322:750–756. [PubMed: 18974356]
122. Rinn JL, et al. Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell*. 2007; 129:1311–1323. [PubMed: 17604720]
123. Wang KC, et al. A long noncoding RNA maintains active chromatin to coordinate homeotic gene expression. *Nature*. 2011; 472:120–124. [PubMed: 21423168]




Figure 1. RNA is a biochemically versatile polymer

a | RNA is particularly well suited for sequence-specific nucleic acid targeting through base pairing interactions over a short region (for example, eight nucleotides). By contrast, proteins require repeat motifs comprising 35–39 amino acids (105–117 base pairs of genomic sequence) to recognize a single RNA base with specificity. Therefore, to recognize eight nucleotides, 280–312 amino acids (840–936 base pairs of genomic sequence) would be required. Compared to the eight base pairs required for an RNA, protein-based nucleic acid recognition requires substantially more genomic sequence¹⁷. **b** | RNA can fold into complex

three-dimensional structures that can specifically bind various ligands, including small molecules and peptides¹⁸. **c** | RNA is suitable for transient expression, because a fully functional RNA can be generated immediately following transcription and processing but can also be rapidly degraded. Together, this allows RNA effectors to be produced in quick pulses. Proteins, however, require additional steps, including mRNA export and translation, to produce a functional peptide. Likewise, both the mRNA and the protein need to be degraded to turn off expression. **d** | RNA is malleable and, therefore, more tolerant of mutations. Although some mutations in protein-coding genes are silent, many are deleterious such as nonsense mutations that generate truncated polypeptides. RNA, however, can tolerate mutations even within the regions responsible for target recognition. **e** | RNA-dependent events can be heritable. For instance, processed pseudogenes were once RNA transcripts that have been genomically integrated. In addition, telomerase uses an RNA template to add telomeric repeats to the ends of chromosomes. ORF, open reading frame; Pol II, RNA polymerase II.




Figure 2. lncRNAs regulate transcription through several mechanisms

a-c | Long non-coding RNAs (lncRNAs) can modulate chromatin through transcription-independent (part **a**) and transcription-dependent mechanisms (parts **b** and **c**). lncRNAs can bind one or more chromatin-modifying complexes and target their activities to specific DNA loci (part **a**). Depending on the nature of the enzymes bound, lncRNA-mediated chromatin modifications can activate or repress gene expression^{22,23,26,27,120}. Chromatin-modifying complexes bound to the RNA polymerase II (Pol II) carboxy-terminal domain (CTD) can modify chromatin during transcription of lncRNAs^{33–35} (part **b**). Transcription of lncRNAs

can also result in chromatin remodelling that can either favour or inhibit the binding of regulatory factors (part **c**). Depending on the nature of the factors that bind during remodelling, gene expression is activated or repressed^{37–40}. **d–g** | lncRNAs can modulate both the general transcription machinery (parts **d** and **e**) as well as specific regulatory factors (parts **f** and **g**). lncRNAs can bind Pol II directly to inhibit transcription⁴⁷ (part **d**). Formation of lncRNA-DNA triplex structures can also inhibit the assembly of the pre-initiation complex⁴⁸ (part **e**). lncRNAs can fold into structures that mimic DNA-binding sites (left) or that generally inhibit or enhance the activity of specific transcription factors (right)^{50–53} (part **f**). lncRNAs can also regulate gene expression by binding specific transport factors to inhibit the nuclear localization of specific transcription factors⁵⁴ (part **g**).




Figure 3. lncRNAs influence mRNA processing and post-transcriptional regulation

a,b | Long non-coding RNAs (lncRNAs) can modulate mRNA processing. Splicing patterns can be influenced by lncRNAs that associate with the pre-mRNA (part **a**). For example, splicing of the first intron of neuroblastoma *MYC* mRNA is prevented by a natural antisense transcript⁶¹. Antisense lncRNAs that associate with an mRNA could direct mRNA editing, perhaps through association of the duplex with ADAR (adenosine deaminase acting on RNA) enzymes that catalyse adenosine to inosine conversion in double-stranded RNA^{63,66} (part **b**). **c-f** | lncRNAs modulate post-transcriptional regulatory events. lncRNAs containing SINEB2 repeat elements can upregulate translation through association with the 5' region of an mRNA⁶⁸ (part **c**). lncRNAs containing Alu repeat elements associate with the Alu elements in the 3' untranslated region (UTR) of an mRNA, and this double-stranded structure can direct Staufen-mediated decay through a pathway that is molecularly similar to nonsense-mediated decay⁷⁰ (part **d**). lncRNAs can mask miRNA-binding sites on a target mRNA to block miRNA-induced silencing through the RNA-induced silencing complex (RISC)⁷² (part **e**). Linear or circular lncRNAs can function as miRNA decoys to sequester miRNAs from their target mRNAs^{74,75} (part **f**).




Figure 4. lncRNAs are involved in various cellular contexts

Long non-coding RNAs (lncRNAs) modulate protein activity by post-translational mechanisms (parts a-c). **a** | Small nucleolar lncRNAs (sno-lncRNAs) generated from the 15q11-q13 locus bind and modulate the activity of the FOX2 alternative splicing factor, and this can inhibit FOX2-mediated splicing⁸⁰. **b** | The highly structured *rncs-1* lncRNA binds Dicer to inhibit the processing of small RNAs⁸¹. **c** | The *gadd7* lncRNA binds and modulates the ability of TDP43 (TAR DNA-binding protein 43) to target and process specific mRNAs⁸⁴. **d** | lncRNAs can act as scaffolds to organize several complexes²⁴. **e** | As the cargo of exosomes that mediate transfer of material between cells, exosomal shuttle RNAs (exRNAs) may act as signalling molecules during cell-cell communication; exosomal cargo includes mRNAs, microRNAs (miRNAs) and lncRNAs¹⁰². **f** | lncRNAs expressed from the switch region of genes encoding antibodies form R-loops to direct class switch recombination via activation-induced deaminase (AID) recruitment¹¹¹.

Table 1

lncRNA-mediated gene expression control

lncRNA	Function	Mechanism	Refs
Regulation of mRNA transcription			
<i>XIST</i>	X inactivation	Chromatin-mediated repression	23,121
<i>HOTAIR</i>	Repression at the <i>HOXD</i> locus	Chromatin-mediated repression	24,122
<i>HOTTIP</i>	Activation at the <i>HOXA</i> locus	Chromatin-mediated activation	123
<i>KCNQ1OT1</i>	Imprinting at the <i>KCNQ1</i> cluster	Chromatin-mediated repression	27
<i>ANRIL</i>	Repression at the <i>INK4b</i> <i>ARF-INK4a</i> locus	Chromatin-mediated repression	26,28
<i>AIRN</i>	Imprinting at the <i>IGF2R</i> cluster	Chromatin-mediated repression, transcription interference	30
<i>ME4</i> antisense	Repression of <i>ME4</i> mRNA	Transcription interference	31
<i>IRT1</i>	Repression of <i>IME1</i> mRNA	Chromatin-mediated repression	32
<i>GAL10</i> lncRNA	Repression of <i>GAL1</i> and <i>CAL10</i> mRNAs	Chromatin-mediated repression	35
<i>PH084</i> antisense	Repression of <i>PH084</i> mRNA	Chromatin-mediated repression	33
<i>ICR1</i>	Repression of <i>FLO11</i> mRNA	Modulation of transcription factor recruitment	37,41
<i>PWR1</i>	Activation of <i>FLO11</i> mRNA	Modulation of transcription factor recruitment	37,41
<i>SRG1</i>	Repression of <i>SER3</i> mRNA	Nucleosome remodelling	38
<i>fbp1</i> ncRNA	Activation of <i>fbp1</i>	Chromatin remodelling	39
<i>UNOCR</i>	Activation of lysozyme mRNA	Nucleosome remodelling	40
Alu repeat-containing RNA	Transcriptional repression during heat shock	Inhibition of Pol II	47
<i>HSR1</i>	Activation of the HSF1 transcription factor	Allosteric activation together with eEF1A	49
Non-coding <i>DHFR</i>	Transcriptional repression of <i>DHFR</i>	Inhibition of pre-initiation complex formation	48
CAS5	Repression of glucocorticoid receptor-mediated transcription	DNA mimicry	50
<i>EVE2</i>	Transcriptional activation of <i>DLX2</i> targets, transcriptional repression of <i>MeCP2</i> targets	Recruitment of <i>DLX2</i> or <i>MeCP2</i>	51,52
<i>CCND1</i> promoter RNA	Repression of <i>CCND1</i> transcription	Allosteric activation of <i>TLS</i>	53
<i>NRON</i>	Repression of NFAT-mediated transcription	Inhibition of transcription factor nucleocytoplasmic shuttling	54
Regulation of mRNA processing			
Neuroblastoma <i>MYC</i> (NAT)	Inhibition of neuroblastoma <i>MYC</i> intron 1 splicing	Unknown mechanism involving the inhibition of splicing via RNA-RNA duplex formation	61
<i>Rev-ErbA</i> alpha	Inhibition of the <i>c-ErbAalpha 2</i> splice isoform	Unknown mechanism involving the inhibition of splicing via RNA-RNA duplex	62

lncRNA	Function	Mechanism	Refs
		formation	
<i>ZEB2</i> (NAT)	Activation of <i>ZEB2</i> translation	Unknown mechanism involving regulated splicing of an IRES-containing intron	59
<i>MALAT1</i>	Ser/Arg splicing factor regulation	Scaffolding of subnuclear domains	64
<i>Sas10</i> mRNA 3'UTR	Repression of <i>Rnp4F</i> mRNA	Unknown mechanism involving RNA editing	66
Modulation of mRNA post-transcriptional regulatory pathways			
Antisense <i>UCHL1</i>	Upregulation of <i>UCHL1</i> protein production	SINE2B element-mediated translational upregulation	68
<i>KCS1</i> antisense	Production of truncated <i>KCS1</i> protein	Unknown mechanism involving base pairing	69
<i>1/2-sbsRNA1</i>	Down-regulation of <i>SERPINE1</i> and <i>FLJ21870</i> mRNAs	Staufen-mediated decay through Alu element base pairing	70
<i>BACE1AS</i>	Up-regulation of <i>BACE1</i>	Stabilization of <i>BACE1</i> mRNA by blocking miRNA-induced repression	71,72
<i>UNCMD1</i>	Control of muscle differentiation through upregulation of <i>MAML1</i> and <i>MEF2C</i> transcription factors	Sequestration of miRNAs	74
<i>HULC</i>	Downregulation of miRNA-mediated repression	Sequestration of miRNAs	75
<i>PTENP1</i> pseudogene	Upregulation of <i>PTEN</i>	Sequestration of miRNAs	79
<i>IPSI</i>	Downregulation of miRNA-mediated repression	Sequestration of miRNAs	76
<i>CDR1as</i>	Downregulation of miRNA-mediated repression	Sequestration of miRNAs	77,78

1/2-sbsRNA1, half-STAU1-binding site RNA1; *AIRN*, antisense of *IGFR2* non-coding RNA; *BACE1AS*, beta-site APP-cleaving enzyme 1 antisense; *CCND1*, cyclin D1; *CDR1as*, CDR1 antisense; *DHFR*, dihydrofolate reductase; *fbp1*, fructose-1,6-bisphosphatase 1; eEF1A, eukaryotic elongation factor 1A; *FLO11*; *GAS5*, growth arrest specific 5; *HOTAIR*, HOX transcript antisense RNA; *HOTTIP*, *HOXA* transcript at the distal tip; *HOX*, homeobox cluster; HSF1, heat shock factor 1; *HSR1*, heat shock RNA1; *HULC*, highly upregulated in liver cancer; *IGF2R*, insulin-like growth factor 2 receptor; *IME*, inducer of meiosis; *IPSI*, INDUCED BY PHOSPHATE STARVATION 1; *IRES*, internal ribosome entry site; *IRT1*, *IME1* regulatory transcript 1; *KCNQ1*, potassium voltage-gated channel, KQT-like subfamily, member 1; *KCNQ1OT1*, *KCNQ1* opposite strand or antisense transcript 1; *LINOCR*, *LPS*-inducible non-coding RNA; lncRNA, long non-coding RNA; *MALAT1*, metastasis associated lung adenocarcinoma transcript 1; *MAML1*, mastermind-like 1; MeCP2, methyl CpG binding-protein 2; *MEF2C*, myocyte enhancer factor 2C; miRNA, microRNA; NAT, natural antisense transcript; ncRNA, non-coding RNA; NFAT, nuclear factor of activated T cells; *NRON*, non-coding repressor of N FAT; Pol II, RNA polymerase II; *PTENP1*, phosphatase and tensin homologue; *Rnp4F*, RNA-binding protein 4F; TLS, translocated in liposarcoma; *UCHL1*, ubiquitin carboxyl-terminal esterase LI; UTR, untranslated region; *XIST*, X inactivation-specific transcript; *ZEB2*, zinc-finger E-box binding homeobox 2.

Table 2

lncRNA-mediated regulation of proteins

lncRNA	Function	Mechanism	Refs
<i>Regulation of protein activity</i>			
CAS5	Repression of glucocorticoid receptor-mediated transcription	DNA mimicry	50
<i>EVE2</i>	Transcriptional activation of DLX2 targets	Activation of DLX2	51,52
<i>CCND1</i> promoter RNA	Repression of <i>CCND1</i> transcription	Allosteric activation of TLS	53
<i>NRON</i>	Repression of NFAT-mediated transcription	Inhibition of transcription factor nucleocytoplasmic shuttling	54
<i>15q11-q13</i> sno-lncRNA	Regulation of alternative splicing	Inhibition of FOX2 function	80
<i>mcs-1</i>	Inhibition of Dicer-mediated repression	Sequestration of Dicer or accessory double-stranded RNA-binding proteins	81
sfRNA	Stabilization of viral and host mRNAs	Inhibition of XRN1-mediated mRNA degradation	82,83
<i>gadd7</i>	Inhibition of TDP43-mediated regulatory events	Sequestration of TDP43	84
<i>Organization of protein complexes</i>			
<i>HOTAIR</i>	Repression at the <i>HOXD</i> locus	Recruitment of PRC2 and LSD1	24
<i>KCNQ1OT1</i>	Imprinting at the <i>KCNQ1</i> cluster	Recruitment of PRC2 and G9A	27
<i>ANRIL</i>	Repression at the <i>INK4b-ARF-INK4a</i> locus	Recruitment of PRC1 and PRC2	26,28
<i>TERC</i>	Addition of telomeric repeats to the ends of chromosomes	Organizational scaffold for telomerase components and template for repeat addition	90
<i>SRPRNA</i>	Directing of proteins to the ER	Organizational scaffold for SRP components	91
<i>NEAT1</i>	Assembly of paraspeckles	Nucleation of subnuclear domains	95–97

CCND1, cyclin D1; ER, endoplasmic reticulum; GAS5, growth arrest specific 5; *HOTAIR*, HOX transcript antisense RNA; HOXD, homeobox D cluster; *KCNQ1*, potassium voltage-gated channel, KQT-like subfamily, member 1; *KCNQ1OT1*, *KCNQ1* opposite strand or antisense transcript 1; LSD1, Lys-specific demethylase 1; NFAT, nuclear factor of activated T cells; *NRON*, non-coding repressor of NFAT; PRC, Polycomb repressive complex; *sfRNA*, subgenomic flavivirus RNA; sno-lncRNA, small nucleolar long non-coding RNA; SRP, signal recognition particle; TDP43, TAR DNA-binding protein 43; *TERC*, telomerase RNA component; TLS, translesion DNA synthesis; XRN1, 5' to 3' exoribonuclease 1.