

The Unstructured N-terminal Region of *Arabidopsis* Group 4 Late Embryogenesis Abundant (LEA) Proteins Is Required for Folding and for Chaperone-like Activity under Water Deficit*

Received for publication, February 6, 2016, and in revised form, March 21, 2016. Published, JBC Papers in Press, March 22, 2016, DOI 10.1074/jbc.M116.720318

Cesar L. Cuevas-Velazquez^{†1}, Gloria Saab-Rincón[§], José Luis Reyes[‡], and Alejandra A. Covarrubias^{‡2}

From the Departamentos de [†]Biología Molecular de Plantas and [§]Ingeniería Celular y Biotecnología, Instituto de Biotecnología, Universidad Nacional Autónoma de México, 62250 Cuernavaca, México

Late embryogenesis abundant (LEA) proteins are a conserved group of proteins widely distributed in the plant kingdom that participate in the tolerance to water deficit of different plant species. *In silico* analyses indicate that most LEA proteins are structurally disordered. The structural plasticity of these proteins opens the question of whether water deficit modulates their conformation and whether these possible changes are related to their function. In this work, we characterized the secondary structure of *Arabidopsis* group 4 LEA proteins. We found that they are disordered in aqueous solution, with high intrinsic potential to fold into α -helix. We demonstrate that complete dehydration is not required for these proteins to sample ordered structures because milder water deficit and macromolecular crowding induce high α -helix levels *in vitro*, suggesting that prevalent conditions under water deficit modulate their conformation. We also show that the N-terminal region, conserved across all group 4 LEA proteins, is necessary and sufficient for conformational transitions and that their protective function is confined to this region, suggesting that folding into α -helix is required for chaperone-like activity under water limitation. We propose that these proteins can exist as different conformers, favoring functional diversity, a moonlighting property arising from their structural dynamics.

Low water availability caused by different environmental conditions such as drought, or low temperatures represents a vulnerable situation for many forms of life, particularly for plants. To contend with and to overcome these adverse environments, numerous complex response mechanisms have been selected in the different species of the plant kingdom. One of the most conserved responses is the accumulation of a group of proteins known as late embryogenesis abundant (LEA)³ pro-

teins (1). LEA proteins have been found in all the orthodox dry seeds (embryos) where they have been searched (1, 2), and they also accumulate in response to water limitation in all vegetative tissues (2, 3). Most LEA proteins show high hydrophilicity, high content of small amino acids, and absence or deficit of hydrophobic residues, properties that are extended to a larger set of proteins called hydrophilins, which have been found in species from the three domains of life and that also accumulate under water deficit (2, 4). The composition of these proteins is also characteristic of a group of proteins known as intrinsically disordered proteins (IDPs) (5, 6). Consistent with the predicted structural disorder for most LEA proteins, structural analyses have confirmed this property for some of them in aqueous solution (7–13). Based on their sequence similarity, LEA proteins have been classified in seven groups or families, each one characterized by the presence of specific sequence motifs (2). In *Arabidopsis thaliana* there are 51 genes encoding LEA proteins from six of the seven families (3). Group 4 LEA (LEA4) proteins are one of the smallest families of LEA proteins in *Arabidopsis* consisting of only three members: AtLEA4-1 (At1g32560), AtLEA4-2 (At2g35300), and AtLEA4-5 (At5g06760) (3, 14). Group 4 LEA proteins are enriched in charged and small amino acid residues, whereas they lack Cys, Phe, and Trp (2, 3, 14). This group is characterized by an N-terminal region ranging from 74 to 78 amino acid residues, containing conserved amino acid sequence motifs. *In silico* analysis predicts that this particular region is able to form an amphipathic α -helix structure. The C-terminal region in this protein family is more variable in sequence and length, and it is predicted to be structurally disordered (2, 14). A phylogenetic analysis of group 4 LEA proteins revealed two subclasses in this family (subgroups 4A and 4B) (14). In *Arabidopsis*, AtLEA4-1 and AtLEA4-2 proteins belong to subgroup 4A, whereas AtLEA4-5 protein fits into subgroup 4B (14).

From the 10 distinctive motifs found in this protein group, the high conservation of motif 2 at the N-terminal region constitutes a signature for this family. The same study also showed that both subgroups emerged from a very early duplication before branching of monocots and dicots, suggesting that this separation gave rise to a subfunctionalization of these subgroups (14). Group 4 LEA proteins and transcripts have been found in dry seeds but also in response to water deficit in vegetative and reproductive tissues (14, 15). Moreover, *Arabidopsis* mutants deficient in group 4 LEA proteins are sensitive to

* This work was partially supported by Grants IN208212 from the Dirección General de Apoyo al Personal Académico/Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica PAPIIT, Universidad Nacional Autónoma de México and Grants 132258 and 221448 from the Consejo Nacional de Ciencia y Tecnología. The authors declare that they have no conflicts of interest with the contents of this article.

The nucleotide sequence(s) reported in this paper has been submitted to the DDBJ/GenBank™/EBI Data Bank with accession number(s) AEE31503.1, AEC09091.1, and AED91062.1.

¹ Supported by a Consejo Nacional de Ciencia y Tecnología doctoral fellowship.

² To whom correspondence should be addressed. Tel.: 52-777-329-1643; Fax: 52-777-313-9988; E-mail: crobles@ibt.unam.mx.

³ The abbreviations used are: LEA, late embryogenesis abundant; IDP, intrinsically disordered proteins; LDH, lactate dehydrogenase; TFE, 2,2,2-trifluoroethanol.

LEA4 Proteins Fold in Response to Water Deficit

water deficit, indicating that these proteins participate in the tolerance to this stress condition (14).

Many different functions have been proposed for group 4 LEA proteins such as membrane protectors, sugar or metal binding, radical scavengers, and protein dehydro- and cryoprotectors (16–20). *A. thaliana* AtLEA4-5 protein was shown to prevent inactivation and conformational changes of reporter enzymes such as lactate dehydrogenase (LDH) and malate dehydrogenase after partial dehydration and freeze/thaw cycles from 1:1 molar ratios, indicating that group 4 LEA proteins have a chaperone-like function to protect other proteins from the effects of water deficit (19, 20).

Studies on animal and bacterial IDPs have shown that these proteins can gain structural order upon binding to a specific partner, interaction that leads to IDP folding (21–25); however, in some other examples, the function of globular chaperones is linked to an order to disorder structural transitions in response to environmental cues such as those imposed by changes in pH or redox state (26, 27). Even though it has been shown that severe dehydration can promote folding of some LEA proteins (8, 11, 13, 28–31), the possibility that the environmental effects caused by mild water deficit in the cell (such as those occurring in vegetative tissues) leads to higher structural order in those IDPs responsive to this stressful environment (*e.g.* LEA proteins and other hydrophilins) and whether this structural changes could be related to their function are still open questions.

In this work, we demonstrate that even though *Arabidopsis* members of subgroups 4A (AtLEA4-2) and 4B (AtLEA4-5) LEA proteins are structurally disordered in solution, low osmotic potentials and macromolecular crowding can induce significant levels of α -helix, particularly in the conserved AtLEA4-5 N-terminal region, whereas the C-terminal region displays high structural disorder. We also show that the AtLEA4-5 N-terminal region is necessary and sufficient for the protective effect of this protein on reporter enzyme activities after freeze-thaw cycles and partial dehydration at low molar ratios. Our data support the hypothesis that cellular environment modulates the structural organization of disordered proteins and that these structural changes are related to their functions.

Experimental Procedures

In Silico Analyses—AtLEA4 proteins were aligned using T-Coffee multiple sequence alignment. Secondary structure prediction was determined using AGADIR helical content predictor (32). Intrinsically disordered tendency was predicted using DISpro (33), PONDR (34), and DISOPRED3 (35).

Plasmid Constructions—ORFs of *AtLEA4-1*, *AtLEA4-2*, and *AtLEA4-5* genes were cloned using cDNA from RNA obtained from *Arabidopsis* dry seeds. AtLEA4-5 ORF was amplified by PCR using specific primers containing at their ends NcoI (5'-AAACCATGGAGTCGATGAAAGAAAC-3') and SalI (5'-GCGGTCGACCCGTTTATCCAGTATATCC-3') restriction sites. This cloning strategy led to a modification in the second codon, which in the recombinant version corresponds to glutamic acid (GAG) instead of glutamine (CAG). The amplicon was cloned into pJET1.2/blunt and subsequently digested with NcoI and SalI for its insertion into pTrc99A vector. To eliminate the Met³³ in AtLEA4-5 ORF used in bacterial cells as an

alternative translation initiation site and responsible of the production of an additional shorter AtLEA4-5 protein, directed mutagenesis of AtLEA4-5 ORF sequence was conducted using the following overlapping primers to exchange Met³³ for a Leu residue: sense (5'-GGAGGAAAAGCGGAGAAGCTGAA-GAC-3') and antisense (5'-GTCTTCAGCTTCTCCGCCTT-TTCTCC-3'). The modified DNA fragment was inserted into pJET1.2/blunt plasmid vector to produce pJET1.2:AtLEA4-5. For protein production, AtLEA4-5 ORF was transferred to pTrc99A plasmid vector by digesting with NcoI and SalI restriction enzymes. The DNA fragments encoding AtLEA4-5 N- and C-terminal regions were obtained from pJET1.2:AtLEA4-5 plasmid, using the following sense and antisense oligonucleotides: 5'-AAACCATGGAGTCGATGAAAGAAAC-3' and antisense 5'-CGCGTTCGACTCAGGTTCCGGCTCCAGC-CGC-3' and sense 5'-AAACCATGGCCGGTTTAGGTTTG-GGGAC-3' and antisense 5'-GCGGTCGACCCGTTTATCC-AGTATATCC-3', respectively, which were also inserted into pTrc99A to generate pTrc99:AtLEA4-5₁₋₇₇ and pTrc99:AtLEA4-5₇₈₋₁₅₈ plasmids, respectively. In all cases, nucleotide sequences were verified accordingly.

Because pTrc99A:AtLEA4-1, pTrc99A:AtLEA4-2, and pTrc99A:AtLEA4-5₇₈₋₁₅₈ did not lead to a successful protein expression in bacteria, instead corresponding ORFs were inserted into the pTYB11 vector to obtain them as intein fusion proteins (IMPACT-CN expression system; New England Biolabs Inc.). To this end, AtLEA4-1 and AtLEA4-2 coding sequences were amplified from pJET1.2 intermediary plasmids using specific oligonucleotides containing SapI and PstI restriction sites: 5'-GGTGGTTGCTCTTCCAACATGCAATCGGCCGAAAC-AGAAG-3' and 5'-GGTGGTCTGCAGTCATTAGTAGTGTGATGATTATGATGTCC-3' for AtLEA4-1 and, 5'-GGTGGTTGCTCTTCCAACATGCAGTCGCGGAAGG-3' and 5'-GGTGGTCTGCAGTCATTAGATCTGTCCC GGCG-3' for AtLEA4-2. To amplify the AtLEA4-5₇₈₋₁₅₈ coding sequence, the oligonucleotides used were 5'-GGTGGTTGCTCTTCCAACACCGGTTTAGGTTTGGGGAC-3' and 5'-GGTGGTCTGCAGTCATTATCCAGTATATCCCCCGC-3'.

Expression and Purification of Recombinant Proteins—Recombinant plasmids derived from pTrc99A or pTYB11 as described above were transformed into *Escherichia coli* BL21(DE3) pLysS competent cells (Promega). Single colonies were inoculated in fresh LB medium containing 100 μ g/ml ampicillin and grown overnight at 37 °C, from which 1 liter of fresh LB medium was inoculated to 0.01 A₆₀₀ and grown at 37 °C to 0.5–0.8 A₆₀₀. At this point, protein expression was induced with 1 mM isopropyl β -D-1-thiogalactopyranoside for 6 h at 25 °C. Cell cultures were harvested by centrifugation. For AtLEA4-5 and AtLEA4-5₁₋₇₇ purification, we used a straightforward method designed for nonacidic recombinant unstructured proteins as described by Campos *et al.* (36). After washing twice with acetone, the protein was resuspended in 10 mM sodium phosphate buffer, pH 7.5, and dialyzed extensively against the same buffer. For intein fused AtLEA4-1, AtLEA4-2, and AtLEA4-5₇₈₋₁₅₈, a different purification procedure was followed. Bacterial pellets were resuspended in lysis buffer (20 mM sodium phosphate, pH 8, 500 mM NaCl, 0.1% Triton X-100) containing one tablet of cOmplete protease inhibitor mix-

ture (Roche) per 50 ml of buffer. The cells were lysed by sonication on ice, and the extract was clarified by centrifugation at $20,000 \times g$ for 30 min at 4 °C. To obtain proteins lacking the intein tag, the clarified extract was loaded onto a chitin column following the procedure described by the manufacturer (IMPACTTM-CN kit). The eluted fractions were analyzed for the presence of the recombinant proteins by SDS-PAGE. In contrast to AtLEA4-2, this analysis showed that AtLEA4-1 and AtLEA4-5_{78–158} are most probably protease-susceptible proteins because we were unable to detect the complete corresponding polypeptides, despite using protease-deficient bacterial strains, as is the case for BL21 and RosettaTM 2(DE3) (Merck-Millipore), strains successfully used to purify a number of recombinant proteins from different organisms. Fractions containing AtLEA4-2 were pooled and extensively dialyzed against 10 mM sodium phosphate buffer, pH 7.5. The purity and identity of the different purified proteins were confirmed by SDS-PAGE and by LC-MS. LC-MS was performed by the Proteomic Facility of the Instituto de Biotecnología/Universidad Nacional Autónoma de México following standard methods using LQT-Orbitrap Velos (Thermo-Fisher) mass spectrometer with nanospray ionization system. Proteins were quantified using their molar extinction coefficient at 280 nm ($\epsilon = 2,980 \text{ M}^{-1} \text{ cm}^{-1}$). Because AtLEA4-5_{1–77} lacks aromatic amino acid residues, its concentration was determined by Bradford assay and verified by SDS-PAGE comparing with known concentrations of AtLEA4-5. Purified proteins were conserved by lyophilization or in aliquots at $-80 \text{ }^\circ\text{C}$ until used. Because it was not possible to obtain whole AtLEA4-5_{78–158}, the complete chemically synthesized polypeptide corresponding to this truncated protein was purchased from Biomatik (Cambridge, Canada) and extensively dialyzed before use. This polypeptide was verified by mass spectrometry and HPLC.

Far UV Circular Dichroism Spectroscopy—AtLEA4-2, AtLEA4-5, and AtLEA4-5_{1–77} recombinant proteins, as well as AtLEA4-5_{78–158} polypeptide were diluted to 0.3 mg/ml and far UV CD spectra were recorded using a Jasco J-715 CD spectropolarimeter (JASCO Analytical Instruments) on a 0.1-cm-path length cell from 190 to 250 nm. Desired temperature was regulated with a Peltier temperature-controlled cell holder (PTC-4235; JASCO). Three spectra were averaged and smoothed to reduce noise. Each spectrum was acquired every 1 nm with 2-s average time per point and 1-nm band pass. Secondary structure estimation was calculated using Dichroweb software (37, 38). The CDSSTR algorithm was used with 4, 7, and SP175 data sets. These assays were reproduced using proteins from at least three independent purification batches.

In Vitro Freeze-Thaw Assay—Freeze-thaw *in vitro* assays were carried out as previously described by Reyes *et al.* (19) with small modifications. Briefly, LDH from rabbit muscle (Roche) was diluted to a final concentration of 250 nM (monomer) with or without the corresponding test protein (AtLEA4-2, AtLEA4-5, AtLEA4-5_{1–77}, AtLEA4-5_{78–158}, or lysozyme) in buffer 25 mM Tris-HCl, pH 7.5. The different test proteins were set to the desired molar ratio from 0.5:1 to 20:1 (test protein: LDH), considering 250 nM as 1:1 molar ratio. Mixtures in a final volume of 100 μl were frozen for 30 s in liquid N₂ and subsequently thawed at 25 °C in a thermomixer (Eppendorf). This

procedure constituted one freeze-thaw cycle, which was repeated up to seven times. After the treatment, LDH activity was measured as reported (19). Enzyme activities for each sample were measured in at least three independent tests (each one with three technical replicates). These experiments were reproduced with proteins from at least three independent purification batches.

In Vitro Partial Dehydration Assay—Partial dehydration *in vitro* assays were performed as previously described by Reyes *et al.* (20) with small modifications. LDH from rabbit muscle (Roche) was diluted to 250 nM (monomer) as the final concentration, in the presence or absence of the corresponding test protein (AtLEA4-2, AtLEA4-5, AtLEA4-5_{1–77}, or AtLEA4-5_{78–158}) in buffer of 25 mM Tris-HCl, pH 7.5. The different test proteins were set to 5:1 molar ratio (1.25 μM). Mixtures in a final volume of 25 μl were placed in a SpeedVac concentrator (Savant Instruments), and water was evaporated to achieve $\geq 98\%$ water loss, keeping constant temperature. The percentage of partial water loss was defined as the amount of water evaporated from the samples by means of weight. Partially dehydrated samples were rehydrated to the initial weight with water, assuring that all solutes were completely resuspended. LDH activity was measured as described above. These assays were reproduced using protein samples from at least three independent purification batches.

Statistical Analyses—Statistical analyses were carried out using one-way analysis of variance test. Significant differences were calculated with Tukey's multiple comparison post-test ($p < 0.01$).

Results

A. thaliana Group 4 Late Embryogenesis Abundant Proteins Are Intrinsically Disordered Proteins in Aqueous Solution—In *A. thaliana*, there are three genes encoding group 4 LEA proteins: AtLEA4-1, AtLEA4-2, and AtLEA4-5 (3, 14). These proteins have low molecular masses between 10.5 and 16.2 kDa and pI values between 8.95 and 9.67. All of them present a 74–78-amino acid long N-terminal domain, highly conserved across all group 4 LEA proteins described so far in the plant kingdom, which contains distinctive motifs for this protein family (Fig. 1A). Because of the amino acid composition of this group of LEA proteins, it was proposed that they are IDPs (2, 12). *In silico* analyses using PONDR (VLXT) (34) (Fig. 1B), DISpro (33), and DISOPRED3 (35) (data not shown) indicated that, despite their sequence similarity, they possess different levels of disorder.

To characterize the LEA4 proteins structural features, we expressed and purified the three recombinant proteins in *E. coli*. AtLEA4-2 and AtLEA4-5 were obtained with more than 95% purity (data not shown). Because AtLEA4-1 was mostly degraded during different expression and purification procedures, the rest of the experiments were performed only with AtLEA4-2 and AtLEA4-5, representing the two LEA4 subgroups, 4A and 4B, respectively (14).

To determine the secondary structure of AtLEA4-2 and AtLEA4-5 in solution, the purified proteins were analyzed by far UV CD. The results obtained corroborated that both proteins are mostly disordered in solution over a wide range of temperatures (Fig. 2, A and B) and pH (data not shown), with

LEA4 Proteins Fold in Response to Water Deficit




FIGURE 1. *In silico* structure prediction of *Arabidopsis* group 4 LEA proteins. *A*, sequence alignment of *Arabidopsis* group 4 LEA proteins showing N-terminal conserved and C-terminal variable regions. *B*, AtLEA4-1 (black), AtLEA4-2 (blue), and AtLEA4-5 (green) structural disorder levels using PONDR predictor. According to this algorithm, proteins with disorder probability values above 0.5 are considered highly disordered. *C*, percentage of α -helix predicted using AGADIR for AtLEA4-1 (black), AtLEA4-2 (blue), and AtLEA4-5 (green).




FIGURE 2. Far UV CD spectra of AtLEA4-2 and AtLEA4-5 in aqueous solution at various temperatures and in different TFE concentrations. *A* and *B*, far UV CD spectra of AtLEA4-2 (*A*) and AtLEA4-5 (*B*) at 10 °C (black), 20 °C (blue), 30 °C (brown), 40 °C (cyan), 50 °C (green), 60 °C (magenta), 70 °C (orange), and 80 °C (purple). Insets in *A* and *B* show $\Delta 10-80$ °C (continuous black line) and $\Delta 20-80$ °C (dashed blue line) difference spectra. *C* and *D*, AtLEA4-2 (*C*) and AtLEA4-5 (*D*) in TFE/water mixtures at 0% (black), 4% (blue), 8% (brown), 12% (cyan), 16% (green), 20% (magenta), 40% (orange), 60% (purple), and 90% (red) TFE. The difference spectra ($\Delta 90-0\%$ TFE) are shown as insets in *C* and *D*. These results were reproduced at least four times in autonomous experiments, using three independent purification batches of both proteins.

the characteristic negative band for random coil structures around 198 nm. Both spectra showed a significant negative signal around 222 nm (typical of α -helix), suggesting that these proteins possess residual α -helix structure. The comparative

analysis of the AtLEA4-2 CD difference spectra obtained at different temperatures ($\Delta 10-80$ °C and $\Delta 20-80$ °C) revealed that this protein is able to form α -helix structures at low temperatures (10 °C), whereas at higher temperatures the protein is

TABLE 1
Percentage of helix, strand, and unordered structures in AtLEA4-2 and AtLEA4-5

Secondary structure content in AtLEA4-2 and AtLEA4-5 proteins was obtained by far UV CD spectrometry and calculated with Dichroweb server.

Treatment	AtLEA4-2			AtLEA4-5		
	Helix	Strand	Unordered	Helix	Strand	Unordered
			%			%
Aqueous solution	5	33	61	5	34	61
4% TFE	7	32	61	6	33	61
8% TFE	16	25	58	6	32	61
12% TFE	21	23	56	17	23	59
16% TFE	35	15	51	18	25	55
20% TFE	42	15	44	32	17	51
40% TFE	57	9	33	49	13	39
60% TFE	62	9	30	57	8	35
90% TFE	80	3	16	88	2	10
10% glycerol	5	34	60	4	34	61
20% glycerol	8	32	61	7	36	57
30% glycerol	18	22	60	7	34	59
40% glycerol	21	23	57	18	24	57
50% glycerol	33	12	53	18	23	59
60% glycerol	43	14	43	34	15	51
70% glycerol	49	9	41	42	11	48
80% glycerol	54	12	33	46	9	45
45% PEG 5000	37	11	52	39	9	53

mostly disordered (Fig. 2A, inset). By contrast, a similar analysis indicated that AtLEA4-5 is mostly disordered under all temperatures tested (Fig. 2B, inset). We did not find evidence of extended helical conformations (e.g. poly-L-proline II) for any of these proteins, such as those found in LEA groups 1, 2, and 6 (10, 39). Altogether, these data demonstrate that AtLEA4 proteins are IDPs with residual α -helix structure in aqueous solution.

AtLEA4-2 and AtLEA4-5 Have the Potential to Acquire High Levels of Ordered Structure—To determine the intrinsic ability of AtLEA4-2 and AtLEA4-5 to attain helicity, CD analyses were performed in the presence of different concentrations of 2,2,2-trifluoroethanol (TFE), a well known α -helix inducer (40, 41). Increasing TFE concentrations promote α -helix formation in both proteins, as revealed by the progressive increase in $[\theta]_{198}$ toward positive values and the transition of the minimum at $[\theta]_{222}$ onward more negative values (Fig. 2, C and D). Difference spectra showed that both proteins are able to gain high helicity levels; however, $\Delta 90-0\%$ TFE indicates that AtLEA4-5 reaches higher α -helix percentage than AtLEA4-2 (Fig. 2, C and D, insets). Assessment of protein secondary structure using Dichroweb (37) indicated that AtLEA4-2 adopts 7% of α -helix in 4% TFE, and it reaches up to 80% α -helix in 90% TFE (Table 1), showing a decrease in its structural disorder from 61% in aqueous solution to 16% in 90% TFE. The α -helix content for AtLEA4-5 increased from 6% to 88% in 4% to 90% TFE, whereas its disordered structure decreased from 61% to 10% (Table 1). Both AtLEA4-2 and AtLEA4-5 CD spectra obtained with 4% to 90% TFE showed isodichroic points (Fig. 2, C and D), indicating that these proteins are able to adopt two structural conformations in equilibrium under these conditions: one mostly unstructured favored in aqueous solution (U_{4-2} and U_{4-5}) and a second one with higher helicity promoted by increasing TFE concentrations (F_{4-2} and F_{4-5}). The fitted straight lines obtained from transition diagrams for AtLEA4-2 and AtLEA4-5 support the formation of two conformers for both proteins

(Fig. 3, E and F), as was indicated by the presence of isodichroic points. These data demonstrate that AtLEA4-2 and AtLEA4-5 possess an intrinsic ability to form α -helical species, which under some conditions are in equilibrium with unfolded conformations ($U_{4-2} \leftrightarrow F_{4-2}$; $U_{4-5} \leftrightarrow F_{4-5}$).

AtLEA4-2 and AtLEA4-5 Fold to α -Helix in Response Low Water Availability and Macromolecular Crowding Induced in Vitro—Previous studies have shown that recombinant LEA proteins from different groups acquire secondary structure, mostly α -helix, when subjected to complete dehydration (8, 11, 13, 28–31). Given the intrinsic potential of AtLEA4-2 and AtLEA4-5 to gain α -helix conformation and because these proteins accumulate even under mild water limitation, we hypothesized that *in vitro* conditions limiting water availability could induce changes in their secondary structure. Addition of increasing glycerol concentrations led to a notorious progressive gain in α -helix structure in both proteins, as shown by the $[\theta]_{198}$ change toward positive values and a deeper minimum at $[\theta]_{222}$ (Fig. 3, A and B). Analysis using Dichroweb estimated a small difference in α -helix content between these two proteins: 54% α -helix for AtLEA4-2 and 46% for AtLEA4-5 at the highest glycerol concentration (80%) (Table 1). We observed the presence of isodichroic points in both cases (Fig. 3, A and B), which was also supported by their corresponding transition diagrams (Fig. 3, E and F), that together with those obtained from TFE treatments showed that AtLEA4-2 and AtLEA4-5 seem to follow the same folding pathway to α -helix under both treatments (Fig. 3, E and F).

Inside living cells, macromolecules are present at very high concentrations (~ 400 g/liter) (42–44), a condition that is typically known as macromolecular crowding (45, 46). This state is further exacerbated in cells under water deficit, reaching macromolecular concentrations up to ~ 900 g/liter upon severe dehydration (42). PEG was used to simulate a crowded environment *in vitro*. The addition of 45% PEG 5000 to AtLEA4-2 or AtLEA4-5 solutions clearly induced changes in their structural conformations (Fig. 3, C and D). Dichroweb estimations indicate 37 and 39% α -helix gains for AtLEA4-2 and AtLEA4-5, respectively (Table 1). Together, these data indicate that AtLEA4-2 and AtLEA4-5 can acquire secondary structure under low water availability or macromolecular crowding *in vitro*, possibly reflecting what occurs in plant cells under water deficit.

The N-terminal Region of AtLEA4 Proteins Is Necessary and Sufficient for the Conformational Changes Induced by Water Deficit—Plant group 4 LEA proteins are characterized by the presence of conserved motifs at their N-terminal region (2, 14). *In silico* analysis predicts that this region of 70, 74, and 77 residues in AtLEA4-1, AtLEA4-2, and AtLEA4-5, respectively, has a higher propensity to adopt α -helical conformations than the C-terminal region (Fig. 1C). To test this prediction, we performed far UV CD experiments using AtLEA4-5 truncated versions: one containing the first 77 amino acids, named AtLEA4-5₁₋₇₇, and a second one corresponding to the 81 amino acid C-terminal region, from residues 78 to 158 (AtLEA4-5₇₈₋₁₅₈) (47).

In contrast to *in silico* predictions, AtLEA4-5₁₋₇₇ behaved as a disordered protein in aqueous solution under all tempera-




FIGURE 3. Far UV CD spectra of AtLEA4-2 and AtLEA4-5 in different concentrations of glycerol or PEG. A and B, far UV CD spectra of AtLEA4-2 (A) and AtLEA4-5 (B) in glycerol/water mixtures at 0% (black), 10% (blue), 20% (brown), 30% (cyan), 40% (green), 50% (magenta), 60% (orange), 70% (purple), and 80% (red) glycerol. Difference spectra ($\Delta 80-0\%$ glycerol) are shown as insets in A and B. C and D, AtLEA4-2 (C) and AtLEA4-5 (D) in aqueous solution (black line) and in PEG/water mixture at 45% PEG 5000 (blue line). Difference spectra ($\Delta 45-0\%$ PEG) are shown as insets in C and D. E and F, transition diagrams for AtLEA4-2 (E) and AtLEA4-5 (F) were obtained using the ellipticity values at 198 and 222 nm from TFE and glycerol titrations. Dashed lines represent the linear fits of the data. These results were reproduced at least four times in autonomous experiments, using three independent purification batches of both proteins.

tures tested (Fig. 4A), with a difference spectrum profile ($\Delta 10-80^\circ\text{C}$) similar to that obtained for the complete AtLEA4-5 (Figs. 2B and 4A, insets). However, the addition of increasing TFE concentrations progressively induced α -helix formation in the truncated protein containing the N-terminal region (Fig. 4B). Likewise, treatments with glycerol and PEG led to the same behavior in this protein (Fig. 4, C and D), reaching up to 55 and 42% α -helix at the highest glycerol and PEG concentrations, respectively (Table 2). Also complete AtLEA4-5 and AtLEA4-2, far UV CD spectra from AtLEA4-5₁₋₇₇ showed isodichroic points when treated with progressively increasing TFE or glycerol concentrations, indicating that this protein is in equilibrium between two states in these conditions: disordered and α -helix conformations (Fig. 4, B and C). The high linear corre-

lation of transition diagrams confirmed this observation (data not shown).

Using the chemically synthesized peptide corresponding to the AtLEA4-5C-terminal region (AtLEA4-5₇₈₋₁₅₈), far UV CD analysis confirmed that AtLEA4-5₇₈₋₁₅₈ is highly disordered in aqueous solution at different temperatures (Fig. 4E). Unlike AtLEA4-2, AtLEA4-5, or AtLEA4-5₁₋₇₇, the difference spectrum $\Delta 10-80^\circ\text{C}$ of AtLEA4-5₇₈₋₁₅₈ indicated the possible formation of poly-L-proline-like structures (Fig. 4E, inset). In contrast to the N-terminal region of these proteins, addition of up to 60% TFE had only minor effects on the structure of AtLEA4-5₇₈₋₁₅₈ (Fig. 4F), inducing 4% to 19% α -helix (Table 2). It was not until a high TFE concentration (90%) was reached that AtLEA4-5₇₈₋₁₅₈ became helical (Fig. 4F), indicating a low




FIGURE 4. **N-terminal conserved region of AtLEA4-5 is necessary and sufficient to induce folding to α -helix under the different conditions tested.** Far UV CD spectra of AtLEA4-5₁₋₇₇ (A–D) and AtLEA4-5₇₈₋₁₅₈ (E–H) under different temperatures (A and E), different TFE (B and F) and glycerol concentrations (C and G), and under 45% PEG 5,000 (D and H). Difference spectra ($\Delta 10-80^\circ\text{C}$; $\Delta 90-0\%$ TFE; $\Delta 80-0\%$ glycerol; $\Delta 45-0\%$ PEG) are shown as insets in each graph. These results were reproduced at least four times in autonomous experiments, using three independent purification batches of AtLEA4-5₁₋₇₇ protein and one batch for AtLEA4-5₇₈₋₁₅₈ polypeptide.

intrinsic competence to acquire ordered structures. This finding was further supported by the results obtained from the addition of glycerol or PEG to AtLEA4-5₇₈₋₁₅₈ solutions, which showed no effect on its structure (Fig. 4, G and H), given that the

negative band at $[\theta]_{222}$ did not show any change and that only a slight increase in the $[\theta]_{198}$ signal was detected. Under the highest glycerol (80%) and PEG (45%) concentrations, AtLEA4-5₇₈₋₁₅₈ only showed 5% α -helix formation (Table 2). Comparison

LEA4 Proteins Fold in Response to Water Deficit

between the difference spectra for AtLEA4-5₁₋₇₇ and AtLEA4-5₇₈₋₁₅₈ in TFE, glycerol and PEG showed the magnitude of the AtLEA4-5₁₋₇₇ folding to α -helix, compared with AtLEA4-5₇₈₋₁₅₈ (Fig. 4, A–H, insets). In accordance, the transition diagram for glycerol titration of AtLEA4-5₇₈₋₁₅₈ showed no linear behavior (data not shown). Together, these data demonstrate that the AtLEA4-5 N-terminal region (AtLEA4-5₁₋₇₇) is necessary and sufficient to drive α -helix conformations in this protein under low water availability or macromolecular crowding conditions.

The N-terminal Region of AtLEA4 Proteins Is Necessary and Sufficient to Prevent Inactivation of Lactate Dehydrogenase Caused by Freeze-Thaw Cycles and Partial Dehydration—Because the motif conservation and ability to fold into α -helical conformations of the AtLEA4-5 are preferentially located at the N-terminal region as compared with its C-terminal region, we

TABLE 2

Percentage of helix, strand, and unordered structures in AtLEA4-5₁₋₇₇ and AtLEA4-5₇₈₋₁₅₈

Secondary structure content in AtLEA4-2 and AtLEA4-5 proteins was obtained by far UV CD spectrometry and calculated with Dichroweb server. NA, not available.

Treatment	AtLEA4-5 ₁₋₇₇			AtLEA4-5 ₇₈₋₁₅₈		
	Helix	Strand	Unordered	Helix	Strand	Unordered
Aqueous solution	6	32	61	3	38	58
4% TFE	6	33	59	4	37	58
8% TFE	7	32	60	4	37	58
12% TFE	18	25	57	5	36	59
16% TFE	22	24	53	5	35	59
20% TFE	36	16	48	5	36	59
40% TFE	59	9	31	8	33	58
60% TFE	65	7	27	19	28	54
90% TFE	80	3	16	49	15	35
10% glycerol	6	33	60	NA	NA	NA
20% glycerol	16	23	62	NA	NA	NA
30% glycerol	19	24	58	6	45	51
40% glycerol	19	23	57	0	53	45
50% glycerol	24	18	57	6	36	56
60% glycerol	39	12	48	4	41	55
70% glycerol	45	9	46	NA	NA	NA
80% glycerol	55	8	35	5	35	58
45% PEG 5000	42	9	50	NA	NA	NA

asked about the competence of these individual segments to protect the reporter enzyme LDH from deleterious effects caused by freeze-thaw cycles and partial dehydration, as was previously shown for the complete protein (19, 20). For this purpose, LDH in the presence or absence of AtLEA4-5, AtLEA4-2, AtLEA4-5₁₋₇₇, or AtLEA4-5₇₈₋₁₅₈ was subjected to *in vitro* freeze-thaw cycles and partial dehydration treatments. AtLEA4-2, the smallest protein of this group, mostly consisting of the N-terminal region (Fig. 1A), showed a similar protective effect on LDH activity as AtLEA4-5 after partial dehydration (Fig. 5A). Interestingly, a comparable protection was produced by AtLEA4-5₁₋₇₇, contrasting with the negligible protective levels showed by the AtLEA4-5 C-terminal region (AtLEA4-5₇₈₋₁₅₈), whose values were rather close to those shown by lysozyme, an unrelated globular protein (Fig. 5, A and B). A similar conclusion can be drawn when different molar ratios of these proteins were used in freeze-thaw *in vitro* assays, where it is evident that the various proteins containing the N-terminal region showed an equivalent protective trend, in opposition to the C-terminal region represented by AtLEA4-5₇₈₋₁₅₈ or to lysozyme (Fig. 5B). This analysis revealed the highest and lowest protective efficiencies for AtLEA4-5 and AtLEA4-5₇₈₋₁₅₈, respectively, among the different proteins tested (Fig. 5B). Because the hydrophilicity index and length are similar between the N-terminal and C-terminal regions, these data are consistent with the hypothesis that the protective activity of these proteins is rather dependent on the conserved motifs present in the N-terminal region and/or on its ability to fold into α -helical conformations.

Discussion

During the last nearly 20 years, research on the so-called IDPs has challenged the classical structure-function paradigm (48). IDPs that either completely lack a well defined three-dimensional structure or contain short regions of disorder, known as intrinsically disordered regions (IDRs) within folded




FIGURE 5. The N-terminal conserved region of AtLEA4-5 is necessary and sufficient to prevent inactivation of LDH after partial dehydration and freeze/thaw cycles *in vitro*. A, remaining LDH enzymatic activity after $\geq 98\%$ water loss induced *in vitro* of samples containing LDH without any additive (control) or with AtLEA4-2, AtLEA4-5, AtLEA4-5₁₋₇₇, and AtLEA4-5₇₈₋₁₅₈ in a 5:1 molar ratio (additive:LDH). B, remaining LDH enzymatic activity after seven freeze-thaw cycles of LDH without any additive or with increasing molar amounts (from 0:1 to 20:1 molar ratio additive:LDH) of AtLEA4-2 (black line with circles), AtLEA4-5 (blue line with squares), AtLEA4-5₁₋₇₇ (red line with up triangles), AtLEA4-5₇₈₋₁₅₈ (magenta line with inverted triangles), and lysozyme (purple line with open circles). Error bars indicate S.E. of three independent tests (with three internal repetitions). Letters indicate significant differences calculated with Tukey's multiple comparison post-test ($p < 0.01$). The data in B were fit to a hyperbola curve, and significant differences were subsequently calculated with Tukey's multiple comparison post-test ($p < 0.01$). These results were reproduced at least four times in autonomous experiments, using three independent purification batches of AtLEA4-2, AtLEA4-5, and AtLEA4-5₁₋₇₇ proteins, and one batch for AtLEA4-5₇₈₋₁₅₈ polypeptide.

domains are highly abundant in eukaryotic proteomes and perform important functions (6). In plants, IDPs participate in developmental control, light perception, transcriptional regulation, and response to abiotic stress (12). LEA proteins are the plant protein family with the highest number of proteins either predicted or characterized as IDPs (12). There is experimental evidence showing that LEA proteins from different groups acquire α -helix after complete drying (8, 11, 13, 28–31). These findings suggest that these proteins form α -helix in the dry seed, but little is known about LEA proteins structural behavior under less severe conditions, such as those present in vegetative tissues under drought. Mouillon *et al.* (49) showed that three *Arabidopsis* group 2 LEA proteins (Cor47, Lti29, and Lti30) are IDPs in solution and that they remain disordered when subjected to low water potential or macromolecular crowding simulated *in vitro* with glycerol and PEG, respectively. Even though Cor47 showed a slight folding into α -helix under high concentrations of glycerol and PEG, the authors propose that dehydrins have evolved to stay disordered under cellular conditions and that disorder might be required to properly fulfill their function (49). A similar effect was also observed experimentally for α -casein, MAP2c, and p21^{Cip1}, three different animal IDPs that show no conformational change when subjected to macromolecular crowding, supporting the idea that the physiological state of these IDPs is also disordered (50).

Even though, most LEA proteins can be considered as IDPs, the analysis of their amino acid sequences indicates variety in their potential to attain different levels of secondary structure (2). Because LEA proteins accumulate not only under severe dehydration, such as that occurring in the dry seed, but also in different tissues under a wide range of water limitation, we investigated the possibility that these proteins could adopt secondary structures under conditions prevailing upon water loss, not necessarily as extreme as those occurring in dry seeds. Group 4 LEA proteins in *Arabidopsis* represented a suitable set of proteins for this purpose, given that there are two protein subtypes (subgroups 4A and 4B): a long variant conformed by two distinctive N-terminal and C-terminal regions (AtLEA4-5, representative of subgroup 4B) and a short one with only the conserved N-terminal region (AtLEA4-2, representative of subgroup 4A) (14). Furthermore, *in silico* analysis showed that in contrast to the C-terminal region, the N-terminal region sequence has the potential to fold into α -helix conformations offering an appropriate system to experimentally compare their structural properties under different environments. In this work, we demonstrate not only that the N-terminal regions of AtLEA4-2 and AtLEA4-5 have an intrinsic capacity to fold into α -helix as shown by the far UV CD spectra in the presence of TFE but also that they gain significant helicity in low water potential and macromolecular crowded solutions. Interestingly, the AtLEA4-5 C-terminal region, which we show is disordered in aqueous solution as predicted, remains disordered under all conditions tested. These results indicate the presence of two functional domains in this LEA protein group. The N-terminal region can become ordered under conditions of water restriction, short of absolute dryness, whereas the persistent disorder of the C-terminal region could indicate a requirement to expose some amino acid residues for a more effective or

additional function. Such possibility could be related to the abundance of His residues in this region, which seems to be involved in their binding to metal ions (18), suggesting multifunctionality in some LEA protein families.

All LEA proteins studied to date from groups 1, 2, 3, 4, and 6 show structural disorder in aqueous solution, but most of them also have an intrinsic potential to acquire helical conformations in the presence of TFE (12). These results suggest that there are conditions where LEA proteins exhibit such structural transformation. For most of these LEA proteins, folding has been detected after extreme dehydration (8, 11, 13, 28–31), but extreme conditions may not be necessary in all cases to induce structural transformation. We show in this work that group 4 LEA proteins reach up to 54 and 39% α -helix under less severe environments regarding water availability and macromolecular crowding, respectively. The levels of α -helix formation for group 4 LEA proteins are significantly higher than those estimated for LEA proteins from other groups under similar treatments (12, 39), indicating that the ability to fold is not necessarily the same for different IDPs or intrinsically disordered regions. This is also supported by the wide range of α -helix formation observed among different LEA proteins upon comparable treatments (9, 12, 39, 49, 51, 52). Other proteins seem to be unable to gain ordered conformations such as Rab18, a group 2 LEA protein (9). This information indicates diversity in their structural plasticity, action mechanisms, and/or in their function.

The results from the protection activity assays demonstrate that the protective role of LEA 4 proteins on LDH under low water availability and/or molecular crowding is confined to the conserved N-terminal region of this protein family. There is no apparent participation of the C-terminal region, which completely lacks this safeguard function. These observations imply that the ability of a LEA4 N-terminal domain to gain helicity is related to this chaperone-like activity, particularly under conditions prevailing in water-deficit environments. Based on *in vitro* evidence, it has been proposed that one mechanism for this activity involves protein-protein interactions (20, 53–55). A comprehensive view brings into consideration the existence of different LEA4 conformers in equilibrium (partially folded or unfolded) under crowded or water-deficit environments, which supports the existence of preformed secondary structural elements, denominated as prestructured motifs that could be implicated in the recognition of different and specific binding partners (56). Our findings strongly suggest that water deficit leads to the stabilization of particular conformations in group 4 LEA proteins that may allow the exposure of different motifs necessary for the binding of their target molecules, hence supporting the idea that LEA proteins of this group function as a structural ensemble, whose dynamism can be modulated by environmental conditions (57–60). This proposed mode of action also exhibits possible binding promiscuity, in consonance with their role as chaperone-like molecules needed during water scarcity. At this point, we cannot discount the possibility that different conformers in these proteins could favor different functions such as protection of proteins and/or membranes, as well as metal binding, a moonlighting property arising from their structural dynamics.

LEA4 Proteins Fold in Response to Water Deficit

In conclusion, the findings reported here indicate that the *in vitro* chaperone-like function of the intrinsically disordered group 4 LEA proteins is closely associated to their ability to adopt ordered structural conformations under prevailing conditions in water-deficit environments. The high correlation found between accumulation under water deficit and intrinsic structural disorder, common features in typical LEA proteins and other hydrophilins (4), suggests a functional advantage for this attribute throughout evolution, not only to maintain structural plasticity that could avoid undesirable consequences under stressful environments, such as dramatic structural modifications leading to a functional breakdown, but also to gain functional and mechanistic diversity given their conformational freedom.

Author Contributions—C. L. C.-V. and A. A. C. designed the experimental strategy. C. L. C.-V. conducted cloning, protein expression, and protein purification. C. L. C.-V. and G. S.-R. performed circular dichroism spectroscopy experiments. C. L. C.-V. and J. L. R. carried out *in vitro* protection assays. A. A. C. supervised research. C. L. C.-V., G. S.-R., and A. A. C. analyzed data. C. L. C.-V. and A. A. C. wrote the article, which was read and approved by all authors.

Acknowledgments—We thank H. Jane Dyson for critical reviewing this manuscript, Rosa M. Solórzano for technical assistance in protein purification, and Francisco Campos for the gift of pTrc99A/AtLEA4-*S*_{1–77} plasmid. We are also grateful to the proteomics (Unidad de Proteómica) and DNA sequencing (Unidad de Síntesis y Secuenciación de ADN) core facilities at the Instituto de Biotecnología of the Universidad Nacional Autónoma de México.

References

1. Bray, E. A. (1997) Plant responses to water deficit. *Trends Plant Sci.* **2**, 48–54
2. Battaglia, M., Olvera-Carrillo, Y., Garcarrubio, A., Campos, F., and Covarrubias, A. A. (2008) The enigmatic LEA proteins and other hydrophilins. *Plant Physiol.* **148**, 6–24
3. Hundertmark, M., and Hinch, D. K. (2008) LEA (late embryogenesis abundant) proteins and their encoding genes in *Arabidopsis thaliana*. *BMC Genomics* **9**, 118
4. Garay-Arroyo, A., Colmenero-Flores, J. M., Garcarrubio, A., and Covarrubias, A. A. (2000) Highly hydrophilic proteins in prokaryotes and eukaryotes are common during conditions of water deficit. *J. Biol. Chem.* **275**, 5668–5674
5. Dyson, H. J., and Wright, P. E. (2005) Intrinsically unstructured proteins and their functions. *Nat. Rev. Mol. Cell Biol.* **6**, 197–208
6. Tompa, P. (2012) Intrinsically disordered proteins: a 10-year recap. *Trends Biochem. Sci.* **37**, 509–516
7. Hughes, S., and Graether, S. P. (2011) Cryoprotective mechanism of a small intrinsically disordered dehydrin protein. *Protein Sci.* **20**, 42–50
8. Hundertmark, M., Popova, A. V., Rausch, S., Seckler, R., and Hinch, D. K. (2012) Influence of drying on the secondary structure of intrinsically disordered and globular proteins. *Biochem. Biophys. Res. Commun.* **417**, 122–128
9. Mouillon, J. M., Gustafsson, P., and Harryson, P. (2006) Structural investigation of disordered stress proteins. Comparison of full-length dehydrins with isolated peptides of their conserved segments. *Plant Physiol.* **141**, 638–650
10. Rivera-Najera, L. Y., Saab-Rincón, G., Battaglia, M., Amero, C., Pulido, N. O., García-Hernández, E., Solórzano, R. M., Reyes, J. L., and Covarrubias, A. A. (2014) A group 6 late embryogenesis abundant protein from common bean is a disordered protein with extended helical structure and oligomer-forming properties. *J. Biol. Chem.* **289**, 31995–32009
11. Shih, M. D., Hsieh, T. Y., Lin, T. P., Hsing, Y. I., and Hoekstra, F. A. (2010) Characterization of two soybean (*Glycine max* L.) LEA IV proteins by circular dichroism and Fourier transform infrared spectrometry. *Plant Cell Physiol.* **51**, 395–407
12. Sun, X., Rikkerink, E. H., Jones, W. T., and Uversky, V. N. (2013) Multifarious roles of intrinsic disorder in proteins illustrate its broad impact on plant biology. *Plant Cell* **25**, 38–55
13. Tolleter, D., Jaquinod, M., Mangavel, C., Passirani, C., Saulnier, P., Manon, S., Teyssier, E., Payet, N., Avelange-Macherel, M. H., and Macherel, D. (2007) Structure and function of a mitochondrial late embryogenesis abundant protein are revealed by desiccation. *Plant Cell* **19**, 1580–1589
14. Olvera-Carrillo, Y., Campos, F., Reyes, J. L., Garcarrubio, A., and Covarrubias, A. A. (2010) Functional analysis of the group 4 late embryogenesis abundant proteins reveals their relevance in the adaptive response during water deficit in *Arabidopsis*. *Plant Physiol.* **154**, 373–390
15. Schmid, M., Davison, T. S., Henz, S. R., Pape, U. J., Demar, M., Vingron, M., Schölkopf, B., Weigel, D., and Lohmann, J. U. (2005) A gene expression map of *Arabidopsis thaliana* development. *Nat. Genet.* **37**, 501–506
16. Dang, N. X., Popova, A. V., Hundertmark, M., and Hinch, D. K. (2014) Functional characterization of selected LEA proteins from *Arabidopsis thaliana* in yeast and *in vitro*. *Planta* **240**, 325–336
17. Hundertmark, M., Dimova, R., Lengefeld, J., Seckler, R., and Hinch, D. K. (2011) The intrinsically disordered late embryogenesis abundant protein LEA18 from *Arabidopsis thaliana* modulates membrane stability through binding and folding. *Biochim. Biophys. Acta* **1808**, 446–453
18. Liu, G., Xu, H., Zhang, L., and Zheng, Y. (2011) Fe binding properties of two soybean (*Glycine max* L.) LEA4 proteins associated with antioxidant activity. *Plant Cell Physiol.* **52**, 994–1002
19. Reyes, J. L., Campos, F., Wei, H., Arora, R., Yang, Y., Karlson, D. T., and Covarrubias, A. A. (2008) Functional dissection of hydrophilins during *in vitro* freeze protection. *Plant Cell Environ.* **31**, 1781–1790
20. Reyes, J. L., Rodrigo, M. J., Colmenero-Flores, J. M., Gil, J. V., Garay-Arroyo, A., Campos, F., Salamini, F., Bartels, D., and Covarrubias, A. A. (2005) Hydrophilins from distant organisms can protect enzymatic activities from water limitation effects *in vitro*. *Plant Cell Environ.* **28**, 709–718
21. Borchers, W., Theillet, F. X., Kätzer, A., Finzel, A., Mishall, K. M., Powell, A. T., Wu, H., Manieri, W., Dieterich, C., Selenko, P., Loewer, A., and Daughdrill, G. W. (2014) Disorder and residual helicity alter p53-Mdm2 binding affinity and signaling in cells. *Nat. Chem. Biol.* **10**, 1000–1002
22. Dyson, H. J., and Wright, P. E. (2002) Coupling of folding and binding for unstructured proteins. *Curr. Opin. Struct. Biol.* **12**, 54–60
23. Radhakrishnan, I., Pérez-Alvarado, G. C., Parker, D., Dyson, H. J., Montminy, M. R., and Wright, P. E. (1997) Solution structure of the KIX domain of CBP bound to the transactivation domain of CREB: a model for activator:coactivator interactions. *Cell* **91**, 741–752
24. Rogers, J. M., Wong, C. T., and Clarke, J. (2014) Coupled folding and binding of the disordered protein PUMA does not require particular residual structure. *J. Am. Chem. Soc.* **136**, 5197–5200
25. Sugase, K., Dyson, H. J., and Wright, P. E. (2007) Mechanism of coupled folding and binding of an intrinsically disordered protein. *Nature* **447**, 1021–1025
26. Reichmann, D., Xu, Y., Cremers, C. M., Ilbert, M., Mittelman, R., Fitzgerald, M. C., and Jakob, U. (2012) Order out of disorder: working cycle of an intrinsically unfolded chaperone. *Cell* **148**, 947–957
27. Tapley, T. L., Körner, J. L., Barge, M. T., Hupfeld, J., Schauerer, J. A., Gafni, A., Jakob, U., and Bardwell, J. C. (2009) Structural plasticity of an acid-activated chaperone allows promiscuous substrate binding. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 5557–5562
28. Goyal, K., Tisi, L., Basran, A., Browne, J., Burnell, A., Zurdo, J., and Tunnacliffe, A. (2003) Transition from natively unfolded to folded state induced by desiccation in an anhydrobiotic nematode protein. *J. Biol. Chem.* **278**, 12977–12984
29. Popova, A. V., Hundertmark, M., Seckler, R., and Hinch, D. K. (2011) Structural transitions in the intrinsically disordered plant dehydration stress protein LEA7 upon drying are modulated by the presence of membranes. *Biochim. Biophys. Acta* **1808**, 1879–1887
30. Shih, M. D., Hsieh, T. Y., Jian, W. T., Wu, M. T., Yang, S. J., Hoekstra, F. A., and Hsing, Y. I. (2012) Functional studies of soybean (*Glycine max* L.) seed

- LEA proteins GmPM6, GmPM11, and GmPM30 by CD and FTIR spectroscopy. *Plant Sci.* **196**, 152–159
31. Shimizu, T., Kanamori, Y., Furuki, T., Kikawada, T., Okuda, T., Takahashi, T., Mihara, H., and Sakurai, M. (2010) Desiccation-induced structuralization and glass formation of group 3 late embryogenesis abundant protein model peptides. *Biochemistry* **49**, 1093–1104
 32. Muñoz, V., and Serrano, L. (1994) Elucidating the folding problem of helical peptides using empirical parameters. *Nat. Struct. Biol.* **1**, 399–409
 33. Cheng, J. L., Sweredoski, M. J., and Baldi, P. (2005) Accurate prediction of protein disordered regions by mining protein structure data. *Data Min. Knowl. Disc.* **11**, 213–222
 34. Romero, P., Obradovic, Z., Li, X., Garner, E. C., Brown, C. J., and Dunker, A. K. (2001) Sequence complexity of disordered protein. *Proteins* **42**, 38–48
 35. Jones, D. T., and Cozzetto, D. (2015) DISOPRED3: precise disordered region predictions with annotated protein-binding activity. *Bioinformatics* **31**, 857–863
 36. Campos, F., Guillén, G., Reyes, J. L., and Covarrubias, A. A. (2011) A general method of protein purification for recombinant unstructured non-acidic proteins. *Protein Expr. Purif.* **80**, 47–51
 37. Lobley, A., Whitmore, L., and Wallace, B. A. (2002) DICHROWEB: an interactive website for the analysis of protein secondary structure from circular dichroism spectra. *Bioinformatics* **18**, 211–212
 38. Whitmore, L., and Wallace, B. A. (2004) DICHROWEB, an online server for protein secondary structure analyses from circular dichroism spectroscopic data. *Nucleic Acids Res.* **32**, W668–W673
 39. Soulages, J. L., Kim, K., Walters, C., and Cushman, J. C. (2002) Temperature-induced extended helix/random coil transitions in a group 1 late embryogenesis-abundant protein from soybean. *Plant Physiol.* **128**, 822–832
 40. Buck, M. (1998) Trifluoroethanol and colleagues: cosolvents come of age: recent studies with peptides and proteins. *Q. Rev. Biophys.* **31**, 297–355
 41. Luo, P., and Baldwin, R. L. (1997) Mechanism of helix induction by trifluoroethanol: a framework for extrapolating the helix-forming properties of peptides from trifluoroethanol/water mixtures back to water. *Biochemistry* **36**, 8413–8421
 42. Ellis, R. J. (2001) Macromolecular crowding: an important but neglected aspect of the intracellular environment. *Curr. Opin. Struct. Biol.* **11**, 114–119
 43. Zimmerman, S. B., and Minton, A. P. (1993) Macromolecular crowding: biochemical, biophysical, and physiological consequences. *Annu. Rev. Biophys. Biomed. Struct.* **22**, 27–65
 44. Zimmerman, S. B., and Trach, S. O. (1991) Estimation of macromolecule concentrations and excluded volume effects for the cytoplasm of *Escherichia coli*. *J. Mol. Biol.* **222**, 599–620
 45. Minton, A. P. (2000) Implications of macromolecular crowding for protein assembly. *Curr. Opin. Struct. Biol.* **10**, 34–39
 46. Minton, A. P. (1997) Influence of excluded volume upon macromolecular structure and associations in “crowded” media. *Curr. Opin. Biotech.* **8**, 65–69
 47. Campos, F., Zamudio, F., and Covarrubias, A. A. (2006) Two different late embryogenesis abundant proteins from *Arabidopsis thaliana* contain specific domains that inhibit *Escherichia coli* growth. *Biochem. Biophys. Res. Commun.* **342**, 406–413
 48. Wright, P. E., and Dyson, H. J. (2015) Intrinsically disordered proteins in cellular signalling and regulation. *Nat. Rev. Mol. Cell Biol.* **16**, 18–29
 49. Mouillon, J. M., Eriksson, S. K., and Harryson, P. (2008) Mimicking the plant cell interior under water stress by macromolecular crowding: disordered dehydrin proteins are highly resistant to structural collapse. *Plant Physiol.* **148**, 1925–1937
 50. Szasz, C. S., Alexa, A., Toth, K., Rakacs, M., Langowski, J., and Tompa, P. (2011) Protein disorder prevails under crowded conditions. *Biochemistry* **50**, 5834–5844
 51. Haaning, S., Radutoiu, S., Hoffmann, S. V., Dittmer, J., Giehm, L., Otzen, D. E., and Stougaard, J. (2008) An unusual intrinsically disordered protein from the model legume *Lotus japonicus* stabilizes proteins *in vitro*. *J. Biol. Chem.* **283**, 31142–31152
 52. McCubbin, W. D., Kay, C. M., and Lane, B. G. (1985) Hydrodynamic and optical-properties of the wheat-germ Em protein. *Can. J. Biochem. Cell Biol.* **63**, 803–811
 53. Cuevas-Velazquez, C. L., Rendón-Luna, D. F., and Covarrubias, A. A. (2014) Dissecting the cryoprotection mechanisms for dehydrins. *Front. Plant Sci.* **5**, 583
 54. Olvera-Carrillo, Y., Luis Reyes, J., and Covarrubias, A. A. (2011) Late embryogenesis abundant proteins: versatile players in the plant adaptation to water limiting environments. *Plant Signal. Behav.* **6**, 586–589
 55. Tompa, P., and Kovacs, D. (2010) Intrinsically disordered chaperones in plants and animals. *Biochem. Cell Biol.* **88**, 167–174
 56. Lee, S. H., Kim, D. H., Han, J. J., Cha, E. J., Lim, J. E., Cho, Y. J., Lee, C., and Han, K. H. (2012) Understanding pre-structured motifs (PreSMos) in intrinsically unfolded proteins. *Curr. Protein Pept. Sci.* **13**, 34–54
 57. Cino, E. A., Karttunen, M., and Choy, W. Y. (2012) Effects of molecular crowding on the dynamics of intrinsically disordered proteins. *PLoS One* **7**, e49876
 58. Mittag, T., Kay, L. E., and Forman-Kay, J. D. (2010) Protein dynamics and conformational disorder in molecular recognition. *J. Mol. Recognit.* **23**, 105–116
 59. Mittag, T., Orlicky, S., Choy, W. Y., Tang, X., Lin, H., Sicheri, F., Kay, L. E., Tyers, M., and Forman-Kay, J. D. (2008) Dynamic equilibrium engagement of a polyvalent ligand with a single-site receptor. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 17772–17777
 60. Uversky, V. N. (2013) Unusual biophysics of intrinsically disordered proteins. *Biochim. Biophys. Acta* **1834**, 932–951