



HHS Public Access

Author manuscript

Am Econ Rev. Author manuscript; available in PMC 2016 May 17.

Published in final edited form as:

Am Econ Rev. 2015 May ; 105(5): 491–495. doi:10.1257/aer.p20151023.

Prediction Policy Problems

Jon Kleinberg,
Cornell

Jens Ludwig,
University of Chicago and NBER

Sendhil Mullainathan, and
Harvard and NBER

Ziad Obermeyer*
Harvard Medical School and Brigham and Women's Hospital

Empirical policy research often focuses on causal inference. Since policy choices seem to depend on understanding the counterfactual—what happens with and without a policy—this tight link of causality and policy seems natural. While this link holds in many cases, we argue that there are also many policy applications where causal inference is not central, or even necessary.

Consider two toy examples. One policy maker facing a drought must decide whether to invest in a rain dance to increase the chance of rain. Another seeing clouds must decide whether to take an umbrella to work to avoid getting wet on the way home? Both decisions could benefit from an empirical duty of rain. But each has different requirements of the estimator. One requires causality: do rain dances cause rain? The other does not, needing only prediction: is the chance of rain high enough to merit an umbrella? We often focus on rain dance like policy problems. But there are many important policy problems umbrella-like. Not only are these prediction problems neglected, machine learning can help us solve them more effectively.

In this paper, we (i) provide a simple framework that clarifies the distinction between causation and prediction; (ii) explain how machine learning adds value over traditional regression approaches in solving prediction problems; (iii) provide an empirical example from health policy to illustrate how improved predictions can generate large social impact; (iv) illustrate how “umbrella” problems are common and important in many important policy domains; and (v) argue that solving these problems produces not just policy impact but also theoretical and economic insights.¹

*Obermeyer: zobermeyer@partners.org.

¹A longer version of this paper—Kleinberg et al. (2015a)—fleshes out each of these points, providing greater detail on the model, the empirical work and a more thorough summary of machine learning.

I. Prediction and Causation

Let Y be an outcome variable (such as rain) which depends in an unknown way on a set of variables X_0 and X . A policy-maker must decide on X_0 (e.g. an umbrella or rain-dance) in order to maximize a (known) payoff function $\pi(X_0, Y)$. Our decision of X_0 depends on the derivative

$$\frac{d\pi(X_0, Y)}{dX_0} = \frac{\partial\pi}{\partial X_0} \underbrace{(Y)}_{\text{prediction}} + \frac{\partial\pi}{\partial Y} \underbrace{\frac{\partial Y}{\partial X_0}}_{\text{causation}}$$

Empirical work can help estimate the two unknowns in this equation: $\frac{\partial Y}{\partial X_0}$ and $\frac{\partial\pi}{\partial X_0}$.

Estimating $\frac{\partial Y}{\partial X_0}$ requires causal inference: answering how much does X_0 affect Y ?

The other term— $\frac{\partial\pi}{\partial X_0}$ —is unknown for a different reason. We know the payoff function but

since its value must be evaluated at Y , knowing the exact value of $\frac{\partial\pi}{\partial X_0}$ requires a prediction Y . We know how much utility umbrellas provide only once we know the level of rain.

Choosing X_0 therefore requires solving both causation and prediction problems. Assume away one of these terms—place an exclusion restriction—and only one problem remains. Rain dances are a pure causal inference problem because rain dances have no direct effect on

payoffs $\frac{\partial\pi}{\partial X_0} = 0$. Umbrellas are a pure prediction problem because umbrellas have no direct effect on rain $\frac{\partial Y}{\partial X_0} = 0$.

This derivative also illustrates two key features of prediction problems. First, the need for

prediction arises exactly because $\frac{\partial\pi}{\partial X_0}$ depends on Y . Prediction is necessary only because the benefit of an umbrella depends on rain. As we illustrate in the final section, this kind of dependency is common for many important policy problems. Second, because only \hat{Y} enters the decision, prediction problems only require low error in \hat{Y} ; they do not require the coefficients to be unbiased or causal.

II. Machine Learning

As a result, standard empirical techniques are not optimized for prediction problems because they focus on unbiasedness. Ordinary least squares (OLS), for example, is only the best linear *unbiased* estimator. To see how it can lead to poor predictions, consider a two variable example where OLS estimation produced $\hat{\beta}_1 = 1 \pm .001$ and $\hat{\beta}_2 = 4 \pm 10$, suggesting a predictor of $x_1 + 4x_2$. But given the noise in $\hat{\beta}_2$, for prediction purposes one would be

tempted to place a smaller (possibly 0) coefficient on x_2 . Introducing this bias could improve prediction by removing noise.

This intuition holds more generally. Suppose we are given a data set D of n points $(y^i, x^i) \sim G$. We must use this data to pick a function $\hat{f} \in \mathcal{F}$ so as to predict the y value of a new data point $(y, x) \sim G$. The goal is to minimize a loss function, which for simplicity we take to be $(y - \hat{f})(x)^2$.

OLS minimizes in-sample error, choosing from \mathcal{F}_{lin} , the set of linear estimators:

$$\hat{f}_{OLS} = \operatorname{argmin}_{f \in \mathcal{F}_{lin}} \sum_{i=1}^n (y^i - f(x^i))^2$$

but for prediction we are not interested in doing well *in sample*: we would like to do well *out of sample*. Ensuring zero bias in-sample creates problems out of sample. To see this,

consider the mean squared error at the new point x , $MSE(x) = E_D \left[(\hat{f}(x) - y)^2 \right]$. This can be decomposed as:

$$E_D \left[(\hat{f}(x) - E_D[\hat{y}_0])^2 \right] + \underbrace{(E_D[\hat{y}_0] - y)^2}_{\text{Bias}}$$

Variance

Because the f varies from sample to sample, it produces variance (the first term). This must be traded off against bias (the first term). By ensuring zero bias, OLS allows no tradeoff.

Machine learning techniques were developed specifically to maximize prediction performance by providing an empirical way to make this bias-variance trade-off (Hastie et al., 2009 provide a useful overview). Instead of minimizing *only* in-sample error, ML techniques minimize:

$$\hat{f}_{ML} = \operatorname{argmin}_{f \in \mathcal{F}} \sum_{i=1}^n (y^i - f(x^i))^2 + \lambda R(f)$$

Here $R(f)$ is a *regularizer* that penalizes functions that create variance. It is constructed such the set of functions $\mathcal{F}_c = \{f | R(f) \leq c\}$ create more variable predictions as c increases. For linear models, larger coefficients allow more variable predictions, so a natural regularizer is $R(f_\beta) = \|\beta\|_d$, which is the LASSO and RIDGE estimators for $d=1$ and 2 respectively. In effect, this minimization now explicitly includes a bias (in-sample error) and variance term ($R(f)$), where λ can be thought of as the price at which we trade off variance to bias. OLS is a special case where we put an infinite (relative) price on bias ($\frac{1}{\lambda} = \infty$).

A key insight of machine learning is that this price λ can be chosen *using the data itself*. Imagine we split the data into f subsets (often called “folds”). For a set of λ , we estimate the

algorithm on $f-1$ of the folds and then see which value of λ produces the best prediction in the f^{th} fold. This cross-validation procedure effectively simulates the bias-variance tradeoff by creating a way to see which λ does best “out of sample”.

These two insights—regularization and empirical choice of the regularization penalty—together also change the kinds of predictors we can consider. First, they allow for “wide” data, to predict even when we have more variables than data points. For example, researchers using language data often have an ten or a hundred times as many variables as data. Second, this allows for far more flexible functional forms. One can include many higher order interaction terms or use techniques such as decision trees which by construction allow for a high degree of interactivity.

Machine learning techniques are in one sense not new: they are a natural offshoot of non-parametric statistics. But they provide a disciplined way to predict \hat{y} which (i) uses the data itself to decide how to make the bias-variance tradeoff and (ii) allows for search over very rich set of variables and functional forms. But everything comes at a cost: one must always keep in mind that because they are tuned for \hat{y} they do not (without many other assumptions) give very useful guarantees for $\hat{\beta}$.

III. Illustrative Application

Osteoarthritis (joint pain and stiffness) is a common and painful chronic illness among the elderly. Replacement of the affected joints, most commonly hips and knees, provide relief each year to around 500,000 Medicare beneficiaries in the US. The medical benefits B are well understood: surgery improves quality of life over the patient’s remaining life expectancy Y . The costs C are both monetary (roughly \$15,000 calculated using 2010 claims data) and non-monetary: surgeries are painful and recovery takes time, with significant disability persisting months afterwards. The benefits accrue over time, so surgery only make sense if someone lives long enough to enjoy them; joint replacement for someone who dies soon afterward is futile—a waste of money and an unnecessary painful imposition on the last few months of life.

The payoff to surgery depends on (eventual) mortality, creating a pure prediction problem. Put differently, the policy challenge is: can we predict which surgeries will be futile using only data available at the time of the surgery? This would allow us save both dollars and disutility for patients.

To study this we example a 20% percent sample of 7.4 million Medicare beneficiaries, 98,090 (1.3%) of which had a claim for hip or knee replacement surgery in 2010.² Of these, 1.4% die in the month after surgery, potentially from complications of the surgery itself, and 4.2% die in the 1–12 months after surgery. This low rate—roughly the average annual mortality rate for all Medicare recipients—seems to suggest on average surgeries are not futile. But the average is misleading because the policy decision is really about whether surgeries on the *predictably riskiest patients* was futile.

²We restricted to fee-for-service beneficiaries with full claims data living in the continental US, and exclude any with joint replacement in 2009 (potentially implying revision surgery or other complication of a prior procedure).

To answer this, we predicted mortality in the 1–12 months after hip or knee replacement using LASSO (see Kleinberg et al., 2015a for full details).³ We used 65,395 observations to fit the models and measured performance on the remaining 32,695 observations. 3,305 independent variables were constructed using Medicare claims dated prior to joint replacement, including patient demographics (age, sex, geography); co-morbidities, symptoms, injuries, acute conditions and their evolution over time; and health-care utilization.

These predictions give us a way to isolate predictably futile surgeries. In Table 1, we sort beneficiaries by predicted mortality risk, showing risk for the riskiest 1%, 2% and so on, *which is highly and predictably concentrated*: for example, the 1% riskiest have a 56% mortality rate, and account for fully 10% of all futile surgeries.⁴

Imagine the dollars from these futile surgeries could instead have been spent on other beneficiaries who would benefit more. To understand the potential savings, we simulated the effect of substituting these riskiest recipients with other beneficiaries who might have benefited from joint replacement procedures under Medicare eligibility guidelines, but did not receive them. To be conservative, rather than comparing to the lowest-risk eligibles, we draw from the median predicted risk distribution of these eligibles, and simulate effects of this replacement in columns (3) and (4). Replacing the riskiest 10th percentile with lower-risk eligibles would avert 10,512 futile surgeries and reallocate the 158 million per year (if applied to the entire Medicare population) to people who benefit from the surgery, at the cost of postponing joint replacement for 38,533 of the riskiest beneficiaries who would not have died.⁵

IV. Prediction Problems Are Common and Important

Our empirical application above highlights how improved prediction using machine learning techniques can have large policy impacts (much like solving causal inference problems have had). There are many other examples as well. In the criminal justice system, for instance, judges have to decide whether to detain or release arrestees as they await adjudication of their case—a decision that depends on a prediction about the arrestee’s probability of committing a crime. Kleinberg et al. (2015b) show that machine learning techniques can dramatically improve upon judges’ predictions and substantially reduce the amount of crime.

Other illustrative examples include: (i) in education, predicting which teacher will have the greatest value add (Rockoff et al., 2011); (ii) in labor market policy, predicting unemployment spell length to help workers decide on savings rates and job search strategies;

³This interval reflects two choices. 1) We excluded deaths in the first month after surgery to focus on prediction of Y rather than the short-term causal effect of X_0 on Y (i.e., operative risk, post-surgical complications). 2) We chose a threshold of 12 months based on studies showing substantial remaining disability 6 months after surgery, but improved clinical outcomes at the 12-month mark (Hamel et al., 2008). Alternatively, a ‘break-even’ threshold could be derived empirically.

⁴One might wonder whether these riskier patients may also be the ones who also stood to benefit the most from the procedure, potentially justifying surgery. However, variables that should correlate with surgery benefit (number of physician visits for hip or knee pain, physical therapy and therapeutic joint injections) do not vary significantly by predicted mortality risk.

⁵The existence of a large pool of low-risk beneficiaries potentially eligible for replacement argues against moral hazard as an explanation for these findings, since physicians who predicted well acting consistent with moral hazard would first exhaust the low-risk pool of patients before operating on higher-risk patients.

(iii) in regulation, targeting health inspections (Kang et al. 2013); (iv) in social policy, predicting highest risk youth for targeting interventions (Chandler et al., 2011); and (v) in the finance sector, lenders identifying the underlying creditworthiness of potential borrowers.

Even this small set of examples are biased by what we *imagine* to be predictable. Some things that seem unpredictable may actually be more predictable than we think using the right empirical tools. As we expand our notion of what is predictable, new applications will arise.

Prediction problems can also generate theoretical insights, for example by changing our understanding of an area. Our empirical application above shows that low-value care is not due just to the standard moral-hazard explanation of health economics but also to mis-prediction. The pattern of discrepancies between human and algorithmic decisions can serve as a behavioral diagnostic about decision making (Kleinberg et al. 2015b). And prediction can shed light on other theoretical issues. For example understanding how people change their behavior as regulators or police change the algorithms they use to target monitoring effort can shed light on the game theory of enforcement.

Prediction policy problems are, in sum, important, common, and interesting, and deserve much more attention from economists than they have received. New advances in machine learning can be adapted by economists to work on these problems, but will require a substantial amount of both theoretical and practical reorientation to yield benefits for those currently engaged in policy studies.

Acknowledgments

For financial support we thank operating grants to the University of Chicago Crime Lab by the MacArthur and McCormick foundations and the National Institutes of Health Common Fund grant DP5 OD012161 to Ziad Obermeyer. Thanks to Marianne Bertrand, Kerwin Charles and Elizabeth Santorella for helpful comments and Brent Cohn, Maggie Makar, and Matt Repka for extremely helpful research assistance. Any errors and all opinions are of course ours alone.

References

- Chandler, Dana, Levitt, Steven, List, John. Predicting and preventing shootings among at-risk youth. *American Economic Review*. 2011; 101(3):288–92.
- Hamel, Mary Beth, Toth, Maria, Legedza, Anna, Rose, Max. Joint replacement surgery in elderly patients with severe osteoarthritis of the hip or knee. *Archives of Internal Medicine*. 2008; 168(13): 1430–40. [PubMed: 18625924]
- Hastie, Trevor, Tibshirani, Robert, Friedman, Jerome. *The Elements of Statistical Learning: Data mining, Inference, and Prediction*. 2nd. Springer-Verlag; 2009.
- Kang, Jun Seok, Kuznetsova, Polina, Choi, Yejin, Luca, Michael. Where not to eat? Improving public policy by predicting hygiene inspections using online reviews. *Empirical methods in natural language processing*. 2013; 22:14–030. 58–77.
- Kleinberg, Jon, Ludwig, Jens, Mullainathan, Sendhil, Obermeyer, Ziad. *Policy prediction problems*. Cambridge, MA: 2015a. NBER Working Paper
- Kleinberg, Jon, Lakkaraju, Himabindu, Leskovec, Jure, Ludwig, Jens, Mullainathan, Sendhil. *Human and machine predictions: An application to bail decisions*. Cambridge, MA: 2015b. NBER Working Paper

Rockoff, Jonah, Jacob, Brian A., Kane, Thomas J., Staiger, Douglas O. Can you recognize an effective teacher when you recruit one? *Education Finance and Policy*. 2011; 6(1):43–74.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Riskiest Joint Replacements

Predicted Mortality Percentile	Observed Mortality Rate	Futile Procedures Averted	Futile Spending (\$ mill.)
1	0.435 (.028)	1984	30
2	0.422 (.028)	3844	58
5	0.358 (.027)	8061	121
10	0.242 (.024)	10512	158
20	0.152 (.020)	12317	185
30	0.136 (.019)	16151	242

Note:

1. We predict 1–12 month mortality using an L_1 regularized logistic regression trained on 65,395 Medicare beneficiaries undergoing joint replacement in 2010, using 3,305 claims-based variables and 51 state indicators. λ was tuned using 10-fold cross-validation in the training set. In columns (1) and (2) we sort a hold-out set of 32,695 by predicted risk into percentiles (column 1) and calculate actual 1–12 month mortality (column 2).
2. Columns (3) and (4) show results of a simulation exercise: we identify a population of eligibles (using published Medicare guidelines: those who had multiple visits to physicians for osteoarthritis and multiple claims for physical therapy or therapeutic joint injections) who did not receive replacement and assign them a predicted risk. We then substitute the high risk surgeries in each row with patients from this eligible distribution for replacement, starting at *median* predicted risk. Column (3) counts the futile procedures averted (i.e., replaced with non-futile procedures) and (4) quantifies the dollars saved in millions by this substitution.