



Published in final edited form as:

J Hepatol. 2016 June ; 64(6): 1378–1387. doi:10.1016/j.jhep.2016.01.035.

The pro-inflammatory effects of miR-155 promote liver fibrosis and alcohol-induced steatohepatitis

Shashi Bala¹, Timea Csak^{1,2}, Banishree Saha¹, James Zatsiorsky¹, Karen Kodys¹, Donna Catalano¹, and Gyongyi Szabo¹

¹Department of Medicine, University of Massachusetts Medical School, Worcester, 01604, Massachusetts, USA

²Brookdale University Hospital and Medical Center, Brooklyn, New York, USA

Abstract

Background & Aims—Alcoholic liver disease (ALD) ranges from fatty liver to inflammation and cirrhosis. miRNA-155 is an important regulator of inflammation. In this study, we describe the in vivo role of miR-155 in ALD.

Methods—Wild type, WT (C57/BL6J) or miR-155 KO and TLR4 KO mice received Lieber-DeCarli diet for 5 weeks. Some mice received corn oil or CCl₄ for 2 or 9 weeks.

Results—We found that miR-155 KO mice are protected from alcohol-induced steatosis and inflammation. The reduction in alcohol-induced fat accumulation in miR-155 KO mice was associated with increased PPRE and PPAR α (miR-155 target) binding and decreased MCP1 production. Treatment with a miR-155 inhibitor increased PPAR γ expression in naive and alcohol treated RAW macrophages. Alcohol increased lipid metabolism genes (FABP4, LXR α , ACC1 and LDLR) in WT mice and this was prevented in KO mice. Alcohol diet induced increase in the number of CD163⁺CD206⁺ infiltrating macrophages and neutrophils in WT was prevented in miR-155 KO mice. Kupffer cells isolated from miR-155 KO mice exhibited predominance of M2 phenotype when exposed to M1 polarized signals and this was due to increased C/EBP β . Profibrotic genes were attenuated in miR-155 KO mice after alcohol diet or CCl₄ treatment. Compared to WT attenuation in CCl₄ induced hydroxyproline and α SMA was observed in KO mice. Finally, we show TLR4 signaling regulates miR-155 as TLR4 KO mice showed no induction of miR-155 after alcohol diet.

Conclusions—Collectively our results demonstrated the role of miR-155 in alcohol-induced steatohepatitis and fibrosis in vivo.

Corresponding Author: Gyongyi Szabo MD, PhD University of Massachusetts Medical School; Department of Medicine, LRB215, 364 Plantation Street, Worcester, MA 01605; Tel: 00-1-508-856-5275, Fax: 00-1-508-856-4770, ; Email: gyongyi.szabo@umassmed.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of interest: None

Author's contributions: Idea conceived and design the experiments: SB, GS. Performed and analyzed the experiments: SB, TC, BS, KK, DC, JZ. Interpreted the data and wrote the paper: SB, and GS. All authors approved the final manuscript.

Keywords

microRNA; inflammation; alcohol; PPAR α ; PPAR γ ; fibrosis

INTRODUCTION

Alcoholic liver disease (ALD) is one of the most common causes of liver diseases in the world [1]. The clinical manifestations of ALD range from simple fatty liver to more severe forms of liver injury, including alcoholic hepatitis, cirrhosis, and hepatocellular carcinoma. Alcoholic hepatitis has high mortality and morbidity with limited treatment options. The pathomechanism of ALD involves complex interactions between the effects of alcohol and its toxic metabolites on various cell types in the liver and gut, induction of reactive oxygen species, and up-regulation of the inflammatory cascade [2–5]. Direct toxic effects of alcohol on hepatocytes, increased intestinal permeability and activation of liver macrophages (Kupffer cells; KCs) by gut-derived lipopolysaccharide (LPS) are major factors contributing to ALD [1–3, 5]. LPS is recognized by the Toll-like receptor (TLR) 4 complex expressed on immune cells as well as parenchymal cells and induces pro-inflammatory cytokine activation [5]. Chronic alcohol exposure also sensitizes liver resident macrophages (KCs) to LPS-induced inflammatory cytokine production [6].

Fatty liver (steatosis) is the first stage of response in the liver to binge drinking or chronic ethanol consumption. Accumulation of lipid products such as triglycerides in hepatocytes leads to lipid superoxidation and oxidative stress, resulting in apoptosis, and hepatic inflammation [7]. microRNAs (miRNAs) are shown to be modulated by alcohol in the liver and in hepatocytes to exert inflammatory and fat accumulation effects [6, 8–10]. Among other miRNAs, miR-155 is a major regulator of immune responses [11]. Various TLR ligands are shown to induce miR-155 in macrophages [11]. Previously we identified miR-155 as an alcohol-induced regulator of increased KC activation and TNF α production in macrophages [6]. In recent years, the role of miR-155 in other cellular processes like fatty acid metabolism and fibrogenic events has also been emerging [12–14]. In a methionine choline-deficient (MCD) diet induced steatohepatitis, we showed that miR-155 targets genes involved in lipid metabolism (Fab4, cpt1a) and early fibrosis (C/EBP β , Smad3) [12].

We earlier demonstrated that miR-155 was increased in both KCs and hepatocytes isolated from alcohol-fed mice. Therefore we hypothesized that miR-155 has a role in alcohol-induced liver injury, inflammation and steatosis. To test this hypothesis, we determined the in vivo role of miR-155 in a mouse model of ALD. Our results revealed that alcohol-induced liver injury, steatosis and inflammation were significantly reduced in miR-155 KO mice compared to wild-type (WT) mice. The reduction in fat accumulation in miR-155 KO mice was associated with increased PPRE and PPAR α binding after alcohol diet. The alcohol-induced increase in neutrophil leukocyte and CD163⁺CD206⁺ (M2) macrophage (profibrotic) infiltration observed in WT mice was prevented in miR-155 KO mice. miR-155 KO mice also showed attenuation in pro-fibrotic genes after alcohol diet as well as in a CCl₄-induced liver fibrosis. Together, these results provide in vivo evidence for the role of miR-155 in ALD and fibrosis.

MATERIALS AND METHODS

Animal studies

Animal studies were approved by the University of Massachusetts Medical School (UMMS) Institutional Animal Use and Care Committee (Worcester, MA). miR-155 KO mice were purchased from Jackson Laboratory (Bar Harbor, Maine, USA) and a breeding colony was maintained in the animal facility of UMMS. Wild type (WT) control mice, C56BL/6 were also obtained from Jackson Laboratory. For chronic ethanol feeding, female mice (n=8–10, 8 weeks old) received 5% (v/v) ethanol (36% ethanol-derived calories) containing Lieber-DeCarli diet or control (pair-fed) diet for 5 weeks. For pair-fed diet, alcohol-derived calories were substituted with dextrin-maltose (Bio-Serv, New Jersey, USA) as described previously [6]. For CCl₄ treatment, WT or miR-155 KO male mice (n=6, 8–10 weeks old) were treated either with corn oil (vehicle control) or CCl₄ (0.6ml/kg i.p.) for 2 or 9 weeks, as described [15]. Mice were sacrificed 72h after last CCl₄ injection. At the end of the experiment, blood was collected by cheek bleeding and mice were sacrificed [6]. Liver tissue was collected and snap frozen for proteins and in RNA later (Qiagen, Germany) for RNA extraction. All samples were stored at –80°C.

Liver cell isolation and Flow cytometry

Primary murine hepatocytes, Kupffer cells (KCs) and liver mononuclear cells (MNCs) were isolated from WT chow fed mice by an enzyme-based tissue digestion method as described [16]. Hepatocytes were plated onto 6 well collagen-coated plates and free-floating cells were removed 3h after plating. Both hepatocytes and MNCs were treated with LPS (100ng/ml) for 6h and processed for RNA extraction. Some hepatocytes were treated with MCP1 (100ng/ml) for 36h and nuclear proteins were used for PPRE binding. KCs isolated from WT or KO mice were treated with or without LPS (100ng/ml), IL-4 (20ng/ml) and IFN γ (20ng/ml) for 18h.

For flow cytometry, MNCs were isolated from perfused livers of WT and miR-155 KO mice after 5 weeks of alcohol diet and processed as described [17]. For cell viability, Live/Dead Fixable Blue Dead staining was used (Life Technologies, NY, USA). Briefly, after blocking for non-specific binding to Fc γ Rs, MNCs ($\sim 10^6$ cells) were stained with CD163, CD11b-FITC, F4/80 PerCP-Cy5.5, and CD206-Alexa 647 (BioLegend, CA, USA) followed by incubation with PE-conjugated secondary antibody for CD163 as described [17]. For neutrophils, the percentage of CD11b⁺ PE and Ly6G^{high} – FITC was determined. Respective isotype-matched control antibodies were used (BioLegend). After washing with FACS buffer, cells were fixed with 1% paraformaldehyde and acquired on a BD LSR II instrument (BD Biosciences, CA, USA). Data was analyzed with FlowJo software (OR, USA).

Additional methods are in the supporting material.

Statistical analysis

Statistical significance was determined using the non-parametric Mann-Whitney test. Data is represented as mean \pm standard error and considered statistically significant at p<0.05.

Results

miR-155 deficiency attenuates chronic alcohol-induced steatosis, liver injury, and oxidative stress in the liver

Inflammation, a major component in alcoholic liver disease, contributes to the perpetuation of liver disease and amplifies steatosis [1]. In our earlier study, we showed induction of miR-155 in the liver, isolated Kupffer cells and hepatocytes from alcohol fed mice [6], however, the functional role of miR-155 in ALD in vivo is unknown. In this study, we determined the role of miRNA-155 using a miR-155 KO mouse model system. Alcohol feeding for 5 weeks in WT mice resulted in hepatic steatosis as evidenced by H&E (Fig.1A), Oil-O-Red staining (Fig.1B), histological scores (Fig.1C) and liver triglyceride levels (Fig. 1D). Compared to WT mice, a significant decrease was observed in fat accumulation in miR-155 KO mice after alcohol feeding (Fig.1A–D). In miR-155 KO mice we found a significant decrease in plasma ALT, a marker of liver injury, compared to WT mice after alcohol feeding (Fig.1E). Alcohol and its metabolites increase the production of reactive oxygen species and enhance peroxidation of lipids, proteins, and DNA [1]. We found that the increased oxidative stress (measured by thiobarbituric acid reactive substances assay) observed in WT mice was prevented in miR-155 KO mice after alcohol diet (Fig.1F), indicating that miR-155 deficiency attenuates alcohol-induced oxidative damage.

Alcohol diet results in increased PPRE and PPAR α DNA binding in miR-155 KO mice

Because we found a decrease in liver fat accumulation in miR-155 KO mice after alcohol diet (Fig.1), we sought to evaluate possible mechanisms involved. Lipid and glucose homeostasis are regulated by peroxisome proliferator-activated receptors (PPARs) that belong to the nuclear hormone receptor superfamily [7, 18]. There are three PPAR subtypes: PPAR α , PPAR γ and PPAR δ [7, 18]. PPAR α regulates alcoholic steatosis via its effects on genes of lipid metabolism, and a decrease in PPAR α has been reported in mouse models of ALD [7, 18, 19]. We extensively performed bioinformatics analysis and found that miR-155 has a seed region at the 3'UTR of PPAR γ gene (www.microrna.org) (Supplementary Fig. 1A). A very recent study identified PPAR α , a direct target of miR-155 [20] and showed that miR-155 regulates inflammation via targeting PPAR α after inducing diffuse alveolar hemorrhage. We found a decrease in PPAR α mRNA after alcohol diet in WT mice and this decrease was prevented in miR-155 KO mice (Fig.2A). We didn't find a significant change in the PPAR γ mRNA after alcohol diet in WT mice, whereas miR-155 KO mice showed increased transcription of PPAR γ (Supplementary Fig.1B). Next we examined peroxisome proliferator-activated receptor response element (PPRE) binding using EMSA. We found a decrease in PPRE binding in WT mice after alcohol diet compared to pair-fed mice (Fig.2B–C). In contrast, alcohol feeding resulted in no decrease in PPRE binding in miR-155 KO mice (Fig.2B–C). PPRE binding was similar between genotypes on PF diet. To determine the binding of PPAR α in miR-155 KO mice after alcohol diet, we performed a supershift assay. Our results (Fig.2B) indicate a decrease in PPAR α binding in alcohol-fed WT mice compared to PF diet fed mice (Fig.2B). Further, we found an increase in PPAR α binding in miR-155 KO mice compared to WT mice after alcohol diet (Fig.2B–C).

To further dissect the role of miR-155 in hepatocytes, we isolated hepatocytes from WT and miR-155 KO mice and treated them with or without MCP1 (100ng/ml) to induce steatosis and evaluated for PPRE binding (Fig.2D). MCP1 treatment resulted in a decrease in PPRE binding in hepatocytes from WT mice, however this decrease was prevented in hepatocytes isolated from miR-155 KO mice (Fig.2D), suggesting a role of miR-155 in steatosis via regulation of PPRE signaling. PPAR γ is highly expressed in macrophages and low in hepatocytes. Since we found that miR-155 has a seed region at the 3'UTR of PPAR γ gene therefore we sought to determine if PPAR γ is a direct target of miR-155 using RAW 264.7 macrophages. Inhibition of miR-155 using a specific miR-155 inhibitor resulted in an increase in PPAR γ gene both in naïve and alcohol treated cells (Fig.2E), suggesting that miR-155 regulates PPAR γ in macrophages.

Since we found that miR-155 regulates PPRE and PPAR γ , we checked for PPAR γ and PPAR α target genes involved in lipid metabolism and fatty acid uptake. Genes such as fatty acid binding protein 4 (FABP4), lipogenesis (LXR α), acetyl-CoA carboxylase 1 [ACC1] and lipid binding transport (LDL receptor) were checked. We found that alcohol induced FABP4 (Fig.2F), LXR α (Fig.2G), ACC1 (Fig.2H), and LDLR (Fig.2I) in the livers of WT mice and not in miR-155 KO mice. Alcohol induced HMGCR mRNA in WT mice and this was prevented in miR-155 KO mice (Fig.2J). No significant changes were observed in genes involved in glucose homeostasis (PCK1 and CEBP α) between genotypes (Supple Fig.1C–D). Other PPAR α target genes such as carnitine palmitoyltransferase 1A (CPT1A) and fatty acid synthase (FAS) were also evaluated. We did not any significant differences between genotypes in these genes after alcohol diet (Supple Fig.1E–F).

Recently MCP1 was shown to promote hepatic steatosis [19, 21]. Therefore we assessed MCP1 expression and found that alcohol increased MCP1 mRNA in WT mice but not in miR-155 KO mice (Fig.2K). After an LPS challenge, alcohol fed WT mice showed significant MCP1 production (Fig.2L and M), and a lower MCP1 induction was present in alcohol fed miR-155 KO (Fig.2L and M). Collectively our results suggest that steatogenic, changes observed in miR-155 KO mice after ethanol diet is due to its effect on PPRE signaling.

Chronic alcohol-induced inflammation and macrophage and neutrophil infiltration are attenuated in miR-155 KO mice

miR-155 regulates genes involved in inflammation [11], and alcohol use causes inflammation, therefore, next we evaluated the markers of inflammation. Chronic alcohol feeding increased TNF α (Fig.3A) and IL-1 β (Fig.3B) protein production in WT mice compared to mice on PF diet. An LPS challenge further increased these cytokines in alcohol-fed WT mice, however, the LPS induced increase in TNF α and IL-1 β proteins was attenuated in alcohol-fed miR-155 KO mice (Fig.3A and B). No significant difference was found in the production of these cytokines between the genotypes that were on PF diet after LPS treatment, supporting a role for miR-155 in the alcohol-induced effects. Macrophage activation markers (CD68 and MIP2) were increased in WT mice, after alcohol feeding, these increases were prevented in miR-155 KO mice (Fig.3C and D).

Recently we showed that alcohol induces both M1 (classical activation) and M2 (alternatively activated profibrotic) phenotype of macrophages in mice [17]. In this context, miR-155 was shown to promote the M1 phenotype of bone marrow derived macrophages [22]. To determine if deficiency of miR-155 KO have any effect on macrophage phenotype after alcohol diet, we performed flow cytometry analysis of the liver immune cells. We found that chronic alcohol diet increased the number of inflammatory monocytes Cd11b⁺ Ly6C⁺ in WT mice, while in miR-155 KO mice alcohol moderately decreased the number of inflammatory monocytes (Fig.3E). Interestingly, alcohol diet also increased the percentage of F4/80⁺ CD163⁺CD206⁺ M2 macrophages in WT mice, but no increase was found in miR-155 KO mice (Fig.3F). miR-155 KO mice on PF diet showed increased % of M2 macrophages. To evaluate how miR-155 is involved in KCs polarization, we isolated KCs from WT and miR-155 KO mice (on chow fed diet) and polarized to M1 and M2 phenotype. As M2 markers, we evaluated for Arginase1, IL-10 and MRC1 levels. LPS or INF γ treatment decreased and IL-4 treatment increased Arginase1 levels in KCs isolated from WT mice (Fig.3G). Compared to WT, KCs isolated from miR-155 KO mice showed a higher increase in Arginase1 levels after LPS or INF γ treatment (Fig.3G). KCs isolated from miR-155 KO mice showed a significant increase in IL-10 mRNA compared to KCs isolated from WT mice after LPS treatment (Fig.3H). No significant differences in MRC1 levels were found between genotypes (Fig.3I). As M1 markers we tested for IL-1 β and iNOS genes. No significant differences were found in IL-1 β mRNA between genotypes after LPS treatment (Fig.3J). Surprisingly, KCs isolated from miR-155 KO mice showed a significant increase in iNOS mRNA compared to WT KCs after LPS or INF γ or IL-4 treatment (Fig. 3K). To determine the possible explanation for this mixed phenotype/polarization, we checked for transcriptional regulators of macrophage polarization.

miR-155 is involved in modulating macrophage polarization via targeting C/EBP β . We found induction of C/EBP β only in KCs isolated from miR-155 KO and not from WT mice after LPS or INF γ treatment (Fig.3L). A higher increase in STAT3 (M2 regulator) was also observed in KCs isolated from miR-155 KO compared to WT mice after LPS or INF γ treatment (Fig.3M). This was specific to C/EBP β as C/EBP α levels were not so different between genotypes (Fig.3N). These results suggest that miR-155 deficient KCs exhibited predominance of M2 phenotype when treated with M1 polarizing stimuli. Next, we attempted to find the explanation for increased iNOS expression in KCs isolated from miR-155 KO after LPS or INF γ treatment. For this, we checked for Hif1 α , a M1 macrophage transcriptional regulator, and found enhanced Hif1 α in KCs isolated from miR-155 KO compared to WT mice after LPS or INF γ treatment (Fig.3O). Collectively, our results suggest that miR-155 plays a role in M1 macrophage polarization. Further, we found that LPS induced whereas IL-4 decreased miR-155 levels, however we did not find any increase in miR-155 after INF γ treatment (Fig.3P). It is possible that kinetics of miR-155 induction with INF γ is different than LPS.

In addition to Kupffer cell/macrophage activation, alcoholic hepatitis is also characterized by neutrophil infiltration to the liver [23, 24]. Neutrophil infiltration has been shown to correlate with the severity of acute alcoholic hepatitis [24, 25]. Alcohol diet resulted in an infiltration of neutrophils (CD11b+Ly6G^{hi}) in the livers of WT mice compared to PF mice (Fig.4A). However, neutrophil infiltration was prevented in miR-155 KO mice after alcohol

diet (Fig. 4A). M2 macrophages are known to play roles in tissue remodeling, and fibrogenesis [22, 26, 27]. Chronic alcohol feeding induces pro-fibrotic gene expression [4]. Because our results indicated no increase in M2 macrophages in miR-155 KO mice after alcohol diet, we next evaluated the expression of pro-fibrotic genes. We found that chronic alcohol feeding resulted in an increase in mRNA levels of TGF β , (Fig.4B), type1 procollagen α (Fig.4C, and vimentin (Fig.4D) in WT mice whereas these increases were either prevented (TGF β and vimentin) or reduced (procollagen 1 α) in miR-155 KO mice (Fig.4 B–D). These results suggested that miR-155 deficiency could be protective from development of liver fibrosis.

Reduction in CCl₄-induced liver fibrosis in miR-155 KO mice

Because our results indicated a decrease in pro-fibrotic genes and M2 macrophages in miR-155 KO mice after alcohol diet, we further evaluated the role of miR-155 in liver fibrosis. The Lieber-DeCarli alcohol mouse model results in a modest increase in early markers of fibrosis [4] while the CCl₄-induced liver fibrosis mouse model induces significant fibrosis [28]. We found that CCl₄ treatment increases hepatic miR-155 expression in WT mice compared to corn oil (Fig.4E). miR-155 levels were not detected in miR-155 KO mice (Fig.4E). CCl₄ treatment for 2 weeks resulted in lymphocyte infiltration and initiation of collagen deposition in the livers of WT mice and an attenuation was observed in miR-155 KO mice (H&E and Sirius red staining (morphometry density units) (Fig.4F). Since CCl₄ induces inflammation, we next evaluated the markers of inflammation and found an increase in the mRNA levels of pro-inflammatory cytokines (TNF α and MCP1) in WT mice and not in miR-155 KO mice after CCl₄ administration (Fig.4G–H). CCl₄ treatment caused the induction of pro-fibrotic genes including α smooth muscle actin (Fig.4I), and TGF β (Fig.4J) and type 1 collagen α (Fig.4K) in WT mice, where as miR-155 KO mice either showed no increases or reduction (Fig.4I–K). Matrix metalloproteinases (MMPs) and metalloproteinase inhibitor (TIMP1) play a crucial role in matrix remodeling [29]. CCl₄ treatment resulted in the induction of the mRNA levels of MMP9 and MMP12 (Fig.4L–M) and TIMP1 (Fig.4N) only in WT mice and not in miR-155 KO mice (Fig.4L–N). Consistent with the mRNA data (Fig.4N), we found that the serum levels of TIMP1 were increased in WT mice after CCl₄ treatment and this increase was prevented in miR-155 KO mice (Fig. 4O). Epithelial-mesenchymal cell transition (EMT) is an important event in the initiation of organ fibrosis. EMT-promoting transcription factor Snail1 has been shown to play a role in liver fibrosis and hepatocyte specific deletion of Snail1 was shown to protect the mice from liver fibrosis [30]. Consistent with previous report [30] we found increase in Snail1 in WT mice after CCl₄ treatment, however this increase in Snail1 was prevented in miR-155 KO mice (Fig.4P).

To show that miR-155 is indeed involved in the development of advanced liver fibrosis we carried out 9 weeks of CCl₄ treatment. Histological scoring of Sirius red staining performed by pathologist expert revealed less bridging fibrosis and perisinusoidal fibrosis in miR-155 KO compared to WT mice (Fig.5A). In concordance with this, less production of hydroxyproline was found in miR-155 KO compared to WT mice (Fig.5B). α smooth muscle actin protein levels were also significantly decreased in miR-155 KO mice compared to WT mice after 9 weeks of CCl₄ treatment (Fig.5C). Collectively, these results suggest a

role of miR-155 in liver fibrosis. Finally, to validate observations from the mouse models, we tested the expression of miR-155 levels in the cirrhotic livers of alcoholic patients. We found a significant induction of miR-155 levels in the alcoholic cirrhotic livers compared to livers without cirrhosis (Fig.5D).

Alcohol induces miR-155 in the liver at the transcriptional level via the TLR4 pathway

Both chronic alcohol and CCl₄ treatments were shown to increase serum LPS levels in mice [31]. Recently, we showed that miR-155 KO mice were protected from alcohol-induced intestinal permeability and inflammation [2]. We found an increase in circulating endotoxin levels in WT mice and not in miR-155 KO mice after alcohol diet (Fig.6A). Upon activation by LPS, KCs release various pro-inflammatory and pro-fibrotic cytokines [6]. The effects of circulating LPS on the kinetics of hepatic miR-155 levels are not known; therefore we next checked the levels of liver miR-155 after a single treatment of LPS in WT mice. Our results indicate that miR-155 levels start to increase 2h after LPS injection and steadily increase over time (Fig.6B). No significant differences were found in miR-155 levels between 6h and 18h after LPS injection (Fig.6B) suggesting that miR-155 levels reached a plateau after 6h of LPS injection. To investigate the cellular source of miR-155 increase after LPS treatment, liver mononuclear cells (MNCs) and hepatocytes isolated from WT mice were treated with or without LPS (100ng/ml). Our results indicate that LPS induces miR-155 only in MNCs but not in hepatocytes (Fig.6C). To corroborate these finding in alcoholic steatohepatitis, we examined the expression of miR-155 in WT mice that were challenged with LPS for 3h after alcohol feeding. We found no significant differences in the induction of miR-155 levels between alcohol and control diet fed mice after LPS challenge (Fig.6D), suggesting that alcohol and LPS may increase miR-155 via the same mechanism.

LPS is recognized by the TLR4 receptor resulting in downstream activation of NF- κ B and pro-inflammatory cytokines. TLR4 signaling plays a crucial role in alcohol-induced effects and mice deficient in TLR4 have attenuation in alcohol-mediated cellular response [32]. Our results indicate a role of TLR4 signaling as no induction in mature miR-155 levels were observed in TLR4 KO mice after alcohol feeding compared to WT mice (Fig.6E). To determine if alcohol has any effect on the transcription of miR-155, we checked primary miR-155 (pri-miR-155) levels. Our results demonstrate a significant increase in pri-miR-155 levels in alcohol-fed WT mice compared to controls (Fig.6F). These results suggest that in vivo alcohol increase miR-155 transcription.

Discussion

microRNAs play a key role in various liver diseases including alcoholic liver disease (ALD) and fibrosis. miR-155 is a major regulator of inflammation and immunity. In this study we defined the functional role of miR-155 in ALD and liver fibrosis using miR-155 KO mice. We found that miR-155 KO mice were protected from alcohol-induced liver injury, oxidative stress, steatosis and inflammation. Our data revealed that miR-155 KO mice exhibited reduced liver fibrosis markers in both alcohol and CCl₄-induced fibrosis. miR-155 is a major mediator of inflammation, however in recent years the role of miR-155 in steatosis and fibrosis has been emerging [12–14] and our results support the role of miR-155 in these

processes. We found that the decrease in PPRE and PPAR α binding observed in WT mice was prevented in miR-155 KO mice after alcohol diet. Our results revealed that miR-155 regulates PPRE signaling in both macrophages and hepatocytes. In primary isolated hepatocytes we found that MCP1 treatment decreased PPRE in WT mice where as this decrease was prevented in miR-155 KO mice suggesting a role of miR-155 in regulation of steatosis via PPRE signaling. In consistent with our results MCP1 treatment was shown to decrease PPRE binding in Huh7 cells [19, 21]. We further identified and validated PPAR γ a direct target miR-155 in macrophages as inhibition of miR-155 using a specific miR-155 inhibitor resulted in an increase in PPAR γ gene in both naïve and alcohol treated cells. Our in vitro findings from hepatocytes and macrophages are in concordance with in vivo findings as we found that alcohol-induced decrease in PPRE binding in WT mice was prevented in miR-155 KO mice.

Ethanol-induced hepatic steatosis is multifunctional and alcohol affects several pathways, including fatty acid uptake, oxidation, and export [7]. Alcohol is known to reduce PPAR α [7], thereby inhibiting fatty acid oxidation in hepatocytes. PPAR α and PPAR γ not only play a pivotal role in hepatic lipid metabolism but also in oxidative stress, inflammatory response, and fibrogenesis [7]. Collectively, our results indicate that steatogenic, inflammatory and fibrogenic changes observed in miR-155 KO mice after ethanol diet are likely due to the effect of miR-155 on PPAR signaling.

miR-155 plays a key role in macrophage polarization and promotes the M1 phenotype of macrophage [22, 33]. While no significant differences were found in the quantity of immune cell infiltration between WT and miR-155 KO mice before and after alcohol feeding, we found that miR-155 deficiency alter the phenotype of infiltrating immune cells. Macrophage and KC activation are major elements in ALD [1]. Studies suggest that miR-155 favors M1 macrophages via its effect on C/EBP- β , SHIP1, and IL13R α 1[11, 22, 33, 34]. Interestingly in our ALD mouse model, the absence of miR-155 prevented an increase in the number of M2 macrophages after alcohol diet, suggesting that miR-155 has a tissue/cell/microenvironment specific role. Consistent with the role of miR-155 in promoting M1 macrophage phenotype we found a higher baseline of M2 macrophages in miR-155 KO mice on control diet compared to WT. Results from our in vitro studies revealed that KCs isolated from miR-155 KO mice (on chow diet) exhibited predominance of M2 phenotype but also had increased M1 phenotype after M1 polarizing signals. This was somewhat unexpected and could be explained by the fact that we found an increase of both C/EBP β (M2 transcriptional regulator) and HIF1 α (M1 transcriptional regulator) after polarization with LPS or INF γ in isolated KCs from miR-155 KO mice. A recent study revealed that iNOS deficient mice had enhanced M1 macrophage differentiation after polarization but without having major effects on M2 macrophages [35]. Our results support the notion that macrophages likely exist as a mixed population of M1/M2 in vivo or as hybrid phenotypes instead of an entirely polarized M1 or M2 phenotype [35].

It is well known that alcoholic steatohepatitis, if sustained, can lead to fibrosis, cirrhosis and hepatocellular carcinoma [1]. CCl $_4$ -induced liver fibrosis is a robust and widely used model of liver fibrosis [28]. Sustained/chronic inflammation leads to fibrosis, therefore, we hypothesized that deficiency of miR-155 will also attenuate CCl $_4$ -induced inflammatory

response and fibrosis. Consistent with findings from the alcohol diet, we found reduced fibrosis in miR-155 KO mice after CCl₄ treatment. Emerging studies support a role of miR-155 in liver fibrosis. Recently, we reported that miR-155 deficiency attenuated MCD diet-induced early liver fibrosis [12]. EMT promoting transcription factor, Snail1 (a putative miR-155 target), plays a role in liver fibrosis and we found that CCl₄-induced increase in Snail1 was prevented in miR-155 KO mice. We found that miR-155 KO mice not only were protected from early fibrosis but also showed attenuation from advanced fibrosis after CCl₄ treatment (9 weeks), revealing a role of miR-155 in promoting liver fibrosis. PPAR γ also acts as an anti-fibrotic gene and we identified PPAR γ as a direct target of miR-155. Also miR-155 targets several genes involved in fibrogenesis (such as SMAD2/5, Snail1, STAT3) it is highly likely that the fibrosis phenotype observed in our study is the cumulative effect of miR-155 on various target genes. Our finding of miR-155 being induced in a mouse model of liver fibrosis is relevant to human disease as we found increased miR-155 levels in the cirrhotic livers of alcoholic patients. Given the critical importance of PPAR signaling in normal lipid and glucose metabolism as well as in the regulation of inflammatory responses, it is highly likely that changes in steatosis, inflammation and fibrosis seen in miR-155 KO mice are due to miR-155's effect on PPAR signaling.

We didn't find differences in alcohol metabolism between genotypes as serum ethanol levels and Cyp2E1 protein levels were comparable between WT and miR-155 KO mice after alcohol diet (Supple Fig. 1G–H). Alcohol intake also increases intestinal permeability, which causes an increase in LPS in the circulation [2]. We found higher endotoxin levels in miR-155 KO mice on pair fed diet after 5 weeks, however this effect was not observed when endotoxin levels were checked after 4 weeks of alcohol diet (Supple Fig. 1I). Circulating LPS is recognized by hepatic macrophages via TLR4 receptor resulting in [6]. The role of the LPS/TLR4 axis is very well known in ALD and TLR4 KO mice were protected from ALD [32]. Our results support a role of TLR4 signaling in ALD as we found that alcohol induces miR-155 via the TLR4 pathway. In summary, our data suggest that miR-155 most likely modulates the pro-inflammatory and pro-fibrotic programs via targeting various signaling cascades that impact not only immune cells but also non-immune cells including hepatocytes. Additionally, our results indicate that the absence of miR-155 not only halts alcohol-induced steatohepatitis but also fibrosis in mouse models.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors greatly appreciate Dr. Jin-Kyu Park help in evaluation and scoring of the histology slides. Human liver samples were provided by the NIH-Funded Liver Tissue Procurement and Cell Distribution System (N01-DK-70004/HHSN26700700004C).

Financial support: This work was supported by grant from National Institute on Alcohol Abuse and Alcoholism-AA020744 to G.S.

List of abbreviations

ALD	alcoholic liver disease
KC	Kupffer cell
LPS	lipopolysaccharide
TLR	toll-like receptor
miRNA	microRNA
WT	wild type
KO	knock out
PPRE	peroxisome proliferator-activated receptor (PPAR) response element
M2	macrophage2
TNF α	tumor necrosis factor α
MCP1	monocyte chemo attractant protein 1
IL-1β	Interleukin-1
TGF β	Transforming growth factor β
LMNCs	liver mononuclear cells
EtOH	ethanol
CCl4	carbon tetrachloride

References

1. Wang HJ, Gao B, Zakhari S, Nagy LE. Inflammation in alcoholic liver disease. *Annu Rev Nutr.* 2012;343–368. [PubMed: 22524187]
2. Lippai D, Bala S, Catalano D, Kodys K, Szabo G. Micro-RNA-155 Deficiency Prevents Alcohol-Induced Serum Endotoxin Increase and Small Bowel Inflammation in Mice. *Alcohol Clin Exp Res.* 2014; 8:2217–2224. [PubMed: 25156614]
3. Petrasek J, Iracheta-Vellve A, Csak T, Satishchandran A, Kodys K, Kurt-Jones EA, et al. STING-IRF3 pathway links endoplasmic reticulum stress with hepatocyte apoptosis in early alcoholic liver disease. *Proc Natl Acad Sci U S A.* 2013; 41:16544–16549. [PubMed: 24052526]
4. Petrasek J, Bala S, Csak T, Lippai D, Kodys K, Menashy V, et al. IL-1 receptor antagonist ameliorates inflammasome-dependent alcoholic steatohepatitis in mice. *J Clin Invest.* 2012; 10:3476–3489. [PubMed: 22945633]
5. Szabo G, Bala S. Alcoholic liver disease and the gut-liver axis. *World J Gastroenterol.* 2010; 11:1321–1329. [PubMed: 20238398]
6. Bala S, Marcos M, Kodys K, Csak T, Catalano D, Mandrekar P, et al. Up-regulation of microRNA-155 in macrophages contributes to increased tumor necrosis factor {alpha} (TNF{alpha}) production via increased mRNA half-life in alcoholic liver disease. *J Biol Chem.* 2011; 2:1436–1444. [PubMed: 21062749]
7. Nan YM, Wang RQ, Fu N. Peroxisome proliferator-activated receptor alpha, a potential therapeutic target for alcoholic liver disease. *World J Gastroenterol.* 2014; 25:8055–8060. [PubMed: 25009377]
8. Dong X, Liu H, Chen F, Li D, Zhao Y. MiR-214 promotes the alcohol-induced oxidative stress via down-regulation of glutathione reductase and cytochrome P450 oxidoreductase in liver cells. *Alcohol Clin Exp Res.* 2014; 1:68–77. [PubMed: 23905773]

9. Meng F, Glaser SS, Francis H, Yang F, Han Y, Stokes A, et al. Epigenetic Regulation of miR-34a Expression in Alcoholic Liver Injury. *Am J Pathol.* 2012
10. Yin H, Hu M, Zhang R, Shen Z, Flatow L, You M. MicroRNA-217 promotes ethanol-induced fat accumulation in hepatocytes by down-regulating SIRT1. *J Biol Chem.* 2012; 13:9817–9826. [PubMed: 22308024]
11. O'Connell RM, Chaudhuri AA, Rao DS, Baltimore D. Inositol phosphatase SHIP1 is a primary target of miR-155. *Proc Natl Acad Sci U S A.* 2009; 17:7113–7118. [PubMed: 19359473]
12. Csak T, Bala S, Lippai D, Kodys K, Catalano D, Iracheta-Vellve A, et al. MicroRNA-155 Deficiency Attenuates Liver Steatosis and Fibrosis without Reducing Inflammation in a Mouse Model of Steatohepatitis. *PLoS One.* 2015; 6:e0129251. [PubMed: 26042593]
13. Miller AM, Gilchrist DS, Nijjar J, Araldi E, Ramirez CM, Lavery CA, et al. MiR-155 has a protective role in the development of non-alcoholic hepatosteatosis in mice. *PLoS One.* 2013; 8:e72324. [PubMed: 23991091]
14. Dai W, Zhao J, Tang N, Zeng X, Wu K, Ye C, et al. MicroRNA-155 attenuates activation of hepatic stellate cell by simultaneously preventing EMT process and ERK1 signalling pathway. *Liver Int.* 2015; 4:1234–1243. [PubMed: 25142507]
15. Bala S, Petrasek J, Mundkur S, Catalano D, Levin I, Ward J, et al. Circulating microRNAs in exosomes indicate hepatocyte injury and inflammation in alcoholic, drug-induced and inflammatory liver diseases. *Hepatology.* 2012
16. Bala S, Csak T, Momen-Heravi F, Lippai D, Kodys K, Catalano D, et al. Biodistribution and function of extracellular miRNA-155 in mice. *Sci Rep.* 2015:10721. [PubMed: 26024046]
17. Saha B, Bala S, Hosseini N, Kodys K, Szabo G. Kruppel-like factor 4 is a transcriptional regulator of M1/M2 macrophage polarization in alcoholic liver disease. *J Leukoc Biol.* 2015
18. Pawlak M, Lefebvre P, Staels B. Molecular mechanism of PPARalpha action and its impact on lipid metabolism, inflammation and fibrosis in non-alcoholic fatty liver disease. *J Hepatol.* 2015; 3:720–733. [PubMed: 25450203]
19. Mandrekar P, Ambade A, Lim A, Szabo G, Catalano D. An essential role for MCP-1 in alcoholic liver injury: Regulation of pro-inflammatory cytokines and hepatic steatosis. *Hepatology.* 2011
20. Zhou S, Wang Y, Meng Y, Xiao C, Liu Z, Brohawn P, et al. In Vivo Therapeutic Success of MicroRNA-155 (miR-155) Antagomir in a Mouse Model of Lupus Alveolar Hemorrhage. *Arthritis Rheumatol.* 2015
21. Nath B, Levin I, Csak T, Petrasek J, Mueller C, Kodys K, et al. Hepatocyte-specific hypoxia-inducible factor-1alpha is a determinant of lipid accumulation and liver injury in alcohol-induced steatosis in mice. *Hepatology.* 2011; 5:1526–1537. [PubMed: 21520168]
22. Cai X, Yin Y, Li N, Zhu D, Zhang J, Zhang CY, et al. Re-polarization of tumor-associated macrophages to pro-inflammatory M1 macrophages by microRNA-155. *J Mol Cell Biol.* 2012; 5:341–343. [PubMed: 22831835]
23. Lemmers A, Moreno C, Gustot T, Marechal R, Degre D, Demetter P, et al. The interleukin-17 pathway is involved in human alcoholic liver disease. *Hepatology.* 2009; 2:646–657. [PubMed: 19177575]
24. Ramaiah SK, Jaeschke H. Role of neutrophils in the pathogenesis of acute inflammatory liver injury. *Toxicol Pathol.* 2007; 6:757–766. [PubMed: 17943649]
25. Bertola A, Park O, Gao B. Chronic plus binge ethanol feeding synergistically induces neutrophil infiltration and liver injury in mice: a critical role for E-selectin. *Hepatology.* 2013; 5:1814–1823. [PubMed: 23532958]
26. Gratchev A, Guillot P, Hakiy N, Politz O, Orfanos CE, Schledzewski K, et al. Alternatively activated macrophages differentially express fibronectin and its splice variants and the extracellular matrix protein betaIG-H3. *Scand J Immunol.* 2001; 4:386–392. [PubMed: 11285119]
27. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol.* 2008; 12:958–969. [PubMed: 19029990]
28. Chiang DJ, Roychowdhury S, Bush K, McMullen MR, Pisano S, Niese K, et al. Adenosine 2A receptor antagonist prevented and reversed liver fibrosis in a mouse model of ethanol-exacerbated liver fibrosis. *PLoS One.* 2013; 7:e69114. [PubMed: 23874883]

29. Hemmann S, Graf J, Roderfeld M, Roeb E. Expression of MMPs and TIMPs in liver fibrosis – a systematic review with special emphasis on anti-fibrotic strategies. *J Hepatol.* 2007; 5:955–975. [PubMed: 17383048]
30. Rowe RG, Lin Y, Shimizu-Hirota R, Hanada S, Neilson EG, Greenson JK, et al. Hepatocyte-derived Snail1 propagates liver fibrosis progression. *Mol Cell Biol.* 2011; 12:2392–2403. [PubMed: 21482667]
31. Seki E, Schnabl B. Role of innate immunity and the microbiota in liver fibrosis: crosstalk between the liver and gut. *J Physiol.* 2012; (Pt 3):447–458. [PubMed: 22124143]
32. Hritz I, Mandrekar P, Velayudham A, Catalano D, Dolganiuc A, Kodys K, et al. The critical role of toll-like receptor (TLR) 4 in alcoholic liver disease is independent of the common TLR adapter MyD88. *Hepatology (Baltimore, Md).* 2008; 4:1224–31.
33. He M, Xu Z, Ding T, Kuang DM, Zheng L. MicroRNA-155 regulates inflammatory cytokine production in tumor-associated macrophages via targeting C/EBPbeta. *Cell Mol Immunol.* 2009; 5:343–352. [PubMed: 19887047]
34. Martinez-Nunez RT, Louafi F, Sanchez-Elsner T. The interleukin 13 (IL-13) pathway in human macrophages is modulated by microrna-155 via direct targeting of interleukin 13 receptor alpha1 (IL13R{alpha}1). *J Biol Chem.* 2010
35. Lu G, Zhang R, Geng S, Peng L, Jayaraman P, Chen C, et al. Myeloid cell-derived inducible nitric oxide synthase suppresses M1 macrophage polarization. *Nat Commun.* 2015:6676. [PubMed: 25813085]




Figure 1. miR-155 deficiency attenuates chronic alcohol-induced steatosis and liver injury
 C57BL/6 wild type (WT) or miR-155 KO mice (n=8–10) were fed with Lieber DeCarli diet containing either 5% ethanol or isocaloric control diet for 5 weeks. Formalin fixed liver sections were stained with Hematoxylin and eosin (A) Oil-O-red stain (B). Slides were observed under light microscope (200X) and representative slides are shown. Histological scores were recorded by a pathology expert (C). Triglyceride levels were measured from the livers (D). ALT levels were measured from plasma (E). Oxidative stress was measured from

whole liver cell lysates by TBARs assay (F). Data is presented as mean \pm SEM. * indicates $p < 0.05$ vs. pair fed (PF) mice and # $p < 0.05$ compared to pair fed KO mice.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2. Alcohol diet results in increased PPRE and PPARα DNA binding in miR-155 KO mice
 WT or miR-155 KO mice were fed with Lieber DeCarli diet as described in methods. Total RNA was used to determine PPARα levels by qPCR (A). 5ug of nuclear proteins were used to detect PPARE binding by EMSA and PPARα antibody was used for supershift assay and density (B). Primary hepatocytes were isolated from WT and miR-155 KO mice (chow fed) and treated them with or without MCP1 (100ng/ml) for 36h. 10ug of nuclear proteins was used to detect PPARE binding by EMSA (D). RAW 264.7 macrophages were transfected with either miR-155 inhibitor or control inhibitor using lipofectamine siRNA reagent as described

in methods. Cells were treated with or without alcohol (50mM), cells were harvested after 48h of transfection and total RNA was used to quantify PPAR γ mRNA levels (E). Expression of lipid metabolism and fatty acid uptake gene expression was evaluated by qPCR. FABP4 (F), LXR α (G), ACC1 (H), and LDLR (I), and HMGCR (J).MCP1 mRNA levels were determined by qPCR (K). 18s was used to normalize Cq values. Whole cell liver lysates (L) and plasma (M) were used to detect MCP1 by ELISA. (D). Data is presented as mean \pm SEM. * indicates $p < 0.05$ vs. pair-fed mice, # $p < 0.05$ vs PF mice treated with LPS. \$ $p < 0.05$ vs PF KO mice.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 3. Chronic alcohol-induced inflammation and macrophage infiltration are attenuated in miR-155 KO mice

WT or miR-155 KO mice were fed with Lieber DeCarli diet as described in methods. Whole cell liver lysates (A and B, left panel) and plasma (A and B, right panel) were used to detect TNF α (A) and IL-1 β (B) by ELISA. RNA was used to determine CD68 (C) and MIP2 (D) mRNA levels by qPCR. MNCs isolated from WT and miR-155 KO mice after alcohol feeding and were stained with combination of antibodies for monocytes, macrophages and neutrophils as described in methods and processed for flow cytometry. % of inflammatory monocytes (E), and M2 macrophages (F) is shown. Kupffer cells were isolated from WT and

KO mice and treated with or without LPS (100ng/ml), IL-4 (20ng/ml) and IFN γ (20ng/ml) for 18h. Total RNA was evaluated for M1 and M2 markers and their regulators. Arginase1 (G), IL-10 (H), MRC1 (I), IL-1 β (J), iNOS (K), C/EBP β (L), STAT3 (M), C/EBP α (N), HiF1 α (O), and miR-155 levels (P). 18s was used to normalize Cq values. Data is presented as mean \pm SEM. * indicates p<0.05 vs. pair fed mice, # p<0.05 vs PF mice treated with LPS. \$ p<0.05 vs PF mice KO mice. ns: non significant.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4. Reduction in CCl₄-induced liver fibrosis in miR-155 KO mice
 WT or miR-155 KO mice were fed with Lieber DeCarli diet. MNCs were stained for neutrophils (A). Liver RNA was used to determine TGFβ (B) procollagen 1α (C) and vimentin (D) mRNA levels by qPCR. 18s was used to normalize Cq values. Data is presented as mean ± SEM. * indicates p<0.05 vs. pair fed mice, ns: non significant. WT or miR-155 KO mice (n=6) were treated either with corn oil or CCl₄ for 2 weeks as described in methods. Total RNA was used to determine miR-155 levels by qPCR (E). Formalin fixed liver sections were stained with Hematoxylin and eosin (F, left panel) and Sirius red stain (F, right panel).

right panel). Slides were observed under light microscope (100X) and representative slides are shown. RNA was used to determine TNF α (G), MCP1 (H), α smooth muscle actin (I), TGF β (J), collagen 1 α (K), MMP9 (L), MMP12 (M), Timp1 (N), Snail1 (P) mRNA levels by qPCR. Timp1 levels were measured from plasma by ELISA (O). 18s or SnoRNA202 was used to normalize Cq values for mRNA, and mouse miR-155 respectively. Data is presented as mean \pm SEM. * indicates p<0.05 vs. corn oil treated mice.




Figure 5. miR-155 promotes liver fibrosis

WT or miR-155 KO mice (n=9) were treated either with corn oil or CCl₄ for 9 weeks. Formalin fixed liver sections were stained with Sirius red stain (A). Slides were observed under light microscope (100X) and representative slides are shown. The amount of hydroxyproline was measured from the liver (B). Western blot analysis for α SMA and density units (C), miR-155 levels were determined from alcoholic cirrhotic livers by qPCR (D). RNU48 was used to normalize Cq values. Data is presented as mean ± SEM. * indicates p<0.05 vs. corn oil treated mice.




Figure 6. Alcohol induces miR-155 in the liver at the transcriptional level via the TLR4 pathway WT or miR-155 KO mice were fed with Lieber DeCarli diet. Endotoxin levels were measured from plasma (A). WT mice were either administered with saline or LPS (2.5mg/kg, ip) for indicated times and miR-155 levels were measured from livers (B). MNCs or hepatocytes were isolated from perfused WT livers as described in methods. Cells were treated with LPS (100ng/ml) for 6h and miR-155 levels were detected (C). WT or miR-155 KO pair- or alcohol- fed mice were injected with LPS (0.5mg/kg, ip) for 3h and livers were isolated and miR-155 levels were detected (D). C57BL/6 WT or TLR4 KO mice (n=6–8) were fed with Lieber DeCarli diet. (E). Primary miR-155 levels were detected using TaqMan gene expression assay. GAPDH was used to normalize Cq values (F). Data is presented as mean ± SEM. * indicates *p*<0.05 vs. pair fed mice (A, D and F). * indicates *p*<0.05 vs. saline treated mice (B) or cells in medium only (C).