



Published in final edited form as:

Avian Dis. 2010 March ; 54(1 Suppl): 466–476. doi:10.1637/8914-043009-Reg.1.

Migration of Waterfowl in the East Asian Flyway and Spatial Relationship to HPAI H5N1 Outbreaks

John Y. Takekawa^{A,K}, Scott H. Newman^B, Xiangming Xiao^C, Diann J. Prosser^D, Kyle A. Spragens^A, Eric C. Palm^A, Baoping Yan^E, Tianxian Li^F, Fumin Lei^G, Delong Zhao^{C,G}, David C. Douglas^H, Sabir Bin Muzaffar^{A,I}, and Weitao Ji^J

^AU.S. Geological Survey, Western Ecological Research Center, San Francisco Bay Estuary Field Station, Vallejo, CA 94592

^BUnited Nations Food and Agriculture Organization, EMPRES Wildlife Unit, Emergency Centre for Transboundary Animal Diseases, Animal Production and Health Division, Rome 00153, Italy

^CDepartment of Botany and Microbiology, Center for Spatial Analysis, University of Oklahoma, Norman, OK 73019

^DU.S. Geological Survey, Patuxent Wildlife Research Center, Beltsville, MD 20705

^EChinese Academy of Sciences, Computer Network Information Center, 4 South 4th Street, Haidian, Beijing 100080, China

^FChinese Academy of Sciences, State Key Laboratory of Virology, Wuhan Institute of Virology, No. 44 Xiao Hong Shan District, Wuchang, Wuhan 430071, China

^GChinese Academy of Sciences, Institute of Zoology, 25 Beisihuanxi Road, Haidian, Beijing 100080, China

^HU.S. Geological Survey, Alaska Science Center, Juneau, AK 99801

^IDepartment of Biology, College of Sciences, United Arab Emirates University, Al Ain, UAE

^JPoyang Lake National Nature Reserve, State Forestry Administration, Nanchang, China

SUMMARY

Poyang Lake is situated within the East Asian Flyway, a migratory corridor for waterfowl that also encompasses Guangdong Province, China, the epicenter of highly pathogenic avian influenza (HPAI) H5N1. The lake is the largest freshwater body in China and a significant congregation site for waterfowl; however, surrounding rice fields and poultry grazing have created an overlap with wild waterbirds, a situation conducive to avian influenza transmission. Reports of HPAI H5N1 in healthy wild ducks at Poyang Lake have raised concerns about the potential of resilient free-ranging birds to disseminate the virus. Yet the role wild ducks play in connecting regions of HPAI H5N1 outbreak in Asia is hindered by a lack of information about their migratory ecology. During 2007–08 we marked wild ducks at Poyang Lake with satellite transmitters to examine the location and timing of spring migration and identify any spatiotemporal relationship with HPAI H5N1 outbreaks. Species included the Eurasian wigeon (*Anas penelope*), northern pintail (*Anas acuta*),

^KCorresponding author: ; Email: john_takekawa@usgs.gov

common teal (*Anas crecca*), falcated teal (*Anas falcata*), Baikal teal (*Anas formosa*), mallard (*Anas platyrhynchos*), garganey (*Anas querquedula*), and Chinese spotbill (*Anas poecilohyncha*). These wild ducks (excluding the resident mallard and Chinese spotbill ducks) followed the East Asian Flyway along the coast to breeding areas in northern China, eastern Mongolia, and eastern Russia. None migrated west toward Qinghai Lake (site of the largest wild bird epizootic), thus failing to demonstrate any migratory connection to the Central Asian Flyway. A newly developed Brownian bridge spatial analysis indicated that HPAI H5N1 outbreaks reported in the flyway were related to latitude and poultry density but not to the core migration corridor or to wetland habitats. Also, we found a temporal mismatch between timing of outbreaks and wild duck movements. These analyses depend on complete or representative reporting of outbreaks, but by documenting movements of wild waterfowl, we present ecological knowledge that better informs epidemiological investigations seeking to explain and predict the spread of avian influenza viruses.

Keywords

Anatidae; waterfowl; Poyang Lake; Brownian bridge movement model; satellite telemetry; highly pathogenic avian influenza; H5N1; East Asian Flyway

Poyang Lake, located 50 km northeast of Nanchang in Jiangxi Province, is the largest freshwater lake in China with a total area of about 3585 km² (39). Over 520,000 birds are found at Poyang Lake during the nonbreeding period (43), many of which are long-distance migrants including over 2700 Siberian cranes (*Grus leucogeranus*), representing 95% of the global population (24,28). Among the 300 bird species recorded, a significant proportion are migratory Anatidae, including the whooper swan (*Cygnus cygnus*), greater white-fronted goose (*Anser albifrons*), greylag goose (*Anser anser*), Chinese spotbill (*Anas poecilohyncha*), Eurasian wigeon (*Anas penelope*), and mallard (*Anas platyrhynchos*) (28,43). Poultry grazing in surrounding croplands maximizes overlap between wild bird and domestic poultry, thereby increasing chances of transmission of avian influenza viruses (AIVs). In addition, a relatively recent farming practice at the Poyang Lake region involves rearing wild waterfowl in captivity, which is preferred by some Chinese consumers over domestic fowl. Migratory swan geese (*Anser cygnoides*) as well as nonmigratory Chinese spotbills are now raised in captivity for sale in markets. Many of these farm-raised waterfowl are allowed to feed or swim in the wetlands, facilitating direct interaction of farmed and free-ranging wild birds.

Areas such as Poyang Lake in which large numbers of domestic ducks and geese are raised in areas with little or no biosafety measures (9) are conducive to the emergence of novel highly pathogenic avian influenza (HPAI) subtypes. For example, phylogenetic analyses of AIVs by Mukhtar *et al.* (30) showed that reassortment of Nanchang and Hokkaido subtypes at Poyang Lake may have resulted in the evolution of HPAI Gs/Gd/1/96 of subtype H5N1. Subsequent introductions of new AIVs from domestic to wild birds are also probable, especially because domestic and wild ducks are intermixed and poultry outnumber wild waterfowl by more than 25 to 1 (43). Evidence for introduction of novel AIVs into wild bird populations exists in the detection of two HPAI H5N1 genotypes ('Z' and 'V') from Poyang Lake between October 2004 and March 2005 by Chen *et al.* (6). While the species sampled

were identified (mallard, falcated teal [*Anas falcata*], and Chinese spotbill), the results were pooled into a mixed “migratory duck” category that combined species with contrasting ecology and behavior, limiting ecological interpretation of these findings (12,31,45). Both Mukhtar *et al.* and Chen *et al.* have implicated wild waterfowl at Poyang Lake in HPAI H5N1 transmission, and both proposed a possible link to Qinghai Lake, where the largest recorded wild bird epizootic occurred in 2005 (6,29). However, little empirical information was available at that time on the migratory movements of wild ducks from the Poyang and Qinghai regions to ecologically establish an association with HPAI H5N1 outbreaks.

In this paper, we present results from eight wild duck species including the Eurasian wigeon (*Anas penelope*), northern pintail (*Anas acuta*), common teal (*Anas crecca*), falcated teal, Baikal teal (*Anas formosa*), mallard, garganey (*Anas querquedula*), and Chinese spotbill, marked with satellite transmitters at Poyang Lake, Wuhan, China during 2007–08. The infection status of birds marked at the onset of the project in March 2007 was diagnosed using both reverse transcriptase-polymerase chain reaction (RT-PCR) and serological testing. We used telemetry locations to develop movement models and utilization distributions of the East Asian Flyway migration routes. We compared these migratory pathways with regional HPAI H5N1 outbreaks reported from this region in poultry and wild birds to determine if there were spatial relationships between wild duck movements and outbreak areas. Finally, we examined relationships between the timing of outbreaks and wild duck life-cycle stages to see if they were correlated.

MATERIALS AND METHODS

Study sites

Ducks were captured at Poyang Lake (29°09'N, 116°13'E) located 50 km northeast of Nanchang in Jiangxi Province, China. Poyang Lake is fed by the Gan and Xiu rivers, and its shallow basin is connected to the Yangtze River. The lake is part of a large alluvial floodplain with seasonal rainfall (between May and September) that expands its area to about 4000 km² (38). Severe flooding has been common since 1950s because of land reclamation, levee construction, and El Niño events (38). The surrounding land use is dominated by rice agriculture, as Poyang Lake is in the center of China's rice production region. The lake basin also supports a rich benthic community of at least 58 invertebrate taxa, which exceeds resources available to waterfowl in the adjacent river systems (41). Satellite telemetry locations for ducks marked at Poyang Lake were used to describe the spring migration range of waterfowl in the East Asian Flyway, extending northward from Poyang Lake to northeast China and southeastern Siberia.

Capture and marking

We captured ducks between March 8 and 20, 2007, and November 27 and December 7, 2007, on the western side of Poyang Lake, China. We captured one duck in a line of monofilament leg nooses, but most ducks were captured by local cooperators with large mistnets erected across open water. Upon capture, birds were placed in individual cloth bags and promptly processed. We recorded species identity, mass, flat wing chord, diagonal or short tarsus (8), sex, and age for each bird.

Ducks were marked with 12 or 18 g solar-powered, 30 g GPS-solar-powered, or 26 g battery-implant Platform Terminal Transmitters (PTT-100 and solar-GPS PTT-100, Microwave Telemetry, Inc., Columbia, MD, USA). Solar transmitters were secured to birds with a teflon harness (Bally Ribbon Mills, Bally, PA), and the implant transmitters were surgically secured in the abdominal cavity (26) by a veterinarian with the antenna protruding dorsally. Transmitter packages averaged <3% of the bird's body weight. Birds were released near capture locations as soon as possible after processing. Procedures for capture, handling, and marking were approved by the USGS Patuxent Wildlife Research Center Animal Care and Use Committee.

Satellite telemetry locations

PTTs were programmed to transmit for 10-hr periods every 2 days, and GPS-PTTs were scheduled to record 6–12 GPS locations each day. Transmissions were received by the Argos satellite tracking system (CLS America Inc., Largo, MD). CLS calculated PTT locations from the perceived Doppler-effect shifts in transmission frequency during a satellite overpass. Each Doppler-derived PTT location was accompanied by a location quality class index. CLS reports a 1-sigma error radius of >1000 m, 350–1000 m, 150–350 m, and 150 m, for location classes indices 0, 1, 2, and 3, respectively. Auxiliary location classes A, B, and Z are not assigned accuracy estimates by CLS. We used a filtering algorithm (D. Douglas, Version 7.03, <http://alaska.usgs.gov/science/biology/spatial/>) to identify and remove implausible auxiliary Doppler locations based on distance moved, movement rate, and turning angle. We used ArcGIS 9.2 (Environmental Systems Research Institute, Inc., Redlands, CA) and Google Earth 5.0 (Google, Mountain View, CA) to plot telemetry locations. Migratory stopover sites were defined as areas where birds moved less than 20 km 24 hr time period.

Environmental data layers

We coanalyzed telemetry locations with a variety of digital thematic maps. For habitat features we used the Moderate-resolution Imaging Spectroradiometer (MODIS)/Terra Land Cover Classification, distributed by the Land Processes Distributed Active Archive Center, U.S. Geological Survey Center for Earth Resources Observation and Science (<http://LPDAAC.usgs.gov>). The MODIS/Terra Land Cover Classification contains a primary thematic that delineates 17 land cover classes defined by the International Geosphere-Biosphere Programme (IGBP). We used a University of Maryland modification of the IGBP scheme (Land Cover Type 2), which we generalized into five broad land cover categories: 1) water, 2) upland or natural vegetation, including barren or sparsely vegetated areas, 3) wetland, 4) cropland or cropland mosaic, and 5) urban.

Poultry density data

Poultry density data were obtained from the United Nations Food and Agriculture Organization (FAO) via the Geonetwork (<http://www.fao.org/geonetwork>). Methodology and sources of the estimates are described in the FAO's Gridded Livestock of the World (44). Briefly, for each country the most recent available livestock census data were converted into densities to produce “observed” data and then disaggregated based on statistical relations with environmental variables in similar agro-ecological zones to produce “predicted” poultry

distributions. The data files were disseminated in raster format (0.0833° resolution), with pixel values containing estimated poultry densities (head/km²).

HPAI H5N1 outbreak data

Information about HPAI H5N1 outbreaks were obtained from the Emergency Prevention System for Transboundary Animal and Plant Pests and Diseases (EMPRES) Database (11) for the period December 2003–February 2009. The 7-yr time frame for outbreaks was used to maximize the number of cases included in statistical analyses, providing a more complete outbreak pattern. Variables associated with each outbreak included date, location (country, administrative region, locality, latitude and longitude), reliability of the field veterinarian's diagnosis and the laboratory approach that was used to confirm viral subtype, and whether the outbreak occurred in wild birds or poultry. We restricted our analyses to records considered reliable according to the FAO EMPRES Program and the World Organisation for Animal Health (OIE).

Spatial analyses

We pooled species into four groups according to body mass: 1) falcated teal, 2) small teal (common teal, garganey, Baikal teal), 3) Eurasian wigeon, and 4) Chinese spotbill. We used all tracking locations that described the spring migration route of each waterfowl group (March to July 2007) for statistical analyses. We applied a fixed kernel home range analysis (19) of each bird's migration route to identify stopover sites. The fixed kernel analysis was performed using Animal Space Use (v. 1.2 Beta) software (20) and applied a least squares cross-validation method to obtain the kernel smoothing parameter. We applied Brownian bridge movement models (21) to create utilization distributions for spring migration pathways of each species group. We used Animal Space Use (v. 1.2) to create the Brownian bridge utilization distributions (BBUDs) and assumed the distribution of location errors was circular normal.

For coanalysis with virus outbreak data, we used the 99% BBUDs for each species group to avoid location outliers. We averaged the BBUDs of all species groups to describe a population-level spring migration route (21) and extracted the upper 60% utilization distribution, partitioned into six 10% intervals (i.e., 40–99%). We included 29 FAO-OIE confirmed HPAI H5N1 outbreak events from 2003 to 2009 involving wild and domestic birds that occurred within the population-level BBUD spring migration corridor. Wild bird mortalities were not analyzed according to taxon because most outbreak cases from this region were identified as “waterbirds” (32). We spatially intersected each HPAI H5N1 outbreak location with the land cover and poultry density maps. An equal number of random locations from within the overall BBUD were also intersected for comparison with the outbreak locations.

Statistical analyses

An information theoretic approach (Akaike Information Criterion [AIC]) was used to compare outbreak and random locations under 21 *a priori* defined logistic regression models. Covariates for the logistic models included latitude, habitat type (water, upland, wetland, cropland, and urban), poultry density, and utilization distribution (BBUD). Models within

2.0 AIC of the top-ranked model were considered to have biological significance (3). All analyses were conducted in Program-R (36) using the “glm” function (family = binomial, link = logit).

We divided the annual cycle of wild ducks into four seasonal stages in their life cycle on the basis of the area, scale of movement, and arrival and departure dates. These stages were 1) breeding and post-breeding (including annual moult): May 1–October 2; 2) fall migration: October 3–November 18; 3) nonbreeding: November 19–February 20; and 4) spring migration: February 21–April 30. We examined the null hypothesis of equal probability of an outbreak within season by comparing likelihood ratios during the four seasonal stages. To evaluate temporal variation, we weighted each PTT-marked duck equally and calculated average poultry exposure estimates throughout the annual cycle. We used a G-statistic (46) to detect if there was a significant difference between the number of observed (reported in EMPRES database) and expected (random distribution in time) outbreaks for each of the four seasonal stages.

AIV testing

Virological analyses were performed on samples collected from birds marked during March 2007, but not performed for samples collected during recapture. We obtained cloacal and tracheal swabs and blood from each bird following standard sampling and transport procedures (10). For all birds three analyses were performed. The first test, a sandwich enzyme-linked immunosorbent assay (ELISA) kit (KeQian Biology Technology Ltd, China), was used to detect the presence of type A influenza virus antigen with optical density recorded at 630 nm wavelength (OD_{630}). An OD_{630} value above 0.23 was regarded as positive. The second test, an rRT-PCR assay, targeted the Matrix gene (13) and was used to screen for type A influenza virus RNA. The third test, an indirect ELISA kit (KeQian Biology Technology), was used to detect the presence of AIV H5 antibody with an OD_{630} value above 0.40 regarded as positive. Tests were conducted according to OIE standards (33). Laboratory analyses were conducted by the Chinese Academy of Sciences, Wuhan Institute of Virology, China.

RESULTS

Capture and marking

We obtained a total of 13,699 locations from 33 ducks marked with PTTs at Poyang Lake in 2007 including the Chinese spotbill (13 adults: 6 males, 7 females), falcated teal (5 adults: 4 males, 1 female; 1 first-year female), Eurasian wigeon (5 adults: 1 male, 4 females), common teal (3 adult males; 1 first-year male), Baikal teal (2 adult males), garganey (1 adult male), northern pintail (1 first-year female), and mallard (1 adult male; Table 1). The total number of locations obtained per duck varied widely (2–909) as did the duration over which signals were received (14–617 days). We obtained few locations for the northern pintail, so it was excluded from analyses. Similarly, the mallard was excluded from analyses because the individual we captured was a resident. The Chinese spotbills were nonmigratory and year-round residents at Poyang Lake. We recovered spring migration tracks for 12 of the other 18 migratory ducks.

Migration from Poyang Lake

None of the birds migrated from the East Asian nonbreeding area at Poyang Lake to the Qinghai Lake breeding area in the Central Asian Flyway (Fig. 1). A single falcated teal migrated northwest toward the Central Asian Flyway, but it stopped 700 km northeast of Qinghai Lake on the eastern edge of the Gobi Desert, a region that presents a large ecological barrier to waterfowl migration.

Falcated teal—We obtained 1557 total locations for migrating falcated teal (Table 1). The falcated teal departed from Poyang Lake from March 28 to April 29. From late March through late April, they used migration sites in the Lower Yangtze River in Anhui and Zhejiang Provinces to the southwest of Shanghai (Table 2). In late April they migrated to sites on Bohai Bay (Fig. 2B) including a stopover site on the south side near Qingdao in Shandong Province. One falcated teal migrated to the northwest, but all other birds migrated along the coast to northern China and southeast Siberia. The core of their spring migration corridor was in the northeastern region of Manchuria in China (Fig. 3A).

Common teal—We obtained 1942 total locations for the species group of small teal that included common teal (4), Baikal teal (2), and garganey (1; Table 1). These teal departed from Poyang Lake from March 26 to May 3. Like the larger falcated teal, they used migration sites in the Lower Yangtze River in Anhui and Zhejiang Provinces to the west of Shanghai (Table 2; Fig. 3B). In late April they migrated to coastal wetland staging areas to the south of Bohai Bay on the Lower Huang River, and to the north of Bohai Bay off the coasts of South Korea, North Korea, and China (Fig. 2B). The small teal migrated farther north than the falcated teal into eastern Siberia south of the Sea of Okhotsk. Their BBUD was similar to that of falcated teal with a more southerly concentration.

Eurasian wigeon—We obtained 1294 total locations for Eurasian wigeon (Table 1). Eurasian wigeon seemed to depart Poyang Lake later than the teal, but their first recorded postdeparture locations were not until late April in northeast China (Figs. 2A, 3C). We obtained fewer locations for the wigeon, so it is difficult to determine if the seemingly late timing of their migration was due to inferior PTT performance (three of the five birds had implant battery-powered transmitters). Eurasian wigeon migrated farther northeast than other ducks, crossing the Sea of Okhotsk and concentrating in the Magadan Oblast of northeast Siberia (Fig. 3C).

Chinese spotbill—In contrast to the teal and Eurasian wigeon, Chinese spotbills are a nonmigratory species (4) that resides year-round in the Poyang Lake region. We obtained 8871 total locations (Table 1). As expected, the Chinese spotbill ducks remained within Jiangxi Province throughout the spring (Fig. 3D).

Spatial relationship of migration pathways and outbreaks

We examined waterfowl movements in relation to land cover, poultry density, and virus outbreaks within a rectangular study area bounded to the northeast at 55°N, 140°E and the southwest at 25°N, 110°E. We averaged the three BBUD isopleths for falcated teal, common and other small teal, and Eurasian wigeon to describe the East Asian Flyway pathway for

wild ducks from Poyang Lake (Fig. 4). Although the migratory corridor extended east to west across most of the Bohai Bay, there was a stopover region used by most ducks in China just north of Bohai Bay. We overlaid the overall BBUD migration pathway on a grid of poultry density and plotted the documented outbreak locations. The AIC analysis indicated that two models fit the data (Table 3): these included the top-ranked model with covariates latitude+poultry density, and the second-ranked model with latitude+poultry density +BBUD. However, a comparative examination of each covariate's contribution to the model (Table 4) indicated that BBUD magnitude was only weakly and inversely related to the outbreak location (*i.e.*, fewer wild bird locations where there were more outbreaks). Thus, we concluded that the top AIC ranked model (latitude+poultry density) was the single most parsimonious model in explaining variation in spatial outbreak locations.

Outbreaks were positively correlated with poultry density and inversely correlated with latitude, indicating that greater frequencies of outbreaks occurred at lower latitudes and with higher poultry densities. When we examined outbreaks in relation to both latitude and date, most confirmed outbreaks occurred at latitudes less than 40°N, and outbreaks began by October before most ducks had reached nonbreeding areas in early November. The spring migration utilization distribution (Fig. 4) had a weak positive relationship with outbreaks, and the χ^2 test indicated that utilization distribution was nonsignificant when compared with the null model (Table 4). None of the land cover class covariates were helpful in explaining the occurrence of outbreak events.

When we examined the outbreaks that occurred within the overall BBUD on a seasonal basis (Fig. 5), we found that the number of observed and expected outbreaks differed by life-cycle period ($G = 19.98$, $\chi^2_{0.05,3} = 7.82$, $P < 0.001$). Outbreaks occurred before many wild ducks had migrated to their nonbreeding areas (Fig. 5). More outbreaks occurred than were expected during the fall (61-day period) and the nonbreeding period (121-day period), whereas fewer occurred during the breeding season (122-day period) and spring migration (61-day period; Fig. 6).

AIV results

In total, 20 birds representing seven species were sampled for AIV, including Eurasian wigeon ($n = 2$), falcated teal ($n = 6$), common teal ($n = 4$), Baikal teal ($n = 2$), mallard ($n = 1$), garganey ($n = 1$), and Chinese spotbill ($n = 4$). The results of the sandwich ELISA and RT-PCR revealed that all 20 ducks were negative for type A influenza virus (Table 5). However, the serology results of the indirect ELISA test for AIV H5 antibody indicated that the only mallard was positive (Table 5). These results indicate that no bird was actively infected by HPAI H5N1; however, the presence of antibody in the blood of the mallard suggested prior exposure to H5N1. All birds that we handled appeared clinically healthy and showed no obvious symptoms of influenza infection.

DISCUSSION

Our ecological studies have been directed at improving understanding of the potential role of wild birds in the transmission of avian influenza. Prior to our satellite telemetry studies, little data were available on migratory movements of wild birds during the nonbreeding season at

Poyang Lake, a region that has been suggested as an important source of HPAI H5N1 (6,30). Because sample size in satellite telemetry projects is limited by the cost of the technology along with battery life and performance, it is often difficult to generalize the sequential movements of individuals to broader migratory patterns of populations. However, application of the recently developed Brownian bridge movement model (21) allowed us to describe the flyway migration corridor in a probabilistic manner. As waterfowl demonstrate a high degree of seasonal fidelity to sites within and between years (1), we believe our BBUD analysis of spring 2007 migration tracks provided a reliable representation of the general migration route for the East Asian Flyway.

We predicted that spatial and temporal concordance of HPAI H5N1 outbreaks with wild duck movements should occur if wild ducks were directly involved in spread of the virus. Lack of correlation between the core flyway corridors of wild birds as indicated by the BBUD analysis and sites of reported outbreaks did not support this hypothesis (Table 3). However, this type of spatial analysis depends on complete or representative reporting of outbreaks. If outbreaks are not fully reported, it could result in erroneous conclusions (22,23). Thus, our analysis highlights the importance of transparency in outbreak reporting to better understand the spread of disease. In addition, our analyses would have been improved with larger samples of the different species. By grouping species with different ecological characteristics, we may have missed individual differences in their movement patterns that may have been spatially correlated more closely to outbreaks.

We found no relationship among HPAI H5N1 outbreaks and land cover characteristics typically associated with waterfowl, such as water or wetland habitats (Table 3). However, a biologically significant relationship between poultry density and sites of HPAI H5N1 outbreaks was detected (Table 4), consistent with modeling studies of the virus in Southeast Asia (14). The relationship of outbreaks with latitude (Fig. 6) could stem from the higher density of poultry farms in southern, compared to northern China. In addition, the relationship of outbreaks to season (Fig. 6) may be due to the higher poultry stocking rates prior to Chinese New Year in January or February, a phenomenon that also governs the incidence of Newcastle disease (16).

Our examination of HPAI H5N1 outbreaks in relation to stages in the annual life cycle of migratory birds suggests a temporal mismatch (Fig. 5). Similar to earlier studies of whooper swans in the East Asian Flyway (26), we found fewer outbreaks than expected during the breeding season, when AIV prevalence typically reaches a maximum in wild bird populations (37). Instead, the peak in observed outbreaks coincided with the nonbreeding season of migratory birds. During the breeding season, birds migrate to latitudes higher than 40°N (Fig. 6), precluding contact with poultry farms. However, the possibility exists that persistence of AIV H5 subtype particles in the environment (2) may be responsible for the lag between the breeding season and the peak in infections two months later. Our data also indicate that migration between nonbreeding and breeding areas is prolonged, with many intervening stopover sites. If AIV particles survive in the environment with increased residence time in colder climates (2), wild birds may act as agents of transmission, “seeding” areas along their migration routes such as rice fields. Larger waterfowl such as geese may defecate as often as every 3.7 minutes (47) and therefore if infected have the

potential to contaminate the environment with high loads of virus. Some duck species, such as the mallard, may shed virus for several days before the onset of clinical signs (25), a period of time in which they could potentially travel several hundred kilometers assuming that the infection did not affect their movements (40). Opportunities for poultry to acquire AIV shed by wild birds at Poyang Lake are created by the practice of herding domestic ducks onto the harvested rice fields to consume waste grains, as well as foraging in wetlands frequented by migratory birds.

Our study identified that most ducks from Poyang Lake migrated across Bohai Bay, a body of water bordering Hebei Province in Northeast China. This area may represent a migratory “thoroughfare” of regional importance for several species of waterbirds such as whooper swans (32), bar-tailed godwits (*Limosa limosa*) (15,35), and swan geese (author’s unpublished data). Studies have identified sites that attract a diversity of wild bird species as important for interspecific transmission of low pathogenic avian influenza virus (7,17,18). Similarly, the convergence of numerous waterbird species from different flyways at Bohai Bay may facilitate exchange of AIVs of different geographic origin. Regional migration stopover sites where wild bird species congregate and intermix are commonplace along other flyways, such as Qinghai Lake in the Central Asian Flyway. These high-density regional stopover areas may serve as catalysts for spread of disease in wild birds (27), just as live poultry or “wet” markets facilitate transmission by bringing together multiple species of domestic fowl drawn from populations of different geographic regions (5,42).

Our tracking results did not show any evidence of a migratory pathway for waterfowl between Poyang Lake in the East Asian Flyway and Qinghai Lake in the Central Asian Flyway (Fig. 1), even when we included all 111 waterbirds marked with satellite transmitters in 2007–08 (author’s unpublished data). In addition to the tracking data for eight species of migratory ducks from Poyang Lake, this included two species (bar-headed geese, ruddy shelduck) involved in the large 2005 outbreak at Qinghai Lake (6,29). Although most waterbirds migrate in a north-south direction, migration of some species may be east-west such as pochards (*Aythya ferina*) (1) and whooper swans (32). It is possible that a larger sample of more species would result in detection of a migratory pathway connecting the East and Central Asian Flyways, but our tracking results to date indicate that, at best, it is not a commonly used route by waterfowl. Alternative possibilities for the movement of HPAI H5N1 from Poyang Lake to Qinghai Lake include that other species may follow that migration route, some species overlap at staging areas such as Dong Ting Lake to the west or Bohai Bay to the north and serve as relay agents, or migration routes from other regions might be involved. For example, our telemetry work in the Central Asian Flyway documented a previously unknown migration route connecting Qinghai Lake to a 2005 outbreak site in Mongolia (34). Another alternative explanation is the trade of poultry or poultry products between these two wetlands. However, because poultry production is largely absent in this migration corridor and we found no connection between the flyways, the movement of HPAI H5N1 from the Poyang Lake and Qinghai Lake regions remains a mystery.

Acknowledgments

This work was supported by the U.S. Geological Survey (Western Ecological Research Center, Patuxent Wildlife Research Center, Alaska Science Center, and Avian Influenza Program), the United Nations Food and Agriculture Organization AGAH-EMPRES Wildlife Unit, National Institutes of Health Fogarty International Center (7R01TW007869-04), the Government of Sweden financial support through donation OSRO/GLO/GO1/SWE, and the Chinese Academy of Sciences (No. 2007FY6210700, INFO-115-D02, KSCX2-YW-N-063, and 2005CB523007). We thank S. Haseltine, R. Kearney, P. Bright, S. Schwarzbach, and J. Howell of USGS and E. Moncada, Q. Gao, V. Martin, F. Guo, J. Lubroth, and J. Domenech of the FAO for support of this project. We thank the staff at Poyang Lake National Nature Reserve, including Liu Guanhua, Manager Zhao, Manager Wu, and Biologist Wu. Ding Changqing and Yin Zuohua (CAS IOZ) provided excellent field assistance, and G. Olsen and B. Siraroonrat assisted with implant surgeries. We are grateful to the Qinghai Lake National Nature Reserve staff, Qinghai Forestry Bureau (S. Li), Chinese Academy of Sciences (X. Hu, L. Hu, N. Kong, Z. Luo), and the U.S. Embassy (W. Chang, D. Jassem) for logistical and field support. We are also grateful to W. Perry for GIS webpage support, C. Hamilton and J. Pinto of FAO for outbreak data from the EMPRES database, and J. Horne for assisting with technical questions on Program Animal Space Use. N. Hill and S. De La Cruz provided helpful to strengthen earlier versions of this manuscript. The use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Abbreviations

AIC	Akaike Information Criterion
AIV	avian influenza virus
BBUD	Brownian bridge utilization distribution
ELISA	enzyme-linked immunosorbent assay
EMPRES	Emergency Prevention System for Transboundary Animal and Plant Pests and Diseases
FAO	Food and Agriculture Organization
GPS	global positioning system
HPAI	highly pathogenic avian influenza
IGBP	International Geosphere-Biosphere Programme
MODIS	Moderate-resolution imaging spectroradiometer
OD	optical density
OIE	Office International des Epizooties
PTT	platform terminal transmitter
RT-PCR	reverse transcriptase-polymerase chain reaction

References

1. Boere, G.; Stroud, D. The flyway concept: what it is and what it isn't. In: Boere, G.; Galbraith, CA.; Stroud, D.; Bridge, LK., editors. *Waterbirds around the world: a global overview of the conservation, management and research of the world's waterbird flyways*. Stationery Office; Edinburgh, Scotland: 2006. p. 940
2. Brown JD, Swayne DE, Cooper RJ, Burns RE, Stallknecht DE. Persistence of H5 and H7 avian influenza viruses in water. *Avian Dis*. 2007; 51:285–289. [PubMed: 17494568]
3. Burnham, KP.; Anderson, DR. *Model selection and multimodel inference: a practical information theoretic approach*. Springer-Verlag; New York: 1998.

4. Carboneras, C. Spot-billed duck. In: del Hoyo, J.; Elliott, A.; Sargatal, J., editors. Handbook of the birds of the world, vol. 1, Ostrich to ducks. Lynx Edicions; Barcelona, Spain: 1992.
5. Cardona C, Yee K, Carpenter T. Are live bird markets reservoirs of avian influenza? *Poult Sci.* 2009; 88:856–859. [PubMed: 19276435]
6. Chen H, Li Y, Li Z, Shi J, Shinya K, Deng G, Qi Q, Tian G, Fan S, Zhao H, Sun Y, Kawaoka Y. Properties and dissemination of H5N1 viruses isolated during an influenza outbreak in migratory waterfowl in western China. *J Virol.* 2006; 80:5976–5983. [PubMed: 16731936]
7. Chen R, Holmes EC. Frequent inter-species transmission and geographic subdivision in avian influenza viruses from wild birds. *Virology.* 2009; 383:156–161. [PubMed: 19000628]
8. Dzubin, A.; Cooch, EG. Measurement of geese: general field methods. California Waterfowl Association; Sacramento: 1992.
9. Food and Agricultural Organization (FAO). Farming systems and poverty: improving farmers' livelihoods in a changing world. Food and Agricultural Organization of the United Nations and World Bank; Rome, Italy: 2001.
10. Food and Agricultural Organization (FAO). Wild birds and avian influenza: an introduction to applied field research and disease sampling techniques. Food and Agricultural Organization of the United Nations and World Bank; Rome, Italy: 2007.
11. Food and Agricultural Organization (FAO). EMPRES-i global animal health information system of FAO's Emergency Prevention Programme for Transboundary Animal Diseases (EMPRES). EMPRES Animal Health Programme, Animal Health Service, Animal Production and Health Division; Rome, Italy: 2009. Available from: <http://empres-i.fao.org/empres-i/home>
12. Feare CJ. The role of wild birds in the spread of HPAI H5N1. *Avian Dis.* 2007; 51:440–447. [PubMed: 17494603]
13. Fouchier RA, Bestebroer TM, Herfst S, Van Der Kemp L, Rimmelzwaan GF, Osterhaus AD. Detection of influenza A viruses from different species by PCR amplification of conserved sequences in the matrix gene. *J Clin Microbiol.* 2000; 38:4096–4101. [PubMed: 11060074]
14. Gilbert M, Xiao X, Pfeiffer DU, Epprecht M, Boles S, Czarneci C, Chaitaweesub P, Kalpravidh W, Minh PQ, Otte MJ, Martin V, Slingenbergh J. Mapping H5N1 highly pathogenic avian influenza risk in Southeast Asia. *Proc Natl Acad Sci U S A.* 2008; 105:4769–4774. [PubMed: 18362346]
15. Gill RE Jr, Tibbitts TL, Douglas DC, Handel CM, Mulcahy DM, Gottschalck JC, Warnock N, McCaffery BJ, Battley PF, Piersma T. Extreme endurance flights by landbirds crossing the Pacific Ocean: ecological corridor rather than barrier? *Proc R Soc B.* 2009; 279:447–457.
16. Higgins, DA.; Shortridge, KF. Newcastle disease in tropical and developing countries. In: Alexander, DJ., editor. Newcastle disease. Springer; New York: 1988. p. 273-302.
17. Hinshaw VS, Webster RG, Turner B. The perpetuation of orthomyxoviruses and paramyxoviruses in Canadian waterfowl. *Can J Microbiol.* 1980; 26:622–629. [PubMed: 7397605]
18. Hinshaw VS, Wood JM, Webster RG, Deibel R, Turner B. Circulation of influenza viruses and paramyxoviruses in waterfowl originating from two different areas of North America. *Bull WHO.* 1985; 63:711–719. [PubMed: 3878741]
19. Horne JS, Garton EO. Likelihood cross-validation versus least squares cross-validation for choosing the smoothing parameter in kernel home-range analysis. *J Wildl Manag.* 2006; 70:641–648.
20. Horne, JS.; Garton, EO. Animal space use 1.2. 2007. [cited 6 Mar 2009]. Available from: http://www.cnr.uidaho.edu/population_ecology/animal_space_use.htm
21. Horne JS, Garton EO, Krone SM, Lewis JS. Analyzing animal movements using Brownian bridges. *Ecology.* 2007; 88:2354–2363. [PubMed: 17918412]
22. John K, Kazwala R, Mfinanga GS. Knowledge of causes, clinical features and diagnosis of common zoonoses among medical practitioners in Tanzania. *BMC Infect Dis.* 2008; 8:162. [PubMed: 19046464]
23. Jorgensen P, an der Heiden M, Kern P, Schoneberg I, Krause G, Alpers K. Underreporting of human alveolar echinococcosis, Germany. *Emerg Infect Dis.* 2008; 14:935–937. [PubMed: 18507906]

24. Kanai Y, Uetaa M, Germogenov N, Nagendran M, Mitad N, Higuchie H. Migration routes and important resting areas of Siberian cranes (*Grus leucogeranus*) between northeastern Siberia and China as revealed by satellite tracking. *Biol Conserv.* 2002; 106:339–346.
25. Keawcharoen J, van Riel D, van Amerongen G, Bestebroer T, Beyer WE, van Lavieren R, Osterhaus AD, Fouchier RA, Kuiken T. Wild ducks as long-distance vectors of highly pathogenic avian influenza virus (H5N1). *Emerg Infect Dis.* 2008; 14:600–607. [PubMed: 18394278]
26. Korschgen CE, Kenow KP, Gendron-Fitzpatrick A, Green WL, Dein FJ. Implanting intra-abdominal radio transmitters with external whip antennas in ducks. *J Wildl Manag.* 1996; 60:132–137.
27. Krapu GL, Reinecke KJ, Jorde DG, Simpson SG. Spring-staging ecology of midcontinent greater white-fronted geese. *J Wildl Manag.* 1995; 59:736–746.
28. Li, ZWD.; Mundkhar, T. Numbers and distribution of waterbirds and wetlands in the Asia-Pacific region: results of the Asian waterbird census 2002–2004. *Wetlands International*; Kuala Lumpur: 2007.
29. Liu J, Xiao H, Lei F, Zhu Q, Qin K, Zhang XW, Zhang XL, Zhao D, Wang G, Feng Y, Ma J, Liu W, Wang J, Gao GF. Highly pathogenic H5N1 influenza virus infection in migratory birds. *Science.* 2005; 309:1206. [PubMed: 16000410]
30. Mukhtar MM, Rasool ST, Song D, Zhu C, Hao Q, Zhu Y, Wu J. Origin of highly pathogenic H5N1 avian influenza virus in China and genetic characterization of donor and recipient viruses. *J Gen Virol.* 2007; 88:3094–3099. [PubMed: 17947535]
31. Muzaffar SB, Ydenberg RC, Jones IL. Avian influenza: an ecological and evolutionary perspective for waterbird scientists. *Waterbirds.* 2006; 29:243–406.
32. Newman SH, Iverson SA, Takekawa JY, Gilbert M, Prosser DJ, Batbayar N, Natsagdorj T, Douglas DC. Migration of whooper swans and outbreak of highly pathogenic avian influenza H5N1 virus in eastern Asia. *PLoS One.* 2009; 4:e5729. [PubMed: 19479053]
33. Office des Internationales Epizooties. *Manual of diagnostic tests and vaccines for terrestrial animals.* World Organization of Animal Health; Paris: 2007. Avian influenza.
34. Prosser DJ, Takekawa JY, Newman SH, Yan B, Douglas DC, Hou Y, Xing Z, Li T, Li Y, Zhao D, Perry WM, Palm EC, Zhang D. Satellite-marked waterfowl reveal migratory connection between H5N1 outbreak areas in China and Mongolia. *Ibis.* 2009; 151:568–576.
35. Qing-Quan B. The finding of Australian satellite tracing Bar-tailed Godwit in Dandong, Liaoning. *China Crane News.* 2008; 12:55–57.
36. R: a language and environment for statistical computing. R Foundation for Statistical Computing; Vienna, Austria: 2008. [cited 6 Mar 2009]. Available from: <http://www.R-project.org>
37. Rundstadler JA, Happ GM, Slemmons RD, Sheng Z-M, Gundlach N, Petrucci M, Senne D, Nolting J, Evers DL, Modrell A, Huson H, Hills S, Rothe T, Marr T, Taubenberger JK. Using RRT-PCR analysis and virus isolation to determine the prevalence of avian influenza virus infections in ducks in Minto Flats State Game Refuge, Alaska, during August 2005. *Arch Virol.* 2007; 152:1901–1910. [PubMed: 17541700]
38. Shankman D, Keim BD, Song H. Flood frequency in China's Poyang Lake region: trends and teleconnections. *Int J Climatol.* 2006; 26:1255–1266.
39. Shankman D, Liang Q. Landscape changes and increasing flood frequency in China's Poyang Lake region. *Prof Geogr.* 2003; 55:434–445.
40. van Gils JA, Munster VJ, Radersma R, Liefhebber D, Fouchier RA, Klaassen M. Hampered foraging and migratory performance in swans infected with low-pathogenic avian influenza A virus. *PLoS ONE.* 2007; 2:e184. [PubMed: 17264886]
41. Wang H-Z, Xu Q-Q, Cui Y-D, Liang Y-L. Macrozoobenthic community of Poyang Lake, the largest freshwater lake of China, in the Yangtze floodplain. *Limnology.* 2007; 8:65–71.
42. Webster RG. Wet markets—a continuing source of severe acute respiratory syndrome and influenza? *Lancet.* 2004; 363:234–236. [PubMed: 14738798]
43. Weitao J, Yun-Bao W. Survey on species and their number of wintering waterbirds in surrounding Poyang Lake area in 2006. *China Crane News.* 2007; 11:37–38.
44. Wint, W.; Robinson, T. *Gridded livestock of the world—2007.* Food and Agriculture Organization; Rome, Italy: 2007.

45. Yasue M, Feare CJ, Bennun L, Fiedler W. The epidemiology of H5N1 avian influenza in wild birds: why we need better ecological data. *BioScience*. 2006; 56:923–929.
46. Zar, JH. *Biostatistical analysis*. 4. Prentice Hall; Upper Saddle River, NJ: 1998.
47. Zhang JX, Liu JJ. Feeding ecology of two wintering geese species at Poyang Lake, China. *Freshwater Ecol*. 1999; 14:439–445.




Fig. 1. General movement paths for more than 111 waterbirds marked with satellite transmitters in the East Asian Flyways depicted in red (this study) and Central Asian Flyways depicted in yellow (author's unpublished data), 2007–08.




Fig. 2. Stopover and staging sites for ducks during the 2007 spring migration including (A) the Yangtze River and (B) Bohai Bay. Stopover sites are represented by 95% fixed kernel home ranges and dates indicate length of stopover periods. Two HPAI H5N1 outbreaks in wild birds from 2003 to 2009 are indicated (○) with the date. Species include Eurasian wigeon (EUWI), common teal (CMTE), falcated teal (FATE), Baikal teal (BATE), and garganey (GARG).




Fig. 3. Brownian bridge utilization distribution for the spring migration of ducks captured and marked at Poyang Lake, China. Groupings include (A) falcated teal, (B) teal (common teal, garganey, Baikal teal), (C) Eurasian wigeon, and (D) Chinese spotbill. Shading from darker to lighter indicates isopleths incorporating 70–99% of total locations.




Fig. 4. Brownian Bridge utilization distribution (BBUD) for ducks from Poyang Lake, China, 2007–08. Shading, from darker to lighter, indicates isopleths incorporating 70–99% of total locations. Darker brown background shading indicates greater density of poultry. Outbreaks of HPAI H5N1 restricted to within the BBUD and occurring 2003–09 are indicated in poultry (▲) and wild birds (○).




Fig. 5.

Comparison of observed (□) and expected (●) HPAI H5N1 outbreak events in the East Asian Flyway by seasonal period in the life cycle of wild ducks. Expected numbers are calculated under the assumption that outbreaks should be proportional to the number of days of a particular life stage. Arrows indicate the direction of the expected from the observed values. Observed and expected values differed ($G_{stat} = 19.98$, $\chi^2_{0.05,3} = 7.82$, $P < 0.001$).




Fig. 6. Relationship between location of ducks and HPAI H5N1 outbreaks according to latitude and seasonal period in the life cycle of wild ducks. Lines indicate the location of all waterfowl, Eurasian wigeon, teal, and outbreaks, where dashed lines indicate missing data. Outbreaks are limited to lower latitudes $<40^{\circ}$ and seem to be detected beginning in September before most wild ducks have migrated south of 40° .

Table 1

Performance of satellite transmitter for ducks marked in 2007–08 at Poyang Lake, China. Species include Baikal teal (BATE), common teal (CMTE), Eurasian wigeon (EUWI), falcated teal (FATE), garganey (GARG), mallard (MALL), northern pintail (NOPI), and Chinese spotbill duck (SBDU). Estimated accuracy of location codes (from best to least, left to right) are described in the Methods.

PTT	Species	Age	Sex	Last message	Working days	Total locations	Location class						
							3	2	1	0	A	B	Z
73290	BATE	A	M	Apr. 17, 2007	29	36	2	1	6	1	11	13	2
74827	BATE	A	M	Mar. 31, 2007	14	9	0	0	1	1	2	5	0
74826	CMTE	A	M	Oct. 1, 2007	198	163	3	12	17	13	43	69	6
74828	CMTE	A	M	Nov. 24, 2008	617	614	8	27	44	118	135	266	16
74829	CMTE	A	M	Jan. 3, 2008	59	291	1	1	1	2	4	13	2
73020	CMTE	J	M	Nov. 3, 2007	231	488	22	74	111	118	70	78	15
39571	EUWI	A	F	Jun. 16, 2008	200	6	0	0	0	0	2	4	0
40217	EUWI	A	F	Nov. 3, 2008	340	221	74	50	19	17	34	27	0
40754	EUWI	A	F	Oct. 17, 2008	323	156	13	26	25	22	28	40	2
73007	EUWI	A	F	Dec. 25, 2007	283	909	56	162	285	218	68	104	16
73031	EUWI	A	M	Jul. 1, 2007	104	2	0	0	0	0	1	1	0
73025	FATE	A	M	Jun. 10, 2008	451	2	0	0	0	0	1	1	0
73034	FATE	A	M	Sep. 12, 2007	179	282	10	23	47	38	73	84	7
73288	FATE	A	M	Nov. 6, 2007	234	222	5	10	19	31	72	80	5
73289	FATE	A	M	Oct. 9, 2008	570	795	15	24	64	146	175	335	36
74830	FATE	A	F	Jul. 31, 2007	134	17	0	1	2	3	6	5	0
73017	FATE	J	F	Jul. 29, 2008	500	239	9	15	39	38	59	75	4
74831	GARG	A	M	Mar. 29, 2008	376	341	23	37	45	41	75	111	9
73294	MALL	A	M	May 4, 2008	414	1002	5	12	20	17	42	106	14
73032	NOPI	J	F	Jan. 24, 2008	54	35	0	1	6	2	9	16	1
45972	SBDU	A	F	Jan. 14, 2009	412	856	20	33	31	32	80	69	15
46082	SBDU	A	F	Dec. 9, 2007	10	19	0	0	1	1	1	4	0
46083	SBDU	A	M	Active	424	2252	50	76	95	71	109	143	15
46126	SBDU	A	F	Dec. 23, 2007	22	58	0	1	0	7	3	4	0
46127	SBDU	A	F	Dec. 19, 2007	18	12	0	0	0	2	2	4	0

PTT	Species	Age	Sex	Last message	Working days	Total locations	Location class						Z
							3	2	1	0	A	B	
46128	SBDU	A	M	Nov. 9, 2008	342	1668	52	63	46	42	80	113	7
46129	SBDU	A	M	Aug. 7, 2008	250	936	17	32	33	32	63	88	13
46130	SBDU	A	M	Aug. 23, 2008	266	916	8	21	31	17	78	113	10
73022	SBDU	A	F	May 13, 2007	62	66	10	7	7	5	12	21	4
73043	SBDU	A	M	Jan. 14, 2008	301	708	15	15	20	12	68	84	14
73044	SBDU	A	F	Apr. 9, 2007	21	76	0	0	2	5	5	7	1
73296	SBDU	A	M	Jan. 2, 2009	655	833	23	28	60	49	67	109	14
74815	SBDU	A	F	Active	422	471	2	6	12	38	72	81	13

Table 2

Spring migration areas in 2007 for wild ducks migrating from Poyang Lake, China. Species include Baikal teal (BATE), common teal (CMTE), falcated teal (FATE), garganey (GARG), and Eurasian wigeon (EUWI).

Date range	Country	Site name	Coordinates (latitude, longitude)	Distance from Poyang Lake (km)	Length of stay (range in days)	Number	Total locations	Species
Mar. 30–May 18	China	Lower Yangtze River	31.11°N, 117.81°E	330	11 (1–37)	7	61	1 BATE, 4 CMTE, 2 FATE, 1 GARG
Apr. 29–Jun. 9	China	Lower Huang River	37.80°N, 118.82°E	1210	10 (1–21)	3	31	2 CMTE, 1 GARG
Apr. 15–Jun. 2	China, N. Korea	North Yellow Sea Coastal Wetlands	39.85°N, 124.26°E	1600	7 (1–27)	7	56	1 BATE, 2 FATE
Apr. 20–Jun. 3	China	Upper Liao River Basin	42.72°N, 123.26°E	2060	12 (1–28)	5	51	2 CMTE, 1 EUWI, 1 FATE, 1 GARG
May 19–Jun. 11	China	Songhua River Wetlands	45.97°N, 112.12°E	2680	7 (1–20)	3	30	2 CMTE, 1 FATE
Apr. 27–May 12	Russia	Lower Amur River Basin	52.94°N, 141.17°E	4540	4 (1–10)	3	7	3 EUWI
May 14–May 29	Russia	Shelikhov Gulf, Sea of Okhotsk	60.26°N, 158.73°E	7000	12 (10–14)	2	41	2 EUWI

Table 3

Best-fitting models under Akaike Information Criterion (AIC) model selection. Outbreak locations from 2003 to 2009 (n = 26) are compared with random points (n = 26). Variables include outbreak latitude (Lat), poultry density (Dens), utilization distribution for satellite-marked wild ducks (BBUD), urban areas (Urban), cropland (Crop), water, and wetlands. Number of parameters (k), AIC score, AIC difference (ΔAIC), likelihood, and AIC weights are reported. Models with biologically significant AIC scores are indicated (*).

Model	Description	k	AIC	ΔAIC	Likelihood	AIC weight
1	Lat+Dens*	3	66.48	0.00	1.00	0.36
2	Lat+Dens+BBUD*	4	67.85	1.38	0.50	0.18
3	Lat+Urban+Crop+BBUD	5	69.04	2.57	0.28	0.10
4	Lat+Crop+Dens+BBUD	5	69.39	2.91	0.23	0.08
5	Lat+Water+Wetland+Dens	5	69.97	3.49	0.17	0.06
6	Lat+Urban+Crop+Dens+BBUD	6	70.05	3.57	0.17	0.06
7	Dens	2	71.62	5.14	0.08	0.03
8	Lat+Water+Urban+Crop+Dens+BBUD	7	71.64	5.16	0.08	0.03
9	Lat	2	71.71	5.23	0.07	0.03
10	Lat+Water+Wetland+Crop +Dens	6	71.85	5.37	0.07	0.03
11	Lat+UD	3	72.59	6.11	0.05	0.02
12	Lat+Water+Wetland+Urban+Crop+Dens+BBUD	8	73.51	7.03	0.03	0.01
13	Lat+Urban+Crop+UD	5	74.14	7.66	0.02	0.01
14	Full model: Lat+Water+Upland+Wetland+Urban+Crop+Dens+BBUD	9	74.20	7.72	0.02	0.01
15	Lat+Water+Wetland+Crop	5	76.70	10.21	0.01	0.00
16	Urban+Crop+Dens+BBUD	5	77.58	11.10	0.00	0.00
17	Null model	0	82.41	15.93	0.00	0.00

Table 4

Top competing AIC models within 2.0 AIC encompassing the variables: intercept, outbreak latitude, poultry density, and wild bird utilization distribution. Statistics include the β coefficients, SE, z -value, significance value, and a χ^2 test comparison to the null model.

Model	Variables	β	SE	z	Pr ($> z $)	Pr ($> \chi^2 $)
1	Intercept	4.3660	2.2050	1.98	0.05	–
	Outbreak latitude	-0.1440	0.0572	-2.52	0.01	<0.001
	Poultry density	0.0001	0.0001	2.23	0.03	0.007
2	Intercept	0.4621	5.4200	0.09	0.93	–
	Outbreak latitude	-0.1572	0.0601	-2.62	0.01	<0.001
	Poultry density	0.0001	0.0001	2.18	0.03	0.007
	Utilization distribution	4.8080	6.1880	0.78	0.44	0.429

Table 5

Summary of virological analyses for the 20 birds marked during March 2007.

PTT	Species	RT-PCR		Sandwich ELISA AIV-Ag (OD ₆₃₀) ^A			Serum
		Tracheal swab	Cloacal swab	Tracheal swab	Cloacal swab	Indirect ELISA AIV-H5-Ab (OD ₆₃₀) ^B	
73290	BATE	Negative	Negative	0.15	0.23	0.30	
74827	BATE	Negative	Negative	0.18	0.21	0.26	
73020	COTE	Negative	Negative	0.23	0.29	0.17	
74826	COTE	Negative	Negative	0.16	0.29	0.32	
74828	COTE	Negative	Negative	0.25	0.26	0.33	
74829	COTE	Negative	Negative	0.19	0.26	0.21	
73007	EUWI	Negative	Negative	0.22	0.18	0.14	
73031	EUWI	Negative	Negative	0.22	0.29	0.38	
73017	FATE	Negative	Negative	0.25	0.31	0.31	
73025	FATE	Negative	Negative	0.23	0.28	0.19	
73034	FATE	Negative	Negative	0.28	0.22	0.27	
73288	FATE	Negative	Negative	0.23	0.23	0.14	
73289	FATE	Negative	Negative	0.18	0.25	0.20	
74830	FATE	Negative	Negative	0.12	0.13	0.20	
74831	GARG	Negative	Negative	0.15	0.21	0.22	
73294	MALL	Negative	Negative	0.27	0.22	0.50 ^C	
73022	SBDU	Negative	Negative	0.24	0.25	0.24	
74043	SBDU	Negative	Negative	0.22	0.28	0.23	
74044	SBDU	Negative	Negative	0.27	0.29	0.22	
73296	SBDU	Negative	Negative	0.20	0.21	0.19	

^A An optical density (at 630 nm) 0.30 was considered positive

^B An optical density (at 630 nm) 0.40 was considered positive.

^C Denotes positive sample.