

SCIENTIFIC REPORTS



OPEN

Genome-wide analysis reveals adaptation to high altitudes in Tibetan sheep

Received: 22 June 2015

Accepted: 09 May 2016

Published: 27 May 2016

Caihong Wei^{1,*}, Huihua Wang^{1,2,3,*}, Gang Liu^{2,*}, Fuping Zhao¹, James W. Kijas⁴, Youji Ma⁵, Jian Lu², Li Zhang¹, Jiaxue Cao¹, Mingming Wu¹, Guangkai Wang¹, Ruizao Liu¹, Zhen Liu¹, Shuzhen Zhang¹, Chousheng Liu² & Lixin Du¹

Tibetan sheep have lived on the Tibetan Plateau for thousands of years; however, the process and consequences of adaptation to this extreme environment have not been elucidated for important livestock such as sheep. Here, seven sheep breeds, representing both highland and lowland breeds from different areas of China, were genotyped for a genome-wide collection of single-nucleotide polymorphisms (SNPs). The F_{ST} and XP-EHH approaches were used to identify regions harbouring local positive selection between these highland and lowland breeds, and 236 genes were identified. We detected selection events spanning genes involved in angiogenesis, energy production and erythropoiesis. In particular, several candidate genes were associated with high-altitude hypoxia, including *EPAS1*, *CRYAA*, *LONP1*, *NF1*, *DPP4*, *SOD1*, *PPARG* and *SOCS2*. *EPAS1* plays a crucial role in hypoxia adaptation; therefore, we investigated the exon sequences of *EPAS1* and identified 12 mutations. Analysis of the relationship between blood-related phenotypes and *EPAS1* genotypes in additional highland sheep revealed that a homozygous mutation at a relatively conserved site in the *EPAS1* 3' untranslated region was associated with increased mean corpuscular haemoglobin concentration and mean corpuscular volume. Taken together, our results provide evidence of the genetic diversity of highland sheep and indicate potential high-altitude hypoxia adaptation mechanisms, including the role of *EPAS1* in adaptation.

The Tibetan Plateau represents 25% of the landmass of China. It is the largest high-altitude area on earth, with an average altitude exceeding 4,500 m¹. Such a high-altitude environment has produced many unique highland species. Compared with neighbouring and lowland populations, the native people and animals have adapted to this habitat, which has a 43% lower partial pressure of oxygen² and 40% higher ultraviolet radiation³. Several animals, such as Tibetan antelopes⁴ and Tibetan Mastiffs^{5,6}, have recently been shown to exhibit adaptation to living in this challenging environment. China has many sheep breeds, such as Tibetan and Mongolian sheep, that are widely distributed from the highlands to lowlands, many of which have evolved over centuries or even millennia^{7,8}. Therefore, they also represent ideal organisms to study plateau adaptability.

To date, the genome sequences of Tibetan antelopes⁴, Tibetan wild boars⁹, yaks¹⁰ and ground tits¹¹ have been generated, and several important pathways and functional categories have been identified, including energy metabolism and oxygen transmission, response to hypoxia, DNA repair and ATPase production. In addition, the mechanisms underlying plateau adaptability have been explored using population surveys of SNP data, successfully identifying candidate genes for genetic adaptation to the Tibetan Plateau. Recent studies have demonstrated that the positively selected haplotypes of *EGLN1* and *PPARA* were significantly associated with the low haemoglobin content of Tibetan people, which is a unique characteristic of this extreme high-altitude population¹². One single-nucleotide polymorphism (SNP) in *EPAS1*, encoding a transcription factor involved in response to

¹Institute of Animal Sciences, Chinese Academy of Agricultural Sciences, National Center for Molecular Genetics and Breeding of Animal, Beijing, People's Republic of China. ²National Animal Husbandry Service, National Center of Preservation & Utilization of Animal Genetic Resources, Beijing, People's Republic of China. ³Institute of apicultural research, Chinese Academy of Agricultural Sciences, Beijing, People's Republic of China. ⁴Livestock Industries, CSIRO, Brisbane, Australia. ⁵College of Animal Science and Technology, Gansu Agriculture University, Lanzhou 730070, People's Republic of China. ^{*}These authors contributed equally to this work. Correspondence and requests for materials should be addressed to C.L. (email: liuchousheng@sina.com) or L.D. (email: lxdu@263.net)

hypoxia, was identified; for this SNP, there was a 78% frequency difference between the Tibetan and Han populations, representing the largest allele frequency difference observed for any human gene to date¹³. Recently, variations in *EPAS1* were identified as high-altitude adaptations in the Tibetan Mastiff^{5,6}. Additionally, other hypoxia-related genes were identified in yaks (*ADAM17*, *ARG2* and *MMP3*), Tibetan antelopes (*ADORA2A*, *CCL2*, *ENG*, *PIK3C2A*, *PKLR*, *ATP12A* and *NOS3*) and Tibetan wild boars (*ALB*, *ECE1*, *GNG2* and *PIK3C2G*)^{4,9,10}. In addition, Qu found 11 candidate positively selected genes (*HIFAN*, *MTOR*, *SRF*, *TXNRD2*, *WNT7B*, *ANGP4*, *ADAM9*, *PSMD2*, *LRRC7*, *MDH1B* and *LRRK1*) associated with a hypoxia response in ground tits¹¹. Almost all of the identified genes mentioned above belong to the list of 247 hypoxia genes that are priority candidates for adaptation to high-altitude hypoxia¹²; however, each species may have different candidate genes. This phenomenon might indicate differences between species in the specific genes selected, even in the same environment. Thus, Tibetan sheep might differ from other species with respect to the molecular mechanisms of high-altitude adaptation. However, there has been no research on the adaptation of Tibetan sheep to the high-altitude plateau.

Data from several amphibians and mammals have indicated that modifications of haemoglobin (Hb) function often play a key role in regulating an adaptive response to high-altitude hypoxia^{14–16}. For example, the haemoglobin concentration of Andean individuals who are native to different elevations is higher than those of Tibetan and sea-level residents^{17,18}. In addition, improved oxygen (O₂) loading and unloading are physiological marks of high-altitude adaptation¹⁹. The O₂-loading capacity is determined by the Hb-O₂ affinity. An increased Hb-O₂ affinity helps maintain pulmonary O₂ loading at a sufficient to maximal level of tissue oxygenation under conditions of extreme hypoxia²⁰. Many highland animals have haemoglobin with high O₂ affinity, such as alpacas²¹, Andean geese²², bar-headed geese^{23,24} and deer mice^{25,26}.

Although many studies have focused on highland humans and animals, the genetic mechanisms underlying the adaptation of domesticated animals to the Tibetan Plateau have been rarely studied. In this study, a genome-wide investigation of the adaptation of Tibetan sheep to the high-altitude plateau was conducted using the Illumina Ovine SNP50K Bead Chip assay. We performed whole-genome SNP sweeps to study the adaptive evolution of high-altitude sheep by analyzing seven breeds. We identified candidate genes using selective sweep mapping and revealed potential genetic mechanisms of high-altitude adaptation in sheep compared with other species. We also more deeply analyzed gene evolution by investigating genes and networks as well as the genetic diversity of *EPAS1*.

Results

Population structure. To assess the relationships between the animals and breeds under investigation, we applied multidimensional scaling (MDS) to analyze 122 individuals genotyped at 20,632 autosomal SNPs. The individuals were representative of seven native Chinese sheep breeds: Hu sheep (HUS), Tong sheep (TON), Large-tailed Han sheep (LTH), Lop sheep (LOP), Tibetan sheep of the Qinghai (TIBQ), Sichuan (TIBS) and Nagqu (TIBN). (Figure 1A, Supplemental Table S1). The first dimension (Component 1) separated domestic individuals into two broad non-overlapping clusters (Fig. 1B). The clusters corresponded well with the geographic origin of each breed within the cluster. Our cluster results showed that the TIBQ, TIBS and TIBN breeds were grouped together as the highland group (G1) and that the HUS, TON, LTH and LOP breeds belonged to the lowland group (G2). To explore further the relationships obtained using the MDS approach, we performed a population structure analysis using the program STRUCTURE 2.3.4²⁷ to obtain a model-based unsupervised hierarchical clustering of the individuals arising from a user-defined number of ancestral populations (*K*). The number of populations was varied from *K* = 2 to 4, and the largest change in the log of the likelihood function (ΔK) was found when *K* = 2 (Fig. S1), which suggests that there is likely a small number of ancestral groups. Additionally, the individuals from low-altitude breeds appeared distinct from those sampled from high-altitude breeds (Fig. 1C, Fig. S2), which was consistent with the MDS results. To further confirm the phylogenetic relationships among the sheep breeds, a neighbour-joining tree²⁸ based on the pairwise genetic distances was constructed. The tree also indicated that high-altitude (TIBQ, TIBS and TIBN) and low-altitude breeds (TON, HUS, LOP and TLH) were split into two distinct branches (Fig. 1D).

Selection signal is correlated with high altitude. F_{ST} , which can reveal a high coefficient of genetic differentiation, has been widely used to identify selective signals among whole-genome SNPs²⁹. We evaluated the population differentiation and calculated the F_{ST} for each SNP between G1 and G2, ultimately identifying 464 SNPs with extremely high values ($F_{ST} > 0.242$, top 1%) from the total tested (46,355 SNPs, Supplemental Table S2). Moreover, we used F_{ST} outlier tests by LOSITAN³⁰ to identify loci with levels of differentiation between the G1 and G2 pairs that were higher or lower than expected under a neutral model, and approximately 96% of the outlier loci were the same as the top 1% F_{ST} value SNPs obtained using the Genepop software³¹ (Supplemental Table S2). Moreover, 40,652 pairwise XP-EHH (Cross Population Extend Haplotype Homozygosity Test) values, which are haplotype-based parameters, were calculated for each SNP³². In total, 2000 unique SNPs were within the top 5% of XP-EHH values (XP-EHH > 0.415) (Supplemental Table S3). The SNPs that were identified by high F_{ST} and XP-EHH values produced a subset of 171 SNPs, which represented 152 candidate genes with strong signatures of selection (Supplemental Table S4). In addition, 70 SNPs with high F_{ST} but null XP-EHH values were included, representing 84 candidate genes (Supplemental Table S5). Finally, genome-wide polymorphisms revealed chromosomal regions that contained 236 genes with evidence of positive selection.

We performed Gene Ontology (GO) enrichment analyses on the highlighted candidate genes in Tibetan sheep populations (Table 1). These genes are involved in response to hypoxia (GO:0001666) and several other biological processes, such as cell morphogenesis involved in neuron differentiation (GO:0048667), axonogenesis (GO:0007409), tissue homeostasis (GO:0001894) and cell maturation (GO:0048469), all of which likely play roles in the local adaptation of Tibetan sheep. To further understand the functions of the candidate genes, we compared them with several genes that are considered likely to be associated with high-altitude adaptation, which led us to




Figure 1. Description of the location and genetic relationships of samples. (A) Sampling and habitats of the seven Tibetan sheep breeds. The map was created using the R package ‘maptools’, version: 0.8–37, URL: <http://r-forge.r-project.org/projects/maptools/>. (B) Plots for the first (Component 1) and second (Component 2) dimensions revealed the clustering of all individuals. (C) Genome-wide admixtures inferred by STRUCTURE 2.3.4. The results from $K = 2$ are shown; (D) Neighbour-joining (NJ) phylogenetic tree for the seven breeds based on pairwise F_{ST} . Each population is represented by a different symbol and colour label: high-altitude breeds are indicated in red, and low-altitude breeds are indicated in blue. The abbreviations for the seven breeds are shown in Supplemental Table S1.

ID	Term	P value	$P_{\text{Bonferroni}}$ value	Associated genes
GO:0048667	cell morphogenesis involved in neuron differentiation	6.60E-05	0.0017	CHN1, CNP, EPHA8, FGFR2, NUMBL, OMG, SOD1, SPTBN4, TBR1
GO:0022829	wide pore channel activity	7.23E-05	0.0017	GJA3, GJB2, VDAC2
GO:0007409	axonogenesis	1.26E-04	0.0029	CHN1, CNP, EPHA8, FGFR2, NUMBL, OMG, SPTBN4, TBR1
GO:0016877	ligase activity, forming carbon-sulfur bonds	5.51E-04	0.0121	ACLX, ACSL6, SLC27A2
GO:0005681	spliceosomal complex	6.95E-04	0.0146	DHX38, HNRNPK, RALY, SMNDC1, TXNL4B, U2AF1
GO:0001666	response to hypoxia	2.71E-02	0.0271	EPAS1, CRYAA, LONP1
GO:0001894	tissue homeostasis	2.71E-02	0.0271	EPAS1, PTHLH, SOD1
GO:0048469	cell maturation	2.71E-02	0.0271	EPAS1, EPHA8, PPARG

Table 1. GO terms enriched with candidate genes for high-altitude adaptation in Tibetan sheep.

prioritize a set of 317 functional candidate genes (Supplemental Table S6). In our Tibetan sheep breeds, we found that the set of candidate loci contained *EPAS1*, *NF1*, *LONP1*, *SOD1* and *DPP4* as well as the homologous genes *PPARG* and *TGFBR3* of the priority candidate genes.

Hypoxia-inducible factors (HIFs) are transcription factors that respond to changes in the available oxygen in the cellular environment under high-altitude conditions. *EPAS1*, also known as HIF2a, is a member of the HIF family that responds to changes in available oxygen in the cellular environment under high-altitude conditions³³. Three candidate genes, *FGFR2*, *NF1* and *RasGef4*, function in the Ras/ERK signalling pathway, which commonly promotes angiogenesis with the HIF pathway under hypoxia³⁴ and may have been selected for this function. Ras is indirectly activated by *FGFR2*³⁵, and *RasGef4*^{36,37} activates Rap1, which regulates the proliferation and migration of human umbilical vein endothelial cells via the ERK and Akt pathways. *NF1* is a negative regulator of the Ras signal transduction pathway³⁸, and the loss of *NF1* results in the activation of the Ras signalling pathway, leading to the aberrant growth of haematopoietic cells³⁹. *DYSF* and *ZEB1* are also target genes of HIFs. The genetic loss of dysferlin (encoded by *DYSF*) caused an abrogated angiogenic response to vascular endothelial growth factor⁴⁰. A study found that *ZEB1* is critical in the regulation of the macrovascular angiogenic response but not that of

microvascular angiogenesis⁴¹. Moreover, under hypoxic conditions, HIF-1 α up-regulates the expression of proteins that induce *TWIST1* and *ZEB1*^{42,43}. However, in different animals, different genes affecting angiogenesis have been detected, such as *DAM17*¹⁰, *NOS3*⁴ and *PLXNA4*⁶ in yaks, Tibetan antelope and Tibetan Mastiffs, respectively.

HIFs were discovered because of their ability to stimulate transcription of the erythropoietin (EPO) gene during hypoxia at high altitudes^{43–45}. Here, *SOCS2* and *LONP1* were also identified as candidate selected genes associated with the primary role of HIFs in promoting the hypoxic EPO response. *SOCS2* regulates EPO-enhanced neuronal differentiation⁴⁶ and, together with *SOCS3*, can regulate EPO signalling in response to hypoxia⁴⁷. Lon peptidase 1 (*LONP1*) is a multifunctional ATP-dependent protease that mainly participates in mitochondrial proteolysis, is involved in heme biosynthesis^{48,49} and can regulate the EPO gene⁵⁰. Under conditions of reduced O₂ availability, HIF-1 α reciprocally regulates *COX4* subunit expression by activating the transcription of the gene encoding *LONP1*⁵¹.

Candidate genes participating in energy metabolism have been identified regularly in animals living in high-altitude areas^{4,10,11,52}. In Tibetan sheep, we identified a few genes associated with energy metabolism that were under positive selection. *DPP4* is a key enzyme involved in glycolysis, where it degrades GLP-1, which is powered by the hydrolysis of ATP, and plays crucial roles in maintaining glucose stability^{53–55}. *PPARG*, another selected gene, reactivates adipogenesis and transcriptionally activates *LPP1* expression, indicating a potential role in the metabolism of phospholipids^{56,57}. Substantial evidence has shown that mitochondrial function is altered in high-altitude adaptations⁵⁸. A change is initiated by reduced oxygen delivery to cells during metabolism, and reactive oxygen species (ROS) are produced by hypoxic mitochondria, which stimulate the activation of HIF-1 and HIF-2^{59,60}. *SOD1*, the anti-oxidant superoxide dismutase 1, which activates ATP production⁶¹ and reduces mitochondrial ROS production⁶², was identified as being under positive selective pressure in Tibetan sheep.

Hypoxia causes an influx of Ca²⁺ and an increase in intracellular Ca²⁺ concentrations by opening store-operated Ca²⁺ (SOC) channels, which results in increased cell contraction⁶³. In our study, *RYR3* was found to encode the ryanodine receptor 3, a calcium channel, and showed selective signatures in a study in Tibetan Mastiffs⁶. The calcineurin genes (*PPP3R1* and *PPP2R1B*), which encode protein phosphatases, play crucial roles in regulating Ca²⁺/calmodulin. Calcineurin can maintain vascular structure and function, and vascular smooth muscle cell proliferation. Calcineurin is a unique calcium/calmodulin-regulated protein phosphatase that functions as a key mediator of the hypertrophic response of the heart. *PPP3R1* (protein phosphatase 3) and *PPP2R1B* (protein phosphatase 2) are regulatory subunits of calcineurin that are important for phosphatase activity⁶⁴.

Tibetan sheep develop low partial oxygen pressure and are exposed to high levels of ultraviolet radiation. Another important candidate selected gene is microphthalmia-associated transcription factor (*MITF*). Hypoxia up-regulates cyclooxygenase-2, leading to the prostaglandin E(2)-mediated loss of *MITF* in cervical stromal cells⁶⁵. In our study, *MITF*, which is associated with melanogenesis, was detected as positively selected in Tibetan sheep. *MITF* was also identified as being subject to high selective pressure in the world's sheep breeds⁶⁶.

HSD11B1L, *HP*, *TGFBR3* and *MSRB3* were also identified as candidate selected genes in Tibetan sheep. *HSD11B1*, which encodes an isoform of the enzyme 11beta-hydroxysteroid dehydrogenase, acts exclusively as an NAD-dependent dehydrogenase in activating cortisol to cortisone⁶⁷, whose activity is down-regulated under high-altitude hypoxia conditions⁶⁸. In Tibetan sheep, another isoform product of *HSD11B1L*, which is believed to act predominantly as an oxo-reductase using NADP(H) as a cofactor to generate cortisone⁶⁹, was identified as a candidate selected gene; this result indicates that 11beta-HSD1a is also involved in the adaptive response to high altitudes. *TGFBR3* is associated with TGF- β , which is an HIF-responsive product that is suspected of playing a role in cancer^{70,71}. *TGFBR3* is involved in regulating cell growth and differentiation and in inflammatory responses^{72–74}. Haptoglobin (Hp; encoded by *HP*) is a plasma glycoprotein, the main biological function of which is to bind free Hb and prevent the loss of iron and subsequent kidney damage following intravascular haemolysis⁷⁵. The gene encoding methionine sulfoxide reductase B3 (*MSRB3*) has been identified as a gene associated with altitude in dogs⁵, although its association with ear types has also been previously reported^{76,77}. In the world's sheep breeds, *MSRB3* on sheep chromosome 3 was noted to develop high selection pressure⁶⁶.

Some particularly extreme values for either F_{ST} or XP-EHH were also noticed (Fig. 2). For F_{ST} , three genes that had undergone strongly selection were reported in previous studies^{66,78}. *PPP1CC* (F_{ST} = 0.78, chromosome 13) is a positional candidate locus for skeletal muscle strength phenotypes⁷⁹. *RXFP2* (F_{ST} = 0.72, chromosome 10) is a candidate gene for sheep horns⁸⁰. *BMP2* (F_{ST} = 0.63, chromosome 13) is associated with body size traits⁸¹. Additionally, for XP-EHH, four genes with strong signatures of selection were detected in the highland group. *WDR92* (XP-EHH = 1.83, chromosome 3) promotes apoptosis induced by tumour necrosis factor- α (TNF- α)⁸². *PNO1* (XP-EHH = 1.62, chromosome 3) is vital to normal cell function⁸³. *PARK2* (XP-EHH = 1.30, chromosome 8) is expressed primarily in the nervous system and is part of the multi-protein E3 ubiquitin ligase complex⁸⁴. *OSR2* (XP-EHH = 1.25, chromosome 9) is a key intrinsic regulator of palatal growth and patterning⁸⁵.

EPAS1 mutations and physiological associations. To further study the functions of the core factor EPAS1 in high-altitude adaption, we measured six haematological parameters in a high-altitude breed Tibetan sheep breed (TIBQ) and a low-altitude sheep breed (LTH). There were no significant differences between males and females within the breeds (Supplemental Table S7). In the high-altitude breeds, the red blood cell (RBC) count, haemoglobin (HGB) concentration, haematocrit (HCT) concentration, mean corpuscular volume (MCV) and mean corpuscular haemoglobin (MCH), but not the mean corpuscular haemoglobin concentration (MCHC), were higher or significantly higher than those of the low-altitude breeds (Supplemental Table S7). This result was similar to that found in dogs⁵. *EPAS1* polymorphisms in native Tibetan people are associated with lower haemoglobin concentrations¹³.




Figure 2. Genome-wide distribution of F_{ST} and XP-EHH values. Red dots represent sites showing significant signals in both the F_{ST} and XP-EHH approaches; black dots represent sites showing significant signal in the F_{ST} approach only. The symbols for candidate genes for adaptation to high-altitude hypoxia in the map are shown in bold and italics.




Figure 3. Genotype-phenotype association with the (A) RBC, (B) MCV and (C) MCH parameters for the 11th SNP in *EPAS1* in the Tibetan breeds. *Indicates a significant difference between the genotypes; **Indicates an extremely significant difference between the genotypes.

To understand the relationship between *EPAS1* and haematological parameters, we designed 10 pairs of primers and detected 12 mutations in the exons of *EPAS1* using Sanger sequencing. Ten mutations were in the coding region, including three non-synonymous mutations, six synonymous mutations, and an AGC insert; two SNPs were in the 3' untranslated regions (Supplemental Table S8). Chi-squared tests showed that the frequencies of the 1st, 2nd, 5th and 11th mutations were significantly different ($P < 0.05$) between the populations (Supplemental Table S9). In the Tibetan breeds, alleles of these four SNPs were tested for association with haematological parameters, and only the 11th SNP showed a significant difference. The CC genotype was associated with a significantly higher RBC count and significantly lower MCV and MCH than the TT genotype (Fig. 3).

Discussion

Tibetan sheep are a hypoxia-tolerant species that live in an extremely inhospitable high-altitude environment, which has high ultraviolet radiation and a low partial pressure of oxygen compared with low-altitude areas⁸⁶. In our study, we genotyped 122 sheep from seven breeds from high and low altitudes using a 50K SNP chip. SNP diversity was examined within the samples of native sheep that were the first to be domesticated by humans. This was the first study to characterize the genetic polymorphisms and evolution of Tibetan sheep.

The examination of a greater number of SNPs allowed STRUCTURE and MDS to robustly detect a distinct geographic pattern within the breeds genotyped. In our study, the seven Chinese indigenous breeds could be divided into two groups (G1 and G2) based on their genetic structure. G1 included Tibetan sheep (TIBQ, TIBS and TIBN) that live in the Qinghai Tibet plateau mountainous area, with an altitude over 3000 m, whereas

EPAS1 plays an important role in high-altitude adaptation in humans and dogs^{5,13}. Tibetans and Tibetan Mastiffs display higher blood flow for oxygen delivery^{5,102}, and this gene is related to the haemoglobin concentration in Andeans¹⁰³ and Tibetans^{13,104–106}. Therefore, we propose that similar mechanisms may have facilitated sheep adaptation to high-altitude hypoxia. In this study, we also found that *EPAS1* had a high positive selection signal in Tibetan sheep breed. One limitation of this study is its relatively small sample size; consequently, the conclusions cannot be extrapolated to all sheep populations. In this study, we examined 12 mutations in the exons of *EPAS1* in detail. However, only four SNPs with different allele frequencies between high- and low-altitude breeds were identified. Mutations in the *EPAS1* coding region and phenotypes are not associated with haematologic parameters; however, in the 3' untranslated regions, one SNP did show an association: the CC genotype was associated with a significantly higher RBC count and significantly lower MCV and MCH compared to the TT genotype in Tibetan sheep. MCV was found to differ among humans living at high altitudes and at sea level, with significantly higher values associated with humans living at higher altitudes¹⁰⁷. These results indicate that increased MCV and MCH are associated with the enhanced ability to carry oxygen under conditions of high-altitude hypoxia. Consistent with this observation, reports of *EPAS1* mutations in human patients have been exclusively associated with haematological phenotypes, especially alterations in MCH¹⁰⁴. Our results indicate that the allele targeted by selection likely confers a functionally relevant adaptation to the hypoxic environment at high altitudes, although the precise physiological mechanism remains to be discovered.

Methods

Ethics Statement. All animals were handled according to the Guidelines for the Biological Studies Animal Care and Use Committee, People's Republic of China. Animal experiments were approved by the Animal Ethics Committee of the Institute of Animal Sciences of Chinese Academy of Agricultural Sciences.

Genotyping and quality control. For the six Chinese sheep breed data set (Supplemental Table S1), blood samples were collected from 12 HUS, 15 TON, 15 LTH, 15 LOP, 14 TIBQ and 14 TIBS sheep. These animals were recently used in the registration and recording system of the National Center for Preservation and Utilization of Genetic Resources of Domestic Animals, National Animal Husbandry Service, Beijing, China. All experimental and surgical procedures were approved by the Biological Studies Animal Care and Use Committee, People's Republic of China. In total, 87 DNA samples were extracted by the IQS DNA extraction method and were genotyped using the Illumina Ovine SNP 50K Bead Chip assay system at Capital Bio Corporation (Beijing, China). The second data set used in this study came from the Ovine HapMap project of the International Sheep Genomics Consortium and is publicly available at <http://www.sheephapmap.org>. The Chinese TIBN breed including 37 individuals was selected in the current study (Supplemental Table S1). We combined the two data sets and obtained 122 individuals and 49,034 SNPs. SNPs that could not pass the following three criteria were removed: (1) SNPs with a minor allele frequency >0.01 ; (2) maximum per-SNP missing rate <0.05 ; and (3) SNPs on chromosome X. After quality control, there were 122 subjects and 46,355 SNPs in the analyzed dataset. Nucleotide diversity π values (Supplemental Table S1) were calculated for each breed using the Bio::PopGen::Statistics package in BioPerl (v1.6.1)¹⁰⁸. Among the 7 breeds, LOP showed the highest diversity ($\pi = 7.84 \times 10^{-6}$), and TIBN had the lowest diversity ($\pi = 7.32 \times 10^{-6}$).

Population analysis. Before analysis, all the SNPs were pruned using the indep-pairwise option, with a non-overlapped window size of 25 SNPs, a step of five SNPs, and a pairwise r^2 threshold of 0.05, resulting in 20,632 independent SNP markers. MDS was performed using PLINK 1.07¹⁰⁹. Pairwise identity-by-state distances were calculated among all the individuals using the 20,632 independent SNP markers, and MDS components were obtained using the mds-plot option, based on the identity-by-state matrix. Population structure was evaluated using STRUCTURE 2.3.4²⁷. All animals were analyzed in triplicate for $K = 2-4$. All analyses were performed with a burn-in length of 20,000, followed by 30,000 MCMC (Markov Chain Monte Carlo) replications for each K -value. The solutions for each K were visualized using DISTRUCT 1.1¹¹⁰.

Priority candidates for adaptation to high-altitude hypoxia. Except for the strongest, clearest selective signals, it is difficult to confidently distinguish true signals from false positives using population genetic data alone¹¹¹. Thus, we generated a set of priority functional candidate loci; 247 genes were obtained from Simonson *et al.*¹², and we searched for 128 genes using his approach, from GO categories, which were associated with hypoxia using the *Bos taurus* background (<http://amigo.geneontology.org/amigo>). The resulting 317 functional candidate loci, including those from *Homo sapiens* and *B. taurus*, are listed in Supplemental Table S6.

Statistical analysis. Before statistical analysis, TIBQ, TIBS and TIBN were clustered into G1, whereas TON, HUS, LOP and TLH were clustered into G2. F_{ST} values per-SNP were calculated using Genepop 4.2.2³¹. The formulae proposed by Weir and Cockerham²⁹ were used to analyze a single locus. The F_{ST} value can theoretically range from 0 to 1. Next, the outlier tests were implemented in LOSITAN, which was run using 50,000 simulations, a 'neutral' mean F_{ST} , confidence intervals of 95% and a false discovery rate of 0.1, using the infinite alleles model³⁰.

To determine if there was selection in G1, we computed the XP-EHH values using haplotype information in the xpehh program from <http://hgdp.uchicago.edu/Software/>. The XP-EHH derives from the idea of extended haplotype homozygosity (EHH), which is defined as the probability that two randomly chosen extended haplotypes carrying a given core haplotype are homozygous¹¹². XP-EHH is used to test whether the site is homozygous in one population but polymorphic in another population through comparison of EHH scores for one core SNP between two populations¹¹². A negative XP-EHH score indicates that selection occurred in the reference population, whereas a positive XP-EHH score indicates that selection occurred in the observed population.

Haplotypes were estimated with fastphase 1.4¹¹³. We used population label information to estimate the phased haplotype background and the following options for each chromosome: -Ku40 -Ki10 -Ki10. Candidate region

annotations were obtained from Ovis 3.1 of the sheep genome from NCBI (ftp://ftp.ncbi.nih.gov/genomes/Ovis_aries/protein/). To perform functional enrichment of the candidate genes, which was required by the ClueGO plugin of Cytoscape 3.2.1¹⁴ using Symbol ID as input parameters, the background organism selected *B. taurus*. P values less than 0.05 after Bonferroni correction for multiple testing were considered statistically significant.

Physiological measurement and association analysis. To verify the accuracy of the experiments, we also sampled Tibetan and Mongolian sheep from different altitudes. Jugular venous blood samples from 30 one-year-old TIBQ sheep (16 males and 14 females) that lived at least 3000 m high in Gansu Province and 31 one-year-old LTH sheep (16 males and 15 females) that lived no higher than 100 m in altitude in Beijing were collected to measure the six haematological parameters: RBC, HGB, HCT, MCV, MCH, and MCHC using a BC-2800Vet Auto Hematology Analyzer (Mindray Co., Ltd, Shenzhen, China). For the association analysis, 10 primer pairs were designed (Supplemental Table S10), and 19 *EPAS1* exons were amplified. Twelve mutations were detected and genotyped by traditional Sanger sequencing. We next used the general linear model with sex as a covariate in SPSS 20.0 to determine the association between haematological parameters and each genotype in the Tibetan breeds. The least square difference test was used for post-hoc analyses.

References

1. Thompson, L. G. *et al.* A high-resolution millennial record of the south asian monsoon from himalayan ice cores. *Science* **289**, 1916–1920 (2000).
2. Peacock, A. J. Oxygen at high altitude. *Brit Med J (Bmj)* **317**, 1063–1066 (1998).
3. Blumthaler, M., Ambach, W. & Ellinger, R. Increase in solar UV radiation with altitude. *J Photoch Photobio B* **39**, 130–134 (1997).
4. Ge, R. L. *et al.* Draft genome sequence of the Tibetan antelope. *Nat Commun* **4**, 1858, doi: 10.1038/ncomms2860 (2013).
5. Gou, X. *et al.* Whole-genome sequencing of six dog breeds from continuous altitudes reveals adaptation to high-altitude hypoxia. *Genome Res.* **24**, 1308–1315, doi: 10.1101/gr.171876.113 (2014).
6. Li, Y. *et al.* Population variation revealed high-altitude adaptation of tibetan mastiffs. *Mol Biol. Evol.* **31**, 1200–1205, doi: 10.1093/molbev/msu070 (2014).
7. Linxin, D. *Animal Genetic Resources In China* (China Agriculture Press, 2011).
8. Cai, D.-W., Han, L., Zhang, X.-L., Zhou, H. & Zhu, H. DNA analysis of archaeological sheep remains from China. *J Archaeol Sci.* **34**, 1347–1355 (2007).
9. Li, M. *et al.* Genomic analyses identify distinct patterns of selection in domesticated pigs and Tibetan wild boars. *Nat Genet* **45**, 1431–1438, doi: 10.1038/ng.2811 (2013).
10. Qiu, Q. *et al.* The yak genome and adaptation to life at high altitude. *Nat Genet* **44**, 946–949, doi: 10.1038/ng.2343 (2012).
11. Qu, Y. *et al.* Ground tit genome reveals avian adaptation to living at high altitudes in the Tibetan plateau. *Nat Commun* **4**, 2071, doi: 10.1038/ncomms3071 (2013).
12. Simonson, T. S. *et al.* Genetic evidence for high-altitude adaptation in Tibet. *Science* **329**, 72–75, doi: 10.1126/science.1189406 (2010).
13. Yi, X. *et al.* Sequencing of 50 human exomes reveals adaptation to high altitude. *Science* **329**, 75–78, doi: 10.1126/science.1190371 (2010).
14. Storz, J. F. Hemoglobin function and physiological adaptation to hypoxia in high-altitude mammals. *J Mammal* **88**, 24–31 (2007).
15. Weber, R. E. High-altitude adaptations in vertebrate hemoglobins. *Resp Physiol Neurobi* **158**, 132–142, doi: 10.1016/j.resp.2007.05.001 (2007).
16. Weber, R. E. & Fago, A. Functional adaptation and its molecular basis in vertebrate hemoglobins, neuroglobins and cytoglobins. *Resp Physiol Neurobi* **144**, 141–159, doi: 10.1016/j.resp.2004.04.018 (2004).
17. Beall, C. M. Two routes to functional adaptation: Tibetan and Andean high-altitude natives. *P Natl Acad Sci. USA* **104**, Suppl 1, 8655–8660, doi: 10.1073/pnas.0701985104 (2007).
18. Beall, C. M. *et al.* An Ethiopian pattern of human adaptation to high-altitude hypoxia. *P Natl Acad Sci. USA* **99**, 17215–17218, doi: 10.1073/pnas.252649199 (2002).
19. Storz, J. F., Scott, G. R. & Cheviron, Z. A. Phenotypic plasticity and genetic adaptation to high-altitude hypoxia in vertebrates. *J Exp Biol.* **213**, 4125–4136 (2010).
20. Storz, J. F. & Moriyama, H. Mechanisms of hemoglobin adaptation to high altitude hypoxia. *High Alt Med Biol.* **9**, 148–157, doi: 10.1089/ham.2007.1079 (2008).
21. Piccinini, M., Kleinschmidt, T., Jurgens, K. D. & Braunitzer, G. Primary structure and oxygen-binding properties of the hemoglobin from guanaco (*Lama guanacoe*, Tylopoda). *Biol. Chem Hoppe Seyler* **371**, 641–648 (1990).
22. Jessen, T. H., Weber, R. E., Fermi, G., Tame, J. & Braunitzer, G. Adaptation of bird hemoglobins to high altitudes: demonstration of molecular mechanism by protein engineering. *P Natl Acad Sci. USA* **88**, 6519–6522 (1991).
23. Liang, Y., Hua, Z., Liang, X., Xu, Q. & Lu, G. The crystal structure of bar-headed goose hemoglobin in deoxy form: the allosteric mechanism of a hemoglobin species with high oxygen affinity. *J Mol Biol.* **313**, 123–137, doi: 10.1006/jmbi.2001.5028 (2001).
24. Zhang, J. *et al.* The crystal structure of a high oxygen affinity species of haemoglobin (bar-headed goose haemoglobin in the oxy form). *J Mol Biol.* **255**, 484–493, doi: 10.1006/jmbi.1996.0040 (1996).
25. Storz, J. F. *et al.* Evolutionary and functional insights into the mechanism underlying high-altitude adaptation of deer mouse hemoglobin. *P Natl Acad Sci. USA* **106**, 14450–14455 (2009).
26. Storz, J. F. *et al.* The molecular basis of high-altitude adaptation in deer mice. *Plos Genet* **3**, e45, doi: 10.1371/journal.pgen.0030045 (2007).
27. Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using multilocus genotype data. *Genetics* **155**, 945–959 (2000).
28. Bruno, W. J., Succi, N. D. & Halpern, A. L. Weighted neighbor joining: a likelihood-based approach to distance-based phylogeny reconstruction. *Mol Biol. Evol.* **17**, 189–197 (2000).
29. Weir, B. S. & Cockerham, C. C. Estimating F-statistics for the analysis of population structure. *Evolution* **38**, 1358–1370, doi: 10.2307/2408641 (1984).
30. Antao, T., Lopes, A., Lopes, R. J., Beja-Pereira, A. & Luikart, G. LOSITAN: a workbench to detect molecular adaptation based on a fast-outlier method. *Bmc Bioinformatics* **9**, 323, doi: 10.1186/1471-2105-9-323 (2008).
31. Roussel, F. genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Mol Ecol Resour* **8**, 103–106, doi: 10.1111/j.1471-8286.2007.01931.x (2008).
32. Sabeti, P. C. *et al.* Positive natural selection in the human lineage. *Science* **312**, 1614–1620, doi: 10.1126/science.1124309 (2006).
33. Pruitt, K. D., Tatusova, T. & Maglott, D. R. NCBI Reference Sequence (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res.* **33**, D501–504, doi: 10.1093/nar/gki025 (2005).
34. Minet, E. *et al.* ERK activation upon hypoxia: involvement in HIF-1 activation. *Febs Lett* **468**, 53–58 (2000).
35. Wilkie, A. O. Cancer drugs to treat birth defects. *Nat Genet* **39**, 1057–1059, doi: 10.1038/ng0907-1057 (2007).

36. Coughlin, J. J., Stang, S. L., Dower, N. A. & Stone, J. C. RasGRP1 and RasGRP3 regulate B cell proliferation by facilitating B cell receptor-Ras signaling. *J Immunol* **175**, 7179–7184 (2005).
37. Yamashita, S. *et al.* CalDAG-GEFIII activation of Ras, R-ras, and Rap1. *J Biol. Chem.* **275**, 25488–25493, doi: 10.1074/jbc.M003414200 (2000).
38. McCormick, F. Ras signaling and NF1. *Curr Opin Genet Dev* **5**, 51–55 (1995).
39. Bollag, G. *et al.* Loss of NF1 results in activation of the Ras signaling pathway and leads to aberrant growth in haematopoietic cells. *Nat Genet* **12**, 144–148, doi: 10.1038/ng0296-144 (1996).
40. Sharma, A. *et al.* A new role for the muscle repair protein dysferlin in endothelial cell adhesion and angiogenesis. *Arterioscler Thromb Vas* **30**, 2196–2204, doi: 10.1161/ATVBAHA.110.208108 (2010).
41. Luo, Z. *et al.* MicroRNA-200C and -150 play an important role in endothelial cell differentiation and vasculogenesis by targeting transcription repressor ZEB1. *Stem Cells* **31**, 1749–1762, doi: 10.1002/stem.1448 (2013).
42. Martin, A. & Cano, A. Tumorigenesis: Twist1 links EMT to self-renewal. *Nat Cell Biol.* **12**, 924–925, doi: 10.1038/ncb1010-924 (2010).
43. Semenza, G. L., Nejfelt, M. K., Chi, S. M. & Antonarakis, S. E. Hypoxia-inducible nuclear factors bind to an enhancer element located 3' to the human erythropoietin gene. *P Natl Acad Sci. USA* **88**, 5680–5684 (1991).
44. Kapitsinou, P. P. *et al.* Hepatic HIF-2 regulates erythropoietic responses to hypoxia in renal anemia. *Blood* **116**, 3039–3048, doi: 10.1182/blood-2010-02-270322 (2010).
45. Semenza, G. L. & Wang, G. L. A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. *Mol Cell Biol.* **12**, 5447–5454 (1992).
46. Wang, L. *et al.* Erythropoietin up-regulates SOCS2 in neuronal progenitor cells derived from SVZ of adult rat. *Neuroreport* **15**, 1225–1229 (2004).
47. Linossi, E. M., Babon, J. J., Hilton, D. J. & Nicholson, S. E. Suppression of cytokine signaling: the SOCS perspective. *Cytokine Growth F R* **24**, 241–248, doi: 10.1016/j.cytogfr.2013.03.005 (2013).
48. Fujita, H. Molecular mechanism of heme biosynthesis. *Tohoku J Exp. Med.* **183**, 83–99 (1997).
49. Tian, Q. *et al.* Lon peptidase 1 (LONP1)-dependent breakdown of mitochondrial 5-aminolevulinic acid synthase protein by heme in human liver cells. *J Biol. Chem.* **286**, 26424–26430, doi: 10.1074/jbc.M110.215772 (2011).
50. Goldberg, M. A., Dunning, S. P. & Bunn, H. F. Regulation of the erythropoietin gene: evidence that the oxygen sensor is a heme protein. *Science* **242**, 1412–1415 (1988).
51. Fukuda, R. *et al.* HIF-1 regulates cytochrome oxidase subunits to optimize efficiency of respiration in hypoxic cells. *Cell* **129**, 111–122, doi: 10.1016/j.cell.2007.01.047 (2007).
52. Cai, Q. *et al.* Genome sequence of ground tit *Pseudopodoces humilis* and its adaptation to high altitude. *Genome Biol.* **14**, R29, doi: 10.1186/gb-2013-14-3-r29 (2013).
53. Das, S. S. *et al.* Regulation of dipeptidyl peptidase 4 production in adipocytes by glucose. *Diabetes Metab Syndr Obes* **7**, 185–194, doi: 10.2147/DMSO.S62610 (2014).
54. Holman, R. R., Paul, S. K., Bethel, M. A., Matthews, D. R. & Neil, H. A. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med.* **359**, 1577–1589, doi: 10.1056/NEJMoa0806470 (2008).
55. Zheng, T. P. *et al.* Increased plasma DPP4 activities predict new-onset atherosclerosis in association with its proinflammatory effects in Chinese over a four year period: A prospective study. *Atherosclerosis* **235**, 619–624, doi: 10.1016/j.atherosclerosis.2014.05.956 (2014).
56. Ren, D., Collingwood, T. N., Rebar, E. J., Wolffe, A. P. & Camp, H. S. PPARgamma knockdown by engineered transcription factors: exogenous PPARgamma2 but not PPARgamma1 reactivates adipogenesis. *Gene Dev* **16**, 27–32, doi: 10.1101/gad.953802 (2002).
57. Yilmaz-Aydogan, H. *et al.* Different effects of PPARA, PPARG and ApoE SNPs on serum lipids in patients with coronary heart disease based on the presence of diabetes. *Gene* **523**, 20–26, doi: 10.1016/j.gene.2013.03.136 (2013).
58. Hoppeler, H., Vogt, M., Weibel, E. R. & Fluck, M. Response of skeletal muscle mitochondria to hypoxia. *Exp. Physiol.* **88**, 109–119 (2003).
59. LaRochelle, W. J. *et al.* PDGF-D, a new protease-activated growth factor. *Nat Cell Biol.* **3**, 517–521, doi: 10.1038/35074593 (2001).
60. Scortegagna, M. *et al.* Multiple organ pathology, metabolic abnormalities and impaired homeostasis of reactive oxygen species in *Epas1*^{-/-} mice. *Nat Genet* **35**, 331–340, doi: 10.1038/ng1266 (2003).
61. Cousse, E. *et al.* G37R SOD1 mutant alters mitochondrial complex I activity, Ca(2+) uptake and ATP production. *Cell Calcium* **49**, 217–225, doi: 10.1016/j.ceca.2011.02.004 (2011).
62. van Patot, M. C. & Gassmann, M. Hypoxia: adapting to high altitude by mutating EPAS-1, the gene encoding HIF-2alpha. *High Alt Med. Biol.* **12**, 157–167, doi: 10.1089/ham.2010.1099 (2011).
63. Wang, Y. X. & Zheng, Y. M. ROS-dependent signaling mechanisms for hypoxic Ca(2+) responses in pulmonary artery myocytes. *Antioxid Redox Sign* **12**, 611–623, doi: 10.1089/ars.2009.2877 (2010).
64. Tang, W. *et al.* Identification of a novel 5-base pair deletion in calcineurin B (PPP3R1) promoter region and its association with left ventricular hypertrophy. *Am Heart J* **150**, 845–851, doi: 10.1016/j.ahj.2004.12.004 (2005).
65. Hari Kishore, A., Li, X. H. & Word, R. A. Hypoxia and PGE(2) regulate MiTF-CX during cervical ripening. *Mol Endocrinol* **26**, 2031–2045, doi: 10.1210/me.2012-1100 (2012).
66. Kijas, J. W. *et al.* Genome-wide analysis of the world's sheep breeds reveals high levels of historic mixture and strong recent selection. *Plos Biol.* **10**, e1001258, doi: 10.1371/journal.pbio.1001258 (2012).
67. Tomlinson, J. W. & Stewart, P. M. Cortisol metabolism and the role of 11beta-hydroxysteroid dehydrogenase. *Best Pract Res. Cl En* **15**, 61–78, doi: 10.1053/beem.2000.0119 (2001).
68. Heiniger, C. D. *et al.* Hypoxia causes down-regulation of 11 beta-hydroxysteroid dehydrogenase type 2 by induction of Egr-1. *Faseb J* **17**, 917–919, doi: 10.1096/fj.02-0582fje (2003).
69. Atanasov, A. G., Nashev, L. G., Schweizer, R. A., Frick, C. & Odermatt, A. Hexose-6-phosphate dehydrogenase determines the reaction direction of 11beta-hydroxysteroid dehydrogenase type 1 as an oxoreductase. *FEBS Lett* **571**, 129–133, doi: 10.1016/j.febslet.2004.06.065 (2004).
70. Kaelin, W. G. The von Hippel-Lindau tumor suppressor protein: roles in cancer and oxygen sensing. *Cold Spring Harb Sym* **70**, 159–166, doi: 10.1101/sqb.2005.70.001 (2005).
71. Kaelin, W. G. Jr. & Ratcliffe, P. J. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell* **30**, 393–402, doi: 10.1016/j.molcel.2008.04.009 (2008).
72. Bonewald, L. F. Regulation and regulatory activities of transforming growth factor beta. *Crit Rev Eukar Gene* **9**, 33–44 (1999).
73. Flanders, K. C. & Wakefield, L. M. Transforming growth factor-(beta)s and mammary gland involution; functional roles and implications for cancer progression. *J Mammary Gland Biol.* **14**, 131–144, doi: 10.1007/s10911-009-9122-z (2009).
74. Letterio, J. J. & Roberts, A. B. Regulation of immune responses by TGF-beta. *Annu Rev Immunol* **16**, 137–161, doi: 10.1146/annurev.immunol.16.1.137 (1998).
75. Beutler, E., Gelbart, T. & Lee, P. Haptoglobin polymorphism and iron homeostasis. *Clin Chem* **48**, 2232–2235 (2002).
76. Boyko, A. R. *et al.* A simple genetic architecture underlies morphological variation in dogs. *Plos Biol.* **8**, e1000451, doi: 10.1371/journal.pbio.1000451 (2010).
77. Vaysse, A. *et al.* Identification of genomic regions associated with phenotypic variation between dog breeds using selection mapping. *Plos Genet* **7**, e1002316, doi: 10.1371/journal.pgen.1002316 (2011).

78. Wei, C. *et al.* Genome-wide analysis reveals population structure and selection in Chinese indigenous sheep breeds. *Bmc Genomics* **16**, 194, doi: 10.1186/s12864-015-1384-9 (2015).
79. Windelinckx, A. *et al.* Identification and prioritization of NUA1 and PPP1CC as positional candidate loci for skeletal muscle strength phenotypes. *Physiol Genomics* **43**, 981–992, doi: 10.1152/physiolgenomics.00200.2010 (2011).
80. Johnston, S. E. *et al.* Genome-wide association mapping identifies the genetic basis of discrete and quantitative variation in sexual weaponry in a wild sheep population. *Mol Ecol* **20**, 2555–2566, doi: 10.1111/j.1365-294X.2011.05076.x (2011).
81. Fan, B. *et al.* Genome-wide association study identifies loci for body composition and structural soundness traits in pigs. *Plos One* **6**, e14726 (2011).
82. Saeki, M. *et al.* Monad, a WD40 repeat protein, promotes apoptosis induced by TNF-alpha. *Biochemical And Biophysical Research Communications* **342**, 568–572, doi: 10.1016/j.bbrc.2006.02.009 (2006).
83. Wang, X. *et al.* Pno1 Tissue-Specific Expression and Its Functions Related to the Immune Responses and Proteasome Activities. *Plos One* **7**, e46093 (2012).
84. Shadrina, M. I. *et al.* Effective quantitative real-time polymerase chain reaction analysis of the parkin gene (PARK2) exon 1–12 dosage. *Bmc Med Genet* **8**, 1 (2007).
85. Lan, Y. *et al.* Odd-skipped related 2 (Osr2) encodes a key intrinsic regulator of secondary palate growth and morphogenesis. *Development* **131**, 3207–3216 (2004).
86. Petropoulos, E. & Timiras, P. Biological effects of high altitude as related to increased solar radiation, temperature fluctuations and reduced partial pressure of oxygen. *Progress in biometeorology. Division A: Progress In Human Biometeorology* **1**, 295 (1974).
87. Huff, C. D., Harpending, H. C. & Rogers, A. R. Detecting positive selection from genome scans of linkage disequilibrium. *Bmc Genomics* **11**, 8, doi: 10.1186/1471-2164-11-8 (2010).
88. Pickrell, J. K. *et al.* Signals of recent positive selection in a worldwide sample of human populations. *Genome Res.* **19**, 826–837, doi: 10.1101/gr.087577.108 (2009).
89. Garraway, L. A. *et al.* Integrative genomic analyses identify MITF as a lineage survival oncogene amplified in malignant melanoma. *Nature* **436**, 117–122, doi: 10.1038/nature03664 (2005).
90. Takeda, K. *et al.* Lipocalin-type prostaglandin D synthase as a melanocyte marker regulated by MITF. *Biochem Bioph Res. Co* **339**, 1098–1106, doi: 10.1016/j.bbrc.2005.11.125 (2006).
91. Peyssonnaud, C., Nizet, V. & Johnson, R. S. Role of the hypoxia inducible factors HIF in iron metabolism. *Cell Cycle* **7**, 28–32 (2008).
92. Chen, L., Endler, A. & Shibasaki, F. Hypoxia and angiogenesis: regulation of hypoxia-inducible factors via novel binding factors. *Exp Mol Med* **41**, 849–857, doi: 10.3858/emmm.2009.41.12.103 (2009).
93. Tanaka, T. *et al.* Cobalt promotes angiogenesis via hypoxia-inducible factor and protects tubulointerstitium in the remnant kidney model. *Lab Invest* **85**, 1292–1307, doi: 10.1038/labinvest.3700328 (2005).
94. Biswas, S. *et al.* Effects of HIF-1alpha and HIF2alpha on Growth and Metabolism of Clear-Cell Renal Cell Carcinoma 786-0 Xenografts. *J Oncol* **2010**, 757908, doi: 10.1155/2010/757908 (2010).
95. Rankin, E. B. *et al.* Hypoxia-inducible factor 2 regulates hepatic lipid metabolism. *Mol Cell Biol.* **29**, 4527–4538, doi: 10.1128/MCB.00200-09 (2009).
96. Shohet, R. V. & Garcia, J. A. Keeping the engine primed: HIF factors as key regulators of cardiac metabolism and angiogenesis during ischemia. *J Mol Med (Berl)* **85**, 1309–1315, doi: 10.1007/s00109-007-0279-x (2007).
97. Tormos, K. V. & Chandel, N. S. Inter-connection between mitochondria and HIFs. *J Cell Mol Med* **14**, 795–804, doi: 10.1111/j.1582-4934.2010.01031.x (2010).
98. Germonpré, M. *et al.* Fossil dogs and wolves from Palaeolithic sites in Belgium, the Ukraine and Russia: osteometry, ancient DNA and stable isotopes. *J Archaeol Sci.* **36**, 473–490 (2009).
99. Zhao, M. *et al.* Mitochondrial genome evidence reveals successful Late Paleolithic settlement on the Tibetan Plateau. *P Natl Acad Sci. USA* **106**, 21230–21235, doi: 10.1073/pnas.0907844106 (2009).
100. Lawson Handley, L. J. *et al.* Genetic structure of European sheep breeds. *Heredity* **99**, 620–631, doi: 10.1038/sj.hdy.6801039 (2007).
101. Zeder, M. A. Animal domestication in the Zagros: a review of past and current research. *Paleorient* **25**, 11–25 (1999).
102. Erzurum, S. C. *et al.* Higher blood flow and circulating NO products offset high-altitude hypoxia among Tibetans. *P Natl Acad Sci. USA* **104**, 17593–17598, doi: 10.1073/pnas.0707462104 (2007).
103. Bigham, A. W. *et al.* Identifying positive selection candidate loci for high-altitude adaptation in Andean populations. *Hum Genomics* **4**, 79–90 (2009).
104. Beall, C. M. *et al.* Natural selection on EPAS1 (HIF2alpha) associated with low hemoglobin concentration in Tibetan highlanders. *P Natl Acad Sci. USA* **107**, 11459–11464, doi: 10.1073/pnas.1002443107 (2010).
105. Peng, Y. *et al.* Genetic variations in Tibetan populations and high-altitude adaptation at the Himalayas. *Mol Biol. Evol.* **28**, 1075–1081, doi: 10.1093/molbev/msq290 (2011).
106. Xu, S. *et al.* A genome-wide search for signals of high-altitude adaptation in Tibetans. *Mol Biol. Evol.* **28**, 1003–1011, doi: 10.1093/molbev/msq277 (2011).
107. Al-Sweedan, S. A. & Alhaj, M. The effect of low altitude on blood count parameters. *Hematol Oncol Stem Cell Ther* **5**, 158–161, doi: 10.5144/1658-3876.2012.158 (2012).
108. Stajich, J. E. *et al.* The Bioperl toolkit: Perl modules for the life sciences. *Genome Res.* **12**, 1611–1618 (2002).
109. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* **81**, 559–575, doi: 10.1086/519795 (2007).
110. Rosenberg, N. A. DISTRUCT: a program for the graphical display of population structure. *Mol Ecol Notes* **4**, 137–138 (2004).
111. Teshima, K. M., Coop, G. & Przeworski, M. How reliable are empirical genomic scans for selective sweeps? *Genome Res.* **16**, 702–712, doi: 10.1101/gr.5105206 (2006).
112. Sabeti, P. C. *et al.* Genome-wide detection and characterization of positive selection in human populations. *Nature* **449**, 913–918 (2007).
113. Scheet, P. & Stephens, M. A fast and flexible statistical model for large-scale population genotype data: applications to inferring missing genotypes and haplotypic phase. *Am J Hum Genet* **78**, 629–644, doi: 10.1086/502802 (2006).
114. Bindea, G. *et al.* ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* **25**, 1091–1093, doi: 10.1093/bioinformatics/btp101 (2009).

Acknowledgements

The authors gratefully acknowledge the International Sheep Genomics Consortium for permitting using Ovine HapMap genotype data. We are also grateful to Prof. Dongmin Liu (CAAS) for reviewing the manuscript. This research was supported by the National Key Technology R&D Program of China (2011BAD28B05-2) and Production and storage of livestock and poultry genetic material and measurement of genetic distance (Agricultural Finance Development (2013), No. 36).

Author Contributions

C.W., H.W. and G.L. drafted the manuscript. Y.M., G.L., J.L., Z.L., R.L. and S.Z. performed the experiments and participated in phenotype data collection. H.W., F.Z., G.W. and J.K. carried out the population genetic studies. L.Z., M.W. and J.C. contributed reagents and materials. H.W. provided the map. C.W., L.D. and C.L. conceived the study, participated in its design and coordination and helped draft the manuscript. All of the authors read and approved the final manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Wei, C. *et al.* Genome-wide analysis reveals adaptation to high altitudes in Tibetan sheep. *Sci. Rep.* **6**, 26770; doi: 10.1038/srep26770 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>