



Published in final edited form as:

*Adv Biol Regul.* 2016 May ; 61: 17–24. doi:10.1016/j.jbior.2015.11.011.

## Membrane and Inhibitor Interactions of Intracellular Phospholipases A<sub>2</sub>

Varnavas D. Mouchlis and Edward A. Dennis\*

Department of Pharmacology and Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, California 92093, USA

### Abstract

Studying phospholipases A<sub>2</sub> (PLA<sub>2</sub>s) is a challenging task since they act on membrane-like aggregated substrates and not on monomeric phospholipids. Multidisciplinary approaches that include hydrogen/deuterium exchange mass spectrometry (DXMS) and computational techniques have been employed with great success in order to address important questions about the mode of interactions of PLA<sub>2</sub> enzymes with membranes, phospholipid substrates and inhibitors. Understanding the interactions of PLA<sub>2</sub>s is crucial since these enzymes are the upstream regulators of the eicosanoid pathway liberating free arachidonic acid (AA) and other polyunsaturated fatty acids (PUFA). The liberation of AA by PLA<sub>2</sub> enzymes sets off a cascade of molecular events that involves downstream regulators such as cyclooxygenase (COX) and lipoxygenase (LOX) metabolites leading to inflammation. Aspirin and other nonsteroidal anti-inflammatory drugs (NSAIDs) work by inhibiting COX, while Zileuton inhibits LOX and both rely on PLA<sub>2</sub> enzymes to provide them with AA. That means PLA<sub>2</sub> enzymes can potentially also be targeted to diminish inflammation at an earlier point in the process. In this review we describe extensive efforts reported in the past to define the interactions of PLA<sub>2</sub> enzymes with membranes, substrate phospholipids and inhibitors using DXMS, molecular docking, and molecular dynamics (MD) simulations.

### Keywords

Phospholipase A<sub>2</sub>; PAPC; catalytic cycle; allosteric regulation; eicosanoid pathway; molecular dynamics simulations; DXMS

### Introduction

Through their catalytic action, phospholipases A<sub>2</sub> (PLA<sub>2</sub>) de-esterify and release free fatty acids from the *sn*-2 position of membrane phospholipids including arachidonic acid (AA) and related polyunsaturated fatty acids (PUFA) (Dennis et al. , 2011). Eicosanoids, which are associated with inflammation, are produced by the downstream enzymes of the

---

To whom correspondence may be addressed. ; Email: edennis@ucsd.edu

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

eicosanoid pathway such as cyclooxygenases (COX), lipoxygenases (LOX), and cytochrome P450s (CYP) that rely on PLA<sub>2</sub>s to provide them with AA and related PUFA (Buczynski et al. , 2009, Dennis and Norris, 2015, Funk, 2001). Existing nonsteroidal anti-inflammatory drugs (NSAID) target the downstream enzymes of the prostaglandin pathway. As the upstream regulators of the eicosanoid pathway, PLA<sub>2</sub> enzymes can be targeted to diminish inflammation at an earlier stage in the process. Besides their implication in the inflammatory process, PLA<sub>2</sub> enzymes are also implicated in various diseases including atherosclerosis, (Karabina et al. , 2010, Rosenson and Hurt-Camejo, 2012) diabetes,(Ayilavarapu et al. , 2010, Bone et al. , 2015) arthritis(Hegen et al. , 2003, Masuda et al. , 2005) and cancer(Nakanishi and Rosenberg, 2006, Park et al. , 2012, Scott et al. , 2010) and inhibition of the enzymatic activity of specific PLA<sub>2</sub>s should be beneficial for the treatment of these diseases (Bone et al. , 2015, Karabina et al. , 2010). Therefore, there has been great interest in understanding the interactions of PLA<sub>2</sub> enzymes with membranes, substrates and inhibitors in order to develop effective new therapeutic agents.

The PLA<sub>2</sub> superfamily consists of 16 groups and many subgroups and each has been classified into one of the six main types based on their characteristics of sequence and structure. The six main types are secreted (sPLA<sub>2</sub>), cytosolic (cPLA<sub>2</sub>), calcium-independent (iPLA<sub>2</sub>), platelet-activating factor acetylhydrolase (PAF-AH) also known as lipoprotein-associated (LpPLA<sub>2</sub>), lysosomal (LPLA<sub>2</sub>), and adipose (AdPLA) (Dennis et al. , 2011). Even though the PLA<sub>2</sub> superfamily constitutes a diverse set of enzymes, they share the same catalytic activity because of convergent evolution. Nevertheless, divergent PLA<sub>2</sub> enzymes have a distinct 3D-structure, binding pocket and catalytic residues as well as a unique, in terms of topology and function, interfacial surface through which they associate with the membrane to extract and bind their phospholipid substrate. Since PLA<sub>2</sub> enzymes have to associate with biological membranes to complete their catalytic cycle, they evolved to act on membrane-like phospholipids such as micelles, membranes and liposomes rather than on monomeric phospholipids. Thus, PLA<sub>2</sub>s follow a special kinetic model which we named “surface dilution kinetics” that was successfully employed to interpret the kinetic characteristics of these enzymes (Carman et al. , 1995).

This short review focuses on the application of hydrogen/deuterium exchange mass spectrometry (DXMS)(Cao et al. , 2013) and computational techniques(Durrant and McCammon, 2011, Sliwoski et al. , 2014) to understanding the mechanism of action and inhibition of Group IVA cytosolic PLA<sub>2</sub> (GIVA cPLA<sub>2</sub>)(Clark et al. , 1991, Sharp et al. , 1991) and Group VIA calcium-independent PLA<sub>2</sub> (GVIA iPLA<sub>2</sub>) (Larsson et al. , 1998). Molecular dynamics (MD) simulations, molecular docking, and other computer-aided design techniques guided by DXMS experimental data have been utilized with great success in understanding the association and interactions of these two intracellular enzymes with membranes in a biphasic water-lipid system as well as with specific phospholipid substrates and inhibitors (Mouchlis et al. , 2015).

### **Structure and function of GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub>**

The 85 kDa human GIVA cPLA<sub>2</sub> (cPLA<sub>2</sub> $\alpha$ , calcium-dependent) contains 749 amino acids and it was cloned and sequenced in 1991 from U937 cells (Clark et al. , 1991, Kramer et al. ,

1991). The crystal structure of the enzyme revealed an N-terminal C2 domain (residues: 13–137), a linker region (residues: 138–143) and a C-terminal  $\alpha/\beta$  hydrolase catalytic domain (residues: 144–749, Fig. 1A) (Dessen et al. , 1999). GIVA cPLA<sub>2</sub> has a deep channel-like binding site that is located in the  $\alpha/\beta$  hydrolase catalytic domain and contains the catalytic dyad of Ser/Asp (Mouchlis et al. , 2015). A peptide region consisting of residues 415–432 is on the top of the binding pocket and is defined as the lid (Dessen et al. , 1999). The lid region is amphipathic showing increased on-exchanged rates in DXMS experimental data upon binding with phospholipid vesicles and inhibitors (Burke et al. , 2009, Burke et al. , 2008). Our recent studies using MD simulations guided by DXMS data showed high flexibility of the lid region during the extraction of a phospholipid molecule from the membrane into the binding pocket (Mouchlis et al. , 2015). GIVA cPLA<sub>2</sub> is regulated by intracellular calcium through a calcium binding site located at the C2 domain that activates the translocation of the enzyme to the phospholipid membrane (Fig. 1A). Using DXMS experiments, we showed that calcium causes conformational changes in the C2 domain to stabilize the C2 domain and facilitate association of the enzyme with the membrane as shown by decreased hydrogen/deuterium (H/D) exchange rates (Hsu et al. , 2008). GIVA cPLA<sub>2</sub> activity is enhanced by binding to phosphatidylinositol-4,5-bisphosphate (PIP<sub>2</sub>) and this activation is calcium-independent (Balsinde et al. , 2000, Leslie and Channon, 1990, Six and Dennis, 2003). Four lysine residues Lys488, Lys541, Lys543, and Lys544, have been shown to constitute a PIP<sub>2</sub> binding site and to be critical for PIP<sub>2</sub>-mediated activation of GVIA cPLA<sub>2</sub> (Das and Cho, 2002, Six and Dennis, 2003). Ceramide 1-phosphate (C1P) which is a phosphorylated sphingolipid has also been suggested to activate GIVA cPLA<sub>2</sub> (Chalfant and Spiegel, 2005, Pettus et al. , 2004). This enzyme exhibits specificity for phospholipids containing AA esterified at the *sn*-2 position (Clark et al. , 1991, Ghosh et al. , 2006).

The human GVIA iPLA<sub>2</sub> (iPLA<sub>2</sub> $\beta$ , PNPLA9, calcium-independent) has a molecular weight of 89.9 kDa and it has longest sequence among the multiple splice variants expressed by the human gene (Larsson Forsell et al. , 1999, Larsson et al. , 1998). The X-ray crystal structure of the enzyme has not been solved yet but homology modeling studies showed that region 121–474 has 51% homology with human ankyrinR (PDB ID: 1N11)(Michaely et al. , 2002) and region 475–806 has 34% homology with patatin (PDB ID: 1OXW) (Rydel et al. , 2003). According to the homology models the structure of GVIA iPLA<sub>2</sub> consists of the ankyrin repeats (residues: 121–383), a linker region (residues 384–474), and an  $\alpha/\beta$  hydrolase catalytic domain (residues: 475–806) (Dennis et al. , 2011, Mouchlis et al. , 2015). No homology template was found for region 1–120 (Mouchlis et al. , 2015). ATP was found to enhance the activity of GVIA iPLA<sub>2</sub>, but it is not a substrate or cofactor for this enzyme (Hazen and Gross, 1991, Lio and Dennis, 1998). GVIA iPLA<sub>2</sub> also contains a catalytic dyad of Ser/Asp and has low specificity for the free fatty acid esterified at the *sn*-2 position (Lio and Dennis, 1998). Besides its phospholipase A<sub>2</sub> activity, GVIA iPLA<sub>2</sub> also exhibits lysophospholipase, transacylase and acyl-CoA thioesterase activity (Jenkins et al. , 2006, Winstead et al. , 2000).

### GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> in eicosanoid biosynthesis

Eicosanoid biosynthesis requires free AA which is provided by PLA<sub>2</sub> enzymes under physiological or inflammatory cell activation. GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> have been implicated in cellular eicosanoid production. GIVA cPLA<sub>2</sub> is activated by Toll-like receptors (LTR), purinergic receptors and other receptors and it is translocated to the perinuclear and endoplasmic reticulum membranes during inflammatory responses (Dennis and Norris, 2015). After its translocation, it releases AA from membrane phospholipids containing esterified AA at the *sn*-2 position leading to the production of pro-inflammatory eicosanoids (Buczynski et al. , 2009, Buczynski et al. , 2007). Our group showed that in macrophages GIVA cPLA<sub>2</sub> may also hydrolyze phospholipids that have other PUFAs such as EPA, DPA and DHA esterified at the *sn*-2 position, which are precursors for anti-inflammatory resolvins (Keyes et al. , 2010, Norris and Dennis, 2012, Norris et al. , 2014). GVIA iPLA<sub>2</sub> constitutively releases a low level of free fatty acids and it is primarily involved in membrane homeostasis and remodeling (Balsinde et al. , 1997, Dennis et al. , 2011) The specificity of GVIA iPLA<sub>2</sub> for the free fatty acid at the *sn*-2 position is not great; thus, its activity may include hydrolysis of AA that is esterified at this position.

Downstream enzymes in the cyclooxygenase (COX) pathway generate prostaglandins (PG), thromboxanes (TX) and lipoxins (LX, Fig.2). In the lipoxygenase (LOX) pathway, the downstream enzymes generate leukotrienes (LTs), hydroxyeicosatetraenoic acids (HETE) and hepxilins (HX, Fig. 2). Finally, in the cytochrome P450 (CYP) epoxy hydrolase pathway, the downstream enzymes generate epoxides and dihydroxy polyunsaturated fatty acids (PUFA, Fig. 2) (Buczynski et al. , 2009, Dennis and Norris, 2015). Pharmaceutical agents that target the prostaglandin pathway have been widely used for decades. Aspirin is the best example among the numerous NSAID that have been commercially available. NSAID inhibit COX-1 and COX-2 enzymes and as a result inhibit prostaglandin biosynthesis (Fig. 2) (Vane, 2002). Zileuton is a 5-LOX inhibitor, which is also a major therapeutic target for the treatment of allergic and asthmatic conditions (Fig. 2) (Rossi et al. , 2010). Other pathways including CYP and sEH have also been considered as pharmaceutical targets for inflammation and inflammatory diseases. Since PLA<sub>2</sub> enzymes are the upstream regulators of the eicosanoid pathway, they should be potential therapeutic targets for inflammation and inflammation-related diseases such as cancer, arthritis and diabetes (Fig. 2).

### Catalytic cycle of GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub>

PLA<sub>2</sub> enzymes are often water-soluble, but act as membrane associated enzymes showing a significant increase in activity when their phospholipid substrate is in a membrane-like form such as micelles, vesicles or liposomes rather than in a monomeric form. Most of the PLA<sub>2</sub> enzymes have to associate with a lipid-water interface through their interfacial surface, and access their phospholipid substrate in order to complete their catalytic cycle which is the hydrolysis of the ester bond at the *sn*-2 position of the phospholipid molecule (Fig. 3). Thus, PLA<sub>2</sub> enzymes can best be described with a special kinetic model called “surface dilution kinetics” (Carman et al. , 1995). This model includes four steps: A) association of the enzyme with the membrane, B) extraction and binding of a phospholipid molecule in the catalytic site, C) hydrolysis of the phospholipid molecule, and D) diffusion of the products

in the membrane, followed by repetition of the cycle (Fig. 3). All of these steps exist in equilibrium. The membrane is a very important factor in the catalytic cycle of these enzymes, and recently we have introduced the idea that the membrane acts as an allosteric ligand shifting the conformation of PLA<sub>2</sub> enzymes from the inactive (in water) to the active (on the membrane surface) form (Mouchlis et al. , 2015). These four steps are described in detail below.

### A. Association of the enzyme with the membrane

GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> associate with membranes to extract and bind their phospholipid substrates even though they are in principal water-soluble enzymes. Therefore, membranes are an important factor for these enzymes. Elucidating their interactions with and conformational changes during the association with membrane is vital in understanding the mechanism of action of these two enzymes.

According to DXMS experimental data, five peptide regions of GIVA cPLA<sub>2</sub> (28–35, 36–39, 258–265, 268–279, and 466–470) exhibited decreased levels of deuteration in the presence of phospholipid vesicles (Burke et al. , 2008). Regions 28–35 and 36–39 are located on the C2 domain and their association with the phospholipid vesicles was found to be calcium-dependent because they exhibited a larger change in exchange in the presence of calcium (Burke et al. , 2008, Hsu et al. , 2008). Region 96–98 of the C2 domain was reported to play a significant role in the association of GIVA cPLA<sub>2</sub> with the membrane (Málková et al. , 2005, Perisic et al. , 1999). Unfortunately, no peptides covering region 96–98 were identified during the digestion of GIVA cPLA<sub>2</sub>, so no DXMS data are available for interactions of that region with phospholipid vesicles (Burke et al. , 2008). Peptide regions 258–265, 268–279, and 466–470 (blue and light blue region in Fig. 4A) are located on the catalytic domain and were used to generate a three-dimensional model of the catalytic domain of GIVA cPLA<sub>2</sub> on the surface of the membrane (Fig. 4A) (Mouchlis et al. , 2015). It is worth mentioning that the C2 domain penetrates the membrane approximately 15 Å, leading to the catalytic domain being near the surface of the membrane. Residues like Trp464 and Met468 penetrate the membrane surface interacting with the hydrophobic tails of the phospholipids, while residues such as Lys273, Lys274, and Arg467 interact with the head groups of the membrane phospholipids (Fig. 4A) (Mouchlis et al. , 2015).

The association of GVIA iPLA<sub>2</sub> with the membrane has also been studied by our group using coarse-grained and atomistic simulations guided by DXMS experimental data (Bucher et al. , 2013, Hsu et al. , 2009). The ankyrin repeats (Fig. 1B) showed no interactions with phospholipid vesicles (Hsu et al. , 2009). MD simulations of the catalytic domain showed that region 708–730 (red region in Fig. 4B) is forming an amphipathic helix after penetrating the membrane with its hydrophilic residues such as Arg710 and Lys719 (Fig. 4B) to interact with the phospholipid head groups, and its hydrophobic residues like Pro711, Pro714, Trp715, Leu717, Val721, and Phe722 to interact with the fatty acid tails of the phospholipids (Bucher et al. , 2013). Our previously published DXMS data and MD simulations indicated that region 708–730 is likely the anchor region from which the enzyme associates with the membrane in contrast to GIVA cPLA<sub>2</sub> that contains the C2 domain which assists membrane association of the enzyme (Bucher et al. , 2013, Burke et al. , 2008, Hsu et al. , 2009).

## B. Extraction and binding of a phospholipid molecule in the catalytic site

Steered molecular dynamics (SMD) simulations guided by DXMS results gave insight into the extraction pathway of a P APC phospholipid molecule from the membrane into the catalytic site of both GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> (Mouchlis et al. , 2015). The direction that P APC was forced into the catalytic site is indicated by the thick black arrow in Figures 4A and 4B. Reference Mouchlis et al. , 2015 includes animations in the supporting information that show the detailed extraction of P APC from the membrane into the catalytic site of both enzymes (Mouchlis et al. , 2015).

During the extraction of the P APC into the catalytic site of GIVA cPLA<sub>2</sub>, three regions that are located near the entrance of the site (261–268, 405–415, and 670–686) were found to play a significant role in the process. Residues like Asn268, Ser408, Gln411, Asn682, and Gln684 were found to participate in hydrogen-bonding with the head group of the P APC molecule while it is extracted into the catalytic site (see Movie S1 in reference Mouchlis et al. , 2015). Arg200 is stabilized by Ser680 and Glu418 when P APC is not in the catalytic site, while P APC enters the site approximately two-thirds of its length; Arg200 interacts with the phosphate group stabilizing its binding. The lid region (cyan region in Fig 4A and 4C) exhibits high flexibility whereas P APC enters the binding pocket (Mouchlis et al. , 2015). Once P APC completely enters the active site and during an MD simulation on the GIVA cPLA<sub>2</sub>-P APC complex, Arg200 and Asn555 interact with the phosphate group while Glu418 and Glu589 interact with the positively charged choline group of the phospholipid head group (Fig. 4C). The oxygen atom of the *sn*-2 carbonyl group is located near the oxyanion hole (Gly197/Gly198) participating in hydrogen-bonding, while its carbon atom is located near catalytic Ser228. Finally the fatty acid chains are placed in the hydrophobic region of the catalytic site with the four double bonds of the *sn*-2 arachidonyl chain participating in  $\pi$ - $\pi$  stacking with the aromatic residues Trp232, Phe295, Phe397, Phe681, and Phe683.

Simulations of the extraction of the P APC molecule into the catalytic site of GVIA iPLA<sub>2</sub> also gave useful information for its extraction pathway (see Movie S2 in reference Mouchlis et al. , 2015). The peptide regions 720–730 (red region in Fig. 4B and 4D) and 640–648 (green region in Fig. 4B and 4D) were found to be involved in the extraction process of P APC. Residues like Thr642 and Tyr643 serve as gate keepers controlling the volume of the GIVA iPLA<sub>2</sub> catalytic site and along with Lys725 assist also the extraction of the P APC through hydrogen-bonding with its phosphate group (Mouchlis et al. , 2015). MD simulations of the GVIA iPLA<sub>2</sub>-P APC complex showed the phosphate group interacts with Lys489, Asn658 and Lys729 while the choline group interacts with Asp484 and Asp733. The oxygen atom of the *sn*-2 carbonyl group is located near the oxyanion hole (Gly486/Gly487) participating in hydrogen-bonding, while its carbon atom is located near catalytic Ser519. An interesting separation of the *sn*-2 from the *sn*-1 fatty acid chain occurs in the case of GVIA iPLA<sub>2</sub> when the substrate binds in the hydrophobic region of its catalytic site. In the hydrophobic region, the fatty acyl chains interact with residues like Val548, Phe549, Phe643, Phe644, Val721, Phe722, Leu770, and Leu778 located on regions that showed a decreased deuteration level in DXMS experimental data.

### C. Hydrolysis of the phospholipid molecule and D. diffusion of the products

Since GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> share a common catalytic dyad of Ser/Asp, they also share a common hydrolytic mechanism (see Fig. S9 in reference Mouchlis et al. , 2015). After the extraction of a phospholipid molecule its binding is stabilized through interactions of its phosphate group with an asparagine and an arginine (GIVA cPLA<sub>2</sub>) or a lysine (GVIA iPLA<sub>2</sub>). Hydrolysis starts with the abstraction of a proton from the catalytic serine by the aspartic acid. The negatively charged serine attacks the carbon atom of the *sn*-2 carbonyl group forming a tetrahedral intermediate which is stabilized by two glycine residues referred to as the oxyanion hole. The tetrahedral intermediate or transition state is then cleaved leading to a serine-acyl enzyme intermediate while the aspartic acid is transferring a proton atom to the lysophospholipid product. In a similar manner, the serine-acyl enzyme intermediate breaks down releasing the free fatty acid (Dessen et al. , 1999, Mouchlis et al. , 2015). The products of the hydrolysis are then diffused back into the membrane and the hydrolytic cycle starts over again. The details of the products diffusion mechanism have not been elucidated yet, but we are planning to study it in future projects.

### Allosteric regulation of GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> by membrane

The association of GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> through their interfacial surface (surface dilution kinetics)(Carman et al. , 1995, Deems et al. , 1975) is the most interesting part of their catalytic cycle (Fig. 3). Our previously published MD simulation supported the notion that the interfacial surface of these two enzymes contains allosteric sites for binding to the membrane (Fig. S8 in reference Mouchlis et al. , 2015). During the association of these enzymes with the membrane, the membrane surface acts as an allosteric ligand shifting the conformational state of the enzyme from the inactive to the active (step A in Fig. 3) in an analogous manner to that described by Changeux (Changeux, 2012, Changeux, 2009, Monod et al. , 1965). In contrast to Changeux's original model, our PLA<sub>2</sub>-membrane models suggest that PLA<sub>2</sub> enzymes have a very large allosteric site or surface interacting with the membrane (Mouchlis et al. , 2015).

### Binding mode of GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> inhibitors

Experimental combined with computational techniques have successfully been used to develop potent and selective inhibitors for PLA<sub>2</sub> enzymes (Mouchlis et al. , 2011a, Mouchlis et al. , 2011b, Mouchlis et al. , 2010a, b, Mouchlis et al. , 2012). MD simulations guided by DXMS experimental data were also successfully used to help in understanding the binding mode of pyrrophenone and a 2-oxoamide in the catalytic site of GIVA cPLA<sub>2</sub>,(Burke et al. , 2009) and a trifluoromethyl ketone in the catalytic site of GVIA iPLA<sub>2</sub> (Hsu et al. , 2013). Pyrrophenone contains a thiazolidinedione ring designed to target Arg200 and a carbonyl group bridging the two benzoyl groups to target the GIVA cPLA<sub>2</sub> catalytic Ser228 and was synthesized in 2001 (Seno et al. , 2001). A 2-oxoamide AX007 contains a carboxylic acid designed to also target Arg200 and a 2-oxoamide to target catalytic Ser228 of GIVA cPLA<sub>2</sub> and was synthesized in 2002 (Kokotos et al. , 2002). The peptide regions 256–265, 268–279, 466–470, 473–478, and 684–689 exhibited decreased deuteration levels upon binding of pyrrophenone and AX007. MD simulations led to enzyme-inhibitor complexes (Fig. 5A and B) that showed interactions of the inhibitors with residues such as Pro263, Leu264, Leu267,

Val272, Tyr275, Trp464, Ile465, Ile469, Met470, and Phe683 (Burke et al. , 2009). Aromatic trifluoromethylketone inhibitors such as PHFK were synthesized by Kokotos' group to target the catalytic Ser519 of GVIA iPLA<sub>2</sub> (Baskakis et al. , 2008, Kokotos et al. , 2010). DXMS experiments showed that peptide regions 483–493, 516–525, 544–549, 631–655, and 773–778 demonstrated a significant decrease in deuteration levels upon binding of PHFK. Docking of PHFK in the catalytic site of GVIA iPLA<sub>2</sub> showed interactions with residues like Thr520, Phe549, Met544, Val548, Phe549, and Leu560 (Fig. 5C).

## Acknowledgments

We would like to thank the National Institutes of Health RO1-GM20501-40 (E.A.D.) for supporting our investigations of phospholipase A<sub>2</sub> over the years.

## References


- Ayilavarapu S, Kantarci A, Fredman G, Turkoglu O, Omori K, Liu H, et al. Diabetes-induced oxidative stress is mediated by Ca<sup>2+</sup>-independent phospholipase A<sub>2</sub> in neutrophils. *J Immunol.* 2010; 184:1507–15. [PubMed: 20053941]
- Balsinde J, Balboa MA, Dennis EA. Antisense inhibition of group VI Ca<sup>2+</sup>-independent phospholipase A<sub>2</sub> blocks phospholipid fatty acid remodeling in murine P388D1 macrophages. *J Biol Chem.* 1997; 272:29317–21. [PubMed: 9361012]
- Balsinde J, Balboa MA, Li WH, Llopis J. Cellular regulation of cytosolic group IV phospholipase A<sub>2</sub> by phosphatidylinositol bisphosphate levels. *J Immunol.* 2000
- Baskakis C, Magrioti V, Cotton N, Stephens D, Constantinou-Kokotou V, Dennis EA, et al. Synthesis of polyfluoro ketones for selective inhibition of human phospholipase A<sub>2</sub> enzymes. *J Med Chem.* 2008; 51:8027–37. [PubMed: 19053783]
- Bone RN, Gai Y, Magrioti V, Kokotou MG, Ali T, Lei X, et al. Inhibition of Ca<sup>2+</sup>-independent phospholipase A<sub>2</sub>β (iPLA<sub>2</sub>β) ameliorates islet infiltration and incidence of diabetes in NOD mice. *Diabetes.* 2015; 64:541–54. [PubMed: 25213337]
- Bucher D, Hsu YH, Mouchlis VD, Dennis EA, McCammon JA. Insertion of the Ca<sup>2+</sup>-independent phospholipase A<sub>2</sub> into a phospholipid bilayer via coarse-grained and atomistic molecular dynamics simulations. *PLoS Comput Biol.* 2013; 9:e1003156. [PubMed: 23935474]
- Buczynski M, Dumlao D, Dennis E. Thematic Review Series: Proteomics. An integrated omics analysis of eicosanoid biology. *J Lipid Res.* 2009; 50:1015–38. [PubMed: 19244215]
- Buczynski MW, Stephens DL, Bowers-Gentry RC, Grkovich A, Deems RA, Dennis EA. TLR-4 and sustained calcium agonists synergistically produce eicosanoids independent of protein synthesis in RAW264. 7 cells. *J Biol Chem.* 2007; 282:22834–47. [PubMed: 17535806]
- Burke J, Babakhani A, Gorfe A, Kokotos G, Li S, Woods V, et al. Location of inhibitors bound to group IVA phospholipase A<sub>2</sub> determined by molecular dynamics and deuterium exchange mass spectrometry. *J Am Chem Soc.* 2009; 131:8083–91. [PubMed: 19459633]
- Burke J, Hsu Y-H, Deems R, Li S, Woods V, Dennis E. A phospholipid substrate molecule residing in the membrane surface mediates opening of the lid region in group IVA cytosolic phospholipase A<sub>2</sub>. *J Biol Chem.* 2008; 283:31227–36. [PubMed: 18753135]
- Cao J, Burke J, Dennis E. Using hydrogen/deuterium exchange mass spectrometry to define the specific interactions of the phospholipase A<sub>2</sub> superfamily with lipid substrates, inhibitors, and membranes. *J Biol Chem.* 2013; 288:1806–13. [PubMed: 23209293]
- Carman G, Deems R, Dennis E. Lipid signaling enzymes and surface dilution kinetics. *J Biol Chem.* 1995; 270:18711–4. [PubMed: 7642515]
- Chalfant CE, Spiegel S. Sphingosine 1-phosphate and ceramide 1-phosphate: expanding roles in cell signaling. *J Cell Sci.* 2005; 118:4605–12. [PubMed: 16219683]
- Changeux J-P. Allostery and the Monod-Wyman-Changeux model after 50 years. *Annu Rev Biophys.* 2012; 41:103–33. [PubMed: 22224598]




- Changeux J-PP. Allosteric receptors: from electric organ to cognition. *Annu Rev Pharmacol Toxicol.* 2009; 50:1–38. [PubMed: 20055696]
- Clark JD, Lin LL, Kriz RW, Ramesha CS, Sultzman LA, Lin AY, et al. A novel arachidonic acid-selective cytosolic PLA<sub>2</sub> contains a Ca<sup>2+</sup>-dependent translocation domain with homology to PKC and GAP. *Cell.* 1991; 65:1043–51. [PubMed: 1904318]
- Das S, Cho W. Roles of catalytic domain residues in interfacial binding and activation of group IV cytosolic phospholipase A<sub>2</sub>. *J Biol Chem.* 2002; 277:23838–46. [PubMed: 11956215]
- Deems RA, Eaton BR, Dennis EA. Kinetic analysis of phospholipase A<sub>2</sub> activity toward mixed micelles and its implications for the study of lipolytic enzymes. *J Biol Chem.* 1975; 250:9013–20. [PubMed: 1194274]
- Dennis E, Cao J, Hsu Y-H, Magrioti V, Kokotos G. Phospholipase A<sub>2</sub> enzymes: physical structure, biological function, disease implication, chemical inhibition, and therapeutic intervention. *Chem Rev.* 2011; 111:6130–85. [PubMed: 21910409]
- Dennis EA, Norris PC. Eicosanoid storm in infection and inflammation. *Nat Rev Immunol.* 2015; 15:511–23. [PubMed: 26139350]
- Dessen A, Tang J, Schmidt H, Stahl M, Clark J, Seehra J, et al. Crystal structure of human cytosolic phospholipase A<sub>2</sub> reveals a novel topology and catalytic mechanism. *Cell.* 1999; 97:349–60. [PubMed: 10319815]
- Durrant JD, McCammon JA. Molecular dynamics simulations and drug discovery. *BMC Biol.* 2011
- Funk CD. Prostaglandins and leukotrienes: advances in eicosanoid biology. *Science.* 2001; 294:1871–5. [PubMed: 11729303]
- Ghosh M, Tucker DE, Burchett SA, Leslie CC. Properties of the Group IV phospholipase A<sub>2</sub> family. *Prog Lipid Res.* 2006; 45:487–510. [PubMed: 16814865]
- Hazen SL, Gross RW. Human myocardial cytosolic Ca<sup>2+</sup>-independent phospholipase A<sub>2</sub> is modulated by ATP. Concordant ATP-induced alterations in enzyme kinetics and mechanism-based inhibition. *Biochem J.* 1991; 280(Pt 3):581–7. [PubMed: 1764021]
- Hegen M, Sun L, Uozumi N, Kume K, Goad ME, Nickerson-Nutter CL, et al. Cytosolic phospholipase A<sub>2</sub>α-deficient mice are resistant to collagen-induced arthritis. *J Exp Med.* 2003; 197:1297–302. [PubMed: 12743172]
- Hsu Y-H, Bucher D, Cao J, Li S, Yang S-W, Kokotos G, et al. Fluoroketone Inhibition of Ca<sup>2+</sup>-Independent Phospholipase A<sub>2</sub> through Binding Pocket Association Defined by Hydrogen/Deuterium Exchange and Molecular Dynamics. *J Am Chem Soc.* 2013; 135:1330–7. [PubMed: 23256506]
- Hsu Y-H, Burke J, Li S, Woods V, Dennis E. Localizing the membrane binding region of Group VIA Ca<sup>2+</sup>-independent phospholipase A<sub>2</sub> using peptide amide hydrogen/deuterium exchange mass spectrometry. *J Biol Chem.* 2009; 284:23652–61. [PubMed: 19556238]
- Hsu Y-HH, Burke JE, Stephens DL, Deems RA, Li S, Asmus KM, et al. Calcium binding rigidifies the C2 domain and the intradomain interaction of GIVA phospholipase A<sub>2</sub> as revealed by hydrogen/deuterium exchange mass spectrometry. *J Biol Chem.* 2008; 283:9820–7. [PubMed: 18211893]
- Jenkins CM, Yan W, Mancuso DJ, Gross RW. Highly selective hydrolysis of fatty acyl-CoAs by calcium-independent phospholipase A<sub>2</sub>β. Enzyme autoacylation and acyl-CoA-mediated reversal of calmodulin inhibition of phospholipase A<sub>2</sub> activity. *J Biol Chem.* 2006; 281:15615–24. [PubMed: 16595686]
- Karabina SA, Gora S, Atout R, Ninio E. Extracellular phospholipases in atherosclerosis. *Biochimie.* 2010; 92:594–600. [PubMed: 20153800]
- Keyes KT, Ye Y, Lin Y, Zhang C. Resolvin E1 protects the rat heart against reperfusion injury. *Am J Physiol Heart Circ Physiol.* 2010
- Kokotos G, Hsu Y-H, Burke J, Baskakis C, Kokotos C, Magrioti V, et al. Potent and selective fluoroketone inhibitors of group VIA calcium-independent phospholipase A<sub>2</sub>. *J Med Chem.* 2010; 53:3602–10. [PubMed: 20369880]
- Kokotos G, Kotsovolou S, Six DA, Constantinou-Kokotou V, Beltzner CC, Dennis EA. Novel 2-Oxoamide Inhibitors of Human Group IVA Phospholipase A<sub>2</sub>. *J Med Chem.* 2002; 45:28912893.

- Kramer RM, Roberts EF, Manetta J, Putnam JE. The  $\text{Ca}^{2+}$ -sensitive cytosolic phospholipase  $\text{A}_2$  is a 100-kDa protein in human monoblast U937 cells. *J Biol Chem*. 1991; 266:5268–72. [PubMed: 2002059]
- Larsson Forsell PK, Kennedy BP, Claesson HE. The human calcium-independent phospholipase  $\text{A}_2$  gene multiple enzymes with distinct properties from a single gene. *Eur J Biochem*. 1999; 262:575–85. [PubMed: 10336645]
- Larsson PK, Claesson HE, Kennedy BP. Multiple splice variants of the human calcium-independent phospholipase  $\text{A}_2$  and their effect on enzyme activity. *J Biol Chem*. 1998; 273:207–14. [PubMed: 9417066]
- Leslie CC, Channon JY. Anionic phospholipids stimulate an arachidonoyl-hydrolyzing phospholipase  $\text{A}_2$  from macrophages and reduce the calcium requirement for activity. *Biochim Biophys Acta*. 1990; 1045:261–70. [PubMed: 2167132]
- Lio YC, Dennis EA. Interfacial activation, lysophospholipase and transacylase activity of group VI  $\text{Ca}^{2+}$ -independent phospholipase  $\text{A}_2$ . *Biochim Biophys Acta*. 1998; 1392:320–32. [PubMed: 9630702]
- Malkova S, Long F, Stahelin RV, Pingali SV, Murray D, Cho W, et al. X-ray reflectivity studies of  $\text{cPLA}_2\alpha$ -C2 domains adsorbed onto Langmuir monolayers of SOPC. *Biophysical journal*. 2005; 89:1861–73. [PubMed: 15994899]
- Masuda S, Murakami M, Komiyama K, Ishihara M, Ishikawa Y, Ishii T, et al. Various secretory phospholipase  $\text{A}_2$  enzymes are expressed in rheumatoid arthritis and augment prostaglandin production in cultured synovial cells. *FEBS J*. 2005; 272:655–72. [PubMed: 15670148]
- Michaely P, Tomchick D, Machius M, Anderson R. Crystal structure of a 12 ANK repeat stack from human ankyrinR. *EMBO J*. 2002; 21:6387–483. [PubMed: 12456646]
- Monod J, Wyman J, Changeux JP. ON THE NATURE OF ALLOSTERIC TRANSITIONS: A PLAUSIBLE MODEL. *J Mol Biol*. 1965; 12:88–118. [PubMed: 14343300]
- Mouchlis VD, Barbayianni E, Mavromoustakos TM, Kokotos G. The application of rational design on phospholipase  $\text{A}_2$  inhibitors. *Curr Med Chem*. 2011a; 18:2566–82. [PubMed: 21568891]
- Mouchlis VD, Bucher D, McCammon JA, Dennis EA. Membranes serve as allosteric activators of phospholipase  $\text{A}_2$ , enabling it to extract, bind, and hydrolyze phospholipid substrates. *PNAS*. 2015; 112:E516–E25. [PubMed: 25624474]
- Mouchlis VD, Magrioti V, Barbayianni E, Cermak N, Oslund RC, Mavromoustakos TM, et al. Inhibition of secreted phospholipases  $\text{A}_2$  by 2-oxoamides based on alpha-amino acids: Synthesis, in vitro evaluation and molecular docking calculations. *Bioorg Med Chem*. 2011b; 19:735–43. [PubMed: 21216150]
- Mouchlis VD, Mavromoustakos TM, Kokotos G. Design of new secreted phospholipase  $\text{A}_2$  inhibitors based on docking calculations by modifying the pharmacophore segments of the FPL67047XX inhibitor. *J Comput-Aided Mol Des*. 2010a; 24:107–15. [PubMed: 20130961]
- Mouchlis VD, Mavromoustakos TM, Kokotos G. Molecular docking and 3D-QSAR CoMFA studies on indole inhibitors of GIIA secreted phospholipase  $\text{A}_2$ . *J Chem Inf Model*. 2010b; 50:1589–601. [PubMed: 20795712]
- Mouchlis VD, Michopoulou V, Constantinou-Kokotou V, Mavromoustakos T, Dennis EA, Kokotos G. Binding conformation of 2-oxoamide inhibitors to group IVA cytosolic phospholipase  $\text{A}_2$  determined by molecular docking combined with molecular dynamics. *J Chem Inf Model*. 2012; 52:243–54. [PubMed: 22196172]
- Nakanishi M, Rosenberg DW. Roles of  $\text{cPLA}_2\alpha$  and arachidonic acid in cancer. *Biochim Biophys Acta*. 2006; 1761:1335–43. [PubMed: 17052951]
- Norris PC, Dennis EA. Omega-3 fatty acids cause dramatic changes in TLR4 and purinergic eicosanoid signaling. *PNAS*. 2012; 109:8517–22. [PubMed: 22586114]
- Norris PC, Gosselin D, Reichart D, Glass CK, Dennis EA. Phospholipase  $\text{A}_2$  regulates eicosanoid class switching during inflammasome activation. *PNAS*. 2014; 111:12746–51. [PubMed: 25139986]
- Park J, Lee C, Jang J-H, Ghim J, Kim Y-J, You S, et al. Phospholipase signalling networks in cancer. *Nat Rev Cancer*. 2012; 12:782–92. [PubMed: 23076158]


- Perisic O, Paterson HF, Mosedale G, Lara-Gonzalez S, Williams RL. Mapping the phospholipid-binding surface and translocation determinants of the C2 domain from cytosolic phospholipase A<sub>2</sub>. *J Biol Chem*. 1999; 274:14979–87. [PubMed: 10329700]
- Pettus BJ, Bielawska A, Subramanian P, Wijesinghe DS, Maceyka M, Leslie CC, et al. Ceramide 1-phosphate is a direct activator of cytosolic phospholipase A<sub>2</sub>. *J Biol Chem*. 2004; 279:11320–6. [PubMed: 14676210]
- Rosenson RS, Hurt-Camejo E. Phospholipase A<sub>2</sub> enzymes and the risk of atherosclerosis. *Eur Heart J*. 2012; 33:2899–909. [PubMed: 22802388]
- Rossi A, Pergola C, Koeberle A. The 5-lipoxygenase inhibitor, zileuton, suppresses prostaglandin biosynthesis by inhibition of arachidonic acid release in macrophages. *Br J Pharmacol*. 2010
- Rydell TJ, Williams JM, Krieger E, Moshiri F, Stallings WC, Brown SM, et al. The crystal structure, mutagenesis, and activity studies reveal that patatin is a lipid acyl hydrolase with a Ser-Asp catalytic dyad. *Biochemistry*. 2003; 42:6696–708. [PubMed: 12779324]
- Scott KF, Sajinovic M, Hein J, Nixdorf S, Galettis P, Liauw W, et al. Emerging roles for phospholipase A<sub>2</sub> enzymes in cancer. *Biochimie*. 2010; 92:601–10. [PubMed: 20362028]
- Seno K, Okuno T, Nishi K, Murakami Y, Yamada K, Nakamoto S, et al. Pyrrolidine inhibitors of human cytosolic phospholipase A<sub>2</sub>. Part 2: synthesis of potent and crystallized 4-triphenylmethylthio derivative 'pyrrophenone'. *Bioorg Med Chem Lett*. 2001; 11:587–90. [PubMed: 11229777]
- Sharp JD, White DL, Chiou XG, Goodson T, Gamboa GC, McClure D, et al. Molecular cloning and expression of human Ca<sup>2+</sup>-sensitive cytosolic phospholipase A<sub>2</sub>. *J Biol Chem*. 1991; 266:14850–3. [PubMed: 1869522]
- Six DA, Dennis EA. Essential Ca<sup>2+</sup>-independent Role of the Group IVA Cytosolic Phospholipase A<sub>2</sub> C2 Domain for Interfacial Activity. *J Biol Chem*. 2003; 278:23842–50. [PubMed: 12672805]
- Sliwoski G, Kothiwale S, Meiler J, Lowe EW. Computational methods in drug discovery. *Pharmacol Rev*. 2014; 66:334–95. [PubMed: 24381236]
- Vane JR. Biomedicine. Back to an aspirin a day? *Science*. 2002; 296:474–5. [PubMed: 11964462]
- Winstead MV, Balsinde J, Dennis EA. Calcium-independent phospholipase A<sub>2</sub>: structure and function. *Biochim Biophys Acta*. 2000; 1488:28–39. [PubMed: 11080674]




**Figure 1.** 3D structure of the enzymes. (A) The X-ray crystal structure of GIVA cPLA<sub>2</sub> (PDB ID 1CJY). (B) The homology model of GVIA iPLA<sub>2</sub>. Reprinted from reference Mouchlis et al. , 2015.




**Figure 2.** Eicosanoid pathway and its therapeutic targets. Downstream regulators such as COX, LOX and CYP rely on PLA<sub>2</sub> enzymes to provide them with AA. Reprinted from Dennis and Norris, 2015.



**Figure 3.**  
The catalytic cycle of GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub>. Reprinted from reference Mouchlis et al. , 2015.



**Figure 4.** Association of the catalytic domain with the membrane bilayer for (A) GIVA cPLA<sub>2</sub> and (B) GVIA iPLA<sub>2</sub>. Substrate binding of a PAPC phospholipid molecule in the catalytic site of (C) GIVA cPLA<sub>2</sub> and (D) GVIA iPLA<sub>2</sub> is shown. Reprinted from reference Mouchlis et al. , 2015.



**Figure 5.** Binding mode of GIVA cPLA<sub>2</sub> and GVIA iPLA<sub>2</sub> inhibitors elucidated using MD simulations guided by DXMS data. (A) Pyrrophenone in GIVA cPLA<sub>2</sub>. (B) 2-Oxoamide in GIVA cPLA<sub>2</sub>. (C) PHFK in GVIA iPLA<sub>2</sub>. Adopted from references 25 and 61.