

Cyclic endogenous estrogen and progesterone vary by mammographic density phenotypes in premenopausal women

Anita Iversen^{a,*}, Hanne Frydenberg^{b,*}, Anne-Sofie Furberg^{a,c}, Vidar G. Flote^b, Sissi Espetvedt Finstad^d, Anne McTiernan^e, Giske Ursin^f, Tom Wilsgaard^a, Peter T. Ellison^g, Grazyna Jasienska^h and Inger Thune^{a,b}

Estrogen and progesterone are key factors in the development of breast cancer, but it remains unclear whether these hormones are associated with mammographic density phenotypes in premenopausal women. We measured percent mammographic density, nondense area, and absolute mammographic density using computer-assisted breast density readings (Maden) from digitized mammograms taken on a scheduled day of the menstrual cycle (day 7–12) among 202 healthy, premenopausal women (Energy Balance and Breast cancer Aspects Study-I). Daily salivary concentrations of 17 β -estradiol and progesterone throughout an entire menstrual cycle and fasting morning serum concentrations of hormones on 3 specific days of the menstrual cycle were assessed. Salivary and serum 17 β -estradiol and progesterone were positively associated with percent mammographic density, we observed by 1 SD increase in overall salivary estradiol (β -value equal to 2.07, $P = 0.044$), luteal salivary progesterone (β -value equal to 2.40, $P = 0.020$). Women with above-median percent mammographic density had a 20% higher mean salivary 17 β -estradiol level throughout the menstrual cycle. The odds ratio for having above-median percent mammographic density (>28.5%) per 1 SD increase in overall salivary 17 β -estradiol was 1.66 (95% confidence interval 1.13–2.45). Women in the top tertile of the overall average daily 17 β -estradiol concentrations had an odds ratio of 2.54

Introduction

Women with higher levels of mammographically measured breast density have a significantly increased risk of developing breast cancer (Pettersson *et al.*, 2014). Absolute mammographic density reflects dense areas of the breast, hypothesized to be composed of epithelial and stromal tissues. Nondense area represents fat tissue and percent mammographic density reflects the relative amounts of fibroglandular and fat tissue (Stone *et al.*, 2010; Boyd *et al.*, 2011; Pettersson *et al.*, 2011). Both percent mammographic density and absolute mammographic density are positively correlated with the number of epithelial cells at risk for malignant transformation.

(confidence interval 1.05–6.16) of above-median percent mammographic density compared with women in the bottom tertile. Our finding of a relationship between estrogen, progesterone, and percent mammographic density and not with other mammographic density phenotypes in premenopausal women is biologically plausible, but needs to be replicated in larger studies. *European Journal of Cancer Prevention* 25:9–18 Copyright © 2015 Wolters Kluwer Health, Inc. All rights reserved.

European Journal of Cancer Prevention 2016, 25:9–18

Keywords: 17 β -estradiol, mammographic density, premenopausal women, progesterone

^aDepartment of Community Medicine, Faculty of Health Sciences, The Arctic University of Norway, ^bDepartment of Oncology, Oslo University Hospital HF, Ullevål, ^cDepartment of Microbiology and Infection Control, University Hospital North Norway, Tromsø, ^dThe Norwegian Directorate of Health, ^eCancer Registry of Norway, Oslo, Norway, ^fFred Hutchinson Cancer Research Center, Seattle, Washington, ^gDepartment of Human Evolutionary Biology, Harvard University, Cambridge, Massachusetts, USA and ^hDepartment of Environmental Health, Jagiellonian University Collegium Medicum, Krakow, Poland

Correspondence Hanne Frydenberg, MD, Department of Oncology, Oslo University Hospital HF, Ullevål, N-0424 Oslo, Norway
Tel: + 47 23026600; fax: + 47 23026601;
e-mail: hanne.frydenberg@medisin.uio.no

*Anita Iversen and Hanne Frydenberg contributed equally to the writing of this article.

Received 17 August 2014 Accepted 23 December 2014

Although a high percent mammographic density is associated with a three-to-six-fold increase in the risk of breast cancer compared with a low percent mammographic density (Pettersson *et al.*, 2014), the absolute dense area is considered to represent the actual target tissue for tumor development as ductal carcinoma *in situ* and invasive breast cancer more often occur in dense areas (Ursin *et al.*, 2005; Gill *et al.*, 2006; Pinto Pereira *et al.*, 2011).

Recently, diverse processes including growth factors, hormones, and interactions among epithelial cells, and the breast microenvironment, including fibroblast and adipocytes, have been shown to influence breast phenotypes (Brower, 2010; Boyd *et al.*, 2011; Pettersson *et al.*, 2014). Various breast cancer risk factors, such as age,

All supplementary digital content is available directly from the corresponding author.

reproductive factors, and BMI, have been associated with different breast tissue compositions (Stone *et al.*, 2010; Boyd *et al.*, 2011; Pettersson *et al.*, 2011). Moreover, estrogen and progesterone promote cellular and epithelial growth in the normal mammary gland (Henderson *et al.*, 1982; Bernstein, 2002; Mctiernan *et al.*, 2009), and are positively associated with an increased risk for breast cancer in both premenopausal and postmenopausal women (Key *et al.*, 2003, 2013). Interestingly, randomized-controlled trials have shown that percent mammographic density increases significantly with administration of combined estrogen–progesterone menopausal hormone therapy (Greendale *et al.*, 2003; Mctiernan *et al.*, 2005), whereas the estrogen receptor antagonist Tamoxifen reduces percent mammographic density (Cuzick *et al.*, 2004; Johansson *et al.*, 2013). However, it is unclear whether estrogen and progesterone levels throughout the menstrual cycle in premenopausal women are associated with mammographic density or vary by mammographic density phenotypes (Boyd *et al.*, 2002; Noh *et al.*, 2006; Yong *et al.*, 2009).

Previously, we observed in the Energy Balance and Breast cancer Aspects (EBBA) study-I that daily cyclic 17 β -estradiol was associated with breast cancer risk factors including age at menarche (Emaus *et al.*, 2008a) and body composition from birth to adult life (Furberg *et al.*, 2005; Jasienska *et al.*, 2006; Emaus *et al.*, 2008b; Finstad *et al.*, 2009; Barrett *et al.*, 2013). Furthermore, in a sub-analysis, we observed a positive association between daily progesterone concentrations and mammographic density in premenopausal women using a modified Wolfe's classification (Furberg *et al.*, 2005).

In the present study, we investigated the associations between salivary and serum sex hormones [estradiol, progesterone, testosterone, dehydroepiandrosterone-sulfate (DHEA-SO₄), follicle-stimulating hormone (FSH), and luteinizing hormone (LH)] and mammographic density phenotypes (percent mammographic density, nondense area, and absolute mammographic density), assessed by the computer-assisted method (Madena; University of Southern California School of Medicine, Los Angeles, California, USA) (Ursin *et al.*, 2003), among premenopausal women from the EBBA-I study.

Materials and methods

Study design, setting, and participants

The women participating in the Norwegian EBBA-I Study (2000–2002) were recruited through local media campaigns. A total of 204 women aged 25–35 years were included, and fulfilled the following criteria: regular menstrual cycles (cycle length: 22–38 days within the previous 3 months), no use of any regular (daily/weekly) medication, no pregnancy or lactation or use of steroid contraceptives over the previous 6 months, and no history of gynecological or chronic disorders (e.g. diabetes, hypo-

hyperthyroidism, polycystic ovary syndrome) (Furberg *et al.*, 2005; Iversen *et al.*, 2011). Two women were excluded because of missing mammographic data, leaving data from 202 premenopausal women available for the present study.

We used a standardized questionnaire to collect information on reproductive history, previous hormone use, and lifestyle habits. The same trained nurse interviewed all participants.

Clinical examination

The participants underwent a clinical examination at three scheduled visits over the course of one menstrual cycle: first visit (days 1–5, early follicular phase), second visit (days 7–12, late follicular phase), and third visit (days 21–25, late luteal phase). They came in on the first possible day after the onset of menstrual bleeding for clinical examinations at the Clinical Research Center, University Hospital of North Norway (UNN), Tromsø. Fasting blood samples were collected and analyzed at the Department of Clinical Chemistry, UNN. Height was measured to the nearest 0.5 cm and weight to the nearest 0.1 kg on an electronic scale, and BMI was calculated in kg/m². At the second visit, participants underwent a full-body scan to estimate total percent body fat using dual-energy X-ray absorptiometry (DPLX-L 2288; Lunar Radiation Corporation, Madison, Wisconsin, USA) (Furberg *et al.*, 2005).

Serum sex steroid hormone assessment procedures

Fasting serum samples were measured at three scheduled visits during the menstrual cycle. Concentrations of estradiol and progesterone were measured using a direct immunometric assay (Immuno-1; Bayer Diagnostics, at the Department of Clinical Chemistry, UNN, Tromsø, Norway) in fresh fasting serum samples at all three visits during the menstrual cycle. The sensitivity for estradiol was 0.01 nmol/l and the coefficient of variation (CV) was 3.9%. The sensitivity and CV for progesterone were 0.13 nmol/l and 5.7%, respectively. Sex hormone-binding globulin was measured using an immunometric method (both Diagnostic Products Corporation, Bierman GmbH, Bad Nauheim, Germany), with a CV of 5–10%. Serum testosterone was measured using an enhanced chemiluminescence immunoassay, using Elecsys 2010 from Roche Diagnostics (Mannheim, Germany), in 153 samples (74%), whereas for the remaining samples, Immuno-1 from Bayer Diagnostics (Tarrytown, New York, USA) was used. The CV was 0.97 in parallel runs of the testosterone assays and no correction formula was used. Serum DHEA-SO₄ was measured using a competitive immunometric assay. LH and FSH were measured in serum samples using Techicon Immuno-1 immunometric assays (Bayer Diagnostics). Both assays were standardized against the WHO 2nd International Standard (for FSH: IRP 78/549 and for LH: IRP 68/40).

The sensitivity of the FSH assay was 0.1 IU/l and the CV was less than 7%. For LH, the assay sensitivity was 0.3 IU/l and the CV was 5–10%.

Daily saliva sampling

The women collected daily morning saliva samples at home for one entire menstrual cycle, and sampling started on the first day of menstrual bleeding according to protocols established at the Reproductive Ecology Laboratory, Harvard University (USA) (Lipson and Ellison, 1989; Ellison and Lipson, 1999; Furberg *et al.*, 2005). Levels of 17 β -estradiol were measured in daily saliva samples from 20 days (reverse cycle day –5 to –24) and levels of progesterone from 14 days (reverse cycle days –1 to –14). ¹²⁵I-labeled RIA kits (#39100; Diagnostic Systems Laboratories, Webster, Texas, USA) were used along with published modifications of the manufacturer's protocols (Furberg *et al.*, 2005). All samples were run in duplicate. All of a participant's samples were run in the same batch, with women assigned to batches randomly. CVs were calculated on the basis of high and low value pools (appropriate to the range of each steroid) included in each assay (Furberg *et al.*, 2005). In the present study, measurements of 17 β -estradiol at the beginning and at the end of the cycles had higher CVs. The sensitivity of the 17 β -estradiol assay (lowest value measurable by assay) was 4 pmol/l. The average intra-assay variability was 9% and the interassay variability ranged from 23% for low pools to 13% for high pools. For progesterone, the sensitivity of the assay was 13 pmol/l. The average intra-assay variability was 10%; the inter-assay variability ranged from 19% for low pools to 12% for high pools.

Alignment of the cycles was based on the identification of the mid-cycle decrease in estradiol (aligned cycle day 0), which provides a good estimate of the day of ovulation (Ellison and Lipson, 1999). Identification of the decrease in the salivary 17 β -estradiol concentration was not satisfactory for 14 women; thus, their cycles were not aligned. The overall average salivary 17 β -estradiol and progesterone were calculated for all women, and additional indices of average hormone concentrations were calculated for 188 women: follicular estradiol index (aligned cycle days –7 to –1), luteal estradiol index (aligned cycle days 0 to +6), mid-menstrual estradiol index (aligned cycle days –7 to +6), and early-mid luteal progesterone index (aligned cycle days 0 to +9).

Mammographic density phenotypes

Bilateral two-view mammograms were obtained from all women during the same menstrual cycle as the serum and salivary sampling were performed, late follicular phase, second visit (days 7–12) at the Center of Breast Imaging, UNN, using a standard protocol (Bjursta *et al.*, 2003; Furberg *et al.*, 2005). Left craniocaudal mammograms were digitized and imported into a computerized

mammographic density assessment program (Madena) (Ursin *et al.*, 1998, 2003), and the breast areas were outlined by a trained research assistant using validated methods (Ursin *et al.*, 1998). The total breast area was defined on the mammographic image using a special outlining tool. The region of interest (ROI) was then outlined. The mammogram reader used a tinting tool to apply yellow tint to areas considered to represent mammographically dense areas. The Madena software estimated the total number of pixels and the number of tinted pixels in the ROI.

Absolute mammographic density represents the number of the tinted pixels within the ROI. The nondense area reflects the total breast area, minus the dense area. Percent mammographic density is the ratio of absolute mammographic density to the total breast area (area of ROI) multiplied by 100. The mammograms were read in four batches, with an equal number of mammograms in each batch. A duplicate reading of 26 randomly selected

Table 1 Descriptive characteristics of the study population: the Norwegian Energy Balance and Breast Cancer Aspects-I study ($n = 202$)^a

Characteristics of the study population	Mean	SD
Age (years)	30.7	3.07
Education (total years)	16.1	3.02
Body composition ^b		
BMI (kg/m ²)	24.4	3.77
Waist circumference (cm)	79.5	9.80
Tissue fat, DEXA scan (%) ^c	34.2	7.62
Reproductive factors		
Parity (% parous)	47.1	–
Age at menarche (years)	13.1	1.36
Cycle length (days)	28.2	3.17
Lifestyle factors		
Energy intake (kJ/day)	8.09	1.90
Previous use of OC (%)	82.7	–
Leisure time (MET h/week)	57.6	88.6
Alcohol (U/week)	2.89	3.38
Current smokers (%)	22.1	–
Serum hormones ^d		
Estradiol (pmol/l)	146.7	61.6
Progesterone (nmol/l)	6.29	4.83
Testosterone (nmol/l)	1.47	0.53
SHBG (nmol/l)	51.9	19.5
DHEA-SO ₄ (mcmol/l)	4.59	2.07
LH (IU/l)	6.28	5.31
FSH (IU/l)	6.34	2.12
Salivary hormones ^e		
Overall average 17 β -estradiol (pmol/l)	17.9	8.79
Overall average progesterone (pmol/l)	130.2	68.3
Mammograms ^c		
Percent mammographic density (%)	29.8	19.0
Absolute mammographic density (cm ²)	34.7	23.4
Nondense area (cm ²)	102.6	66.2
Total mammographic density (cm ²)	137.3	62.5

All analyses have used descriptive statistics.

DEXA, dual-energy X-ray; DHEA-SO₄, dehydroepiandrosterone-sulfate; FSH, follicle-stimulating hormone; LH, luteinizing hormone; MET, metabolic equivalents; OC, oral contraceptives; SHBG, sex hormone-binding globulin.

^aNumbers may vary because of missing information.

^bMeasurements at days 1–5 after the onset of the menstrual cycle.

^cMammograms and total tissue fat (DEXA) were taken days 7–12 (mid-cycle phase).

^dSerum samples in early follicular phase: days 1–5 after the onset of the menstrual cycle.

^eDaily saliva samples throughout an entire menstrual cycle.

Table 2 The association by 1 SD higher level of sex steroid hormones and mammographic density phenotypes in premenopausal women ($n = 202$)^a using multivariable linear regression models

Sex steroid hormones	Percent density (%)		Absolute density (cm ²)		Nondense area (cm ²)	
	β -Value (95% CI)	P-value	β -Value (95% CI)	P-value	β -Value (95% CI)	P-value
Estradiol						
Saliva samples (pmol/l)						
Overall average by 1 SD (8.79) ^b	2.07 (0.05 to 4.08)	0.044	1.32 (-1.87 to 4.52)	0.415	-0.26 (-6.22 to 5.70)	0.932
Mid-menstrual by 1 SD (8.98) ^c	1.61 (-0.45 to 3.68)	0.125	0.52 (-2.66 to 3.70)	0.747	1.12 (-5.05 to 7.29)	0.721
Follicular by 1 SD (9.58) ^d	1.94 (-0.15 to 4.02)	0.069	1.19 (-2.04 to 4.41)	0.469	0.40 (-5.87 to 6.66)	0.901
Luteal by 1 SD (9.22) ^e	1.15 (-0.89 to 3.19)	0.267	-0.18 (-3.30 to 2.95)	0.912	1.73 (-4.34 to 7.80)	0.575
AUC by 1 SD (133.0) ^f	1.63 (-0.43 to 3.70)	0.125	0.58 (-2.60 to 3.76)	0.719	0.96 (-5.21 to 7.14)	0.759
Serum samples (nmol/l)						
Early follicular by 1 SD (0.06) ^g	0.41 (-1.55 to 2.37)	0.680	0.27 (-2.82 to 3.36)	0.863	1.71 (-4.06 to 7.47)	0.560
Late follicular by 1 SD (0.35) ^h	0.08 (-1.91 to 2.07)	0.936	-0.97 (-4.13 to 2.18)	0.543	-2.83 (-8.65 to 3.00)	0.339
Late luteal by 1 SD (0.20) ⁱ	2.78 (0.74 to 4.80)	0.008	2.12 (-1.13 to 5.36)	0.200	-7.30 (-13.30 to -1.31)	0.014
Progesterone						
Saliva samples (pmol/l)						
Overall average by 1 SD (68.3) ^b	1.75 (-0.22 to 3.71)	0.081	1.44 (-1.67 to 4.55)	0.363	-3.53 (-9.31 to 2.55)	0.230
Early-mid luteal by 1 SD (73.5) ^j	2.40 (0.39 to 4.41)	0.020	2.13 (-0.98 to 5.24)	0.178	-4.93 (-10.95 to 1.10)	0.108
AUC by 1 SD (717.7) ^f	1.91 (-0.11 to 3.93)	0.064	1.56 (-1.56 to 4.68)	0.325	-3.61 (-9.65 to 2.44)	0.241
Serum samples (nmol/l)						
Early follicular by 1 SD (6.29) ^g	-0.62 (-2.59 to 1.35)	0.536	-0.10 (3.22 to 3.02)	0.948	1.79 (-4.02 to 7.61)	0.544
Late follicular by 1 SD (7.59) ^h	2.94 (1.02 to 4.86)	0.003	4.28 (1.22 to 7.33)	0.006	-1.59 (-7.33 to 4.18)	0.589
Late luteal by 1 SD (20.11) ⁱ	0.93 (-1.16 to 3.02)	0.380	0.61 (-2.70 to 3.91)	0.718	-2.89 (-9.05 to 3.28)	0.357
Androgens						
Serum samples ^g						
Testosterone, by 1 SD (0.53 nmol/l)	0.79 (-1.23 to 2.82)	0.441	0.16 (-3.05 to 3.36)	0.922	-3.32 (-9.19 to 2.55)	0.266
DHEA-SO ₄ , by 1 SD (2.07 mcmmol/l)	1.55 (-0.46 to 3.56)	0.131	1.96 (-1.22 to 5.14)	0.225	-2.85 (-8.70 to 3.01)	0.339
SHBG, by 1 SD (19.53 nmol/l) ^g	0.60 (-1.53 to 2.72)	0.581	-0.13 (-3.48 to 3.22)	0.938	-0.02 (-0.35 to 0.31)	0.891
Gonadotrophins						
Serum samples (IU/l) ^g						
LH, early follicular by 1 SD (5.32)	0.08 (-1.90 to 2.06)	0.936	-0.48 (-3.58 to 2.63)	0.763	-0.55 (-6.33 to 5.24)	0.852
FSH, early follicular by 1 SD (2.13)	0.68 (-1.33 to 2.70)	0.505	1.24 (-1.93 to 4.41)	0.442	-5.76 (-11.61 to 0.97)	0.054

All analyses have used multivariable linear regression models, and are adjusted for age (continuous), BMI (continuous), parity (categorical), previous oral contraceptives (categorical), and current smokers (categorical).

AUC, area under curve; CI, confidence interval; DHEA-SO₄, dehydroepiandrosterone-sulfate; FSH, follicle-stimulating hormone; LH, luteinizing hormone; SHBG, sex hormone-binding globulin.

^aNumbers may vary because of missing information.

^bDaily saliva samples throughout an entire menstrual cycle.

^cDaily saliva samples in the mid-menstrual phase: aligned cycle day -7, +6.

^dDaily saliva samples in the follicular phase: aligned cycle day -7, -1.

^eDaily saliva samples in the luteal phase: aligned cycle day 0, +6.

^fAUC estimated from aligned cycle days -10, +9.

^gSerum samples in the early follicular phase: day 1-5 after onset of menstrual cycle.

^hSerum samples in the late follicular phase: day 7-12 after onset of menstrual cycle.

ⁱSerum samples in the late luteal phase: day 21-25 after onset of menstrual cycle.

^jDaily saliva samples in the luteal phase: aligned cycle day 0, +9.

mammograms from two of the batches showed a Pearson's correlation coefficient of 0.97. The reader was blinded to all the characteristics of the study population.

Ethical considerations

All the participants signed an informed consent form and the Regional Committee for Medical Research Ethics and the Norwegian Data Protection Agency approved the study.

Statistical analysis

The associations between sex steroid hormones in both saliva and serum and mammographic density were analyzed using multivariable linear and logistic regression models. Mammographic density outcome variables were used as both continuous and dichotomized variables representing lower and higher density using median values as cut-off points: percent mammographic density (28.5%), nondense area (84.7 cm²), and absolute

mammographic density (32.4 cm²). Both mammographic and hormone variables were approximately normally distributed, enabling data analyses by parametric tests.

On the basis of suggested biological mechanisms influencing levels of estradiol and progesterone or mammographic density, several models were used, including a variety of potentially confounding variables. Age, BMI, parity, previous oral contraceptives (OC) use, and current smoking were included as covariates in the final models. The adjusted β -values and odds ratio (ORs) of having above-median mammographic density were estimated according to a 1 SD higher level of ovarian hormones. The area under the curve for estradiol and progesterone was calculated for each participant with an aligned cycle using the trapezium rule (Matthews *et al.*, 1990).

To study in detail how measures of mammographic density vary among premenopausal women in groups of 'low', 'medium', and 'high' cyclic endogenous estrogen

and progesterone levels, we used tertiles (T1–T3). Furthermore, we used adjusted linear mixed models for repeated measures to study variations in daily salivary 17 β -estradiol and salivary progesterone across the menstrual cycle according to low and high (median split) levels of mammographic density phenotypes. The Toeplitz covariance structure yielded the best fit to the data and was used in all models.

All statistical tests were two-sided using a 5% significance level. Statistical analyses were carried out using SPSS version 21.0 (IBM Corporation, Armonk, New York, USA).

Results

Patient characteristics

There were 204 healthy participants; their mean age was 30.7 years. The mean (median) percent mammographic density was 29.8% (28.5%), nondense area 102.6 cm² (84.7 cm²), and the absolute mammographic density was 34.7 cm² (32.4 cm²) (Table 1).

Association between hormones and mammographic density

Percent mammographic density was positively associated with salivary estrogen and progesterone after adjustments. Both a 1 SD increase in the overall average salivary estradiol ($\beta=2.07$, $P=0.044$) and late luteal serum

estradiol ($\beta=2.78$, $P=0.008$) were positively associated with percent mammographic density after adjusting for age, BMI, parity, smoking, and previous OC use in linear regression analysis (Table 2). Furthermore, a one SD increase in luteal salivary progesterone ($\beta=2.40$, $P=0.020$), and late follicular serum progesterone ($\beta=2.94$, $P=0.003$) were positively associated with percent mammographic density (Table 2). No associations were observed between gonadotropins (LH, FSH) or androgens (testosterone, DHEA-SO₄) and percent mammographic density, and no associations were observed between salivary 17 β -estradiol, or progesterone with either absolute mammographic density or nondense area (Table 2). Late follicular serum progesterone was the only hormone positively associated with absolute mammographic density ($\beta=4.28$, $P=0.006$) (Table 2).

In adjusted logistic regression analysis, a 1 SD increase in salivary 17 β -estradiol in all menstrual phases was associated with statistically significant 50–60% higher odds of above-median percent mammographic density (i.e. >28.5%) (Table 3). In addition, a 1 SD increase in late follicular serum progesterone was associated with an OR of 1.72 [95% confidence interval (CI) 1.00–2.95] for having above-median percent mammographic density. No associations were found between the dichotomized (above-median) nondense area or absolute mammographic density and

Table 3 Odds ratio with 95% confidence interval for above-median percent mammographic density (>28.5%) by 1 SD higher level of ovarian hormones among premenopausal women (n = 202)^a

Endogenous ovarian hormones	Percent mammographic density above median (>28.5%)	
	OR (95% CI)	P-value
Estradiol		
Saliva samples (pmol/l)		
Overall average by 1 SD (8.79 pmol/l) ^b	1.66 (1.13 to 2.45)	0.010
Mid-menstrual by 1 SD (8.98 pmol/l) ^c	1.66 (1.11 to 2.48)	0.013
Follicular by 1 SD (9.58 pmol/l) ^d	1.69 (1.12 to 2.55)	0.012
Luteal by 1 SD (9.22 pmol/l) ^e	1.55 (1.06 to 2.29)	0.025
AUC by 1 SD (133.0 pmol/l \times cycle) ^f	1.67 (1.11 to 2.50)	0.013
Serum samples (nmol/l)		
Early follicular by 1 SD (0.06 nmol/l) ^g	0.94 (0.67 to 1.32)	0.707
Late follicular by 1 SD (0.35 nmol/l) ^h	1.31 (0.91 to 1.90)	0.153
Late luteal by 1 SD (0.20 nmol/l) ⁱ	7.61 (1.85 to 31.23)	0.005
Progesterone		
Saliva samples (pmol/l)		
Overall by 1 SD (68.27 pmol/l) ^b	1.32 (0.91 to 1.90)	0.144
Early-mid luteal by 1 SD (73.5 pmol/l)	1.48 (0.99 to 2.20)	0.055
AUC by 1 SD (717.7 pmol/l) ^f	1.33 (0.91 to 1.94)	0.146
Serum samples (nmol/l)		
Early follicular by 1 SD (6.29 nmol/l) ^g	0.78 (0.51 to 1.20)	0.255
Late follicular by 1 SD (7.59 nmol/l) ^h	1.72 (1.00 to 2.95)	0.051
Late luteal by 1 SD (20.11 nmol/l) ⁱ	1.39 (0.96 to 2.02)	0.084

All analyses have used multivariable logistic regression models, and are adjusted for age (continuous), BMI (continuous), parity (categorical), previous oral contraceptives (categorical), and current smokers (categorical).

AUC, area under curve; CI, confidence interval; OR, odds ratio.

^aNumbers may vary because of missing information.

^bDaily saliva samples throughout an entire menstrual cycle.

^cDaily saliva samples in the mid-menstrual phase: aligned cycle days –7, +6.

^dDaily saliva samples in the follicular phase: aligned cycle days –7, –1.

^eDaily saliva samples in the luteal phase: aligned cycle days 0, +6.

^fAUC estimated from aligned cycle days –10, +9.

^gSerum samples in the early follicular phase: day 1–5 after onset of menstrual cycle.

^hSerum samples in the late follicular phase: day 7–12 after onset of menstrual cycle.

ⁱSerum samples in the late luteal phase: day 21–25 after onset of menstrual cycle.

levels of estradiol or progesterone (Supplementary Table 1).

Women in the top tertile of the overall average salivary 17 β -estradiol concentration had a 2.5 times higher odds of above-median percent mammographic density ($\geq 28.5\%$) compared with women in the bottom tertile [T3 vs. T1: OR 2.54 (95% CI 1.05–6.16) $P_{\text{trend}}=0.037$] (Table 4). Similarly, associations were observed for mid-menstrual, luteal, and area under the curve measures of salivary 17 β -estradiol. Women in the two top tertiles of salivary luteal progesterone had a 3.4 times higher odds of above-median percent mammographic density compared with women in the bottom tertile [T2 vs. T1: OR 3.46 (95% CI 1.34–8.92), T3 vs. T1: OR 3.37 (95% CI 1.31–8.68), $P_{\text{trend}}=0.014$] (Table 4).

Associations between repeated measures of hormones and mammographic density

In the mixed linear regression models, we found that women with above-median percent mammographic density had 20% higher mean salivary 17 β -estradiol levels throughout the menstrual cycle compared with women with below-median percent mammographic density ($P=0.011$) (Fig. 1a). No associations were observed between salivary estradiol and progesterone with either absolute mammographic density or nondense area (Fig. 1b–f).

Discussion

In this study of premenopausal women, positive associations were observed between salivary and serum estradiol and progesterone, and percent mammographic density, but no clear associations were observed between these hormones and other mammographic density phenotypes. Women in the top tertiles of overall average daily 17 β -estradiol and luteal progesterone concentrations had about two to three times higher odds of having above-median percent mammographic density compared with women in the bottom tertile.

These results are unique, but partly supported (Noh *et al.*, 2006; Yong *et al.*, 2009). A positive association was observed between serum estradiol concentrations ($n=192$) (Yong *et al.*, 2009) and serum concentration of progesterone ($n=204$) (Noh *et al.*, 2006), and percent mammographic density among older premenopausal women. In addition, urinary estrogen metabolites were associated with both percent and absolute mammographic density among premenopausal women (Walker *et al.*, 2009). However, the associations observed between endogenous hormones and mammographic density phenotypes were weaker or inconclusive when BMI was included as a covariate (Boyd *et al.*, 2002; Tamimi *et al.*, 2005; Johansson *et al.*, 2008; Walker *et al.*, 2009). Thus, the associations observed between endogenous hormones and different mammographic density phenotypes

Table 4 Odds ratios for higher percent mammographic density (>28.5%) associated with endogenous ovarian hormones by tertiles of salivary 17 β -estradiol and progesterone ($n=202$)^a

Endogenous ovarian hormones	T1	T2	T3	P_{trend}
Tertiles of salivary 17 β -estradiol (T1–T3) (pmol/l)				
Overall ^b	1.0 (ref)	2.14 (0.93 to 4.97)	2.54 (1.05 to 6.16)	0.037
Mid-menstrual ^c	1.0 (ref)	1.83 (0.74 to 4.49)	2.44 (0.97 to 6.15)	0.057
Luteal ^d	1.0 (ref)	1.83 (0.74 to 4.53)	2.70 (1.07 to 6.80)	0.034
AUC ^e	1.0 (ref)	1.95 (0.78 to 4.85)	2.40 (0.95 to 6.05)	0.063
Tertiles of salivary progesterone (T1–T3) (pmol/l)				
Overall ^f	1.0 (ref)	2.02 (0.84 to 4.83)	2.36 (1.00 to 5.58)	0.054
Early-mid luteal ^g	1.0 (ref)	3.46 (1.34 to 8.92)	3.37 (1.31 to 8.68)	0.014
AUC ^h	1.0 (ref)	2.63 (1.03 to 6.72)	3.16 (1.25 to 8.02)	0.017

All analyses have used multivariable logistic regression models, and have adjusted for age (continuous), BMI (continuous), parity (categorical) previous oral contraceptives (categorical), and current smokers (categorical). AUC, area under curve; CI, confidence interval; OR, odds ratio; T1–T3, tertiles of 17 β -estradiol and progesterone in the saliva.

^aNumbers may vary because of missing information.

^bDaily saliva estradiol throughout an entire menstrual cycle. T1 ≤ 13.0 pmol/l, T2 13.0–20.6 pmol/l, T3 > 20.60 pmol/l.

^cDaily saliva estradiol in the mid-menstrual phase: aligned cycle days –7, +6.

T1 ≤ 13.1 pmol/l, T2 13.1–20.1 pmol/l, T3 > 20.1 pmol/l.

^dDaily saliva estradiol in the luteal phase: aligned cycle days 0, +6.

T1 ≤ 12.3 pmol/l, T2 12.3–20.1 pmol/l, T3 > 20.1 pmol/l

^eAUC estradiol estimated from aligned cycle days –10, +9. T1 ≤ 191.5 pmol/l

\times cycle, T2 191.5–307.0 pmol/l \times cycle, T3 > 307.0 pmol/l \times cycle.

^fDaily saliva progesterone throughout an entire menstrual cycle. T1 ≤ 96.2 pmol/l,

T2 96.2–146.3 pmol/l, T3 > 146.3 pmol/l.

^gDaily saliva progesterone in luteal phase: aligned cycle day 0, +6.

T1 ≤ 104.9 pmol/l, T2 104.9–161.9 pmol/l, T3 > 161.9 pmol/l.


^hAUC progesterone estimated from aligned cycle days –10, +9.

T1 ≤ 1005.0 pmol/l \times cycle, T2 1005.0–1494.0 pmol/l \times cycle, T3 > 1494.0 pmol/l \times cycle.

have been divergent (Boyd *et al.*, 2002; Walker *et al.*, 2009).

The present study provides novel information by including both fasting serum levels of 17 β -estradiol and progesterone from 3 scheduled days, and daily salivary levels of both hormones over an entire menstrual cycle. Previous studies have measured only blood hormones at discrete periods (Boyd *et al.*, 2002; Noh *et al.*, 2006; Walker *et al.*, 2009; Yong *et al.*, 2009; Maskarinec *et al.*, 2012). Our study provides a more accurate estimate of likely exposure to estradiol and progesterone across the entire menstrual cycle (Bellem *et al.*, 2011). However, we observed no correlations between serum and salivary hormones among all the participants, which is consistent with the results of others (Lu *et al.*, 1999) as a correlation between salivary and serum concentrations within the individual was observed, but not in total. This may be explained by the fact that serum hormones are dependent on the protein-binding capacity, which differs markedly among individuals. Moreover, the serum levels in one individual cannot be predicted from salivary concentrations in others (Ellison and Lipson, 1999; Lu *et al.*, 1999). Interestingly, almost all of our observed associations were in relation to estrogen and progesterone and percent mammographic density, and not with other mammographic phenotypes. Recently, a meta-analysis of 13 case–control

Fig. 1



Adjusted mean salivary ovarian hormones by high (red line) and low (blue line) percent mammographic density (a, c), absolute mammographic density (b, e), and nondense area (c, f) among 202 premenopausal women. Note that all analyses have used linear mixed models for repeated measures, adjusted for age (continuous), BMI (kg/m²) (continuous), parity (categorical), previous oral contraceptives (categorical), and current smokers (categorical). Mean salivary estradiol throughout the menstrual cycle: (a) Percent mammographic density: ≤ 28.5%, 16.3 pmol/l; > 28.5%, 20.4 pmol/l ($P=0.011$). (b) Absolute mammographic density: ≤ 32.4 cm², 17.4 pmol/l; > 32.4 cm², 19.4 pmol/l ($P=0.148$). (c) Nondense area: ≤ 84.7 cm², 18.9 pmol/l; > 84.7 cm², 17.8 pmol/l ($P=0.501$). Mean salivary progesterone level throughout the menstrual cycle: (d) percent mammographic density: ≤ 28.5%, 121.8 pmol/l; > 28.5%, 140.7 pmol/l ($P=0.118$). (e) Absolute mammographic density: ≤ 32.4 cm², 122 pmol/l; > 32.4 cm², 140 pmol/l ($P=0.092$). (f) Nondense area: ≤ 84.7 cm², 134.9 pmol/l; > 84.7 cm², 126.3 pmol/l ($P=0.490$).

studies including both premenopausal and postmenopausal women found that percent mammographic density was a stronger risk factor for breast cancer than absolute mammographic density, potentially suggesting that the ratio of fibroglandular to fat tissue may be important in relation to the development of breast cancer (Pettersson *et al.*, 2014). Recently, the importance of both percent and absolute mammographic density has been considered, in addition to established risk factors, in a predictive model for breast cancer (Rauh *et al.*, 2012). However, a recent study found absolute breast density to be a better breast cancer risk marker in women with an unfavorable metabolic profile as percent density was correlated negatively with nondense area and adiposity (Schetter *et al.*, 2014). However, the women in our study had a mean BMI of 24.4 kg/m².

Mammographic density reflects proliferation of epithelial and stromal cells, as well as the cumulative exposure of the breast to different mitogens including sex hormones (Boyd *et al.*, 2011), and in the 'Pike model', breast tissue aging is hypothesized to reflect reproductive factors and the cumulative hormone exposure. The rate of breast tissue aging is most rapid at the time of menarche, slows with each pregnancy, and slows further in the menopausal phase (Pike *et al.*, 1983); the same pattern has been observed in mammographic density (Boyd *et al.*, 2007; Boyd, 2013). Interestingly, local estrogen production in the breast, rather than circulating estrogen levels, has been suggested to be more relevant to breast density in postmenopausal women (Pettersson *et al.*, 2014), whereas one may hypothesize that percent mammographic density may be a better marker in premenopausal women.

Whether percent mammographic density or a specific threshold of percent mammographic density in early adulthood is predictive of breast cancer risk later in life remains unclear. However, previous studies in premenopausal (Van Gils *et al.*, 2000) and postmenopausal (Yaghjian *et al.*, 2013) women have found a two-fold to three-fold increase in breast cancer risk for women with percent mammographic density above 25%. These observations support comparison of groups of women with above-versus below-median percent mammographic density, as we did in our study.

There are several strengths, but also limitations to our study. Fasting serum samples were collected at three scheduled visits during the menstrual cycle, and salivary measurements of unbound estradiol and progesterone concentrations were collected daily over an entire menstrual cycle, following strict procedures (Lipson and Ellison, 1989; Ellison and Lipson, 1999; Gann *et al.*, 2001; Furberg *et al.*, 2005; Jasienska and Jasienski, 2008). Using noninvasive daily salivary samples, we could measure the free biologically active forms of estrogen and progesterone, which are considered ideal measures among premenopausal women (Ellison and Lipson, 1999; Bellem *et al.*, 2011). In addition, we measured the total sex steroid hormone, and could

capture the total sex steroid hormone exposure throughout the entire menstrual cycle. All mammograms were assessed within a narrow time frame (between days 7 and 12), thereby avoiding the bias of variation in mammographic density during the menstrual cycle (Morrow *et al.*, 2010). We used a validated computer-assisted method to quantify mammographic density (Ursin *et al.*, 1998), and all mammograms were read by one experienced blinded reader. The study population was homogeneous and included healthy women aged 25–35 years from the same cultural background.

However, our sample size was small, the study design was cross-sectional, and because of ethical concerns, we could only obtain one measure of mammographic density, and therefore could not measure changes in density patterns across the menstrual cycle. Finally, all serum samples from each woman were not measured in the same batches, which could have potentially introduced some error. However, serum concentrations of estradiol and progesterone in premenopausal women are high, with lower CVs, and therefore there is generally less likelihood for measurement error compared with values in postmenopausal women. Each woman's salivary estradiol and progesterone, however, were assayed in the same batch. These similar results observed in salivary and serum concentrations suggested that the associations using serum values are valid. The salivary samples were only measured during one menstrual cycle, and we could not capture the intercycle variations (Chatterton *et al.*, 2005). However, among stable-weight women, marked changes in hormonal levels from cycle to cycle are not expected, thus lowering the required number of measured cycles. Even a single cycle per woman would be sufficient to provide adequate statistical power (Jasienska and Jasienski, 2008).

Conclusion

Our study provides novel data linking endogenous sex hormones throughout the menstrual cycle to percent mammographic density in particular. The present observations are biologically plausible, and may be of potential clinical interest. However, larger studies including estrogen and progesterone across the menstrual cycle in various populations are needed to define the clinical implications of these findings.

Acknowledgements

The authors thank the participants in the EBBA-I study. They are grateful to Gunn Kristin Knudsen, Heidi Jakobsen, Anna-Kirsti Kvitnes, Sissel Andersen, and Nils Bjurstam, and the Clinical Research Department and Center of Breast Imaging, University Hospital of North Norway, for professional assistance in the data collection. This work was supported by the Foundation for the Norwegian Health and Rehabilitation Organization (59010-2000, 59010-2001, 59010-2002); the Norwegian

Cancer Society (05087, TP 49 258); the Aakre Foundation (5695-2000, 5754-2002); the Northern Norway Regional Health Authority (SFP-563-06); the Norwegian research Council (213997); and a PhD grant from University of Tromsø.

Conflicts of interest

There are no conflicts of interest.

References

- Barrett ES, Thune I, Lipson SF, Furberg AS, Ellison PT (2013). A factor analysis approach to examining relationships among ovarian steroid concentrations, gonadotrophin concentrations and menstrual cycle length characteristics in healthy, cycling women. *Hum Reprod* **28**:801–811.
- Bellem A, Meiyappan S, Romans S, Einstein G (2011). Measuring estrogens and progestagens in humans: an overview of methods. *Gend Med* **8**:283–299.
- Bernstein L (2002). Epidemiology of endocrine-related risk factors for breast cancer. *J Mammary Gland Biol Neoplasia* **7**:3–15.
- Bjurstam N, Björneld L, Warwick J, Sala E, Duffy SW, Nyström L, *et al.* (2003). The Gothenburg breast screening trial. *Cancer* **97**:2387–2396.
- Boyd NF (2013). Mammographic density and risk of breast cancer. *Am Soc Clin Oncol Educ Book*. doi: 10.1200/EdBook_AM.2013.33.e57.
- Boyd NF, Stone J, Martin LJ, Jong R, Fishell E, Yaffe M, *et al.* (2002). The association of breast mitogens with mammographic densities. *Br J Cancer* **87**:876–882.
- Boyd NF, Guo H, Martin LJ, Sun L, Stone J, Fishell E, *et al.* (2007). Mammographic density and the risk and detection of breast cancer. *N Engl J Med* **356**:227–236.
- Boyd NF, Martin LJ, Yaffe MJ, Minkin S (2011). Mammographic density and breast cancer risk: current understanding and future prospects. *Breast Cancer Res* **13**:223.
- Brower V (2010). Homing in on mechanisms linking breast density to breast cancer risk. *J Natl Cancer Inst* **102**:843–845.
- Chatterton RT Jr, Mateo ET, Hou N, Rademaker AW, Acharya S, Jordan VC, Morrow M (2005). Characteristics of salivary profiles of oestradiol and progesterone in premenopausal women. *J Endocrinol* **186**:77–84.
- Cuzick J, Warwick J, Pinney E, Warren RM, Duffy SW (2004). Tamoxifen and breast density in women at increased risk of breast cancer. *J Natl Cancer Inst* **96**:621–628.
- Ellison PT, Lipson SF (1999). Salivary estradiol – a viable alternative? *Fertil Steril* **72**:951–952.
- Emaus A, Espetvedt S, Veierød MB, Ballard-Barbash R, Furberg AS, Ellison PT, *et al.* (2008a). 17-beta-estradiol in relation to age at menarche and adult obesity in premenopausal women. *Hum Reprod* **23**:919–927.
- Emaus A, Veierød MB, Furberg AS, Espetvedt S, Friedenreich C, Ellison PT, *et al.* (2008b). Physical activity, heart rate, metabolic profile, and estradiol in premenopausal women. *Med Sci Sports Exerc* **40**:1022–1030.
- Finstad SE, Emaus A, Potischman N, Barrett E, Furberg AS, Ellison PT, *et al.* (2009). Influence of birth weight and adult body composition on 17beta-estradiol levels in young women. *Cancer Causes Control* **20**:233–242.
- Furberg AS, Jasienska G, Bjurstam N, Torjesen PA, Emaus A, Lipson SF, *et al.* (2005). Metabolic and hormonal profiles: HDL cholesterol as a plausible biomarker of breast cancer risk. The Norwegian EBBA Study. *Cancer Epidemiol Biomarkers Prev* **14**:33–40.
- Gann PH, Giovanazzi S, Van Horn L, Branning A, Chatterton RT Jr (2001). Saliva as a medium for investigating intra- and interindividual differences in sex hormone levels in premenopausal women. *Cancer Epidemiol Biomarkers Prev* **10**:59–64.
- Gill JK, Maskarinec G, Pagano I, Kolonel LN (2006). The association of mammographic density with ductal carcinoma in situ of the breast: the Multiethnic Cohort. *Breast Cancer Res* **8**:R30.
- Greendale GA, Reboussin BA, Slone S, Wasilauskas C, Pike MC, Ursin G (2003). Postmenopausal hormone therapy and change in mammographic density. *J Natl Cancer Inst* **95**:30–37.
- Henderson BE, Ross RK, Pike MC, Casagrande JT (1982). Endogenous hormones as a major factor in human cancer. *Cancer Res* **42**:3232–3239.
- Iversen A, Thune I, McTiernan A, Emaus A, Finstad SE, Flote V, *et al.* (2011). Ovarian hormones and reproductive risk factors for breast cancer in premenopausal women: the Norwegian EBBA-I study. *Hum Reprod* **26**:1519–1529.
- Jasienska G, Jasienski M (2008). Interpopulation, interindividual, intercycle, and intracycle natural variation in progesterone levels: a quantitative assessment and implications for population studies. *Am J Hum Biol* **20**:35–42.
- Jasienska G, Ziolkiewicz A, Lipson SF, Thune I, Ellison PT (2006). High ponderal index at birth predicts high estradiol levels in adult women. *Am J Hum Biol* **18**:133–140.
- Johansson H, Gandini S, Bonanni B, Mariette F, Guerrieri-Gonzaga A, Serrano D, *et al.* (2008). Relationships between circulating hormone levels, mammographic percent density and breast cancer risk factors in postmenopausal women. *Breast Cancer Res Treat* **108**:57–67.
- Johansson H, Bonanni B, Gandini S, Guerrieri-Gonzaga A, Cazzaniga M, Serrano D, *et al.* (2013). Circulating hormones and breast cancer risk in premenopausal women: a randomized trial of low-dose tamoxifen and fenretinide. *Breast Cancer Res Treat* **142**:569–578.
- Key TJ, Appleby PN, Reeves GK, Roddam A, Dorgan JF, Longcope C, *et al.* (2003). Body mass index, serum sex hormones, and breast cancer risk in postmenopausal women. *J Natl Cancer Inst* **95**:1218–1226.
- Key TJ, Appleby PN, Reeves GK, Travis RC, Alberg AJ, Barricarte A, *et al.* (2013). Sex hormones and risk of breast cancer in premenopausal women: a collaborative reanalysis of individual participant data from seven prospective studies. *Lancet Oncol* **14**:1009–1019.
- Lipson SF, Ellison PT (1989). Development of protocols for the application of salivary steroid analysis to field conditions. *Am J Hum Biol* **1**:249–255.
- Lu Y, Bentley GR, Gann PH, Hodges KR, Chatterton RT (1999). Salivary estradiol and progesterone levels in conception and nonconception cycles in women: evaluation of a new assay for salivary estradiol. *Fertil Steril* **71**:863–868.
- Maskarinec G, Heak S, Morimoto Y, Custer L, Franke AA (2012). The relation of urinary estrogen metabolites with mammographic densities in premenopausal women. *Cancer Epidemiol* **36**:e310–e316.
- Matthews JN, Altman DG, Campbell MJ, Royston P (1990). Analysis of serial measurements in medical research. *BMJ* **300**:230–235.
- McTiernan A, Martin CF, Peck JD, Aragaki AK, Chlebowski RT, Pisano ED, *et al.* (2005). Estrogen-plus-progestin use and mammographic density in postmenopausal women: Women's Health Initiative randomized trial. *J Natl Cancer Inst* **97**:1366–1376.
- McTiernan A, Chlebowski RT, Martin C, Peck JD, Aragaki A, Pisano ED, *et al.* (2009). Conjugated equine estrogen influence on mammographic density in postmenopausal women in a substudy of the women's health initiative randomized trial. *J Clin Oncol* **27**:6135–6143.
- Morrow M, Chatterton RT Jr, Rademaker AW, Hou N, Jordan VC, Hendrick RE, Khan SA (2010). A prospective study of variability in mammographic density during the menstrual cycle. *Breast Cancer Res Treat* **121**:565–574.
- Noh JJ, Maskarinec G, Pagano I, Cheung LW, Stanczyk FZ (2006). Mammographic densities and circulating hormones: a cross-sectional study in premenopausal women. *Breast* **15**:20–28.
- Pettersson A, Hankinson SE, Willett WC, Lagiou P, Trichopoulos D, Tamimi RM (2011). Nondense mammographic area and risk of breast cancer. *Breast Cancer Res* **13**:R100.
- Pettersson A, Graff RE, Ursin G, Santos Silva ID, McCormack V, Baglietto L, *et al.* (2014). Mammographic density phenotypes and risk of breast cancer: a meta-analysis. *J Natl Cancer Inst* **106**. doi: 10.1093/jnci/dju078.
- Pike MC, Krailo MD, Henderson BE, Casagrande JT, Hoel DG (1983). 'Hormonal' risk factors, 'breast tissue age' and the age-incidence of breast cancer. *Nature* **303**:767–770.
- Pinto Pereira SM, McCormack VA, Hipwell JH, Record C, Wilkinson LS, Moss SM, *et al.* (2011). Localized fibroglandular tissue as a predictor of future tumor location within the breast. *Cancer Epidemiol Biomarkers Prev* **20**:1718–1725.
- Rauh C, Hack CC, Häberle L, Hein A, Engel A, Schrauder MG, *et al.* (2012). Percent mammographic density and dense area as risk factors for breast cancer. *Geburtshilfe Frauenheilkd* **72**:727–733.
- Schetter SE, Hartman TJ, Liao J, Richie JP, Prokopczyk B, DuBrock C, *et al.* (2014). Differential impact of body mass index on absolute and percent breast density: implications regarding their use as breast cancer risk biomarkers. *Breast Cancer Res Treat* **146**:355–363.
- Stone J, Ding J, Warren RM, Duffy SW, Hopper JL (2010). Using mammographic density to predict breast cancer risk: dense area or percentage dense area. *Breast Cancer Res* **12**:R97.
- Tamimi RM, Hankinson SE, Colditz GA, Byrne C (2005). Endogenous sex hormone levels and mammographic density among postmenopausal women. *Cancer Epidemiol Biomarkers Prev* **14** (Pt 1):2641–2647.
- Ursin G, Astrahan MA, Salane M, Parisky YR, Pearce JG, Daniels JR, *et al.* (1998). The detection of changes in mammographic densities. *Cancer Epidemiol Biomarkers Prev* **7**:43–47.

- Ursin G, Ma H, Wu AH, Bernstein L, Salane M, Parisky YR, *et al.* (2003). Mammographic density and breast cancer in three ethnic groups. *Cancer Epidemiol Biomarkers Prev* **12**:332–338.
- Ursin G, Hovanessian-Larsen L, Parisky YR, Pike MC, Wu AH (2005). Greatly increased occurrence of breast cancers in areas of mammographically dense tissue. *Breast Cancer Res* **7**:R605–R608.
- van Gils CH, Hendriks JH, Otten JD, Holland R, Verbeek AL (2000). Parity and mammographic breast density in relation to breast cancer risk: indication of interaction. *Eur J Cancer Prev* **9**:105–111.
- Walker K, Fletcher O, Johnson N, Coupland B, McCormack VA, Folkard E, *et al.* (2009). Premenopausal mammographic density in relation to cyclic variations in endogenous sex hormone levels, prolactin, and insulin-like growth factors. *Cancer Res* **69**:6490–6499.
- Yaghjian L, Colditz GA, Rosner B, Tamimi RM (2013). Mammographic breast density and subsequent risk of breast cancer in postmenopausal women according to the time since the mammogram. *Cancer Epidemiol Biomarkers Prev* **22**:1110–1117.
- Yong M, Atkinson C, Newton KM, Aiello Bowles EJ, Stanczyk FZ, Westerlind KC, *et al.* (2009). Associations between endogenous sex hormone levels and mammographic and bone densities in premenopausal women. *Cancer Causes Control* **20**:1039–1053.