

SCIENTIFIC REPORTS



OPEN

Exogenous N addition enhances the responses of gross primary productivity to individual precipitation events in a temperate grassland

Received: 20 November 2015

Accepted: 10 May 2016

Published: 06 June 2016

Qun Guo¹, Zhong-min Hu¹, Sheng-gong Li¹, Gui-rui Yu¹, Xiao-min Sun¹, Ling-hao Li², Nai-shen Liang³ & Wen-ming Bai²

Predicted future shifts in the magnitude and frequency (larger but fewer) of precipitation events and enhanced nitrogen (N) deposition may interact to affect grassland productivity, but the effects of N enrichment on the productivity response to individual precipitation events remain unclear. In this study, we quantified the effects of N addition on the response patterns of gross primary productivity (GPP) to individual precipitation events of different sizes (P_{size}) in a temperate grassland in China. The results showed that N enrichment significantly increased the time-integrated amount of GPP in response to an individual precipitation event (GPP_{total}), and the N-induced stimulation of GPP increased with increasing P_{size} . N enrichment rarely affected the duration of the GPP response, but it significantly stimulated the maximum absolute GPP response. Higher foliar N content might play an important role in the N-induced stimulation of GPP. GPP_{total} in both the N-addition and control treatments increased linearly with P_{size} with similar P_{size} intercepts (approximately 5 mm, indicating a similar lower P_{size} threshold to stimulate the GPP response) but had a steeper slope under N addition. Our work indicates that the projected larger precipitation events will stimulate grassland productivity, and this stimulation might be amplified by increasing N deposition.

Comprising approximately 40% of the global land cover¹, grasslands are extraordinarily sensitive to the alteration of precipitation regimes in the context of global climate change². Grassland productivity is profoundly constrained by precipitation^{2–4}, and this constraint is characterized by features such as the total amount of precipitation and its timing, seasonal distribution, frequency, the dry spells between events, etc^{5–8}. As an important aspect of altered precipitation regimes, the size of individual precipitation events (P_{size}) has been predicted to dramatically change in the future; e.g., there will be a trend toward less frequent but more intense precipitation events in arid and semiarid regions^{9,10}. Additionally, lines of evidence have indicated that the increase in atmospheric nitrogen (N) deposition due to human activities is having serious impacts on ecosystems^{11–13}. Therefore, elucidating the interactive effects of altered precipitation regimes and N enrichment on grassland productivity is particularly important for predicting the response of ecosystems to future global climate change.

Recent studies have demonstrated that, globally, ecosystem productivity is co-limited by precipitation and N, and it is usually stimulated by the addition of water or N^{14–16}. However, whether N enrichment could enhance productivity response to precipitation remains controversial. Studies in temperate grasslands have found that N enrichment has little effect on productivity in response to water addition^{17,18}, and other studies have reported that the stimulation of productivity with N addition could depend on external environment conditions^{15,16,19,20}. To

¹Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China. ²State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China. ³Global Carbon Cycle Research Section Center for Global Environmental Research (CGER), National Institute for Environmental Studies (NIES), Tsukuba 305-8506, Japan. Correspondence and requests for materials should be addressed to Z.-m.H. (email: huzm@igsnr.ac.cn) or S.-g.L. (email: lisg@igsnr.ac.cn)




Figure 1. Seasonal gross primary productivity (GPP) and soil water content (SWC) dynamics of the control and N-addition treatments in 2012 (a) and 2013 (b) in a temperate grassland in Inner Mongolia, China. The six selected precipitation events are indicated by arrows above the precipitation event bars. DOY: day of year; GPP_N : GPP of the N-addition treatment; GPP_{CK} : GPP of the control; GPP_{Eddy} : GPP derived from the eddy covariance system.

date, most studies have generally focused on the effects of N addition on productivity in response to total annual precipitation (e.g., ref. 16), but availability of soil moisture is intermittent in arid and semi-arid areas. Therefore, to know how pulses of moisture following individual precipitation events affect productivity and how N addition regulate the productivity response to individual precipitation events is essential to an advanced understanding of the interactive effects between precipitation and N enrichment.

Several parameters are used in the literature to characterize the productivity-response pattern to various sizes of individual precipitation events, such as the maximum productivity response, the duration of the response, and the time-integrated amount of productivity response to a given individual precipitation event^{21,22}. Furthermore, the productivity response is constrained by two precipitation event threshold sizes, i.e., the lower threshold (R^L), which triggers the productivity response, and the upper threshold (R^U), above which the response levels off^{23,24}. Previous studies have shown that N enrichment deeply influences vegetation properties (e.g., photosynthesis or plant growth rate) and soil moisture^{13,18}, which are also important to the determination of the above parameters^{21,22,24}. However, the extent to which N enrichment affects these parameters and the effects that changed parameters will exert on the patterns of productivity in response to individual precipitation events remains unclear.

In this study, through experiments that manipulated N availability, we quantified the effects of N addition on the responses of gross primary productivity (GPP) to different sizes of individual precipitation events in a temperate steppe in Inner Mongolia, China. A multichannel automated measurement system was employed to estimate GPP at high temporal resolution, and our main objectives were as follows. First, we needed to clarify the extent to which N enrichment affected the time-integrated amount of GPP response to individual precipitation events and how the N-induced GPP response varied with the size of individual precipitation events. Second, we quantified the effects of N addition on the parameters that characterize the patterns of the GPP response to individual precipitation events, i.e., its maximum potential and duration, the lower P_{size} threshold initiating the GPP response and the upper threshold that achieves the maximum response (saturation). Finally, we determined how these parameters impact the total GPP response and the underlying mechanisms.

Results

Seasonal gross primary productivity and soil moisture dynamics. We found that the GPP measured by the multichannel automated measurement system in this study was highly consistent with that from an eddy covariance system (GPP_{Eddy}), indicating that our GPP-estimation methods were reliable (Fig. 1). The control GPP (GPP_{CK} in terms of mean \pm standard errors) during the two measurement periods (from day of year (DOY) 137 to 262 in 2012 and from DOY 157 to 249 in 2013) was $303.2 \pm 25.5 \text{ g C m}^{-2}$ ($n = 3$) in 2012 and $289.8 \pm 5.7 \text{ g C m}^{-2}$ ($n = 3$) in 2013. The GPP values of the N-addition treatments (GPP_N in terms of mean \pm standard errors) were $395.94 \pm 32.9 \text{ g C m}^{-2}$ ($n = 3$) in 2012 and $365.75 \pm 40.1 \text{ g C m}^{-2}$ ($n = 3$) in 2013, which were significantly higher (30.6% and 26.2%, respectively) than those of the control ($n = 3$, $p < 0.05$).

Seasonally, both GPP_{CK} and GPP_N were closely related to the increase in soil moisture, but GPP_N was higher than GPP_{CK} on most days (Fig. 1). Generally, both GPP_{CK} and GPP_N first synchronously increased and then subsequently decreased after precipitation events. Although GPP_{CK} and GPP_N were more similar before rainfall,




Figure 2. Daily GPP and SWC dynamics after different sizes of precipitation events in a temperate grassland in Inner Mongolia, China. Rainfalls occurring over consecutive days were considered to be one precipitation event (e.g., a 21.9-mm precipitation event was composed of rainfalls over three consecutive days). The y-axes of all of the subpanels are scaled the same for GPP, SWC, and precipitation.

GPP_N was significantly higher than GPP_{CK} after rainfall, and this increase in GPP after rainfall was dependent on the size of the precipitation events (Fig. 2).

Effects of N addition on the GPP responses to different sizes of precipitation events. The duration of the GPP response to an individual precipitation event (τ_R), the maximum absolute GPP increase induced by the event (GPP_{max}), and the time-integrated amount of GPP increase during the response period (GPP_{total}) all significantly linearly increased with increasing P_{size} in both the control and N-addition treatments ($n = 6$, $p < 0.01$, R^2 ranged from 0.86 to 0.94) (Fig. 3a–c). When P_{size} ranged from 4.7 to 40.2 mm, GPP_{total} increased from 2.0 to 53.8 g C m⁻² on the control and from 2.4 to 66.6 g C m⁻² under the N-addition treatments. The GPP_{max} varied from 1.4 to 6.4 g C m⁻² d⁻¹ for the control and 0.9 to 8.7 g C m⁻² d⁻¹ for N-addition treatments, whereas the τ_R values were much more similar in the control and N-addition treatments, ranging from 5 to 18 d and 5 to 17 d, respectively.

Except for the τ_R - P_{size} relationship (whose slopes were both 0.33), the slopes of the linear regressions of both GPP_{total} - P_{size} and GPP_{max} - P_{size} were steeper under the N-addition treatment than the control (Fig. 3a–c). The slope of the GPP_{max} - P_{size} relationship in the N-addition treatment (0.21, ranging from 0.14 to 0.28) tended to be larger than that of the control (0.14, ranging from 0.09 to 0.21) ($n = 6$, $p = 0.09$). Although the slope of the GPP_{total} - P_{size} regression (1.72, ranging from 1.32 to 2.36) did not significantly increase due to N addition ($n = 6$, $p = 0.22$), it was still much larger than that of the control (1.39, ranging from 1.06 to 1.93). However, the lower precipitation event threshold size (R^L) required to induce a GPP response, i.e., the P_{size} intercept of the GPP_{total} - P_{size} regression, was much more similar in both the N-addition treatments (4.92 ± 1.85 mm) and the control (5.23 ± 1.88 mm) ($p > 0.1$). The upper precipitation event threshold (R^U), at which GPP response leveled off, was not observed in this study.




Figure 3. The variations in the time-integrated amount of the GPP response during the response period (GPP_{total}) (a), the duration of the GPP response to a precipitation event (τ_R) (b), and the maximum absolute increase in GPP induced by the event (GPP_{max}) (c) along with the size of individual precipitation events (P_{size}) in a temperate grassland in Inner Mongolia.




Figure 4. Variations of the differences in GPP_{total} (ΔGPP_{total}) and GPP_{max} (ΔGPP_{max}) between the control and N-addition treatments with the size of precipitation events (P_{size}) in a temperate grassland in Inner Mongolia.

The difference in GPP_{total} between the N-addition treatments and the control increased significantly with increasing P_{size} (Fig. 4a), while the difference of GPP_{max} also increased (Fig. 4b). There was no remarkable difference in τ_R ($n = 6$, $p > 0.1$), indicating that the N-induced stimulation of GPP_{total} is mainly due to the increased GPP_{max} .

Leaf area index and foliar N content. The leaf area index (LAI) in 2012, 2013, and 2014 exhibited a hump-shaped pattern of seasonal variation (Fig. 5a). LAI was very similar among years in the early growing season (from DOY130 to 190) but differed considerably throughout the rest of the growing season ($n = 13$, $p = 0.04$), being much lower in 2014 due to a long dry spell (21 days) (Fig. 5a). Lacking vegetation cover data for each treatment in 2012 and 2013, we used the vegetation cover data obtained in 2014 to approximate the status of the vegetation cover in 2012 and 2013 based on the LAI similarity in the early growing season (i.e., the study period),




Figure 5. Leaf area index (LAI) over three years (a) and vegetation cover in 2014 (b). The panel inset in (b) shows the difference in vegetation cover (Δ cover) between the N-addition (N, empty circle) and the control (CK, solid circle) treatments in 2014. LAI was derived from normalized difference vegetation index (NDVI) data from the Moderate Resolution Imaging Spectroradiometer (8-d time resolution and 250-m spatial resolution).




Figure 6. Comparison of the foliar N content among different years and treatments. An asterisk above the error bars indicates a significant difference between treatments ($p < 0.001$). Values showed in the figure are averaged with standard errors from 8 plots of each treatment.

and then we evaluated the vegetation cover differences between the treatments in 2012 and 2013. The vegetation cover increased at first and reached its peak in mid-August (Fig. 5b). There was no difference in vegetation cover between the N-addition treatments and the control during the early growing season ($n = 8$, $p = 0.91$), but a larger difference appeared from the middle to the end of the growing season ($n = 5$, $p = 0.41$).

No significant (the control) or only slight (the N-addition treatments) inter-annual variations in foliar N content were observed across the three years (Fig. 6). Foliar N content in the N-addition treatments was higher than that of the control ($n = 8$, $p < 0.001$); i.e., foliar N content in the N-addition treatments was 43%, 30%, and 34% higher than that in the control in 2012, 2013, and 2014, respectively (Fig. 6).

Discussion

The time-integrated amount of the GPP response to an individual precipitation event (GPP_{total}) greatly increased due to N addition in the temperate steppe of Inner Mongolia, and this stimulation increased with the size of the precipitation event. Previous studies have shown that exogenous N addition boosts the productivity response to the amount of annual precipitation^{15,16}, and our observations provide evidence that N-induced stimulation of productivity could occur in response to an individual precipitation event. In arid and semi-arid ecosystems, such as the steppe in this study, productivity is primarily limited by water availability^{25–27}, so ecosystems are only limited by nitrogen when the soil water content reaches a critical level²⁸. Thus, we may postulate that N addition-induced stimulation of the GPP response to individual precipitation events may be more likely when there is a relatively long-lasting period of ample soil moisture. Soil moisture pulses due to larger precipitation events that percolate deep into the soil profile may serve as an important water source that can be maintained for much longer periods^{29,30}, leading to more benefits of plant growth (GPP) from greater N release, so the stimulation of GPP_{total}

due to N addition increased with P_{size} . It has been predicted that precipitation events will tend to be larger in size but fewer in number in the context of global climate change¹⁰, which may alleviate the water limitation imposed on vegetation, especially in arid and semi-arid environments, and thus promote plant growth (GPP)^{29–31}. This stimulation of productivity under altered precipitation regimes will likely be further augmented in the context of increasing N deposition in the future.

For any individual precipitation event, the increase in GPP_{total} by N enrichment could be primarily ascribed to the enhancement of the maximum absolute GPP response induced by the event (GPP_{max}) rather than the duration of the GPP response (τ_R). Under the given soil moisture conditions due to an individual event, N enrichment will stimulate GPP_{max} in two ways. Firstly, the photosynthetic capacity may be enhanced by increased foliar N content, and previous studies have shown that the increase in foliar N content resulting from N enrichment is one of the main reasons for the enhancement of photosynthesis^{32–35}. Secondly, N enrichment relieves N limitation in ecosystems and subsequently increases plant growth and vegetation cover¹⁸, thus facilitating higher carbon sequestration per unit area. Our results showed that N addition did boost foliar N content relative to the control, whereas leaf area index was much more similar in both the control and N-addition treatments. Due to the relatively low temperatures during the study period (i.e., the early growing season), plants grew slowly, so a significant increase in vegetation cover due to N addition was not observed. However, obviously higher vegetation cover was observed in the N-addition treatments than in the control during the warmer mid-late growing season, indicating that an increase in vegetation cover might also contribute to stimulating GPP_{max}. Previous studies have revealed that N enrichment could affect soil moisture by impacting vegetation cover¹⁸, and it has been indicated that τ_R is constrained by the duration of higher soil moisture²². Therefore, N enrichment could affect τ_R by modulating soil moisture resulting from rainfall. Nevertheless, there was no striking difference in vegetation cover between the control and N-addition treatments, leading to a similar τ_R .

N enrichment had minor impacts on the lower P_{size} threshold (R^L) that triggers a detectable GPP-response. Previous studies have demonstrated that the productivity response starts at an approximately 5-mm P_{size} threshold in typical steppes of Inner Mongolia^{36,37}, which is similar to the observation in this study. As plant growth is primarily constrained by water availability, it appears that the steppe is more limited by soil moisture rather than N availability when the P_{size} is relatively small, e.g., less than 5 mm, implying that N enrichment has a minor or negligible impact on R^L . When the P_{size} was larger than R^L , GPP_{total} increased linearly with P_{size} in both treatments because of the increasing relief from water limitation and hence a greater plant response²². Previous studies have proposed that a plant's physiological activity cannot exceed some maximum value due to functional or structural constraints, and furthermore, soil moisture would be saturated after precipitation events of a certain size²⁴. Therefore, there should be an upper P_{size} threshold (R^U), above which no additional GPP response occurs. R^U was not observed in this study because such sizes of individual precipitation events, especially with a P_{size} larger than 50 mm, were not observed, but R^U deserves further examination in the future.

In conclusion, based on a field experiment consisting of N addition and continuous measurements of net ecosystem CO₂ exchange with a multichannel automated measurement system, we quantified the effects of N addition on the responses of GPP to individual precipitation events in a temperate steppe in Inner Mongolia, China. Our results showed that N enrichment increased the time-integrated amount of the GPP response to an individual precipitation event through the enhanced maximum absolute GPP response after the event, and the stimulation of GPP induced by N addition increased with the size of the precipitation events (P_{size}). N enrichment rarely affected the lower P_{size} threshold for inducing a detectable GPP-response. Our work has important implications for theoretical modeling to obtain an advanced understanding of the response of productivity to different sizes of precipitation events and increased N deposition in grassland ecosystems.

Methods

Study site. This study was conducted at the Duolun Restoration Ecology Research Station of the Institute of Botany, Chinese Academy of Sciences, which is located in a typical steppe in Inner Mongolia, China (42°02'N, 116°117'E; 1,324 m a.s.l.). The dominant species in this grassland include *Stipa krylovii* Roshev., *Artemisia frigid* Willd., *Potentilla acaulis* L., *Cleistogenes squarrosa* (Trin.) Keng., *Allium bidentatum* Fisch. ex Prokh., and *Agropyron cristatum* (L.) Gaertn. Mean annual precipitation (MAP) is approximately 385.5 mm with the majority (over 80%) falling in the growing season from May to September. Annual mean temperature is 2.1 °C with the highest monthly mean occurring in July (18.9 °C) and the lowest in January (−17.5 °C). The soil type is chestnut; the soil pH value is 7.12; and the mass-based nitrogen and phosphorus content is 0.17% and 0.28%, respectively. The length of the growing season is approximately 150 d, and the average plant height is 0.4 m. During the two years of the experimental period, both the amount of precipitation (285 mm in 2012 and 262 mm in 2013) and the median precipitation event size (6.3 mm in 2012 and 7.7 mm in 2013) were similar. The aboveground biomass estimated by the harvest method was 151.3 g m^{−2} in 2012 and 143.6 g m^{−2} in 2013, respectively, indicating that the climate and vegetation cover did not differ greatly between 2012 and 2013.

Design of the manipulative experiments. The experiment employed a complete randomized block design with two treatments, including a control (0 g N m^{−2} yr^{−1}) and one N-addition level (10 g N m^{−2} yr^{−1}), the rate of which is similar to the critical threshold for the N-induced increase in aboveground biomass³⁸. A total of four parallel blocks (each block consisted of 4 3-m × 4-m plots) were established. Two of the four plots in each block were set as control plots, and the remaining two plots were used for the N addition treatment. There were buffer zones of at least 3 m between blocks and intervals of 2 m between the plots within each block. The manipulative N-addition experiments began in 2012 and have continued to this day.

Urea (CO(NH₂)₂) was scattered on the N-addition plots twice before the first rainfall of each month during the growing season, from May to June, each year (2012 and 2013). If there were no rains in the first half of each month, the urea solution would be added to the N-addition plots, and an equal amount of water would be added

to the control plots. However, during our two-year experiment period, this situation did not occur. N addition was performed on May 11 and June 3 in 2012 and May 3 and June 5 in 2013.

Measurements of gross primary productivity and meteorological variables. We measured net ecosystem CO₂ exchange between the atmosphere and grassland with a multichannel automated measurement system developed by Dr. Liang at the National Institute for Environmental Studies of Japan. The system is comprised of 12 transparent chambers and a control box, and six chambers were used in this study, i.e., three for the control and three for the N-addition treatments. The main components of the control box are an infrared gas analyzer (IRGA, Li-840, Li-Cor Inc., Lincoln, USA) and a data logger (CR3000, Campbell Scientific, Inc., Logan, UT, USA), and detailed diagram of this multichannel automated measurement system can be found in ref. 22. During the measurements, 12 chambers are sequentially closed by a homemade relay board controlled by the data logger and then the air in the closed chamber is circulated through the IRGA by a microdiaphragm pump (CM-50, Enomoto Ltd., Tokyo, Japan). The sampling period for each chamber is 150 s (i.e., it takes 0.5 h to measure all of the 12 chambers), and the data logger monitors the CO₂ concentration output signals from the IRGA at a rate of 1 Hz and records their averages at 10-s intervals. To remove the interference from impure air and erratic air pressure, we excluded the data from the first 10 s and the last 10 s. We obtained one measurement from each chamber in half an hour, and the net ecosystem exchange of CO₂ (NEE) was calculated from the CO₂ exchange rate during the measurement periods (i.e., 130 s):

$$NEE = (V \times P(1 - W)/(R \times S \times T)) \times (u_c/u_t), \quad (1)$$

where V is the volume of the chamber (m³); P is the air pressure (Pa); W is the water vapor mole fraction; R is the universal gas constant (8.314 Pa m³ mol⁻¹ K⁻¹); S is the base area of the chamber (m²); T is the air temperature in the chamber (K); and u_c/u_t is the rate of increase in the CO₂ mole fraction (mmol mol⁻¹ s⁻¹) in the chamber calculated by the least squares method.

Because nighttime NEE only represents ecosystem respiration (R_e), we used the relationship between nighttime NEE and soil temperature to estimate daytime R_e (R_{ed})³⁹. Furthermore, soil water content (SWC) was also considered when evaluating the R_{ed} because our study was carried out in a dry environment⁴⁰. In this study, R_{ed} was finally estimated as follows:

$$R_{ed} = R_{e,ref} \times (b_1 + b_2SWC) \times ((T_s - T_{ref})/10), \quad (2)$$

where $R_{e,ref}$ is the ecosystem respiration at the reference temperature (T_{ref} , 10°C) and optimal soil moisture; T_s is the soil temperature; b_1 and b_2 are constants evaluated from the relationship between nighttime R_e and temperature and SWC.

Finally, GPP was indirectly estimated as follows:

$$GPP = R_{ed} - NEE, \quad (3)$$

where the negative NEE denotes a carbon flux into the steppe, and a positive NEE denotes the reverse. Because our GPP estimation methods are similar to those from an eddy covariance system, which we also have in the same field experiment, we compared the results between the two systems to evaluate the uncertainty of the GPP estimation in this study.

There are two thermocouples in each chamber to measure the air temperature and the soil temperature (at a depth of 5 cm), which are recorded by a data logger with the same time resolution as that of the CO₂ flux. Volumetric soil water content at the depths of 5 cm, 20 cm, and 40 cm were measured and calculated simultaneously at half-hourly and daily intervals with a meteorological measurement system near the treatments. Finally, we acquired daily GPP and soil water content data from DOY 137 to DOY 262 in 2012 and from DOY 157 to DOY 249 in 2013. The missing growing season data were mostly due to the rough conditions when the experiments must be suspended or the failure of the multichannel automated measurement system.

Leaf samples of five dominant species in each plot were collected in mid-August to determine foliar N content, and the N content of the five species was averaged for each plot. Vegetation cover for the two treatments was estimated by means of SamplePoint software⁴¹ from the quadrat photos taken every week in 2014. Vegetation cover data were missing because the quadrat photos were not available in 2012 and 2013.

Data analysis. In the present study, an individual precipitation event was defined as a series of one or more consecutive days of rainfall followed by at least one day without rainfall.

The GPP observed from 3 transparent chambers was averaged to evaluate the GPP for each treatment. We denoted GPP prior to the precipitation event as GPP_{base} , which was averaged over the three days before the event to better represent pre-rainfall GPP. Then, we evaluated the GPP response to an individual precipitation event in both the control and N-addition treatments in terms of its duration, maximum, and time-integrated amount. (1) The duration of the GPP response (τ_R) was calculated as the number of days required for GPP to return to its pre-precipitation event level, i.e., the GPP_{base} . If the daily GPP did not reach the GPP_{base} due to the occurrence of the next precipitation event, we used a linear extrapolation method for the estimation. (2) The maximum absolute GPP increase (GPP_{max}) induced by the precipitation event was calculated as the maximal difference between daily GPP after rainfall and GPP_{base} . (3) The time-integrated amount of the GPP response during the response period (duration) (GPP_{total}) was calculated as

$$GPP_{total} = \sum_{i=1}^{\tau_R} (GPP_i - GPP_{base}). \quad (4)$$

where subscript i is the i th day after the occurrence of a precipitation event.

Based on the regression between GPP_{total} and P_{size} , we also evaluated the lower and upper event size threshold that stimulated a detectable GPP-response or at which no additional gain in the GPP response would result from the increasing size of the event, e.g., the intercept of a linear function (lower threshold) or the critical thresholds of a logistic curve (lower and upper thresholds).

The GPP response to an individual precipitation event may be confounded by other factors, e.g., antecedent soil water prior to the precipitation event⁴², leaf area index, and photosynthetic capacity at different growth stages^{21,24,43}. To precisely evaluate the GPP response to an individual precipitation event, several criteria must be satisfied. First, antecedent SWC should be very low (mostly lower than the average SWC, i.e., $0.088 \text{ cm}^3 \text{ cm}^{-3}$ in this study), and the inter-pulse period should be longer than 5 days to minimize the effect of the previous precipitation events, which may shadow or confound the GPP response to the current precipitation event of interest. Second, we normalized the GPP-response (i.e., GPP_{max} , GPP_{total} , but not τ_R because τ_R is mostly affected by the duration of SWC) according to LAI. LAI was evaluated by normalized difference vegetation index (NDVI) data with models developed by ref. 44 ($LAI = 0.106e^{4.064NDVI}$, $R^2 = 0.94$). NDVI data were from the Moderate Resolution Imaging Spectroradiometer (MODIS) with 8-d time resolution and 250-m spatial resolution (<http://daac.ornl.gov/MODIS/modis.html>). Finally, the GPP response to a precipitation event might vary at different growth stages because of varying photosynthetic capacities (assuming a stronger response in the middle of the growing season than early or late in the growing season), and for the purpose of comparison, we chose the neighboring precipitation events to minimize the possible effects of vegetation phenology.

In total, six independent precipitation events from DOY 155 to 188 were selected for the related analysis (7.7, 40.2, and 21.9-mm precipitation events on DOY 155, 164, and 176 in 2012 and 4.7, 19.6, and 11-mm precipitation events on DOY 159, 168, and 188 in 2013). Hence, our results only reflected the GPP response during the early growing season.

Statistical analysis. Normality of data was tested by the Kolmogorov-Smirnov test. One-way ANOVA was employed to estimate the differences in the variables of interest (e.g., GPP, GPP_{total} , LAI, vegetation cover, foliar N content, etc.) between the two treatments or the different years. Comparison of slope differences between the two treatments was performed using the R software package (version 3.1.2, an open source software program), and the slope comparison was implemented by the “smart” package with “slope.com” command. A *t*-test was used to estimate whether these differences were significant, and the significance level was set at 0.05.

References

1. Reynolds, J. F. *et al.* Global desertification: Building a science for dryland development. *Science* **316**, 847–851 (2007).
2. Knapp, A. K. & Smith, M. D. Variation among biomes in temporal dynamics of aboveground primary production. *Science* **291**, 481–484 (2001).
3. Huxman, T. E. *et al.* Convergence across biomes to a common rain-use efficiency. *Nature* **429**, 651–654 (2004).
4. Michaletz, S. T., Cheng, D. L., Kerkhoff, A. J. & Enquist, B. J. Convergence of terrestrial plant production across global climate gradients. *Nature* **512**, 39–43 (2014).
5. Brookshire, E. N. J. & Weaver, T. Long-term decline in grassland productivity driven by increasing dryness. *Nat Commun* **6**, 1–7 (2015).
6. Craine, J. M. *et al.* Timing of climate variability and grassland productivity. *P Natl Acad Sci USA* **109**, 3401–3405 (2012).
7. Hovenden, M. J., Newton, P. C. & Wills, K. E. Seasonal not annual rainfall determines grassland biomass response to carbon dioxide. *Nature* **511**, 583–586 (2014).
8. Knapp, A. K. *et al.* Rainfall variability, carbon cycling, and plant species diversity in a mesic grassland. *Science* **298**, 2201–2205 (2002).
9. Easterling, D. R. *et al.* Climate extremes: Observations, modeling, and impacts. *Science* **289**, 2068–2074 (2000).
10. IPCC. *Climate Change 2013: The Physical Science Basis* (eds. Stocker, T. F. *et al.*) Ch. 12, 1076–1087 (Cambridge University Press, 2013).
11. Xia, J. Y., Niu, S. L. & Wan, S. Q. Response of ecosystem carbon exchange to warming and nitrogen addition during two hydrologically contrasting growing seasons in a temperate steppe. *Glob Change Biol* **15**, 1544–1556 (2009).
12. Humbert, J. Y., Dwyer, J. M., Andrey, A. & Arlettaz, R. Impacts of nitrogen addition on plant biodiversity in mountain grasslands depend on dose, application duration and climate: a systematic review. *Glob Change Biol* **22**, 110–120 (2016).
13. Field, C. & Mooney, H. A. *The photosynthesis–nitrogen relationship in wild plants* (ed. Givnish, T. J.) 25–55 (Cambridge University Press, 1986).
14. Austin, A. T. *et al.* Water pulses and biogeochemical cycles in arid and semiarid ecosystems. *Oecologia* **141**, 221–235 (2004).
15. Harpole, W. S., Potts, D. L. & Suding, K. N. Ecosystem responses to water and nitrogen amendment in a California grassland. *Glob Change Biol* **13**, 2341–2348 (2007).
16. Hooper, D. U. & Johnson, L. Nitrogen limitation in dryland ecosystems Responses to geographical and temporal variation in precipitation. *Biogeochemistry* **46**, 247–293 (1999).
17. Lu, X. T., Dijkstra, F. A., Kong, D., Wang, Z. W. & Han, X. G. Plant nitrogen uptake drives responses of productivity to nitrogen and water addition in a grassland. *Sci Rep-UK* **4**, doi: 10.1038/srep04817 (2014).
18. Niu, S. L. *et al.* Non-Additive Effects of Water and Nitrogen Addition on Ecosystem Carbon Exchange in a Temperate Steppe. *Ecosystems* **12**, 915–926 (2009).
19. Lee, M., Manning, P., Rist, J., Power, S. A. & Marsh, C. A global comparison of grassland biomass responses to CO₂ and nitrogen enrichment. *Philos Trans R Soc Lond B Biol Sci* **365**, 2047–2056 (2010).
20. Kinugasa, T., Tsunekawa, A. & Shinoda, M. Increasing nitrogen deposition enhances post-drought recovery of grassland productivity in the Mongolian steppe. *Oecologia* **170**, 857–865 (2012).
21. Huxman, T. E. *et al.* Precipitation pulses and carbon fluxes in semiarid and arid ecosystems. *Oecologia* **141**, 254–268 (2004).
22. Guo, Q. *et al.* Responses of gross primary productivity to different sizes of precipitation events in a temperate grassland ecosystem in Inner Mongolia, China. *J Arid Land* **8**, 36–46 (2016).
23. Noy-Meir, I. Desert ecosystems: environment and producers. *Annu Rev Ecol Evol Syst* **4**, 23–51 (1973).
24. Ogle, K. & Reynolds, J. F. Plant responses to precipitation in desert ecosystems: integrating functional types, pulses, thresholds, and delays. *Oecologia* **141**, 282–294 (2004).
25. Hu, Z. M., Fan, J. W., Zhong, H. P. & Yu, G. R. Spatiotemporal dynamics of aboveground primary productivity along a precipitation gradient in Chinese temperate grassland. *Sci China Ser D* **50**, 754–764 (2007).
26. Hu, Z. M. *et al.* Precipitation-use efficiency along a 4500-km grassland transect. *Global Ecol Biogeogr* **19**, 842–851 (2010).

27. Bai, Y. F. *et al.* Primary production and rain use efficiency across a precipitation gradient on the mongolia plateau. *Ecology* **89**, 2140–2153 (2008).
28. Yahdjian, L., Gherardi, L. & Sala, O. E. Nitrogen limitation in arid-subhumid ecosystems: A meta-analysis of fertilization studies. *J Arid Environ* **75**, 675–680 (2011).
29. Knapp, A. K. *et al.* Consequences of more extreme precipitation regimes for terrestrial ecosystems. *BioScience* **58**, 811–821 (2008).
30. Guo, Q. *et al.* Contrasting responses of gross primary productivity to precipitation events in a water-limited and a temperature-limited grassland ecosystem. *Agr Forest Meteorol* **214–215**, 169–177 (2015).
31. Thomey, M. L. *et al.* Effect of precipitation variability on net primary production and soil respiration in a Chihuahuan Desert grassland. *Glob Change Biol* **17**, 1505–1515 (2011).
32. Bai, X. F., Cheng, J. H., Zheng, S. X., Zhan, S. X. & Bai, Y. F. Ecophysiological responses of *Leymus chinensis* to nitrogen and phosphorus additions in a typical steppe. *J Plant Ecol* **38**, 103–115 (2014).
33. Hikosaka, K. Interspecific difference in the photosynthesis-nitrogen relationship: patterns, physiological causes, and ecological importance. *J Plant Res* **117**, 481–494 (2004).
34. Mao, W. *et al.* Biomass allocation response of species with different life history strategies to nitrogen and water addition in sandy grassland in Inner Mongolia. *J Plant Ecol* **38**, 125–133 (2014).
35. Takashima, T., Hikosaka, K. & Hirose, T. Photosynthesis or persistence: nitrogen allocation in leaves of evergreen and deciduous Quercus species. *Plant Cell Environ* **27**, 1047–1054 (2004).
36. Hao, Y. B., Wang, Y. F., Mei, X. R. & Cui, X. Y. The response of ecosystem CO₂ exchange to small precipitation pulses over a temperate steppe. *Plant Ecol* **209**, 335–347 (2010).
37. Zhang, B. *et al.* Effects of rainfall amount and frequency on vegetation growth in a Tibetan alpine meadow. *Climatic Change* **118**, 197–212 (2013).
38. Bai, Y. F. *et al.* Tradeoffs and thresholds in the effects of nitrogen addition on biodiversity and ecosystem functioning: evidence from inner Mongolia Grasslands. *Glob Change Biol* **16**, 358–372 (2010).
39. Lloyd, J. & Taylor, J. A. On the temperature-dependence of soil respiration. *Funct Ecol* **8**, 315–323 (1994).
40. Reichstein, M. *et al.* Ecosystem respiration in two Mediterranean evergreen Holm Oak forests: drought effects and decomposition dynamics. *Funct Ecol* **16**, 27–39 (2002).
41. Booth, D. T., Cox, S. E. & Berryman, R. D. Point sampling digital imagery with ‘SamplePoint’. *Environ Monit Assess* **123**, 97–108 (2006).
42. Potts, D. L. *et al.* Antecedent moisture and seasonal precipitation influence the response of canopy-scale carbon and water exchange to rainfall pulses in a semi-arid grassland. *New Phytol* **17**, 849–860 (2006).
43. Reynolds, J. F., Kemp, P. R., Ogle, K. & Fernandez, R. J. Modifying the ‘pulse-reserve’ paradigm for deserts of North America: precipitation pulses, soil water, and plant responses. *Oecologia* **141**, 194–210 (2004).
44. Hu, Z. M. *et al.* Partitioning of evapotranspiration and its controls in four grassland ecosystems: Application of a two-source model. *Agr Forest Meteorol* **149**, 1410–1420 (2009).

Acknowledgements

This study was jointly supported by the National Natural Science Foundation of China (31400425, 31570437, 41301043, 31420103917), Youth Innovation Promotion Association, CAS (2015037), the Funding for talented young scientists of IGSNRR (2013RC203), and National Key Project of Scientific and Technical Supporting Program (2013BAC03B03). We would like to thank Dr. Meng-xun Zhu for her help in field observation and data processing.

Author Contributions

Z.-m.H., Q.G. and S.-g.L. designed the study, Z.-m.H. and Q.G. compiled the data and produced the results. Z.-m.H., Q.G., S.-g.L., G.-r.Y. and X.-m.S. designed the field experiments, and L.-h.L., N.-s.L. and W.-m.B. provided supplementary data. All of the authors contributed to the interpretation of the results and the writing.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Guo, Q. *et al.* Exogenous N addition enhances the responses of gross primary productivity to individual precipitation events in a temperate grassland. *Sci. Rep.* **6**, 26901; doi: 10.1038/srep26901 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>