

Does the Method and Timing of Intravenous Ketamine Administration Affect Postoperative Morphine Requirement After Major Abdominal Surgery?

Feryal Biçer, Zeynep Eti, Kemal Tolga Saraçoğlu, Koray Altun, Fevzi Yılmaz Göğüş
Department of Anaesthesiology, Marmara University Faculty of Medicine, İstanbul, Turkey

Objective: Single intravenous bolus administration and postoperative or perioperative infusions are the most preferred methods of ketamine. Nevertheless, there is no clear explanation on the ideal ketamine administration method. In this study, we aimed to compare the effects of the most common ketamine administration methods and administration time on postoperative opioid consumption.

Methods: Fifty-two patients undergoing colectomy for colon cancer were randomly assigned into four groups. Group 1 was the control group. Group 2 received only a single intravenous bolus dose of 0.5 mg kg⁻¹ ketamine at induction. Group 3 received 0.5 mg kg⁻¹ intravenous ketamine bolus at induction and perioperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹. Group 4 received a bolus of 0.5 mg kg⁻¹ intravenous ketamine at induction and perioperative and postoperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹. Postoperatively, visual analogue scale pain scores, side effects, and morphine consumption were recorded.

Results: There was no statistically significant difference in postoperative pain scores. Total morphine consumption was found to be significantly lower in Group 4 compared to the other groups (p=0.03, p=0.004, p=0.03, respectively). During the 1st, 2nd, and 4th hours in the postoperative period, patient-controlled analgesia morphine consumption was significantly lower in Group 4 compared to the control group (p<0.01).

Conclusion: Preoperative single-bolus dose or intraoperative low-dose ketamine infusion does not decrease postoperative morphine consumption; however, per- and postoperative 48-hour ketamine infusion has a significant effect in decreasing morphine consumption without decreasing the incidence of side effects in patients undergoing major abdominal surgery.

Key Words: Ketamine, postoperative analgesia, colectomy, morphine

Introduction

Postoperative pain occurs through several anatomical and biochemical pathways. When the N-methyl-D-aspartate receptors (NMDAR) in the central nervous system are stimulated with afferent nociceptive inputs, the neuronal sensitization process is activated, and thus, pain perception increases. NMDAR activation does not only increase cellular response to painful stimulus, it also reduces the neuronal sensitivity against opioid receptor agonists (1). Activation of NMDAR is also the main mechanism of the hyperalgesia that is induced by opioids (2). Besides inhibiting central sensitization, using NMDAR antagonists with opioids also prevents tolerance that may develop against opioid analgesia (3).

Multimodal analgesia aims to improve efficiency by blocking different nociceptive pathways; therefore, when agents are administered in this way, their dose requirement and side effects can be reduced accordingly. Ketamine, which is a non-competitive NMDAR antagonist, has analgesic characteristics when it is used in subanaesthetic doses. However, several studies hypothesizing that subanaesthetic doses of ketamine may reduce the postoperative opioid requirement have reported controversial results. It has been reported that a low dose of ketamine administered intravenously (IV) as a part of multimodal analgesia after cesarean section did not have an additional postoperative analgesic effect (4). It has been indicated that adding ketamine to remifentanyl infusion as an adjuvant does not lead to a decrease in pain severity after cholecystectomy, time to first analgesic requirement, and morphine consumption during the first 24 hours (2). However, some randomized and controlled studies have concluded that ketamine decreases postoperative opioid consumption and incidence of opi-

oid-related side effects (5-9). These controversial outcomes may have been resulted from the methodology of the studies, heterogeneity of study populations, differences in patient-controlled analgesia (PCA) protocols or pain measurement methods, and the method of ketamine administration (10). There are several ways to administer ketamine, but the intravenous route has been reported to be the optimal method (11). Single intravenous bolus administration and postoperative or perioperative infusions are the most preferred methods. Nevertheless, meta-analyses did not bring a clear explanation on the ideal ketamine administration method (12, 13).

This prospective, randomized, double-blind study aimed to compare the effects of the most common ketamine administration methods-i.e., single bolus or infusion-and administration time (pre-induction, continuous infusion during preoperative and/or postoperative period) on postoperative opioid consumption.

Methods

After Marmara University Medical School ethics committee approval and patients' written informed consent, 52 patients, aged 30-70 years, American Society of Anesthesiology physical status I or II, undergoing colectomy for colon cancer under general anaesthesia were randomly assigned into four groups in this prospective, randomized, double-blinded, placebo-controlled study. Group allocation was determined by drawing white, closed envelopes containing group numbers. The patients, the anaesthesia team, and the team managing the postoperative analgesia were blinded to the study groups. A different anaesthesiologist prepared the solutions. Group 1 was the control group and received peri- and postoperative saline infusion. Group 2 received only a single-bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction and saline infusion peri- and postoperatively. Group 3 received 0.5 mg kg⁻¹ ketamine bolus IV at induction and perioperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹ and postoperative saline infusion. Group 4 received a bolus of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative and postoperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹.

All patients were instructed about the use of a patient-controlled analgesia (PCA) device and a 100-mm visual analog scale (VAS) the day before the operation. Exclusion criteria were patients' refusal, history of psychiatric disorders, altered mental status, alcohol abuse, opioid abuse or chronic opioid treatment, uncontrolled hypertension, renal or hepatic insufficiency, inability to use a PCA device, pregnancy, allergy to ketamine or morphine, and history of seizure or intracranial hypertension.

All patients were premedicated with 0.07 mg kg⁻¹ midazolam and 0.5 mg atropine intramuscularly administered 45 min before surgery. A central venous catheter, peripheral venous catheters, an arterial catheter, and urinary Foley catheters were inserted. Patients were continuously monitored for heart rate (ECG), oxygen saturation (SpO₂), end-tidal CO₂ pressure, invasive arterial blood pressure, and central venous pressure.

General anaesthesia was induced with thiopental sodium 5 mg kg⁻¹ and vecuronium 0.1 mg kg⁻¹ IV. Endotracheal intubation was performed in all patients. Anaesthesia was maintained with 1 MAC isoflurane and 70% N₂O in oxygen. Residual neuromuscular blockade was reversed with neostigmine 0.03 mg kg⁻¹ and atropine 0.015 mg kg⁻¹ IV.

Postoperatively in the recovery room, pain intensity was assessed using a 0-100-mm VAS, and IV morphine bolus was administered to keep the VAS score ≤30. Then, IV PCA with 1 mg mL⁻¹ morphine solution was started to deliver a 1.5-cc bolus dose with a 10-min lockout time. PCA was continued for 48 h after surgery. No other analgesics were used during the study. During follow-up, in case of patients' VAS ≥30, 2 mL morphine was loaded from a PCA device.

Postoperatively at 1, 2, 4, 8, 12, 24, 36, and 48 Hours, VAS scores, side effects (nausea and vomiting, pruritus, sedation with a 4-point scale, respiratory depression, psychomimetic side effects, such as hallucination and nightmares), and morphine consumption were recorded. Ondansetron 4 mg IV was used for the treatment of nausea and vomiting.

Statistical analysis

A reduction of 30% in postoperative morphine consumption was considered clinically significant. In order to obtain a power of 80% and significance level of 5%, the sample size for each study group was calculated as 13 patients or more. Demographic characteristics of the patients were compared statistically by one-way analysis of variance (ANOVA), while time-dependent variables were compared by two-way analysis of variance, and non-parametric data were analyzed by Kruskal-Wallis test. A *p*<0.05 was considered statistically significant, and the Tukey-Kramer test was used as the post hoc test.

Results

The demographic characteristics of the patients did not differ between groups (Table 1). There was no statistically significant difference in postoperative pain scores between the groups (Table 2). There was no significant difference in the amount of initial morphine requirement in the recovery room (Table 3).

Table 1. Patient characteristics (mean±SD)

	Group 1 (n=13)	Group 2 (n=13)	Group 3 (n=13)	Group 4 (n=13)
Age (year)	60.87±7.9	56.77±9.0	56.81±9.7	57.1±15.7
Weight (kg)	66.12±14.1	73.55±12.2	78.45±11.8	67.4±10.4
Male: female	4:4	5:4	7:4	4:6

p>0.05. Group 1: control group. Group 2: single-bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction. Group 3: bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹. Group 4: a bolus of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative and postoperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹

Table 2. Postoperative VAS scores (mm) (mean±SD)

	Group 1 (control, n=13)	Group 2 (n=13)	Group 3 (n=13)	Group 4 (n=13)	p
1 st hour	41.00±22.26	51.22±27.81	40.00±22.00	38.40±30.38	0.7075
2 nd hour	35.62±17.08	48.55±29.50	34.72±15.40	37.10±26.76	0.5411
4 th hour	22.25±12.89	37.22±25.64	23.72±11.74	25.60±31.05	0.4689
6 th hour	14.87±5.96	23.44±13.23	22.09±8.12	23.50±28.85	0.6893
8 th hour	16.75±6.54	16.55±7.03	16.82±7.58	11.00±28.85	0.8170
12 th hour	14.14±6.12	13.44±3.00	17.45±6.15	11.00±7.12	0.1101
24 th hour	12.87±5.71	12.22±6.03	13.45±8.23	14.90±6.98	0.8551
36 th hour	11.50±7.98	14.55±7.10	10.82±9.63	16.70±13.07	0.5231
48 th hour	10.37±5.99	13.55±13.39	11.82±10.62	7.77±7.88	0.6323

p>0.05. Group 1: control group. Group 2: single-bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction. Group 3: bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹. Group 4: a bolus of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative and postoperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹

Table 3. Morphine loading dose and total morphine consumption (mg) (mean±SD)

	Group 1 (n=13)	Group 2 (n=13)	Group 3 (n=13)	Group 4 (n=13)
Morphine bolus dose	8.25±2.91	5.66±3.04	8.81±2.85	8.70±6.16
Total PCA morphine consumption	101.87±53.31	115.61±43.48	98.37±49.16	61.77±35.38*

*p<0.05 compared with other groups. Group 1: control group. Group 2: single-bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction. Group 3: bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹. Group 4: a bolus of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative and postoperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹

Table 4. Side effects n (%)

	Group 1 n=13	Group 2 n=13	Group 3 n=13	Group 4 n=13
Nausea-vomiting	1 (7.7)	3 (23)	1 (7.7)	1 (7.7)
Nightmares	1 (7.7)	0 (0)	3 (23)	4 (30.7)
Hallucination	-	-	-	-

p>0.05. Group 1: control group. Group 2: single-bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction. Group 3: bolus dose of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹. Group 4: a bolus of 0.5 mg kg⁻¹ ketamine IV at induction and perioperative and postoperative ketamine infusion at a rate of 0.25 mg kg⁻¹ h⁻¹

and 4th hours in the postoperative period, PCA morphine consumption was significantly lower in Group 4 compared to the control group (p<0.01). The incidence of nightmares was recorded in 7.7%, 0%, 23%, and 30.7% of patients in each group, respectively (Table 4) (p>0.05). No hallucinations were reported in any of the groups, and there was no statistically significant difference in the incidence of nightmares, nausea and vomiting, pruritus, or sedation scores. Respiratory depression was not recorded in any patients.

Discussion

In this clinical study, we observed that low-dose ketamine decreased postoperative opioid requirements only when administered as an IV bolus at induction and IV infusion during the perioperative and postoperative 48 hours, without decreasing the incidence of opioid side effects.

Total PCA-morphine consumption was found to be significantly lower in Group 4 compared to the other groups (p=0.03, p=0.004, p=0.03, respectively). However, no significant difference was found between Group 1, Group 2, and Group 3 regarding total morphine consumption. During the 1st, 2nd,

Many studies conducted about the analgesic effect of IV low-dose ketamine have revealed controversial results. IV bolus ketamine has been reported to decrease morphine consumption in patients having severe acute pain due to trauma (14, 15). Argiriadou et al. (16) concluded that compared to a placebo group, intraoperative ketamine infusion was more effective in post-thoracotomy pain management. As ketamine provides early knee mobilization, it has been reported as a useful adjuvant for perioperative multimodal analgesia in patients undergoing arthroplasty (17). Another randomized, placebo-controlled study stated the opioid-sparing effect of preemptive IV bolus ketamine (18). On the other hand, Lopez-Alvarez et al. (19) indicated that morphine consumption and postoperative pain scores were higher in the group in which ketamine was administered. However, in this study, ketamine was only administered at induction; then, the patients received remifentanyl infusion during the surgery. Sveticic et al. (20) concluded that combining ketamine with PCA-morphine administration did not provide any additional benefits; so, they did not suggest ketamine usage in routine practice. In this study, patients received PCA with a bolus of morphine plus ketamine 1.5 mg each after major orthopaedic surgery. This result may be related to the sole use of ketamine in the postoperative period. In a meta-analysis of 35 randomized, controlled studies conducted in children, all of the included studies, except 2 of them, planned to examine the effects of single-bolus dose ketamine or single bolus plus intraoperative IV infusion (21). Ketamine was infused for 24 hours in one of the two studies mentioned above. The patients in this study received postoperative IV paracetamol, rectal morniflumate, and IV nalbuphine infusion. However, the pain management in the postoperative period is recommended to evaluate using the PCA method, because IV PCA provides superior pain relief compared with conventional on-demand analgesia (22, 23). In the second study, morphine consumption within 72 hours was evaluated using IV PCA without postoperative ketamine infusion, and it was found that perioperative infusion affected neither postoperative pain scores nor opioid consumption.

Such controversial outcomes may have resulted from different factors, such as heterogeneity of the study population and differences in PCA protocols or pain measurement methods; however, the most important reason for these controversial results is the methodological differences in ketamine administration. Previous studies have focused on ketamine regarding its administration time (during preoperative, perioperative, or postoperative period), use of different doses, and administration method, such as IV, intra-articular, or intramuscular. However, there have not been any studies analyzing both administration time and methods regarding ketamine. The current study examined the effects of using single preoperative IV low-dose ketamine bolus

and administering IV infusion during the peri- and postoperative periods on total morphine consumption as a part of multimodal analgesia. No statistical difference was found between the groups regarding pain scores during the study period. Similar pain scores were achieved with significantly lower total morphine consumption in Group 4 than the other groups. There was no difference in the PCA-morphine consumption at the postoperative 1st and 2nd hours between the group having a single-bolus dose and the one receiving peri- and postoperative ketamine infusion. However, the group having peri- and postoperative ketamine infusion consumed only 46% of the total morphine consumed by the group that received single-bolus dose ketamine. This can be attributed to the short duration of action of ketamine. Therefore, administering IV single-bolus dose ketamine did not affect the opioid consumption except for the first 2 postoperative hours. Ketamine, which is a short-acting anaesthetic agent, can be adequately effective only if it is administered as IV infusion.

In the present study, there was no increase in the incidence of behavioral, psychotomimetic, or neurological complications in patients receiving ketamine. In the postoperative period, even patients having ketamine infusion for 48 hours did not develop hallucinations. Moreover, nightmare incidence did not show a significant difference between groups. This may have resulted from use of low-dose ketamine-not from duration of infusions.

The limitation of our study is the number of patients in the groups (13 patients in each group). Although this is enough to detect a 30% decrease in morphine consumption, it may be different for the incidence of side effects. The incidence of hallucinations, which is insignificantly higher in Group 4, may be significant with a higher number of patients.

Conclusion

According to the results of the present study, the administration method of low-dose ketamine as an adjuvant in postoperative multimodal analgesia affects opioid consumption. Preoperative single-bolus dose or intraoperative low-dose ketamine infusion does not decrease postoperative morphine consumption; however, peri- and postoperative 48-hour ketamine infusion has a significant effect in decreasing morphine consumption without increasing the incidence of side effects in patients undergoing major abdominal surgery.

Ethics Committee Approval: Ethics committee approval was received for this study from the ethics committee of Marmara University Faculty of Medicine.

Informed Consent: Written informed consent was obtained from patients who participated in this study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - Z.E.; Design - Z.E., F.B.; Supervision - Z.E., F.Y.G.; Funding - Z.E., F.B.; Materials - Z.E., F.B., K.T.S.; Data Collection and/or Processing - F.B., K.A.; Analysis and/or Interpretation - Z.E., F.B., K.A.; Literature Review - Z.E., F.B., K.T.S.; Writer - Z.E., F.B., K.T.S.; Critical Review - Z.E., F.Y.G.; Other - F.B., Z.E., K.T.S., K.A., F.Y.G.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

References

- Sveticic G, Farzanegan F, Zmoos P, Zmoos S, Eichenberger U, Curatolo M. Is the combination of morphine with ketamine better than morphine alone for postoperative intravenous patient-controlled analgesia? *Anesth Analg* 2008; 106: 287-93. [\[CrossRef\]](#)
- Cunha Leal P, Kimiko Sakata R, Salomão R, Jun Sadatsune E, Machado Issy A. Assessment of the effect of ketamine in combination with remifentanyl on postoperative pain. *Rev Bras Anesthesiol* 2013; 63: 178-82. [\[CrossRef\]](#)
- Gharai B, Jafari A, Aghamohammadi H, Kamranmanesh M, Poorzamani M, Elyassi H, et al. Opioid-sparing effect of preemptive bolus low-dose ketamine for moderate sedation in opioid abusers undergoing extracorporeal shock wave lithotripsy: a randomized clinical trial. *Anesth Analg* 2013; 116: 75-80. [\[CrossRef\]](#)
- Bauchat JR, Higgins N, Wojciechowski KG, McCarthy RJ, Toledo P, Wong CA. Low-dose ketamine with multimodal postcesarean delivery analgesia: a randomized controlled trial. *Int J Obstet Anesth* 2011; 20: 3-9. [\[CrossRef\]](#)
- Menkiti ID, Desalu I, Kushimo OT. Low-dose intravenous ketamine improves postoperative analgesia after caesarean delivery with spinal bupivacaine in African parturients. *Int J Obstet Anesth* 2012; 21: 217-21. [\[CrossRef\]](#)
- Remérand F, Le Tendre C, Baud A, Couvret C, Pourrat X, Favard L, et al. The early and delayed analgesic effects of ketamine after total hip arthroplasty: a prospective, randomized, controlled, double-blind study. *Anesth Analg* 2009; 109: 1963-71. [\[CrossRef\]](#)
- Karcioglu M, Davarci I, Tuzcu K, Bozdogan YB, Turhanoglu S, Aydogan A, et al. Addition of ketamine to propofol-alfentanil anesthesia may reduce postoperative pain in laparoscopic cholecystectomy. *Surg Laparosc Endosc Percutan Tech* 2013; 23: 197-202. [\[CrossRef\]](#)
- Pacreu S, Fernández Candil J, Moltó L, Carazo J, Fernández Galinski S. The perioperative combination of methadone and ketamine reduces post-operative opioid usage compared with methadone alone. *Acta Anaesthesiol Scand* 2012; 56: 1250-6. [\[CrossRef\]](#)
- Bell RE, Dahl JB, Moore RA, Kalso E. Perioperative ketamine for acute postoperative pain. *Cochrane Database Syst Rev* 2006; CD004603.
- Weinbroum AA. Non-opioid IV adjuvants in the perioperative period: pharmacological and clinical aspects of ketamine and gabapentinoids. *Pharmacol Res* 2012; 65: 411-29. [\[CrossRef\]](#)
- Marland S, Ellerton J, Andolfatto G, Strapazzon G, Thomassen O, Brandner B, et al. Ketamine: Use in Anesthesia. *CNS Neurosci Ther* 2013; 19: 381-9. [\[CrossRef\]](#)
- Liu Y, Zheng Y, Gu X, Ma Z. The efficacy of NMDA receptor antagonists for preventing remifentanyl-induced increase in postoperative pain and analgesic requirement: a meta-analysis. *Minerva Anesthesiol* 2012; 78: 653-67.
- Dale O, Somogyi AA, Li Y, Sullivan T, Shavit Y. Does intraoperative ketamine attenuate inflammatory reactivity following surgery? A systematic review and meta-analysis. *Anesth Analg* 2012; 115: 934-43. [\[CrossRef\]](#)
- Galinski M, Dolveck F, Combes X, Limoges V, Smail N, Pommier V, et al. Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007; 25: 385-90. [\[CrossRef\]](#)
- Jennings PA, Cameron P, Bernard S, Walker T, Jolley D, Fitzgerald M, et al. Morphine and ketamine is superior to morphine alone for out-of-hospital trauma analgesia: a randomized controlled trial. *Ann Emerg Med* 2012; 59: 497-503. [\[CrossRef\]](#)
- Argiriadou H, Papagiannopoulou P, Foroulis CN, Anastasiadis K, Thomaidou E, Papakonstantinou C, et al. Intraoperative infusion of S(+)-ketamine enhances post-thoracotomy pain control compared with perioperative parecoxib when used in conjunction with thoracic paravertebral ropivacaine infusion. *J Cardiothorac Vasc Anesth* 2011; 25: 455-61. [\[CrossRef\]](#)
- Adam F, Chauvin M, Du Manoir B, Langlois M, Sessler DI, Fletcher D. Small-dose ketamine infusion improves postoperative analgesia and rehabilitation after total knee arthroplasty. *Anesth Analg* 2005; 100: 475-80. [\[CrossRef\]](#)
- Gharai B, Jafari A, Aghamohammadi H, Kamranmanesh M, Poorzamani M, Elyassi H, et al. Opioid-sparing effect of preemptive bolus low-dose ketamine for moderate sedation in opioid abusers undergoing extracorporeal shock wave lithotripsy: a randomized clinical trial. *Anesth Analg* 2013; 116: 75-80. [\[CrossRef\]](#)
- López-Álvarez S, Mayo-Moldes M, Zaballos M, Iglesias BG, Blanco-Dávila R. Esmolol versus ketamine-remifentanyl combination for early postoperative analgesia after laparoscopic cholecystectomy: a randomized controlled trial. *Can J Anaesth* 2012; 59: 442-8. [\[CrossRef\]](#)
- Sveticic G, Farzanegan F, Zmoos P, Zmoos S, Eichenberger U, Curatolo M. Is the combination of morphine with ketamine better than morphine alone for postoperative intravenous patient-controlled analgesia? *Anesth Analg* 2008; 106: 287-93. [\[CrossRef\]](#)
- Dahmani S, Michelet D, Abback PS, Wood C, Brasher C, Nivoche Y, et al. Ketamine for perioperative pain management in children: a meta-analysis of published studies. *Paediatr Anaesth* 2011; 21: 636-52. [\[CrossRef\]](#)

22. Pöpping DM, Zahn PK, Van Aken HK, Dasch B, Boche R, Pogatzki-Zahn EM. Effectiveness and safety of postoperative pain management: a survey of 18 925 consecutive patients between 1998 and 2006 (2nd revision): a database analysis of prospectively raised data. *Br J Anaesth* 2008; 101: 832-40. [\[CrossRef\]](#)
23. Palmer PP, Miller RD. Current and developing methods of patient-controlled analgesia. *Anesthesiol Clin* 2010; 28: 587-99. [\[CrossRef\]](#)