

Field-Friendly Test for Monitoring Multiple Immune Response Markers during Onset and Treatment of Exacerbated Immunity in Leprosy

Paul L. A. M. Corstjens,^a Anouk van Hooij,^b Elisa M. Tjon Kon Fat,^a Susan J. F. van den Eeden,^b Louis Wilson,^b Annemieke Geluk^b

Department of Molecular Cell Biology, Leiden University Medical Center, Leiden, The Netherlands^a; Department of Infectious Diseases, Leiden University Medical Center, Leiden, The Netherlands^b

Acute inflammatory reactions represent the major cause of irreversible neuropathy in leprosy. These tissue-destroying episodes have considerable overlap with acute immunological complications (flares) in several chronic (autoimmune) diseases that similarly warrant early detection. However, the lack of diagnostic tests impedes early diagnosis of these reactions. Here, we evaluated a user-friendly multiplex lateral flow assay for the simultaneous detection of IP-10 and anti-phenolic glycolipid I antibodies for longitudinally monitoring early onset and treatment of leprosy reactions.

Leprosy, an infectious disease caused by *Mycobacterium leprae*, still poses a major health threat in developing countries. Additionally, leprosy represents an intriguing model of human immunoregulatory disease, since its interindividual variability in clinical manifestations closely parallels the ability of the host to establish effective immunity to *M. leprae* (1, 2). This has resulted in leprosy being the first disease for which researchers identified HLA-disease association (3), human regulatory cells (4), and the Th1/Th2 concept for human T cells (5).

Although leprosy can be treated effectively with multidrug therapy (MDT), it can be complicated by acute inflammatory episodes, called leprosy reactions, that may occur before, during, and after the completion of MDT (6). Two distinct types of reactions are distinguished: reversal reactions (RRs) and erythema nodosum leprosum (ENL). These immunological complications occur in up to 50% of leprosy patients and represent the major cause of irreversible neurological damage and consequent anatomical deformities. Prompt diagnosis and treatment aid recovery from inflammatory nerve damage and reduce the risk of permanent disability considerably (7). However, if diagnosis and treatment are delayed for more than 6 months after symptom initiation, neuropathy is likely to be permanent (8). Tests for the early detection of leprosy reactions may make significant differences in clinical outcomes, especially when the tests are user-friendly and robust.

Previous work has shown that gamma-interferon (IFN- γ)-inducible protein 10 (IP-10) is a useful biomarker for the detection of *M. tuberculosis* infection (9) or to indicate *M. leprae* exposure (10, 11). Moreover, increased IP-10 serum levels are part of the biomarker profile characterizing the early onset of RRs (12, 13). Levels of IP-10 decline again during antireactional therapy (13), similar to what has been described during tuberculosis treatment (14). With respect to the humoral immune response, IgM directed against the *M. leprae*-specific phenolic glycolipid I (PGL-I), although not informative for the identification of RR onset (13), represents a useful biomarker for monitoring the efficacy of treatment of leprosy (reactions), since IgM levels drop when reactions are effectively subdued (13). More importantly, since a significant percentage of new patients in many areas where leprosy is endemic initially seek care because of reactions and only then are diagnosed

with leprosy, the detection of anti-PGL-I IgM serves as a confirmation of leprosy diagnosis.

Areas where leprosy is endemic are often short of sophisticated laboratories, which makes it imperative to develop diagnostic tests that are suitable for use in field settings. The combination of the user-friendly rapid lateral flow assay (LFA) format with the fluorescent quantitative up-converting phosphor (UCP) reporter technology has previously demonstrated usefulness for detecting and monitoring a variety of analytes (15–17). Additionally, besides having high sensitivity due to the complete absence of background fluorescence, UCP-LF test strips can be stored as permanent records, allowing reanalysis in a reference laboratory. To develop diagnostic assays for application in resource-limited areas, we previously developed UCP-LFAs for single detection of cytokines (IFN- γ , IP-10, interleukin 10 [IL-10], CCL4) as well as antibodies against *M. leprae* PGL-I for the diagnosis of nonreactive leprosy and tuberculosis (18–20). Generally, the performance of one biomarker can be significantly enhanced by using a custom-made grouping of independent biomarkers called a biomarker profile or signature.

In this study, we combined previous findings to evaluate the application of a multiplex UCP-LFA format for monitoring RR onset and treatment in leprosy patients. For this purpose, a UCP-LFA measuring IP-10 and anti-PGL-I IgM simultaneously was used to analyze serum samples from patients with borderline lepromatous leprosy collected prospectively in Bangladesh ($n = 4$ [13]), Brazil ($n = 3$ [13]), Nepal ($n = 2$ [13, 21]), and The Netherlands ($n = 1$ [22]). Newly diagnosed leprosy patients without

Received 31 January 2016 Returned for modification 22 February 2016

Accepted 23 March 2016

Accepted manuscript posted online 30 March 2016

Citation Corstjens PLAM, van Hooij A, Tjon Kon Fat EM, van den Eeden SJF, Wilson L, Geluk A. 2016. Field-friendly test for monitoring multiple immune response markers during onset and treatment of exacerbated immunity in leprosy. *Clin Vaccine Immunol* 23:515–519. doi:10.1128/CVI.00033-16.

Editor: R. S. Abraham, Mayo Clinic

Address correspondence to Annemieke Geluk, a.geluk@lumc.nl.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.




FIG 1 Comparison between single and multiplex IP-10 UCP-LFAs. UCP-LFAs for detection of IP-10 were performed in a single (A) or multiplex (B) format as described previously (18) using sera from 10 leprosy patients at leprosy diagnosis before MDT in the absence of any clinical sign of reactions and at least 3 months before reaction (before RR), at diagnosis of reaction before steroids (RR), or after MDT and reactional treatment, at least 1 month after the end of steroid use (after RR) (13). Differences in cytokine concentrations between test groups were analyzed with the Wilcoxon matched-pairs signed-rank test for nonparametric distribution using GraphPad Prism version 5.01 for Windows (GraphPad Software, San Diego, CA, USA). The dotted line indicates the limit of detection (LOD) (0.17). The y axis indicates the ratio of the relative fluorescence units (RFUs) measured at the respective test and flow-control lines. The statistical significance level used was $P \leq 0.05$. (C) For correlations, the Spearman r was calculated using GraphPad Prism version 5.01.

reactions at recruitment were entered into the study after informed consent was obtained. Ethical approval of the study protocol was obtained through appropriate ethics committees: the Ethical Review Committee of ICDDR,B (no. PR-10032 and no. PR-2007-069), the Brazilian National Council of Ethics in Research (CONEP) and UFU Research Ethics Committee (no. 499/2008), and the Nepal Health Research Council (NHR no. 751). Leprosy was diagnosed based on clinical, bacteriological, and histological observations and classified by skin biopsy specimens according to Ridley and Jopling (23). For analysis by the UCP-LFA, leprosy patients who developed reactions during MDT were tested using samples obtained at three different time points: (i) without clinical signs of reactions ≥ 3 months before RR, (ii) at RR diagnosis, before steroid treatment, and (iii) after RR, ≥ 1 month after ending steroid treatment. Concentrations of antibodies against PGL-I and IP-10 were measured in all sera using a single UCP-LFA

for either IP-10 or anti-PGL-I IgM and a multiplex UCP-LFA for both markers. Simultaneous detection of IP-10 and anti-PGL-I IgM was performed following a two-phase protocol described for single analyte detection (18, 24, 25). The protocol included a pre-flow incubation (60 min, 37°C, 900 rpm) of 10 μ l 100-fold-diluted sample with 90 μ l LF assay buffer containing 100 ng of the UCP^{αIP-10} conjugate and 100 ng of the UCP^{αIgM} conjugate (18).

The serum levels of IP-10 measured with the multiplex UCP-LFA at RR onset differed between the patients but were all significantly higher than those in the absence of reactions ($P = 0.045$ [Fig. 1]). Similarly, IP-10 concentrations were significantly reduced after treatment ($P = 0.027$). On the other hand, the anti-PGL-I IgM levels detected with the multiplex UCP-LFA did not identify the onset of RRs, which is in agreement with our previous findings (13). Seven patients were positive for anti-PGL-I IgM (range, 0.5 to 1.87) at diagnosis, but at RR onset, only two patients,

TABLE 1 Clinical parameters of leprosy patients with RR^a

Country	Symbol ^b	BI ^c	Type	Sex	Age (yrs)	Before RR ^d		At RR		After RR	
						PGL-I	IP-10	PGL-I	IP-10	PGL-I	IP-10
Bangladesh	■	2+	BL	Male	36	0.86	0.03	0.53	0.23	0.14	0.13
Bangladesh	+	2+	BL	Female	39	0.09	0.07	0.15	0.20	0.09	0.03
Bangladesh	○	0	BL	Male	36	0.84	0.12	0.64	0.87	0.50	0.15
Bangladesh	▼	2+	BL	Male	32	1.18	0.10	1.70	0.22	0.38	0.04
Brazil	●	3.2+	BL	Male	25	1.17	0.14	1.85	0.67	0.57	0.19
Brazil	△	2.42+	BL	Male	40	0.50	0.00	0.15	0.40	0.07	0.31
Brazil	*	4.28+	BL	Male	68	1.87	0.18	0.89	0.61	0.14	0.21
Nepal	⊙	0.25+	BL	Female	40	0.07	0.05	0.05	0.12	0.03	0.00
Nepal	★	0	BL	Male	33	0.10	0.14	0.07	0.49	0.03	0.07
Netherlands	⊠	5+	BL	Male	17	1.72	0.07	1.09	0.45	0.78	0.06

^a Values for the ratio of the relative fluorescence units (RFUs) measured at the respective test and flow-control lines of single UCP-LFA for anti-PGL-I IgM and IP-10 are shown.

^b Symbols used correspond to the symbols used for each individual in Fig. 1 and 2.

^c BI, bacterial index.

^d RR, reversal reaction.




FIG 2 Comparison between single and multiplex PGL-I UCP-LFAs. UCP-LFAs for detection of anti-PGL-I IgM antibodies were performed in a single (A) or multiplex (B) format as described previously (18) using sera from 10 leprosy patients (see Fig. 1). Differences in IgM levels between test groups were analyzed with the Wilcoxon matched-pairs signed-rank test for nonparametric distribution using GraphPad Prism version 5.01 for Windows (GraphPad Software). The dotted line indicates the LOD (0.11). The y axis indicates the ratio of the relative fluorescence units (RFUs) measured at test and flow-control lines. The statistical significance level used was $P \leq 0.05$. (C) For correlations, the Spearman r was calculated using GraphPad Prism version 5.01.

males from Bangladesh (aged 32 years; bacterial index [BI], ≥ 2) and Brazil (aged 25 years; BI, ≥ 3.2), showed increased antibody levels. These patients were not different from the others in BI or age (Table 1). However, serology clearly allowed monitoring of treatment efficacy for patients who were seropositive at RR onset, since levels were significantly reduced after treatment ($P = 0.0195$). Our data show that single and multiplex UCP-LFAs correlated well for IP-10 ($r = 0.91$ [Fig. 1C]) and anti-PGL-I IgM ($r = 0.96$ [Fig. 2C]), demonstrating no relevant interference between the two biomarkers.

Biomarkers as reliable correlates of disease complications and response to therapy represent essential tools for early diagnosis of disease states in chronic infections. In areas where leprosy is endemic, leprosy reactions are frequently misdiagnosed due to a lack of expertise. The evaluation of this multiplex UCP-LFA shows that this quantitative test, when used for intraindividual monitoring, can aid health care workers in the early diagnosis of reactional episodes, allowing timely treatment. In view of the frequent recurrence of these episodes, this test is particularly useful for monitoring treatment efficacy.

Since acute inflammatory (delayed hypersensitivity) reactions are a frequently occurring, tissue-destroying phenomenon in chronic (infectious) diseases as well as in autoimmune diseases, similar multiplex IP-10 UCP-LFAs adapted, e.g., for the detection of antibodies against tumor necrosis factor alpha (TNF- α) (25) could also be applied to diagnose similar episodes in other diseases, such as rheumatoid arthritis (26), psoriasis (27), and Crohn's disease, (28) for which conscientious, personalized treatment monitoring is also vital.

ACKNOWLEDGMENTS

We gratefully acknowledge the patients for their participation and Sayera Banu (IDDRB, Dhaka), Deanna Hagge (MLR, Kathmandu), Isabela Goulart (National Reference Center for Sanitary Dermatology and Leprosy Uberlandia), and Colette van Hees (EMC, Rotterdam) for organizing the recruitment of leprosy patients.

We declare no financial or commercial conflicts of interest.

FUNDING INFORMATION

This work, including the efforts of Annemieke Geluk, was funded by Order of Malta Grants for Leprosy Research (MALTALAP). This work, including the efforts of Annemieke Geluk, was funded by The Heiser Program for Research in Leprosy and Tuberculosis of the New York Community Trust (P13-000392). This work, including the efforts of Annemieke Geluk, was funded by The Q. M. Gastmann-Wichers Foundation. This work, including the efforts of Annemieke Geluk, was funded by The Netherlands Leprosy Relief Foundation (NLR) (ILEP 702.02.68, ILEP 7.01.02.48, and ILEP 701.02.49). This work, including the efforts of Annemieke Geluk and Paul Corstjens, was funded by European and Developing Countries Clinical Trials Partnership (EDCTP) (IP-2009-32040).

REFERENCES

- Scollard DM, Adams LB, Gillis TP, Krahenbuhl JL, Truman RW, Williams DL. 2006. The continuing challenges of leprosy. *Clin Microbiol Rev* 19:338–381. <http://dx.doi.org/10.1128/CMR.19.2.338-381.2006>.
- Geluk A. 2013. Biomarkers for leprosy: would you prefer T (cells)? *Lepr Rev* 84:3–12.
- Ottenhoff TH, Gonzalez NM, de Vries RR, Convit J, van Rood JJ. 1984. Association of HLA specificity LB-E12 (MB1, DC1, MT1) with lepromatous leprosy in a Venezuelan population. *Tissue Antigens* 24:25–29.
- Ottenhoff TH, Elferink DG, Klatser PR, de Vries RR. 1986. Cloned suppressor T cells from a lepromatous leprosy patient suppress *Mycobacterium leprae* reactive helper T cells. *Nature* 322:462–464. <http://dx.doi.org/10.1038/322462a0>.
- Modlin RL. 1994. Th1-Th2 paradigm: insights from leprosy. *J Invest Dermatol* 102:828–832. <http://dx.doi.org/10.1111/1523-1747.ep12381958>.
- Scollard DM, Martelli CM, Stefani MM, Maroja MF, Villahermosa L, Pardillo F, Tamang KB. 2015. Risk factors for leprosy reactions in three endemic countries. *Am J Trop Med Hyg* 92:108–114. <http://dx.doi.org/10.4269/ajtmh.13-0221>.
- Lockwood DN, Saunderson P. 2012. Nerve damage in leprosy: a continuing challenge for scientists, clinicians and service providers. *Int Health* 4:77–85. <http://dx.doi.org/10.1016/j.inhe.2011.09.006>.
- Ranque B, Nguyen VT, Vu HT, Nguyen TH, Nguyen NB, Pham XK, Schurr E, Abel L, Alcais A. 2007. Age is an important risk factor for onset and sequelae of reversal reactions in Vietnamese patients with leprosy. *Clin Infect Dis* 44:33–40. <http://dx.doi.org/10.1086/509923>.
- Ruhwald M, Dominguez J, Latorre I, Losi M, Richeldi L, Pasticcini MB, Mazzolla R, Goletti D, Butera O, Bruchfeld J, Gaines H, Gerogianni I, Tuuminen T, Ferrara G, Eugen-Olsen J, Ravn P. 2011. A multicentre evaluation of the accuracy and performance of IP-10 for the diagnosis of infection with *M. tuberculosis*. *Tuberculosis (Edinb)* 91:260–267. <http://dx.doi.org/10.1016/j.tube.2011.01.001>.
- Geluk A, Bobosha K, van der Ploeg-van Schip JJ, Spencer JS, Banu S, Martins SB, Cho SN, Franken KL, Kim HJ, Bekele Y, Uddin MK, Abdul HS, Aseffa A, Pessolani MC, Pereira GM, Dockrell HM, Ottenhoff TH. 2012. New biomarkers with relevance to leprosy diagnosis applicable in areas hyperendemic for leprosy. *J Immunol* 188:4782–4791. <http://dx.doi.org/10.4049/jimmunol.1103452>.
- Bobosha K, Tang ST, van der Ploeg-van Schip JJ, Bekele Y, Martins MV, Lund O, Franken KL, Khadge S, Pontes MA, Goncalves HS, Hussien J, Thapa P, Kunwar CB, Hagge DA, Aseffa A, Pessolani MC, Pereira GM, Ottenhoff TH, Geluk A. 2012. *Mycobacterium leprae* virulence-associated peptides are indicators of exposure to *M. leprae* in Brazil, Ethiopia and Nepal. *Mem Inst Oswaldo Cruz* 107(Suppl 1):S112–S123.
- Scollard DM, Chaduvula MV, Martinez A, Fowlkes N, Nath I, Stryjewska BM, Kearney MT, Williams DL. 2011. Increased CXC ligand 10 levels and gene expression in type 1 leprosy reactions. *Clin Vaccine Immunol* 18:947–953. <http://dx.doi.org/10.1128/CVI.00042-11>.
- Khadge S, Banu S, Bobosha K, van der Ploeg-van Schip JJ, Goulart IM, Thapa P, Kunwar CB, van Meijgaard KE, van den Eeden SJ, Wilson L, Kabir S, Dey H, Goulart LR, Lobato J, Carvalho W, Bekele Y, Franken KL, Aseffa A, Spencer JS, Oskam L, Ottenhoff TH, Hagge DA, Geluk A. 2015. Longitudinal immune profiles in type 1 leprosy reactions in Bangladesh, Brazil, Ethiopia and Nepal. *BMC Infect Dis* 15:477. <http://dx.doi.org/10.1186/s12879-015-1128-0>.
- Wergeland J, Pullar N, Assmus J, Ueland T, Tonby K, Feruglio S, Kvale D, Damas JK, Aukrust P, Mollnes TE, Dyrhol-Riise AM. 2015. IP-10 differentiates between active and latent tuberculosis irrespective of HIV status and declines during therapy. *J Infect* 70:381–391. <http://dx.doi.org/10.1016/j.jinf.2014.12.019>.
- van Dam GJ, de Dood CJ, Lewis M, Deelder AM, van Lieshout L, Tanke HJ, van Rooyen LH, Corstjens PL. 2013. A robust dry reagent lateral flow assay for diagnosis of active schistosomiasis by detection of *Schistosoma* circulating anodic antigen. *Exp Parasitol* 135:274–282. <http://dx.doi.org/10.1016/j.exppara.2013.06.017>.
- Zuiderwijk M, Tanke HJ, Sam NR, Corstjens PL. 2003. An amplification-free hybridization-based DNA assay to detect *Streptococcus pneumoniae* utilizing the up-converting phosphor technology. *Clin Biochem* 36:401–403. [http://dx.doi.org/10.1016/S0009-9120\(03\)00057-2](http://dx.doi.org/10.1016/S0009-9120(03)00057-2).
- Corstjens PL, Chen Z, Zuiderwijk M, Bau HH, Abrams WR, Malamud D, Sam NR, Tanke HJ. 2007. Rapid assay format for multiplex detection of humoral immune responses to infectious disease pathogens (HIV, HCV, and TB). *Ann N Y Acad Sci* 1098:437–445. <http://dx.doi.org/10.1196/annals.1384.016>.
- Bobosha K, Tjon Kon Fat EM, van den Eeden SJ, Bekele Y, van der Ploeg-van Schip JJ, de Dood CJ, Dijkman K, Franken KL, Wilson L, Aseffa A, Spencer JS, Ottenhoff TH, Corstjens PL, Geluk A. 2014. Field-evaluation of a new lateral flow assay for detection of cellular and humoral immunity against *Mycobacterium leprae*. *PLoS Negl Trop Dis* 8:e2845. <http://dx.doi.org/10.1371/journal.pntd.0002845>.
- Corstjens PL, Zuiderwijk M, Tanke HJ, van der Ploeg-van Schip JJ, Ottenhoff TH, Geluk A. 2008. A user-friendly, highly sensitive assay to detect the IFN-gamma secretion by T cells. *Clin Biochem* 41:440–444. <http://dx.doi.org/10.1016/j.clinbiochem.2007.12.015>.
- Corstjens PL, Tjon Kon Fat EM, de Dood CJ, van der Ploeg-van Schip JJ, Franken KL, Chegou NN, Sutherland JS, Howe R, Mihret A, Kassa

- D, van der Vyver M, Sheehama VJ, Simukonda F, Mayanja-Kizza H, Ottenhoff TH, Walzl G, Geluk A. 2015. Multi-center evaluation of a user-friendly lateral flow assay to determine IP-10 and CCL4 levels in blood of TB and non-TB cases in Africa. *Clin Biochem* 49:22–31. <http://dx.doi.org/10.1016/j.clinbiochem.2015.08.013>.
21. Mayboroda OA, van Hooij A, Derks R, van den Eeden SJ, Dijkman K, Khadge S, Thapa P, Kunwar CB, Hagge DA, Geluk A. 2016. Exploratory urinary metabolomics of type 1 leprosy reactions. *Int J Infect Dis* 45:46–52. <http://dx.doi.org/10.1016/j.ijid.2016.02.012>.
 22. Geluk A, van Meijgaarden KE, Wilson L, Bobosha K, van der Ploeg-van Schip JJ, van den Eeden SJ, Quinten E, Dijkman K, Franken KL, Haisma EM, Haks MC, van Hees CL, Ottenhoff TH. 2014. Longitudinal immune responses and gene expression profiles in type 1 leprosy reactions. *J Clin Immunol* 34:245–255. <http://dx.doi.org/10.1007/s10875-013-9979-x>.
 23. Ridley DS, Jopling WH. 1966. Classification of leprosy according to immunity. A five-group system. *Int J Lepr Other Mycobact Dis* 34:255–273.
 24. Corstjens PL, de Dood CJ, van der Ploeg-van Schip JJ, Wiesmeijer KC, Riuttamaki T, van Meijgaarden KE, Spencer JS, Tanke HJ, Ottenhoff TH, Geluk A. 2011. Lateral flow assay for simultaneous detection of cellular- and humoral immune responses. *Clin Biochem* 44:1241–1246. <http://dx.doi.org/10.1016/j.clinbiochem.2011.06.983>.
 25. Corstjens PL, Fidler HH, Wiesmeijer KC, de Dood CJ, Rispens T, Wolbink GJ, Hommes DW, Tanke HJ. 2013. A rapid assay for on-site monitoring of infliximab trough levels: a feasibility study. *Anal Bioanal Chem* 405:7367–7375. <http://dx.doi.org/10.1007/s00216-013-7154-0>.
 26. Hanaoka R, Kasama T, Muramatsu M, Yajima N, Shiozawa F, Miwa Y, Negishi M, Ide H, Miyaoka H, Uchida H, Adachi M. 2003. A novel mechanism for the regulation of IFN-gamma inducible protein-10 expression in rheumatoid arthritis. *Arthritis Res Ther* 5:R74–R81. <http://dx.doi.org/10.1186/ar616>.
 27. Levis WR, Martiniuk F. 2013. Psoriasis and leprosy are teaching us T-cell plasticity. *J Drugs Dermatol* 12:1082.
 28. Grant AV, Alter A, Huong NT, Orlova M, Thuc NV, Ba NN, Thai VH, Abel L, Schurr E, Alcais A. 2012. Crohn's disease susceptibility genes are associated with leprosy in the Vietnamese population. *J Infect Dis* 206:1763–1767. <http://dx.doi.org/10.1093/infdis/jis588>.