

Erratum

J Vet Intern Med 2016;30:689
10.1111/jvim.13912

Erratum for Comparison of Efficacy of Long-term Oral Treatment with Telmisartan and Benazepril in Cats with Chronic Kidney Disease (pages 1479–1487)

This article corrects:

Comparison of Efficacy of Long-term Oral Treatment with Telmisartan and Benazepril in Cats with Chronic Kidney Disease (pages 1479–1487).

U. Sent, R. Gössl, J. Elliott, H. M. Syme and T. Zimmering

Article first published online: 16 OCT 2015 | DOI: 10.1111/jvim.13639

In Table 2 the number 7 is missing in the Telmisartan group, IRIS 2b, substaging UP/C >0.4.

Table 2. Telmisartan and Benazepril group (intention-to-treat population, n = 112 each), IRIS stage at baseline including substaging.

Telmisartan (n = 112) ^a	IRIS Stage 2a (n = 36)	IRIS Stage 2b (n = 48)	IRIS Stage 3 (n = 21)	IRIS Stage 4 (n = 3) ^b
Substaging by SBP				
<150 mmHg	17	30	6	1
150–159 mmHg	11	5	5	1
160–179 mmHg	8	13	10	0
≥180 mmHg	0	0	0	0
Substaging by UP/C ^a				
<0.2	1	1	0	0
0.2–1.4	26	40	13	0
>0.4	9	7	8	2
Benazepril (n = 112)	IRIS Stage 2a (n = 47)	IRIS Stage 2b (n = 44)	IRIS Stage 3 (n = 19)	IRIS Stage 4(n = 2)
Substaging by SBP				
<150 mmHg	24	18	8	1
150–159 mmHg	7	9	5	1
160–179 mmHg	16	16	6	0
≥180 mmHg	0	1	0	0
Substaging by UP/C				
<0.2	0	1	0	0
0.2–0.4	42	31	15	1
>0.4	5	12	4	1

^aIRIS stage 1 (n = 4) not shown.

^bOne missing value for IRIS 4.