

Classifications in Brief

Kellgren-Lawrence Classification of Osteoarthritis

Mark D. Kohn BA, Adam A. Sassoon MD, Navin D. Fernando MD

Received: 15 September 2015 / Accepted: 1 February 2016 / Published online: 12 February 2016
© The Association of Bone and Joint Surgeons® 2016

History

Osteoarthritis (OA) ranks globally among the 50 most common sequelae of diseases and injuries, affecting over 250 million people or 4% of the world's population [39]. Of the global disease burden for OA, knee OA constitutes 83% [39]. A detailed analysis of Medicare beneficiaries reported the TKA annual utilization rate ranging from 287,006 in 2006 to 301,956 in 2010 [26]. The demand for TKA is expected to grow exponentially over the coming decades with epidemiological data suggesting a 673% increase in the United States by 2030, representing 3.48 million procedures annually [22].

As a polymorphic disease with a variety of clinical presentations, OA is challenging to rigorously define. A commonly encountered definition of OA describes "...a heterogeneous group of conditions that leads to joint symptoms and signs which are associated with defective integrity of articular cartilage, in addition to related changes in the underlying bone and at the joint margins" [1]. The pathogenesis of OA is poorly understood but is thought to include a complex interplay among mechanical,

biochemical, cellular, genetic, and immunologic phenomena [7]. Several attempts to develop diagnostic criteria for OA were previously undertaken and incorporate patient-reported joint pain in conjunction with consistent radiographic findings [1, 6, 24]. OA generally can be subcategorized into primary (idiopathic) and secondary OA [1, 25]. Common causes of secondary OA include post-traumatic, dysplastic, infectious, inflammatory, or biochemical etiologies that are relatively well understood. Although the etiology of primary OA remains largely undefined, genetic factors, age-related physiological changes, ethnicity, and biomechanical factors likely play an important role [16].

Plain radiography remains a mainstay in the diagnosis of OA. The first formalized attempts at establishing a radiographic classification scheme for OA were described by Kellgren and Lawrence (KL) in 1957 [19]. After studying rheumatism in coal miners at the Bedford Colliery in North West England [18], Kellgren investigated the inter- and intraobserver reliability of radiographic changes of rheumatism observed in the hand [17]. After concluding that there was wide disagreement among different observers, KL endeavored to establish a classification scheme with an associated set of standardized radiographs for OA of diarthrodial joints. They proposed a five-grade classification scheme and examined plain radiographs of eight joints including the distal interphalangeal joint (DIP), metacarpophalangeal joint (MCP), first carpometacarpal joint (CMC), wrist, cervical spine, lumbar spine, hips, and knees to calculate the inter- and intraobserver reliability of each [19]. They found that the tibiofemoral joint of the knee had the highest interobserver correlation coefficient of $r = 0.83$ (range of all joints studied, 0.10–0.83) as well as the second highest intraobserver correlation coefficient of

Each author certifies that he or she, or a member of his or her immediate family, has no funding or commercial associations (eg, consultancies, stock ownership, equity interest, patent/licensing arrangements, etc) that might pose a conflict of interest in connection with the submitted article.

All ICMJE Conflict of Interest Forms for authors and *Clinical Orthopaedics and Related Research*® editors and board members are on file with the publication and can be viewed on request.

M. D. Kohn, A. A. Sassoon, N. D. Fernando (✉)
University of Washington, 10330 Meridian Avenue N, Suite 270,
Seattle, WA 98133, USA
e-mail: navinf@uw.edu

$r = 0.83$ (range of all joints studied, 0.42–0.88) among the diarthrodial joints they examined [19]. These early results would predict the future application of their classification scheme to the knee specifically. Currently, the KL classification is the most widely used clinical tool for the radiographic diagnosis of OA [5].

Purpose

The KL classification has been commonly used as a research tool in epidemiological studies of OA, including landmark articles by Felson et al. [10] in the Framingham Osteoarthritis Study, and Bagge et al. [3] assessing osteoarthritis in European populations. The KL classification was also used in the development of atlases of radiographic features of OA, including the work done by Scott et al. [33].

The Kellgren and Lawrence classification may also assist healthcare providers with a treatment algorithm to guide clinical decision-making, specifically defining which patients may benefit most from surgical management. Furthermore, some insurers currently require providers to include documentation of the KL classification to receive approval for a TKA [37, 38].

Despite its common use, all research and clinical efforts using the KL classification depend critically on rigorous validation and continuous reevaluation of the schema's relevance to patient-centered outcomes.

Description of the Kellgren-Lawrence Classification System

Based on the data presented in their original work, the KL classification is typically applied specifically within the context of knee OA. The KL classification was originally described using AP knee radiographs. Each radiograph was assigned a grade from 0 to 4, which they correlated to increasing severity of OA, with Grade 0 signifying no presence of OA and Grade 4 signifying severe OA [19] (Fig. 1). Additionally, KL provided detailed radiographic descriptions of OA (Table 1). Although it is unclear from the original paper whether the radiographic descriptions were presented with the intent of demonstrating a linear disease progression of OA that begins with the formation of osteophytes and culminates in the altered shape of bone ends, other authors have criticized the KL system on the basis of this assumption [5, 27].

Validation

In their original paper, KL acknowledged variability in the radiographic evaluation of OA and described the inter- and

intraobserver reliability of their system [19] (Table 2). In particular, the authors comment that estimates of OA prevalence in all joints examined vary considerably between observers ($\pm 31\%$ deviation from the mean number of diagnoses of OA) and less so within the same observer ($\pm 5\%$ deviation from the mean number of diagnoses of OA). The authors report the inter- and intraobserver reliability correlation coefficient of the knee to both be 0.83 and comment that, “A significant correlation between the two observers was obtained for all joints except the wrist” and “Two readings by the same observer gave only a slightly better correlation on the reading of individual x rays...” [19]. It is unlikely, however, that the authors used the term “significant” within the same statistical context that the word carries in the literature today because no p values or confidence intervals were reported for their data. To determine the inter- and intraobserver correlation coefficients, KL used two observers to evaluate a series of 510 radiographs of eight joints including the DIP, MCP, first CMC, wrist, cervical spine, lumbar spine, hips, and knees from 85 patients aged 55 to 64 years selected randomly from an urban population. The authors did not explicitly report how many radiographs from each joint were used in their calculations or mention the qualifications or training of the two observers who graded the radiographs.

More recently, Wright [40] reevaluated the interobserver reliability of the KL system in addition to five other radiographic classification schemes used to grade knee OA (Table 3). The group used radiographs from 632 patients enrolled in the Multicenter ACL Revision Study (MARS) consortium, which consisted of cohorts of patients from 83 surgeons from 52 sites. The investigators used three independent and blinded observers (specific qualifications and training not explicitly stated by authors) to grade weight-bearing AP and/or Rosenberg radiographs (depending on availability) with six radiographic classification schemes. In addition, their radiographic findings were compared with arthroscopic evidence of tibiofemoral chondral disease. They reported that the KL system was the most studied among the different classification systems and had an interobserver reliability intraclass correlation coefficient of 0.51 to 0.89 (considered “moderate” to “very good” based on the author's provided definitions) from studies since the original KL article [40]. The investigators provided their own interobserver reliability intraclass correlation coefficients using the Rosenberg radiograph 0.54 (95% confidence interval [CI], 0.48–0.59) and AP radiographs 0.38 (95% CI, 0.33–0.43), which they characterized as moderate and poor, respectively [40]. The authors provide an explanation for this wide range in interobserver reliability by suggesting that differences in technique, population age group, and degree of OA likely contributed




Fig. 1A–D AP radiographs of the knee presented in the original Kellgren–Lawrence article [19]. **(A)** Representative knee radiograph of KL classification Grade 1, which demonstrates doubtful narrowing of the joint space with possible osteophyte formation. **(B)** Representative knee radiograph of KL classification Grade 2, which demonstrates possible narrowing of the joint space with definite osteophyte formation. **(C)** Representative knee radiograph of KL classification Grade 3, which demonstrates definite narrowing of joint

space, moderate osteophyte formation, some sclerosis, and possible deformity of bony ends. **(D)** Representative knee radiograph of KL classification Grade 4, which demonstrates large osteophyte formation, severe narrowing of the joint space with marked sclerosis, and definite deformity of bone ends. Reprinted with permission from Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthritis. *Ann Rheum Dis.* 1957;16:494–502.

Table 1. Radiologic features of osteoarthritis as described by Kellgren and Lawrence [19]

Formation of osteophytes on joint margin or on the tibial spines
Periarticular ossicles (primarily with regard to DIP and PIP joints)
Narrowing of joint cartilage associated with sclerosis of subchondral bone
Small pseudocystic areas with sclerotic walls situated usually in the subchondral bone
Altered shape of bone ends, particularly in the head of the femur

Modified with permission from Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthritis. *Ann Rheum Dis.* 1957;16:494–502. DIP = distal interphalangeal; PIP = posterior interphalangeal.

to the differing results from the various studies. It is also likely that variable interpretations of the KL classification by observers in different studies had a considerable effect on the reliability data they reported.

Table 2. Inter- and intraobserver correlation coefficients of the Kellgren and Lawrence classification in various joints [19]

Joint examined	Interobserver correlation coefficient (r)	Intraobserver correlation coefficient (r)
First carpometacarpal joint	0.78	0.81
Wrist	0.10	0.62
Cervical spine	0.57	0.66
Dorsolumbar spine	0.52	0.42
Hips	0.40	0.75
Knees	0.83	0.83
Metacarpophalangeal	Not done	0.88

Modified with permission from Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthritis. *Ann Rheum Dis.* 1957;16:494–502.

In the 1987 study of the Framingham population by Felson et al. [10], the investigators had two academically based bone and joint radiologists examine standing AP

Table 3. Grading scales for the radiographic osteoarthritis classification systems [40]

Scale	Grade and characteristics
Kellgren-Lawrence [19, 30]	0: No JSN or reactive changes 1: Doubtful JSN, possible osteophytic lipping 2: Definite osteophytes, possible JSN 3: Moderate osteophytes, definite JSN, some sclerosis, possible bone-end deformity 4: Large osteophytes, marked JSN, severe sclerosis, definite bone ends deformity
IKDC [14, 15]	A: No JSN* B: > 4-mm joint space; small osteophytes, slight sclerosis, or femoral condyle flattening C: 2- to 4-mm joint space D: < 2-mm joint space
Fairbank [9, 36]	0: Normal 1: Squaring of tibial margin 2: Flattening of femoral condyle, squaring and sclerosis of tibial margin 3: JSN, hypertrophic changes, or both 4: > 75% JSN with secondary feature
Brandt et al. [4]	0: < 25% JSN without secondary features (subchondral sclerosis, geodes, and osteophytes) 1: < 25% JSN with secondary features of 25%–50% JSN without secondary features 2: 25%–50% JSN with secondary features or 50%–75% JSN without secondary features 3: 50%–75% JSN with secondary features or > 75% JSN without secondary features 4: > 75% JSN with secondary features
Ahlbäck [11]	0: Normal 1: JSN [†] (with or without subchondral sclerosis) 2: Obliteration of joint space 3: Bone defect/loss < 5 mm 4: Bone defect and/or loss 5–10 mm
Jäger-Wirth [29, 32]	0: No arthrosis 1: Initial arthrosis, small osteophytes, minimal JSN 2: Moderate arthrosis, approximately 50% JSN 3: Medium-grade arthrosis 4: Heavy arthrosis

Reprinted with permission from Wright RW. Osteoarthritis classification scales: interobserver reliability and arthroscopic correlation. *J Bone Joint Surg Am.* 2014;96:1145–1151; * JSN = joint space narrowing; IKDC = International Knee Documentation Committee; [†] joint space narrowing is < 3 mm of the joint space or < 50% of the other compartment.

radiographs of the knee of 1424 elderly patients (ages 63–94 years old, mean = 73 years) and reported an interrater intraclass correlation coefficient of 0.85 (“very good”) using the KL classification. Although the study provided unique population-based data using highly trained observers, the authors admit that their population lacked ethnic diversity by having no black, Hispanic, or non-European ethnic groups. Given the growing ethnic diversity of the US population, it may be difficult to extrapolate the results found in this study to the broader population. In later works by Scott et al. [33], the authors used two skeletal radiologists and two rheumatologists to examine 30 standing AP knee radiographs randomly selected by an investigator not involved in the readings from the Baltimore Longitudinal Study of Aging and reported an interreader intraclass correlation coefficient of 0.68 (“good”) and intrareader intraclass correlation coefficient of 0.87 (“very good”). The authors do not provide detailed ethnic demographic information about their study population but report that their subjects consisted of 25 men and five women (ages 42–84 years old, mean age of men = 67 years, mean age of women = 71 years). Both studies admit to selecting a similar number of radiographs from each KL classification grade for calculating their intraclass correlation coefficients, which may provide more stable predictions of reliability scores but may not be as readily applicable to larger populations. Another study by Gossec et al. [12] used 50 radiographs selected from 1759 radiographs from five databases of trials or cohort studies to evaluate the interreader and intrareader intraclass correlation coefficient of standing, extended knee radiographs using two trained rheumatologists and reported that the KL classification had an interreader intraclass correlation coefficient of 0.72 (95% CI, 0.38–0.86) (“good”) and an intrareader intraclass correlation coefficient of 0.72 (95% CI, 0.55–0.83) (“good”). No criteria for the selection of the 50 radiographs from the collection of 1759 were explicitly stated by the authors and it is not clear if the selection included relatively equal numbers of radiographs from each KL classification grade. Given the differing demographic makeup of each population studied, including differences in ethnic group, percentage of male versus female, and age as well as the differing selection criteria for the radiographs examined by the observers, which themselves varied in number, qualifications, and training, it is understandable that the range of interobserver intraclass correlation coefficients for the KL classification cited by Wright [40] (0.51–0.89) is as wide as it is.

Although independent validation of the KL classification has been examined by multiple authors, less is known about the KL classification’s diagnostic accuracy, that is the degree to which the radiographic findings actually reflect the physical state of the joint; the best work on this




Fig. 2 Diagram of how the Rosenberg radiograph would be set up and performed [28]. Reprinted with permission from Rosenberg TD, Paulos LE, Parker RD, Coward DB, Scott SM. The forty-five-degree posteroanterior flexion weight-bearing radiograph of the knee. *J Bone Joint Surg Am.* 1988;70:1479–1483.

suggests that using the Rosenberg view (the 45° posteroanterior flexion weightbearing radiograph; Fig. 2 [28]) results in higher interrater reliability (0.54; 95% CI, 0.48–0.59) and better correlation to arthroscopic evidence of OA (Spearman rho 0.42; 95% CI, 0.33–0.49) than does using the AP radiograph (interrater reliability 0.38; 95% CI, 0.33–0.43; Spearman rho 0.30; 95% CI, 0.23–0.38) [40]. Those authors argue that this may be because the Rosenberg radiograph provides better visualization of the femoral condyles in midflexion, a common site of articular surface degeneration.

Limitations

Despite the wide application of the KL classification, the system has several noted limitations. Perhaps the most widely argued criticism of the KL system is its application to disease progression and insensitivity to change. The KL system has been criticized by Spector and Cooper [35] for assuming a linear radiographic progression of OA, starting with osteophyte formation, proceeding to joint space narrowing (JSN), and terminating in deformation of articular surfaces. However, as the authors point out, JSN in the absence of osteophyte formation cannot be measured in the KL system, which becomes problematic in patients with

knee pain and radiographically evident loss of cartilage but a lack of osteophytes on their knee radiographs. Although this occurrence is thought to be less common, the presence of marginal osteophytes likely represents a hypertrophic response to mechanical stress in conjunction with enchondral ossification, which may occur along a spectrum depending on individual physiology [21]. Other classification systems have attempted to segregate the radiological findings by performing individual evaluations of the joint space for the medial and lateral compartments separately and doing likewise again for osteophyte formation in each location [2, 5, 27, 34]. However, these classification systems have not been adopted widely, likely as a result of challenges in standardizing definitions and an inherent difficulty in discriminating JSN and osteophyte formation. In addition, Günther and Sun [13] described inferior interobserver intraclass correlation coefficients for medial compartment (0.62) and lateral compartment (0.47) JSN as well as medial femorotibial, lateral femorotibial, and tibial spine osteophyte formation (0.75, 0.74, and 0.63, respectively) as compared with the overall KL classification score (0.81) when using three observers, one of whom was an experienced orthopaedic surgeon and two of whom were orthopaedic residents. As discussed by Emrani et al. [8], classifications based on JSN may be preferable to the KL system for monitoring progression of OA, whereas the KL system may be better for assessing severity of osteoarthritic disease. A criticism of multiple classification systems (including the KL) is the lack of recognition of patellofemoral arthritis as a distinct or contributory radiographic factor.

Another criticism of the KL system is the inconsistency in its original description by the authors and variable applications of the classification in subsequent studies. In their original paper, KL provided simple descriptions of each grade, “none, doubtful, minimal, moderate, severe,” along with radiographic features considered evidence of OA [19] (Table 1); however, they never explicitly specified which radiographic features correspond to which grade. Years after their original article, Grade 2 was changed to “the presence of definite osteophytes with minimal joint space narrowing” [20]. After that, Grade 2 was changed again to “definite osteophytes but the joint space is unimpaired” [23]. These conflicting descriptions, although they were not present in the initial paper or made by its original authors, have nonetheless led to substantial confusion among investigators and inconsistent application.

Schiphof et al. [30] examined epidemiological cohort studies between 1966 and 2006 that incorporated the original KL criteria and found five different descriptions of the KL grading system. Interestingly, some cohort studies contained inconsistencies within the same article. They later examined the impact of these differences in descriptions of the KL criteria [31] by having two trained readers

examine the weightbearing, extended, AP knee radiographs of 3071 people using the five unique descriptions. The authors calculated reproducibility, agreement using the κ statistic, and association with patient-reported knee complaints. They determined that the reproducibility of three of the descriptions was “good” (weighted $\kappa = 0.66, 0.69, 0.63$), whereas the reproducibility of the original description and a fifth description was “moderate” and “poor” (weighted $\kappa = 0.41$ and 0.35 , respectively). The authors determined the agreement between the original KL criteria and the four alternatives to be “moderate” (weighted κ approximately 0.50). The correlation to patient-reported knee complaints was the strongest with the original description (odds ratio [OR], 2.2 [95% CI, 1.9 – 2.7], 4.3 [95% CI, 3.2 – 5.7], 18.3 [95% CI, 7.1 – 47.2] at Grade > 1 , Grade > 2 , Grade > 3 , respectively); however, the differences in OR between the original and alternative descriptions were small and sometimes not significant [31]. Additionally, the authors admit that patient-reported knee complaints could be influenced by the presence of patellofemoral OA, which is not taken into account with the KL system and must be contextualized when interpreting this data. The authors conclude their article with recommendations for using the original description to differentiate patients categorized as Grade > 2 (definite/mild OA) from patients categorized as Grade < 2 (none/possible OA) and to use several of the alternative descriptions to differentiate patients categorized as Grade 0 (no OA) from patients categorized as Grade 1 (possible OA). Given these results, it is clear that discrepancies in the description and application of the KL system are present in the literature; however, the exact impact of these differences on patient care is unclear and requires further investigation.

Conclusions and Uses

Although the KL system has limitations, it remains widely used in clinical practice and in research. Like any radiographic classification tool, the KL system is used ideally in conjunction with a thorough clinical assessment. Altman et al. [1] proposed criteria combining the medical history, physical examination, laboratory as well as radiographic tests to diagnose knee OA, an approach that provides a more comprehensive assessment of a patient’s disease state in comparison to the KL system alone.

Radiographic classification systems like KL seek to standardize the interpretation of studies that many clinicians order during an initial assessment of a patient presenting with clinical findings suggestive of knee OA. In their original paper, Kellgren and Lawrence intended to create a standard reference for the radiographic evaluation of OA for the purposes of field surveys and clinical trials.

Although the KL system has been validated with respect to inter- and intraobserver reliability and a recent article suggests high diagnostic accuracy [40], further research applications should likely focus on the development of treatment algorithms based on classification grade. Such algorithms may better guide clinical decision-making through an evidence-based approach.

References

- Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, Christy W, Cooke TD, Greenwald R, Hochberg M, et al. Development of criteria for the classification and reporting of osteoarthritis. Classification of osteoarthritis of the knee. Diagnostic and Therapeutic Criteria Committee of the American Rheumatism Association. *Arthritis Rheum.* 1986;29:1039–1049.
- Altman RD, Gold GE. Atlas of individual radiographic features in osteoarthritis, revised. *Osteoarthritis Cartilage.* 2007;15(Suppl A):A1–56.
- Bagge E, Bjelle A, Valkenburg HA, Svanborg A. Prevalence of radiographic osteoarthritis in two elderly European populations. *Rheumatol Int.* 1992;12:33–38.
- Brandt KD, Fife RS, Braunstein EM, Katz B. Radiographic grading of the severity of knee osteoarthritis: relation of the Kellgren and Lawrence grade to a grade based on joint space narrowing, and correlation with arthroscopic evidence of articular cartilage degeneration. *Arthritis Rheum.* 1991;34:1381–1386.
- Braun HJ, Gold GE. Diagnosis of osteoarthritis: imaging. *Bone.* 2012;51:278–288.
- Brooks P, Hochberg M. Outcome measures and classification criteria for the rheumatic diseases. A compilation of data from OMERACT (Outcome Measures for Arthritis Clinical Trials), ILAR (International League of Associations for Rheumatology), regional leagues and other groups. *Rheumatology (Oxford).* 2001;40:896–906.
- Creamer P, Hochberg MC. Osteoarthritis. *Lancet.* 1997;350:503–508.
- Emrani PS, Katz JN, Kessler CL, Reichmann WM, Wright EA, McAlindon TE, Losina A. Joint space narrowing and Kellgren-Lawrence progression in knee osteoarthritis: an analytic literature synthesis. *Osteoarthritis Cartilage.* 2008;16:873–882.
- Fairbank TJ. Knee joint changes after meniscectomy. *J Bone Joint Surg Br.* 1948;30:664–670.
- Felson DT, Naimark A, Anderson J, Kazis L, Castelli W, Meenan RF. The prevalence of knee osteoarthritis in the elderly. The Framingham Osteoarthritis Study. *Arthritis Rheum.* 1987;30:914–918.
- Galli M, De Santis V, Tafuro L. Reliability of the Ahlbäck classification of knee osteoarthritis. *Osteoarthritis Cartilage.* 2003;11:580–584.
- Gossec L, Jordan JM, Mazza SA, Lam MA, Suarez-Almazor ME, Renner JB, Lopez-Olivo MA, Hawker G, Dougados M, Maillefert JF; OARSI-OMERACT task force. 'total articular replacement as outcome measure in OA.' Comparative evaluation of three semi-quantitative radiographic grading techniques for knee osteoarthritis in terms of validity and reproducibility in 1759 X-rays: report of the OARSI-OMERACT task force. *Osteoarthritis Cartilage.* 2008;16:742–748.
- Günther KP, Sun Y. Reliability of radiographic assessment in hip and knee osteoarthritis. *Osteoarthritis Cartilage.* 1999;7:239–246.
- Hefti F, Müller W, Jakob RP, Stäubli HU. Evaluation of knee ligament injuries with the IKDC form. *Knee Surg Sports Traumatol Arthrosc.* 1993;1:226–234.
- Irrgang JJ, Anderson AF, Boland AL, Harner CD, Neyret P, Richmond JC, Shelbourne KD. Responsiveness of the International Knee Documentation Committee Subjective Knee Form. *Am J Sports Med.* 2006;34:1567–1573.
- Johnson VL, Hunter DJ. The epidemiology of osteoarthritis. *Best Pract Res Clin Rheumatol.* 2014;28:5–15.
- Kellgren JH. Radiological signs of rheumatoid arthritis; a study of observer differences in the reading of hand films. *Ann Rheum Dis.* 1956;15:55–60.
- Kellgren JH, Lawrence JS. Rheumatism in miners. II. X-ray study. *Br J Ind Med.* 1952;9:197–207.
- Kellgren JH, Lawrence JS. Radiological assessment of osteoarthritis. *Ann Rheum Dis.* 1957;16:494–502.
- Kellgren JH, Lawrence JS. *The Epidemiology of Chronic Rheumatism. Atlas of Standard Radiographs.* Vol 2. Oxford, UK: Blackwell Scientific; 1963.
- Kijowski R, Blankenbaker D, Stanton P, Fine J, De Smet A. Arthroscopic validation of radiographic grading scales of osteoarthritis of the tibiofemoral joint. *AJR Am J Roentgenol.* 2006;187:794–799.
- Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am.* 2007;89:780–785.
- Lawrence JS. *Rheumatism in Populations.* London, UK: Heinemann; 1977:99–100.
- Lequesne M. Clinical features, diagnostic criteria, functional assessments and radiological classifications of osteoarthritis (excluding the spine). *Rheumatology.* 1982;7:1–10.
- Menkes CJ. Radiographic criteria for classification of osteoarthritis. *J Rheumatol Suppl.* 1991;27:13–15.
- Nwachukwu BU, McCormick F, Provencher MT, Roche M, Rubash HE. A comprehensive analysis of Medicare trends in utilization and hospital economics for total knee and hip arthroplasty from 2005 to 2011. *J Arthroplasty.* 2015;30:15–18.
- Roemer FW, Crema MD, Trattinig S, Guermazi A. Advances in imaging of osteoarthritis and cartilage. *Radiology.* 2011;260:332–354.
- Rosenberg TD, Paulos LE, Parker RD, Coward DB, Scott SM. The forty-five-degree posteroanterior flexion weight-bearing radiograph of the knee. *J Bone Joint Surg Am.* 1988;70:1479–1483.
- Scheller G, Sobau C, Bülow JU. Arthroscopic partial lateral meniscectomy in an otherwise normal knee: Clinical, functional, and radiographic results of a long-term follow-up study. *Arthroscopy.* 2001;17:946–952.
- Schiphof D, Boers M, Bierma-Zeinstra SM. Differences in descriptions of Kellgren and Lawrence grades of knee osteoarthritis. *Ann Rheum Dis.* 2008;67:1034–1036.
- Schiphof D, de Klerk BM, Kerkhof HJ, Hofman A, Koes BW, Boers M, Bierma-Zeinstra SM. Impact of different descriptions of the Kellgren and Lawrence classification criteria on the diagnosis of knee osteoarthritis. *Ann Rheum Dis.* 2011;70:1422–1427.
- Schroeder-Boersch H, Töws P, Jani L. [Reproducibility of radiologic markers of osteoarthritis. Evaluating a score for radiologic classification of degenerative changes in osteoarthritis of the knee joint] [in German]. *Z Orthop Ihre Grenzgeb.* 1998;136:293–297.
- Scott WW, Lethbridge-Cejku M, Reichle R, Wigley FM, Tobin JD, Hochberg MC. Reliability of grading scales for individual radiographic features of osteoarthritis of the knee. The Baltimore longitudinal study of aging atlas of knee osteoarthritis. *Invest Radiol.* 1993;28:497–501.

34. Spector T, Cooper C, Cushnighan J, Hart DJ, Dieppe PA. *A Radiographic Atlas of Knee Osteoarthritis*. London, UK: Springer Verlag; 1992:1–15.
35. Spector TD, Cooper C. Radiographic assessment of osteoarthritis in population studies: whither Kellgren and Lawrence? *Osteoarthritis Cartilage*. 1993;1:203–206.
36. Tapper EM, Hoover NW. Late results after meniscectomy. *J Bone Joint Surg Am*. 1969;51:517–526, passim.
37. TMMP 311–Knee Arthroplasty–Total & Partial. Triad Healthcare Musculoskeletal Health Services website. Available at: <http://www.triadhealthcareinc.com/providers/PDFview.aspx?fld=Cur%5CCurrent%20Policies&fn=TMMP%20311%20-%20Knee%20Arthroplasty%20-%20Total%20and%20Partial.pdf>. February 20, 2015. Accessed December 3, 2015.
38. Total knee and total hip replacement bundle. The Bree Collaborative website. Available at: http://www.breecollaborative.org/wp-content/uploads/tkrthr_bundle.pdf. November 21, 2013. Accessed December 3, 2015.
39. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S, Aboyans V, Abraham J, Ackerman I, Aggarwal R, Ahn SY, Ali MK, Alvarado M, Anderson HR, Anderson LM, Andrews KG, Atkinson C, Baddour LM, Bahalim AN, Barker-Collo S, Barrero LH, Bartels DH, Basáñez MG, Baxter A, Bell ML, Benjamin EJ, Bennett D, Bernabé E, Bhalla K, Bhandari B, Bikbov B, Bin Abdulhak A, Birbeck G, Black JA, Blencowe H, Blore JD, Blyth F, Bolliger I, Bonaventure A, Boufous S, Bourne R, Boussinesq M, Braithwaite T, Brayne C, Bridgett L, Brooker S, Brooks P, Brugha TS, Bryan-Hancock C, Bucello C, Buchbinder R, Buckle G, Budke CM, Burch M, Burney P, Burstein R, Calabria B, Campbell B, Canter CE, Carabin H, Carapetis J, Carmona L, Cella C, Charlson F, Chen H, Cheng AT, Chou D, Chugh SS, Coffeng LE, Colan SD, Colquhoun S, Colson KE, Condon J, Connor MD, Cooper LT, Corriere M, Cortinovis M, de Vaccaro KC, Couser W, Cowie BC, Criqui MH, Cross M, Dabhadkar KC, Dahiya M, Dahodwala N, Damsere-Derry J, Danaei G, Davis A, De Leo D, Degenhardt L, Dellavalle R, Delossantos A, Denenberg J, Derrett S, Des Jarlais DC, Dharmaratne SD, Dherani M, Diaz-Torne C, Dolk H, Dorsey ER, Driscoll T, Duber H, Ebel B, Edmond K, Elbaz A, Ali SE, Erskine H, Erwin PJ, Espindola P, Ewoigbokhan SE, Farzadfar F, Feigin V, Felson DT, Ferrari A, Ferri CP, Fèvre EM, Finucane MM, Flaxman S, Flood L, Foreman K, Forouzanfar MH, Fowkes FG, Franklin R, Fransen M, Freeman MK, Gabbe BJ, Gabriel SE, Gakidou E, Ganatra HA, Garcia B, Gaspari F, Gillum RF, Gmel G, Gosselin R, Grainger R, Groeger J, Guillemin F, Gunnell D, Gupta R, Haagsma J, Hagan H, Halasa YA, Hall W, Haring D, Haro JM, Harrison JE, Havmoeller R, Hay RJ, Higashi H, Hill C, Hoen B, Hoffman H, Hotez PJ, Hoy D, Huang JJ, Ibeanusi SE, Jacobsen KH, James SL, Jarvis D, Jasrasaria R, Jayaraman S, Johns N, Jonas JB, Karthikeyan G, Kassebaum N, Kawakami N, Keren A, Khoo JP, King CH, Knowlton LM, Kobusingye O, Koranteng A, Krishnamurthi R, Laloo R, Laslett LL, Lathlean T, Leasher JL, Lee YY, Leigh J, Lim SS, Limb E, Lin JK, Lipnick M, Lipshultz SE, Liu W, Loane M, Ohno SL, Lyons R, Ma J, Mabweijano J, MacIntyre MF, Malekzadeh R, Mallinger L, Manivannan S, Marcenes W, March L, Margolis DJ, Marks GB, Marks R, Matsumori A, Matzopoulos R, Mayosi BM, McAnulty JH, McDermott MM, McGill N, McGrath J, Medina-Mora ME, Meltzer M, Mensah GA, Merriman TR, Meyer AC, Miglioli V, Miller M, Miller TR, Mitchell PB, Mocumbi AO, Moffitt TE, Mokdad AA, Monasta L, Montico M, Moradi-Lakeh M, Moran A, Morawska L, Mori R, Murdoch ME, Mwaniki MK, Naidoo K, Nair MN, Naldi L, Narayan KM, Nelson PK, Nelson RG, Nevitt MC, Newton CR, Nolte S, Norman P, Norman R, O'Donnell M, O'Hanlon S, Olives C, Omer SB, Ortblad K, Osborne R, Ozgediz D, Page A, Pahari B, Pandian JD, Rivero AP, Patten SB, Pearce N, Padilla RP, Perez-Ruiz F, Perico N, Pesudovs K, Phillips D, Phillips MR, Pierce K, Pion S, Polanczyk GV, Polinder S, Pope CA 3rd, Popova S, Porrini E, Pourmalek F, Prince M, Pullan RL, Ramaiah KD, Ranganathan D, Razavi H, Regan M, Rehm JT, Rein DB, Remuzzi G, Richardson K, Rivara FP, Roberts T, Robinson C, De León FR, Ronfani L, Room R, Rosenfeld LC, Rushton L, Sacco RL, Saha S, Sampson U, Sanchez-Riera L, Sanman E, Schwebel DC, Scott JG, Segui-Gomez M, Shahraz S, Shepard DS, Shin H, Shivakoti R, Singh D, Singh GM, Singh JA, Singleton J, Sleet DA, Sliwa K, Smith E, Smith JL, Stapelberg NJ, Steer A, Steiner T, Stolk WA, Stovner LJ, Sudfeld C, Syed S, Tamburlini G, Tavakkoli M, Taylor HR, Taylor JA, Taylor WJ, Thomas B, Thomson WM, Thurston GD, Tleyjeh IM, Tonelli M, Towbin JA, Truelsen T, Tsilimbaris MK, Ubeda C, Undurraga EA, van der Werf MJ, van Os J, Vavilala MS, Venketasubramanian N, Wang M, Wang W, Watt K, Weatherall DJ, Weinstock MA, Weintraub R, Weisskopf MG, Weissman MM, White RA, Whiteford H, Wiersma ST, Wilkinson JD, Williams HC, Williams SR, Witt E, Wolfe F, Woolf AD, Wulf S, Yeh PH, Zaidi AK, Zheng ZJ, Zonies D, Lopez AD, Murray CJ, AlMazroa MA, Memish ZA. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2163–2196.
40. Wright RW. Osteoarthritis classification scales: interobserver reliability and arthroscopic correlation. *J Bone Joint Surg Am*. 2014;96:1145–1151.