



2016 Laparoscopic Surgery: Global view

Preoperative, intraoperative and postoperative risk factors for anastomotic leakage after laparoscopic low anterior resection with double stapling technique anastomosis

Kenji Kawada, Yoshiharu Sakai

Kenji Kawada, Yoshiharu Sakai, Department of Surgery, Graduate School of Medicine, Kyoto University, Kyoto 606-8507, Japan

Author contributions: Kawada K wrote the paper; Sakai Y contributed critical revision of the manuscript for important intellectual content.

Conflict-of-interest statement: No potential conflicts of interest exist.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Kenji Kawada, MD, PhD, Department of Surgery, Graduate School of Medicine, Kyoto University, 54 Shogoin-Kawara-cho, Sakyo-ku, Kyoto 606-8507, Japan. kkawada@kuhp.kyoto-u.ac.jp
Telephone: +81-75-3667595
Fax: +81-75-3667642

Received: April 7, 2016
Peer-review started: April 8, 2016
First decision: May 27, 2016
Revised: May 30, 2016
Accepted: June 13, 2016
Article in press: June 13, 2016
Published online: July 7, 2016

Abstract

Anastomotic leakage (AL) is one of the most devastating

complications after rectal cancer surgery. The double stapling technique has greatly facilitated intestinal reconstruction especially for anastomosis after low anterior resection (LAR). Risk factor analyses for AL after open LAR have been widely reported. However, a few studies have analyzed the risk factors for AL after laparoscopic LAR. Laparoscopic rectal surgery provides an excellent operative field in a narrow pelvic space, and enables total mesorectal excision surgery and preservation of the autonomic nervous system with greater precision. However, rectal transection using a laparoscopic linear stapler is relatively difficult compared with open surgery because of the width and limited performance of the linear stapler. Moreover, laparoscopic LAR exhibits a different postoperative course compared with open LAR, which suggests that the risk factors for AL after laparoscopic LAR may also differ from those after open LAR. In this review, we will discuss the risk factors for AL after laparoscopic LAR.

Key words: Risk factor; Laparoscopic low anterior resection; Anastomotic leakage

© **The Author(s) 2016.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Recently, many studies have reported that laparoscopic rectal surgery is becoming popular and exhibits favorable outcomes compared with open surgery. However, the anastomotic leakage (AL) rate after laparoscopic low anterior resection (LAR) is yet about 10%, and AL remains a huge challenge despite many surgical and technological advances. Here we review the current literature published with respect to the risk factors for AL after laparoscopic LAR.

Kawada K, Sakai Y. Preoperative, intraoperative and postoperative risk factors for anastomotic leakage after laparoscopic low

anterior resection with double stapling technique anastomosis. *World J Gastroenterol* 2016; 22(25): 5718-5727 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v22/i25/5718.htm> DOI: <http://dx.doi.org/10.3748/wjg.v22.i25.5718>

INTRODUCTION

Laparoscopic surgery for colon cancer was introduced in the 1990s, and has shown promising results. Laparoscopic low anterior resection (LAR) for rectal cancer is technically more difficult than laparoscopic colectomy because of the difficulties related to a narrow pelvic space. A higher incidence of positive circumferential margins after laparoscopic LAR was reported in an initial randomized controlled trial (RCT)^[1], but an increasing number of studies have shown that laparoscopic surgery for rectal cancer provides surgical safety and oncological outcomes equivalent to open surgery^[2-6]. Recent large-scale RCTs such as COLOR II^[7] and COREAN^[8] have reported favorable outcomes for laparoscopic surgery compared with open surgery for rectal cancer.

The double stapling technique (DST) has greatly facilitated intestinal reconstruction, especially for anastomosis after LAR. Anastomotic leakage (AL) is one of the most devastating complications after rectal cancer surgery. AL impairs not only short-term outcomes (morbidity, mortality, length of hospital stay, and financial cost) but also long-term oncological outcomes (survival and local recurrence)^[9-11]. Therefore, it is important to identify the patients who are at high risk of AL for improving overall outcomes. Despite technical improvements and instrumental developments, recent studies have reported that the AL rate ranges from 3% to 19%^[9,12-15]; the most commonly reported rate is approximately 10%-13% from recent large population databases in the United States^[12] and Japan^[15]. AL after rectal resection is influenced by many factors including not only surgical factors but also medical factors related to the systemic conditions in patients.

Several risk factors, including age, sex, intraoperative bleeding, obesity, preoperative chemoradiotherapy, protective diverting stoma, pelvic drainage, tumor size, tumor location and the level of anastomosis have been reported to be risk factors for AL after open LAR^[16-21]. In contrast, only a few studies have examined risk factors for AL after laparoscopic LAR^[22-31] (Table 1). In addition, the rates of protective diverting stoma, preoperative chemoradiotherapy (CRT), and total mesorectal excision (TME) in each study were not consistent, which might produce different results. Several studies reported that laparoscopic surgery and open surgery for rectal cancer did not differ in terms of the AL rate^[1,2,4]. Laparoscopic rectal surgery provides an excellent operative field in a narrow pelvic space, and enables

the preservation of autonomic nerves more precisely. However, rectal transection using a laparoscopic linear stapler is relatively difficult when compared with open surgery because of the width and limited performance of the linear stapler. The devices and techniques used for laparoscopic LAR are different from those used for open LAR. Moreover, laparoscopic LAR exhibits a different postoperative course compared with open LAR, including less blood loss, faster recovery of peristalsis, faster initiation of oral intake, and shorter hospital stay. Notably, multicenter, prospective and cohort studies using propensity score matching analysis have reported that risk factors for AL after laparoscopic or robotic LAR are different from those after open LAR^[30,31]. Factors related to technical difficulty such as male sex, previous abdominal surgery, lower location of tumor and the use of more than 2 cartridges for rectal transection were found to be significant only in laparoscopic or robotic LAR groups^[31].

In this review, we will discuss the risk factors for AL after laparoscopic LAR. Risk factors are categorized into (1) preoperative; (2) intraoperative; and (3) postoperative factors. The identification of high-risk patients has great clinical relevance and ultimately improves patient outcomes. Although more prospective studies are needed, this review provides major insight into identifying important risk factors for AL after laparoscopic LAR.

PREOPERATIVE RISK FACTORS

Male gender

Males have a narrow pelvis, which makes rectal dissection and anastomosis more difficult and more prone to surgical complications. In fact, male gender has been reported as an increased risk factor for AL after open LAR^[16,17,19-21] as well as laparoscopic LAR^[28,30,31]. The influence of androgen-related differences in the intestinal microcirculation may be involved^[32].

Body mass index

Some studies have shown that obesity measured by body mass index (BMI) can increase the risk of AL^[33-35]. Yamamoto *et al.*^[27] reported that BMI was independently predictive for AL after laparoscopic LAR. In place of BMI, waist circumference and waist/hip ratio may predict the risk of AL^[36]. In addition, measuring visceral fat area may be more sensitive than BMI in predicting AL after laparoscopic surgery^[37].

Preoperative radiotherapy/chemoradiotherapy

Preoperative radiotherapy (RT) with or without concomitant chemotherapy is generally recommended for patients with locally advanced rectal cancer followed by TME surgery. It is accepted that these therapeutic modalities can reduce the local recurrence

Table 1 Selected studies to investigate the risk factors for Anastomotic leakage after laparoscopic low anterior resection

Ref.	Year	Sample size	AL rate	Tumor Location ¹	Covering stoma	Risk factors
Ito <i>et al</i> ^[22]	2008	180	5.0%	R, RS	+	Anastomosis level, multiple stapler firings
Kim <i>et al</i> ^[23]	2009	270	6.3%	R, RS, S	+	Tumor location
Huh <i>et al</i> ^[24]	2010	223	8.5%	R	-	Tumor location, operation time
Choi <i>et al</i> ^[25]	2010	156	10.3%	R, RS	-	Anastomosis level, operation time
Akiyoshi <i>et al</i> ^[26]	2011	363	3.6%	R, RS	+	Tumor location, abdominal drain
Yamamoto <i>et al</i> ^[27]	2012	111	5.4%	R	+	BMI
Park <i>et al</i> ^[28]	2013	1187	6.3%	R, RS	-	Male, stage, transfusion, tumor location preoperative CRT, multiple stapler firings
Kawada <i>et al</i> ^[29]	2014	154	12.3%	R	-	Tumor size, precompression before stapler firings
Katsuno <i>et al</i> ^[30]	2015	209	15.3%	R	+	Male
Kim <i>et al</i> ^[31]	2016	1154	6.7%	R	+	Male, smoking, alcohol intake, previous abdominal surgery, operation time, tumor location, multiple stapler firings

¹R: Rectum; RS: Rectosigmoid colon; S: Sigmoid colon.

rate^[38-40]. Although effective in targeting cancer cells, RT has a wide array of detrimental effects on intestinal tissue and wound healing and has long been believed to be a risk factor for AL. There are many retrospective studies that have reported the relationship between preoperative RT and AL^[20,21,28]. However, prospective trials and cohort studies have shown contradictory results. The MRC CR07 RCT^[41] reported that there was no difference in AL between preoperative RT and selective postoperative CRT. A Dutch TME trial^[42] reported that there was no significant difference in AL rates (TME plus preoperative RT vs TME alone). A recent report using propensity score matching analysis have also reported that preoperative CRT does not increase the risk of AL after LAR^[43]. Most surgeons perform a temporary protective diverting stoma to minimize the consequences of AL in patients who have received preoperative CRT or RT.

Preoperative chemotherapy

Preoperative chemotherapy is a well-known risk factor for AL^[13]; however, the mechanism underlying this association is poorly understood. Recent use of antiangiogenic agents also increases the risk of AL. The first studies examining bevacizumab (Avastin), a humanized anti-vascular endothelial growth factor antibody, reported several patients with bowel perforation^[44,45]. The mechanism of this perforation is proposed to be arterial microthromboembolic disease leading to bowel ischemia. The same mechanism can cause AL. Bevacizumab has a half-life of 20 days, and the manufacturer recommends stopping its treatment at least 4 wk before surgery.

Antibiotics

A meta-analysis of eight RCTs reported that combining preoperative intestinal decontamination with oral antibiotics and perioperative intravenous antibiotics reduced postoperative infection including AL, com-

pared with use of intravenous antibiotics alone^[46]. Notably, a recent RCT showed that intravenous plus oral antibiotics (cefmetazole, kanamycin and metronidazole) significantly reduced the risk of surgical site infection (SSI) compared with intravenous antibiotics alone (7.3% vs 12.8%, $P = 0.028$), while no significant difference was seen in the rate of AL^[47]. Further studies are required to elucidate the effect of preoperative oral antibiotics on AL.

Medications

Although it is assumed that impaired healing with corticosteroid use would affect the AL rate, it is difficult to find an absolute correlation. Prolonged use of corticosteroids can be a risk factor for AL, particularly when combined with other immunosuppressive drugs^[48-50]. A recent systematic review reported that the AL rate after lower gastrointestinal surgery was 6.8% in the corticosteroid group compared with 3.3% in the non-corticosteroid group, although the duration and dose of corticosteroid treatment were heterogeneous^[51]. A meta-analysis with six RCTs reported that perioperative use of nonsteroidal anti-inflammatory drugs (NSAIDs) had no statistically significant effect on the AL rate^[52]. However, non-selective NSAIDs and non-selective cyclooxygenase (COX) 2 inhibitors were reported to be associated with a higher AL rate^[53]. Therefore, NSAIDs should be used with caution in the postoperative period. In general, the postoperative pain after laparoscopic surgery is less than that after open surgery, which may result in the decreased usage of NSAIDs and decreased rate of AL in laparoscopic surgery.

Other factors, such as smoking and alcohol, have also been reported to be risk factors for AL after LAR^[31,54-57]. The effect of smoking might be secondary to ischemia caused by smoking-related microvascular disease. Large quantities of alcohol consumption might be a surrogate for poor nutritional status.

Mechanical bowel preparation

Mechanical bowel preparation (MBP) is performed before colorectal surgery to reduce massive bowel contents, which can be a source of colorectal AL and infectious bacterial pathogens. However, the routine use of MBP is being abandoned gradually, because some RCT studies and meta-analyses have concluded that omitting MBP before surgery has fewer postoperative morbidities including AL and SSI^[58-61]. The practice of omitting MBP is further promoted because MBP causes some discomforts to patients, such as nausea, vomiting, dehydration and electrolyte abnormalities. However, recent some studies from the United States databases have reported that combining MBP and oral antibiotics results in a significantly lower incidence of AL, incisional SSI and hospital readmission compared with no preoperative bowel preparation in colorectal surgery^[62-64]. Moreover, regarding the long-term effect of MBP, the 10-year cancer-specific survival rate was recently reported to be significantly better in MBP group than in non-MBP group^[65,66]. Further studies are required to elucidate the effect of MBP on AL.

INTRAOPERATIVE RISK FACTORS

Level of anastomosis

It is widely accepted that the risk of AL increases with more distal anastomosis. Although it is well accepted that a low anastomosis has a higher incidence of AL, the mechanism remains unknown. It is hypothesized that the height of the anastomosis or the tumor location can reflect technical difficulties of LAR, resulting in local tissue trauma, increased tension, or poor blood supply. A number of studies reported that lower anastomosis level is an important risk factor for AL after open LAR^[16-21] as well as laparoscopic LAR^[23-26,28,31].

Surgical technique and multiple stapler firings

Surgical technique has a substantial impact on post-operative complications including AL. In laparoscopic LAR, optimal port placement is important to reduce the number of linear stapler firings for rectal transection in a narrow pelvis. The use of multiple staplers (*e.g.*, ≥ 3 cartridges) for rectal transection is a major cause of AL after laparoscopic LAR^[22,23,26,28,29,31]. When the number of stapler cartridges increases, there is a concern that an increased number of stapler firings can lead to small defects between the staple lines and, in turn, cause AL. Therefore, laparoscopic surgeons need to make efforts to reduce the number of linear stapler firings to two or less. Several different techniques have been proposed to reduce AL. Ito *et al.*^[22] reported that vertical rectal transection through an additional suprapubic site was useful for avoiding multiple stapler firings and decreasing the AL rate. Kuroyanagi *et al.*^[67] reported that rectal transection was performed using two cartridges in most cases, with coordinated

operator-assistant movement, and that removal of the crossing point of staple lines was important to delete the potential source of AL. In a clinical setting, we previously analyzed whether the remnant crossing point could increase the AL rate, and found that it was not significantly associated with AL^[29]. Therefore, we assume that surgeons do not need to persist in removal of the crossing point, especially when the crossing point is placed near the edge of the rectal stump and so removal of the crossing point is technically difficult.

It is notable that intracorporeal^[68] or transanal^[69] reinforcing sutures could effectively reduce AL after LAR, but the results of these studies are not conclusive. DST is inevitably associated with bilateral intersecting staple lines at the rectal stump, so-called dog ears. The dog ears are the weak spots associated with potential AL^[70]. Recently, a combined laparoscopic LAR and eversion technique without dog ear formation was reported to be useful to reduce AL for mid and distal rectal cancer^[71].

Precompression before stapler firings

We previously reported that a sufficient amount of precompression time before stapler firings resulted in reduced intestinal wall thickness and proper staple formation in animal models^[72-74]. In addition, we recently reported that precompression before stapler firings and tumor size (≥ 5.0 cm) were associated with AL after laparoscopic LAR in a clinical setting, and that precompression before stapler firing tended to reduce the AL occurring in the early postoperative period^[29]. Precompression time before stapler firings and proper cartridge selection according to the wall thickness are critical to achieve secure staple formation.

Diameter of circular stapler

Kim *et al.*^[23] reported the association between a larger diameter circular stapler and increased rates of AL. They speculated that a larger diameter circular stapler made the distal rectum more distended. A distended rectum with thinned rectal wall may cause inadequate blood supply to the anastomosis site. We previously analyzed whether the diameter of a circular stapler could affect the AL rate, and found that it was not significantly associated with AL^[29]. Further studies are required to elucidate the effect of diameter of a circular stapler.

Tumor characteristics

Tumor size is a well-known risk factor for AL after laparoscopic LAR^[13,29,30]. A bulky tumor could adversely affect the ease of rectal transection and anastomosis in the limited pelvic space. Some studies demonstrated that tumor size greater than 5 cm was independently predictive of AL^[13,29]. Advanced stage is also a risk factor for AL after laparoscopic LAR^[28].

Blood supply

Despite the multifactorial etiology of AL, insufficient perfusion and technical factors are considered to play a substantial role in the development of AL^[9,75-77]. For this reason, surgeons often assess intestinal perfusion by several clinical checks, such as the color of the bowel wall, palpable pulsation, and bleeding from marginal arteries. These checks are subjective and based on the surgeon's experience, and may well lead to misinterpretations even by experienced surgeons^[78]. In recent years, near-infrared (NIR) fluorescence technology with indocyanine green (ICG) has been the most promising method that provides a real-time assessment of intestinal perfusion^[79-83]. The first study to use fluorescence imaging for colorectal surgery was published by Kudzus *et al.*^[79]. They reported that fluorescence imaging resulted in a proximal change of the initially planned transection line in 13.9% (28/201), and that intraoperative fluorescence imaging reduced AL by 4% compared with a control group (7.5% vs 3.5%). These data have been confirmed by Jafari *et al.*^[80] during robotic-assisted laparoscopic rectal surgery. Moreover, a multi-institutional prospective study, PILLAR-II, recently reported that fluorescence imaging changed surgical plans in 8% (11/139), and that the AL rate was 1.4% (2/139) in laparoscopic left-sided/anterior resection^[82]. In addition, Sherwinter *et al.*^[84] evaluated the intraluminal aspect of the anastomosis transanally after DST construction using a transanal NIR imaging system. The assessment of rectal stump perfusion by transanal ICG imaging can be a promising method, although further studies are needed to correlate this technique to the clinical outcome. However, another recent another report stated that the intraoperative fluorescence imaging does not reduce the AL rate in colorectal surgery from a case-matched retrospective study with the use of historical control subjects^[85]. Because of the limited number of patients and the likely multifactorial nature of AL, it is hard to draw robust conclusions concerning the beneficial effect of fluorescence imaging on the AL rate.

The concept of high ligation of the inferior mesenteric artery to achieve optimal oncological results suppresses the vascular supply from the left colic artery, and vascularization of the proximal colon is dependent on marginal vessels from the middle colic artery. The preservation of the left colonic artery in laparoscopic LAR was reported to be associated with lower risk of AL^[86]. With the progressive increase in the aging population, vascular disease can also be a factor contributing to insufficient blood supply, even in the case of low ligation.

Blood loss

Blood loss greater than 100 mL and blood transfusion are independent risk factors for AL^[28-30,87,88], but it is unclear whether this is a specific manifestation due to

blood loss or whether blood loss is a surrogate for poor operative technique or challenging surgery.

Operation time

Although operation time is well known to be one of the risk factors for AL after laparoscopic LAR^[23-25,28,29,31], the experienced skill of the surgeon is also thought to act as a confounding variable. In patients with severe obesity, narrow pelvis, bulky tumor, and in cases with adverse intraoperative events, the operation time is prolonged. When the operation time is long, bacterial exposure and tissue damage can increase, which may cause inflammation and ultimately increase AL.

Anastomotic tension

Many surgeons assume that sufficient mobilization of the splenic flexure is necessary to lower anastomotic tension, especially when the anastomotic site is very low. Minimal anastomotic tension is thought to be one of the requirements of proper surgical technique; yet, this concept remains largely hypothetical. To our knowledge, there have been no experimental studies investigating the role of tension during an intestinal anastomosis. Lack of data likely stems from the difficulty in designing studies that investigate anastomotic tension in a clinical setting.

POSTOPERATIVE RISK FACTORS

Diverting stoma

Fecal diversion is one of the most widely used methods to prevent AL. However, there is still debate as to whether the creation of a diverting stoma (DS) can reduce AL. Some randomized controlled trials reported that DS could reduce the rate of symptomatic AL^[89,90], while a recent large multicenter cohort study using propensity score matching analysis indicated that DS was not associated with symptomatic AL^[14]. A considerable number of retrospective studies also described the beneficial effect of DS on AL^[16,20,91], while some studies stated that the creation of a DS did not reduce the rate of AL^[21,92]. It is generally agreed that the creation of a DS can reduce the incidence of the severe complications of AL, including fecal peritonitis and septicemia. We need to bear in mind that even a temporarily-intended stoma can induce dehydration and renal impairment^[93]. Moreover, re-operation for reversal of stoma may also be associated with morbidity and even death^[94].

Transanal drainage tube

The safety and efficacy of transanal drainage tube (TDT) placement to decrease the risk of AL after rectal cancer surgery has not been validated. In theory, TDT decreases the intraluminal pressure around the anastomotic site, and protects the anastomosis from watery stool and flatus when intestinal motility improves. There are only a few reports to investigate

whether TDT can prevent AL after open LAR, but the results are inconsistent, with some studies indicating favorable outcomes^[95-97], while other studies reported unfavorable outcomes^[98]. Moreover, it has been reported that TDT can reduce the rate of AL after laparoscopic LAR^[99]. There are slight differences in each study such as material and diameter of TDT, length of TDT insertion and duration of TDT placement. A standardized procedure for TDT should be validated and further investigation is required to elucidate its usefulness. With regard to reducing the intraluminal pressure around the anastomotic site, the concept of creating a DS is nearly the same as that for TDT. However, a DS increases patient discomfort and overall cost, and requires further surgery for closure of the DS. If the efficacy of prevention of AL is nearly equal for both procedures, it follows that TDT is superior to DS for this reason.

TDT can also be useful to cure localized peritonitis related to AL. Several reports have stated that the TDT is effective for localizing AL and controlling sepsis following LAR^[100,101]. Shrinkage of an abscess by a TDT inserted into the cavity can localize inflammation, which results in a reduced incidence of re-operation.

Abdominal drains

The use of an abdominal drain has been debated widely in terms of early detection of complications as well as preventing AL. After TME surgery, a large presacral space in which a hematoma or seroma may develop constitutes a nidus for bacterial growth, which may extend to the anastomosis and cause AL. Pelvic drainage can prevent this process and help to control AL. A systematic review including several RCTs reported no significant difference in the rate of AL, concluding there was insufficient evidence to support routine drainage^[102]. However, a recent meta-analysis indicated a reduction of AL rate with pelvic drainage^[103]. Akiyoshi *et al*^[26] reported that the presence of an abdominal drain was an independent predictive factor for AL after laparoscopic LAR. The current evidence does not support drainage of a colonic anastomosis, but the LAR case for abdominal drains is less clear.

Intestinal microbes

The human intestinal microbiome is thought to play a key role in the pathogenesis of obesity, gastrointestinal malignancies, and Crohn's disease^[104]. Recently, the role of microflora in anastomotic healing is attracting more attention^[105]. One powerful modality contributing to major alterations in composition and virulence of the gastrointestinal microflora is radiation. The susceptibility to RT-induced diarrhea could be linked to differential initial microbial colonization^[106]. In a rat model of LAR, Olivas *et al*^[107] demonstrated that the combination of preoperative RT and intestinal inoculation with *Pseudomonas aeruginosa*

resulted in a higher rate of AL, whereas radiation alone or *Pseudomonas aeruginosa* alone did not cause AL. In an AL rat model, it has been recently reported that *Enterococcus faecalis* contributes to the pathogenesis of AL through collagen degradation and matrix metalloproteinase 9 (MMP9) activation in host intestinal tissues, and that either elimination of *Enterococcus faecalis* through direct topical antibiotics or pharmacological suppression of MMP9 could prevent AL^[108]. Patients undergoing colectomy are at a unique risk of *Clostridium difficile* because of the additional physical disruption of the colonic microflora. The impact of postoperative *Clostridium difficile* infection is being increasingly reported with overall worse outcome after colon resection^[109]. It was reported that postoperative diarrhea or high stoma output regardless of *Clostridium difficile* infection could increase significantly more superficial surgical site infections including AL^[110], which may indicate the interaction between AL and the intraluminal pressure increased by postoperative diarrhea. Further investigation focusing on intestinal microbes could be important for uncovering the elusive causes of AL.

CONCLUSION

AL remains a huge challenge despite many surgical and technological advances. Our review identified several risk factors for AL after laparoscopic LAR, all of which are readily available in clinical settings. Continued high-quality research is of paramount importance to reduce the risk and subsequent effects associated with AL.

REFERENCES

- 1 **Guillou PJ**, Quirke P, Thorpe H, Walker J, Jayne DG, Smith AM, Heath RM, Brown JM. Short-term endpoints of conventional versus laparoscopic-assisted surgery in patients with colorectal cancer (MRC CLASICC trial): multicentre, randomised controlled trial. *Lancet* 2005; **365**: 1718-1726 [PMID: 15894098 DOI: 10.1016/S0140-6736(05)66545-2]
- 2 **Zhou ZG**, Hu M, Li Y, Lei WZ, Yu YY, Cheng Z, Li L, Shu Y, Wang TC. Laparoscopic versus open total mesorectal excision with anal sphincter preservation for low rectal cancer. *Surg Endosc* 2004; **18**: 1211-1215 [PMID: 15457380 DOI: 10.1007/s00464-003-9170-1]
- 3 **Lelong B**, Bege T, Esterni B, Guiramand J, Turrini O, Moutardier V, Magnin V, Monges G, Pernoud N, Blache JL, Giovannini M, Delpero JR. Short-term outcome after laparoscopic or open restorative mesorectal excision for rectal cancer: a comparative cohort study. *Dis Colon Rectum* 2007; **50**: 176-183 [PMID: 17180257 DOI: 10.1007/s10350-006-0751-7]
- 4 **Lujan J**, Valero G, Hernandez Q, Sanchez A, Frutos MD, Parrilla P. Randomized clinical trial comparing laparoscopic and open surgery in patients with rectal cancer. *Br J Surg* 2009; **96**: 982-989 [PMID: 19644973 DOI: 10.1002/bjs.6662]
- 5 **Milsom JW**, de Oliveira O, Trencheva KI, Pandey S, Lee SW, Sonoda T. Long-term outcomes of patients undergoing curative laparoscopic surgery for mid and low rectal cancer. *Dis Colon Rectum* 2009; **52**: 1215-1222 [PMID: 19571696 DOI: 10.1007/DCR0b013e3181a73e81]
- 6 **Arezzo A**, Passera R, Scozzari G, Verra M, Morino M. Laparoscopy for rectal cancer reduces short-term mortality and morbidity: results

- of a systematic review and meta-analysis. *Surg Endosc* 2013; **27**: 1485-1502 [PMID: 23183871 DOI: 10.1007/s00464-012-2649-x]
- 7 **Bonjer HJ**, Deijen CL, Abis GA, Cuesta MA, van der Pas MH, de Lange-de Klerk ES, Lacy AM, Bemelman WA, Andersson J, Angenete E, Rosenberg J, Fuerst A, Haglund E. A randomized trial of laparoscopic versus open surgery for rectal cancer. *N Engl J Med* 2015; **372**: 1324-1332 [PMID: 25830422 DOI: 10.1056/NEJMoa1414882]
 - 8 **Jeong SY**, Park JW, Nam BH, Kim S, Kang SB, Lim SB, Choi HS, Kim DW, Chang HJ, Kim DY, Jung KH, Kim TY, Kang GH, Chie EK, Kim SY, Sohn DK, Kim DH, Kim JS, Lee HS, Kim JH, Oh JH. Open versus laparoscopic surgery for mid-rectal or low-rectal cancer after neoadjuvant chemoradiotherapy (COREAN trial): survival outcomes of an open-label, non-inferiority, randomised controlled trial. *Lancet Oncol* 2014; **15**: 767-774 [PMID: 24837215 DOI: 10.1016/S1470-2045(14)70205-0]
 - 9 **Kingham TP**, Pachter HL. Colonic anastomotic leak: risk factors, diagnosis, and treatment. *J Am Coll Surg* 2009; **208**: 269-278 [PMID: 19228539 DOI: 10.1016/j.jamcollsurg.2008.10.015]
 - 10 **Branagan G**, Finnis D. Prognosis after anastomotic leakage in colorectal surgery. *Dis Colon Rectum* 2005; **48**: 1021-1026 [PMID: 15789125 DOI: 10.1007/s10350-004-0869-4]
 - 11 **Mirnezami A**, Mirnezami R, Chandrakumaran K, Sasapu K, Sagar P, Finan P. Increased local recurrence and reduced survival from colorectal cancer following anastomotic leak: systematic review and meta-analysis. *Ann Surg* 2011; **253**: 890-899 [PMID: 21394013 DOI: 10.1097/SLA.0b013e3182128929]
 - 12 **Kang CY**, Halabi WJ, Chaudhry OO, Nguyen V, Pigazzi A, Carmichael JC, Mills S, Stamos MJ. Risk factors for anastomotic leakage after anterior resection for rectal cancer. *JAMA Surg* 2013; **148**: 65-71 [PMID: 22986932 DOI: 10.1001/2013.jamasurg.2]
 - 13 **Qu H**, Liu Y, Bi DS. Clinical risk factors for anastomotic leakage after laparoscopic anterior resection for rectal cancer: a systematic review and meta-analysis. *Surg Endosc* 2015; **29**: 3608-3617 [PMID: 25743996 DOI: 10.1007/s00464-015-4117-x]
 - 14 **Shiomi A**, Ito M, Maeda K, Kinugasa Y, Ota M, Yamaue H, Shiozawa M, Horie H, Kuriu Y, Saito N. Effects of a diverting stoma on symptomatic anastomotic leakage after low anterior resection for rectal cancer: a propensity score matching analysis of 1,014 consecutive patients. *J Am Coll Surg* 2015; **220**: 186-194 [PMID: 25529899 DOI: 10.1016/j.jamcollsurg.2014.10.017]
 - 15 **Matsubara N**, Miyata H, Gotoh M, Tomita N, Baba H, Kimura W, Nakagoe T, Simada M, Kitagawa Y, Sugihara K, Mori M. Mortality after common rectal surgery in Japan: a study on low anterior resection from a newly established nationwide large-scale clinical database. *Dis Colon Rectum* 2014; **57**: 1075-1081 [PMID: 25101603 DOI: 10.1097/DCR.0000000000000176]
 - 16 **Peeters KC**, Tollenaar RA, Marijnen CA, Klein Kranenbarg E, Steup WH, Wiggers T, Rutten HJ, van de Velde CJ. Risk factors for anastomotic failure after total mesorectal excision of rectal cancer. *Br J Surg* 2005; **92**: 211-216 [PMID: 15584062 DOI: 10.1002/bjs.4806]
 - 17 **Rullier E**, Laurent C, Garrelon JL, Michel P, Saric J, Parneix M. Risk factors for anastomotic leakage after resection of rectal cancer. *Br J Surg* 1998; **85**: 355-358 [PMID: 9529492 DOI: 10.1046/j.1365-2168.1998.00615.x]
 - 18 **Yeh CY**, Changchien CR, Wang JY, Chen JS, Chen HH, Chiang JM, Tang R. Pelvic drainage and other risk factors for leakage after elective anterior resection in rectal cancer patients: a prospective study of 978 patients. *Ann Surg* 2005; **241**: 9-13 [PMID: 15621985]
 - 19 **Jung SH**, Yu CS, Choi PW, Kim DD, Park IJ, Kim HC, Kim JC. Risk factors and oncologic impact of anastomotic leakage after rectal cancer surgery. *Dis Colon Rectum* 2008; **51**: 902-908 [PMID: 18408971 DOI: 10.1007/s10350-008-9272-x]
 - 20 **Eriksen MT**, Wibe A, Norstein J, Haffner J, Wiig JN. Anastomotic leakage following routine mesorectal excision for rectal cancer in a national cohort of patients. *Colorectal Dis* 2005; **7**: 51-57 [PMID: 15606585 DOI: 10.1111/j.1463-1318.2004.00700.x]
 - 21 **Matthiessen P**, Hallböök O, Andersson M, Rutegård J, Sjö Dahl R. Risk factors for anastomotic leakage after anterior resection of the rectum. *Colorectal Dis* 2004; **6**: 462-469 [PMID: 15521937 DOI: 10.1111/j.1463-1318.2004.00657.x]
 - 22 **Ito M**, Sugito M, Kobayashi A, Nishizawa Y, Tsunoda Y, Saito N. Relationship between multiple numbers of stapler firings during rectal division and anastomotic leakage after laparoscopic rectal resection. *Int J Colorectal Dis* 2008; **23**: 703-707 [PMID: 18379795 DOI: 10.1007/s00384-008-0470-8]
 - 23 **Kim JS**, Cho SY, Min BS, Kim NK. Risk factors for anastomotic leakage after laparoscopic intracorporeal colorectal anastomosis with a double stapling technique. *J Am Coll Surg* 2009; **209**: 694-701 [PMID: 19959036 DOI: 10.1016/j.jamcollsurg.2009.09.021]
 - 24 **Huh JW**, Kim HR, Kim YJ. Anastomotic leakage after laparoscopic resection of rectal cancer: the impact of fibrin glue. *Am J Surg* 2010; **199**: 435-441 [PMID: 19481197 DOI: 10.1016/j.amjsurg.2009.01.018]
 - 25 **Choi DH**, Hwang JK, Ko YT, Jang HJ, Shin HK, Lee YC, Lim CH, Jeong SK, Yang HK. Risk factors for anastomotic leakage after laparoscopic rectal resection. *J Korean Soc Coloproctol* 2010; **26**: 265-273 [PMID: 21152228 DOI: 10.3393/jksc.2010.26.4.265]
 - 26 **Akiyoshi T**, Ueno M, Fukunaga Y, Nagayama S, Fujimoto Y, Konishi T, Kuroyanagi H, Yamaguchi T. Incidence of and risk factors for anastomotic leakage after laparoscopic anterior resection with intracorporeal rectal transection and double-stapling technique anastomosis for rectal cancer. *Am J Surg* 2011; **202**: 259-264 [PMID: 21871980 DOI: 10.1016/j.amjsurg.2010.11.014]
 - 27 **Yamamoto S**, Fujita S, Akasu T, Inada R, Moriya Y, Yamamoto S. Risk factors for anastomotic leakage after laparoscopic surgery for rectal cancer using a stapling technique. *Surg Laparosc Endosc Percutan Tech* 2012; **22**: 239-243 [PMID: 22678320 DOI: 10.1097/SLE.0b013e31824fbb56]
 - 28 **Park JS**, Choi GS, Kim SH, Kim HR, Kim NK, Lee KY, Kang SB, Kim JY, Lee KY, Kim BC, Bae BN, Son GM, Lee SI, Kang H. Multicenter analysis of risk factors for anastomotic leakage after laparoscopic rectal cancer excision: the Korean laparoscopic colorectal surgery study group. *Ann Surg* 2013; **257**: 665-671 [PMID: 23333881 DOI: 10.1097/SLA.0b013e31827b8ed9]
 - 29 **Kawada K**, Hasegawa S, Hida K, Hirai K, Okoshi K, Nomura A, Kawamura J, Nagayama S, Sakai Y. Risk factors for anastomotic leakage after laparoscopic low anterior resection with DST anastomosis. *Surg Endosc* 2014; **28**: 2988-2995 [PMID: 24853855 DOI: 10.1007/s00464-014-3564-0]
 - 30 **Katsuno H**, Shiomi A, Ito M, Koide Y, Maeda K, Yatsuoka T, Hase K, Komori K, Minami K, Sakamoto K, Saida Y, Saito N. Comparison of symptomatic anastomotic leakage following laparoscopic and open low anterior resection for rectal cancer: a propensity score matching analysis of 1014 consecutive patients. *Surg Endosc* 2016; **30**: 2848-2856 [PMID: 26487228 DOI: 10.1007/s00464-015-4566-2]
 - 31 **Kim CW**, Baek SJ, Hur H, Min BS, Baik SH, Kim NK. Anastomotic Leakage After Low Anterior Resection for Rectal Cancer Is Different Between Minimally Invasive Surgery and Open Surgery. *Ann Surg* 2016; **263**: 130-137 [PMID: 25692355 DOI: 10.1097/SLA.0000000000001157]
 - 32 **Ba ZF**, Yokoyama Y, Toth B, Rue LW, Bland KI, Chaudry IH. Gender differences in small intestinal endothelial function: inhibitory role of androgens. *Am J Physiol Gastrointest Liver Physiol* 2004; **286**: G452-G457 [PMID: 14563675 DOI: 10.1152/ajpgi.00357.2003]
 - 33 **Volk A**, Kersting S, Held HC, Saeger HD. Risk factors for morbidity and mortality after single-layer continuous suture for ileocolonic anastomosis. *Int J Colorectal Dis* 2011; **26**: 321-327 [PMID: 20697722 DOI: 10.1007/s00384-010-1040-4]
 - 34 **Biondo S**, Parés D, Kreisler E, Ragué JM, Fracalvieri D, Ruiz AG, Jaurrieta E. Anastomotic dehiscence after resection and primary anastomosis in left-sided colonic emergencies. *Dis Colon Rectum* 2005; **48**: 2272-2280 [PMID: 16228841 DOI: 10.1007/s10350-005-0159-9]
 - 35 **Senagore AJ**, Delaney CP, Madboulay K, Brady KM, Fazio VW. Laparoscopic colectomy in obese and nonobese patients. *J Gastrointest Surg* 2003; **7**: 558-561 [PMID: 12763416 DOI: 10.1016/S1091-255X(02)00124-5]
 - 36 **Kartheuser AH**, Leonard DF, Penninckx F, Paterson HM, Brandt

- D, Remue C, Bugli C, Dozois E, Mortensen N, Ris F, Tiret E. Waist circumference and waist/hip ratio are better predictive risk factors for mortality and morbidity after colorectal surgery than body mass index and body surface area. *Ann Surg* 2013; **258**: 722-730 [PMID: 24096768 DOI: 10.1097/SLA.0b013e3182a6605a]
- 37 **Watanabe J**, Tatsumi K, Ota M, Suwa Y, Suzuki S, Watanabe A, Ishibe A, Watanabe K, Akiyama H, Ichikawa Y, Morita S, Endo I. The impact of visceral obesity on surgical outcomes of laparoscopic surgery for colon cancer. *Int J Colorectal Dis* 2014; **29**: 343-351 [PMID: 24297037 DOI: 10.1007/s00384-013-1803-9]
- 38 **Bosset JF**, Collette L, Calais G, Mineur L, Maingon P, Radosevic-Jelic L, Daban A, Bardet E, Beny A, Ollier JC. Chemotherapy with preoperative radiotherapy in rectal cancer. *N Engl J Med* 2006; **355**: 1114-1123 [PMID: 16971718 DOI: 10.1056/NEJMoa060829]
- 39 **Roh MS**, Colangelo LH, O'Connell MJ, Yothers G, Deutsch M, Allegra CJ, Kahlenberg MS, Baez-Diaz L, Ursiny CS, Petrelli NJ, Wolmark N. Preoperative multimodality therapy improves disease-free survival in patients with carcinoma of the rectum: NSABP R-03. *J Clin Oncol* 2009; **27**: 5124-5130 [PMID: 19770376 DOI: 10.1200/JCO.2009.22.0467]
- 40 **Heald RJ**, Husband EM, Ryall RD. The mesorectum in rectal cancer surgery--the clue to pelvic recurrence? *Br J Surg* 1982; **69**: 613-616 [PMID: 6751457 DOI: 10.1002/bjs.1800691019]
- 41 **Sebag-Montefiore D**, Stephens RJ, Steele R, Monson J, Grieve R, Khanna S, Quirke P, Couture J, de Metz C, Myint AS, Bessell E, Griffiths G, Thompson LC, Parmar M. Preoperative radiotherapy versus selective postoperative chemoradiotherapy in patients with rectal cancer (MRC CR07 and NCIC-CTG C016): a multicentre, randomised trial. *Lancet* 2009; **373**: 811-820 [PMID: 19269519 DOI: 10.1016/S0140-6736(09)60484-0]
- 42 **Marijnen CA**, Kapiteijn E, van de Velde CJ, Martijn H, Steup WH, Wiggers T, Kranenbarg EK, Leer JW. Acute side effects and complications after short-term preoperative radiotherapy combined with total mesorectal excision in primary rectal cancer: report of a multicenter randomized trial. *J Clin Oncol* 2002; **20**: 817-825 [PMID: 11821466 DOI: 10.1200/JCO.20.3.817]
- 43 **Chang JS**, Keum KC, Kim NK, Baik SH, Min BS, Huh H, Lee CG, Koom WS. Preoperative chemoradiotherapy effects on anastomotic leakage after rectal cancer resection: a propensity score matching analysis. *Ann Surg* 2014; **259**: 516-521 [PMID: 23598382 DOI: 10.1097/SLA.0b013e31829068e5]
- 44 **Heinzerling JH**, Huerta S. Bowel perforation from bevacizumab for the treatment of metastatic colon cancer: incidence, etiology, and management. *Curr Surg* 2006; **63**: 334-337 [PMID: 16971205 DOI: 10.1016/j.cursur.2006.06.002]
- 45 **Saif MW**, Elfiky A, Salem RR. Gastrointestinal perforation due to bevacizumab in colorectal cancer. *Ann Surg Oncol* 2007; **14**: 1860-1869 [PMID: 17356952 DOI: 10.1245/s10434-006-9337-9]
- 46 **Roos D**, Dijkstra LM, Tijssen JG, Gouma DJ, Gerhards MF, Oudemans-van Straaten HM. Systematic review of perioperative selective decontamination of the digestive tract in elective gastrointestinal surgery. *Br J Surg* 2013; **100**: 1579-1588 [PMID: 24264779 DOI: 10.1002/bjs.9254]
- 47 **Hata H**, Yamaguchi T, Hasegawa S, Nomura A, Hida K, Nishitai R, Yamanokuchi S, Yamanaka T, Sakai Y. Oral and Parenteral Versus Parenteral Antibiotic Prophylaxis in Elective Laparoscopic Colorectal Surgery (JMTO PREV 07-01): A Phase 3, Multicenter, Open-label, Randomized Trial. *Ann Surg* 2016; **263**: 1085-1091 [PMID: 26756752 DOI: 10.1097/SLA.0000000000001581]
- 48 **Golub R**, Golub RW, Cantu R, Stein HD. A multivariate analysis of factors contributing to leakage of intestinal anastomoses. *J Am Coll Surg* 1997; **184**: 364-372 [PMID: 9100681]
- 49 **Sliker JC**, Komen N, Mannaerts GH, Karsten TM, Willemsen P, Murawska M, Jeekel J, Lange JF. Long-term and perioperative corticosteroids in anastomotic leakage: a prospective study of 259 left-sided colorectal anastomoses. *Arch Surg* 2012; **147**: 447-452 [PMID: 22249852 DOI: 10.1001/archsurg.2011.1690]
- 50 **Konishi T**, Watanabe T, Kishimoto J, Nagawa H. Risk factors for anastomotic leakage after surgery for colorectal cancer: results of prospective surveillance. *J Am Coll Surg* 2006; **202**: 439-444 [PMID: 16500248 DOI: 10.1016/j.jamcollsurg.2005.10.019]
- 51 **Eriksen TF**, Lassen CB, Gögenur I. Treatment with corticosteroids and the risk of anastomotic leakage following lower gastrointestinal surgery: a literature survey. *Colorectal Dis* 2014; **16**: O154-O160 [PMID: 24215329 DOI: 10.1111/codi.12490]
- 52 **Burton TP**, Mittal A, Soop M. Nonsteroidal anti-inflammatory drugs and anastomotic dehiscence in bowel surgery: systematic review and meta-analysis of randomized, controlled trials. *Dis Colon Rectum* 2013; **56**: 126-134 [PMID: 23222290 DOI: 10.1097/DCR.0b013e31825fe927]
- 53 **Gorissen KJ**, Benning D, Berghmans T, Snoeijs MG, Sosef MN, Hulsewe KW, Luyer MD. Risk of anastomotic leakage with nonsteroidal anti-inflammatory drugs in colorectal surgery. *Br J Surg* 2012; **99**: 721-727 [PMID: 22318712 DOI: 10.1002/bjs.8691]
- 54 **Bertelsen CA**, Andreasen AH, Jørgensen T, Harling H. Anastomotic leakage after anterior resection for rectal cancer: risk factors. *Colorectal Dis* 2010; **12**: 37-43 [PMID: 19175624 DOI: 10.1111/j.1463-1318.2008.01711.x]
- 55 **Richards CH**, Campbell V, Ho C, Hayes J, Elliott T, Thompson-Fawcett M. Smoking is a major risk factor for anastomotic leak in patients undergoing low anterior resection. *Colorectal Dis* 2012; **14**: 628-633 [PMID: 21749605 DOI: 10.1111/j.1463-1318.2011.02718.x]
- 56 **Kruschewski M**, Rieger H, Pohlen U, Hotz HG, Buhr HJ. Risk factors for clinical anastomotic leakage and postoperative mortality in elective surgery for rectal cancer. *Int J Colorectal Dis* 2007; **22**: 919-927 [PMID: 17260142 DOI: 10.1007/s00384-006-0260-0]
- 57 **Sørensen LT**, Jørgensen T, Kirkeby LT, Skovdal J, Vennits B, Wille-Jørgensen P. Smoking and alcohol abuse are major risk factors for anastomotic leakage in colorectal surgery. *Br J Surg* 1999; **86**: 927-931 [PMID: 10417567 DOI: 10.1046/j.1365-2168.1999.01165.x]
- 58 **Contant CM**, Hop WC, van't Sant HP, Oostvogel HJ, Smeets HJ, Stassen LP, Neijenhuis PA, Idenburg FJ, Dijkhuis CM, Heres P, van Tets WF, Gerritsen JJ, Weidema WF. Mechanical bowel preparation for elective colorectal surgery: a multicentre randomised trial. *Lancet* 2007; **370**: 2112-2117 [PMID: 18156032 DOI: 10.1016/S0140-6736(07)61905-9]
- 59 **Slim K**, Vicaut E, Launay-Savary MV, Contant C, Chipponi J. Updated systematic review and meta-analysis of randomized clinical trials on the role of mechanical bowel preparation before colorectal surgery. *Ann Surg* 2009; **249**: 203-209 [PMID: 19212171 DOI: 10.1097/SLA.0b013e318193425a]
- 60 **Van't Sant HP**, Weidema WF, Hop WC, Oostvogel HJ, Contant CM. The influence of mechanical bowel preparation in elective lower colorectal surgery. *Ann Surg* 2010; **251**: 59-63 [PMID: 20009750 DOI: 10.1097/SLA.0b013e3181c0e75c]
- 61 **Dahabreh IJ**, Steele DW, Shah N, Trikalinos TA. Oral Mechanical Bowel Preparation for Colorectal Surgery: Systematic Review and Meta-Analysis. *Dis Colon Rectum* 2015; **58**: 698-707 [PMID: 26200685 DOI: 10.1097/DCR.0000000000000375]
- 62 **Scarborough JE**, Mantyh CR, Sun Z, Migaly J. Combined Mechanical and Oral Antibiotic Bowel Preparation Reduces Incisional Surgical Site Infection and Anastomotic Leak Rates After Elective Colorectal Resection: An Analysis of Colectomy-Targeted ACS NSQIP. *Ann Surg* 2015; **262**: 331-337 [PMID: 26083870 DOI: 10.1097/SLA.0000000000001041]
- 63 **Moghadamyeghaneh Z**, Hanna MH, Carmichael JC, Mills SD, Pigazzi A, Nguyen NT, Stamos MJ. Nationwide analysis of outcomes of bowel preparation in colon surgery. *J Am Coll Surg* 2015; **220**: 912-920 [PMID: 25907871 DOI: 10.1016/j.jamcollsurg.2015.02.008]
- 64 **Kim EK**, Sheetz KH, Bonn J, DeRoo S, Lee C, Stein I, Zarinsefat A, Cai S, Campbell DA, Englesbe MJ. A statewide colectomy experience: the role of full bowel preparation in preventing surgical site infection. *Ann Surg* 2014; **259**: 310-314 [PMID: 23979289 DOI: 10.1097/SLA.0b013e3182a62643]
- 65 **Jung B**, Pählman L, Nyström PO, Nilsson E. Multicentre randomized clinical trial of mechanical bowel preparation in elective colonic resection. *Br J Surg* 2007; **94**: 689-695 [PMID: 17514668 DOI: 10.1002/bjs.5816]

- 66 **Collin Å**, Jung B, Nilsson E, Pählman L, Folkesson J. Impact of mechanical bowel preparation on survival after colonic cancer resection. *Br J Surg* 2014; **101**: 1594-1600 [PMID: 25204295 DOI: 10.1002/bjs.9629]
- 67 **Kuroyanagi H**, Oya M, Ueno M, Fujimoto Y, Yamaguchi T, Muto T. Standardized technique of laparoscopic intracorporeal rectal transection and anastomosis for low anterior resection. *Surg Endosc* 2008; **22**: 557-561 [PMID: 18193475 DOI: 10.1007/s00464-007-9626-9]
- 68 **Maeda K**, Nagahara H, Shibutani M, Ohtani H, Sakurai K, Toyokawa T, Muguruma K, Tanaka H, Amano R, Kimura K, Sugano K, Ikeya T, Iseki Y, Hirakawa K. Efficacy of intracorporeal reinforcing sutures for anastomotic leakage after laparoscopic surgery for rectal cancer. *Surg Endosc* 2015; **29**: 3535-3542 [PMID: 25673349 DOI: 10.1007/s00464-015-4104-2]
- 69 **Baek SJ**, Kim J, Kwak J, Kim SH. Can trans-anal reinforcing sutures after double stapling in lower anterior resection reduce the need for a temporary diverting ostomy? *World J Gastroenterol* 2013; **19**: 5309-5313 [PMID: 23983434 DOI: 10.3748/wjg.v19.i32.5309]
- 70 **Roumen RM**, Rahusen FT, Wijnen MH, Croiset van Uchelen FA. "Dog ear" formation after double-stapled low anterior resection as a risk factor for anastomotic disruption. *Dis Colon Rectum* 2000; **43**: 522-525 [PMID: 10789750 DOI: 10.1007/BF02237198]
- 71 **Zhuo C**, Liang L, Ying M, Li Q, Li D, Li Y, Peng J, Huang L, Cai S, Li X. Laparoscopic Low Anterior Resection and Eversion Technique Combined With a Nondog Ear Anastomosis for Mid- and Distal Rectal Neoplasms: A Preliminary and Feasibility Study. *Medicine (Baltimore)* 2015; **94**: e2285 [PMID: 26683958 DOI: 10.1097/MD.0000000000002285]
- 72 **Nakayama S**, Hasegawa S, Nagayama S, Kato S, Hida K, Tanaka E, Itami A, Kubo H, Sakai Y. The importance of precompression time for secure stapling with a linear stapler. *Surg Endosc* 2011; **25**: 2382-2386 [PMID: 21184102 DOI: 10.1007/s00464-010-1527-7]
- 73 **Nakayama S**, Hasegawa S, Hida K, Kawada K, Sakai Y. Obtaining secure stapling of a double stapling anastomosis. *J Surg Res* 2015; **193**: 652-657 [PMID: 25277356 DOI: 10.1016/j.jss.2014.08.044]
- 74 **Hasegawa S**, Nakayama S, Hida K, Kawada K, Sakai Y. Effect of Tri-Staple Technology and Slow Firing on Secure Stapling Using an Endoscopic Linear Stapler. *Dig Surg* 2015; **32**: 353-360 [PMID: 26228297 DOI: 10.1159/000437216]
- 75 **Vignali A**, Gianotti L, Braga M, Radaelli G, Malvezzi L, Di Carlo V. Altered microperfusion at the rectal stump is predictive for rectal anastomotic leak. *Dis Colon Rectum* 2000; **43**: 76-82 [PMID: 10813128 DOI: 10.1007/BF02237248]
- 76 **Sheridan WG**, Lowndes RH, Young HL. Tissue oxygen tension as a predictor of colonic anastomotic healing. *Dis Colon Rectum* 1987; **30**: 867-871 [PMID: 3677962 DOI: 10.1007/BF02555426]
- 77 **Kologlu M**, Yorganci K, Renda N, Sayek I. Effect of local and remote ischemia-reperfusion injury on healing of colonic anastomoses. *Surgery* 2000; **128**: 99-104 [PMID: 10876192 DOI: 10.1067/msy.2000.107414]
- 78 **Karliczek A**, Harlaar NJ, Zeebregts CJ, Wiggers T, Baas PC, van Dam GM. Surgeons lack predictive accuracy for anastomotic leakage in gastrointestinal surgery. *Int J Colorectal Dis* 2009; **24**: 569-576 [PMID: 19221768 DOI: 10.1007/s00384-009-0658-6]
- 79 **Kudszus S**, Roesel C, Schachtrupp A, Höer JJ. Intraoperative laser fluorescence angiography in colorectal surgery: a noninvasive analysis to reduce the rate of anastomotic leakage. *Langenbecks Arch Surg* 2010; **395**: 1025-1030 [PMID: 20700603 DOI: 10.1007/s00423-010-0699-x]
- 80 **Jafari MD**, Lee KH, Halabi WJ, Mills SD, Carmichael JC, Stamos MJ, Pigazzi A. The use of indocyanine green fluorescence to assess anastomotic perfusion during robotic assisted laparoscopic rectal surgery. *Surg Endosc* 2013; **27**: 3003-3008 [PMID: 23404152 DOI: 10.1007/s00464-013-2832-8]
- 81 **Hellan M**, Spinoglio G, Pigazzi A, Lagares-Garcia JA. The influence of fluorescence imaging on the location of bowel transection during robotic left-sided colorectal surgery. *Surg Endosc* 2014; **28**: 1695-1702 [PMID: 24385249 DOI: 10.1007/s00464-013-3377-6]
- 82 **Jafari MD**, Wexner SD, Martz JE, McLemore EC, Margolin DA, Sherwinter DA, Lee SW, Senagore AJ, Phelan MJ, Stamos MJ. Perfusion assessment in laparoscopic left-sided/anterior resection (PILLAR II): a multi-institutional study. *J Am Coll Surg* 2015; **220**: 82-92.e1 [PMID: 25451666 DOI: 10.1016/j.jamcollsurg.2014.09.015]
- 83 **Degett TH**, Andersen HS, Gögenur I. Indocyanine green fluorescence angiography for intraoperative assessment of gastrointestinal anastomotic perfusion: a systematic review of clinical trials. *Langenbecks Arch Surg* 2016; Epub ahead of print [PMID: 26968863 DOI: 10.1007/s00423-016-1400-9]
- 84 **Sherwinter DA**, Gallagher J, Donkar T. Intra-operative transanal near infrared imaging of colorectal anastomotic perfusion: a feasibility study. *Colorectal Dis* 2013; **15**: 91-96 [PMID: 22632448 DOI: 10.1111/j.1463-1318.2012.03101.x]
- 85 **Kin C**, Vo H, Welton L, Welton M. Equivocal effect of intraoperative fluorescence angiography on colorectal anastomotic leaks. *Dis Colon Rectum* 2015; **58**: 582-587 [PMID: 25944430 DOI: 10.1097/DCR.0000000000000320]
- 86 **Hinoi T**, Okajima M, Shimomura M, Egi H, Ohdan H, Konishi F, Sugihara K, Watanabe M. Effect of left colonic artery preservation on anastomotic leakage in laparoscopic anterior resection for middle and low rectal cancer. *World J Surg* 2013; **37**: 2935-2943 [PMID: 24005279 DOI: 10.1007/s00268-013-2194-3]
- 87 **Leichtle SW**, Mouawad NJ, Welch KB, Lampman RM, Cleary RK. Risk factors for anastomotic leakage after colectomy. *Dis Colon Rectum* 2012; **55**: 569-575 [PMID: 22513436 DOI: 10.1097/DCR.0b013e3182423c0d]
- 88 **Mäkelä JT**, Kiviniemi H, Laitinen S. Risk factors for anastomotic leakage after left-sided colorectal resection with rectal anastomosis. *Dis Colon Rectum* 2003; **46**: 653-660 [PMID: 12792443 DOI: 10.1007/s10350-004-6627-9]
- 89 **Matthiessen P**, Hallböök O, Rutegård J, Simert G, Sjö Dahl R. Defunctioning stoma reduces symptomatic anastomotic leakage after low anterior resection of the rectum for cancer: a randomized multicenter trial. *Ann Surg* 2007; **246**: 207-214 [PMID: 17667498 DOI: 10.1097/SLA.0b013e3180603024]
- 90 **Chude GG**, Rayate NV, Patris V, Koshariya M, Jagad R, Kawamoto J, Lygidakis NJ. Defunctioning loop ileostomy with low anterior resection for distal rectal cancer: should we make an ileostomy as a routine procedure? A prospective randomized study. *HepatoGastroenterology* 2008; **55**: 1562-1567 [PMID: 19102343]
- 91 **Lefebure B**, Tuech JJ, Bridoux V, Costaglioli B, Scotte M, Teniere P, Michot F. Evaluation of selective defunctioning stoma after low anterior resection for rectal cancer. *Int J Colorectal Dis* 2008; **23**: 283-288 [PMID: 17768630 DOI: 10.1007/s00384-007-0380-1]
- 92 **Enker WE**, Merchant N, Cohen AM, Lanouette NM, Swallow C, Guillem J, Paty P, Minsky B, Weyrauch K, Quan SH. Safety and efficacy of low anterior resection for rectal cancer: 681 consecutive cases from a specialty service. *Ann Surg* 1999; **230**: 544-552; discussion 552-554 [PMID: 10522724 DOI: 10.1097/00000658-199910000-00010]
- 93 **Gessler B**, Haglind E, Angenete E. A temporary loop ileostomy affects renal function. *Int J Colorectal Dis* 2014; **29**: 1131-1135 [PMID: 25026894 DOI: 10.1007/s00384-014-1949-0]
- 94 **Sharma A**, Deeb AP, Rickles AS, Iannuzzi JC, Monson JR, Fleming FJ. Closure of defunctioning loop ileostomy is associated with considerable morbidity. *Colorectal Dis* 2013; **15**: 458-462 [PMID: 22974343 DOI: 10.1111/codi.12029]
- 95 **Xiao L**, Zhang WB, Jiang PC, Bu XF, Yan Q, Li H, Zhang YJ, Yu F. Can transanal tube placement after anterior resection for rectal carcinoma reduce anastomotic leakage rate? A single-institution prospective randomized study. *World J Surg* 2011; **35**: 1367-1377 [PMID: 21437746 DOI: 10.1007/s00268-011-1053-3]
- 96 **Zhao WT**, Hu FL, Li YY, Li HJ, Luo WM, Sun F. Use of a transanal drainage tube for prevention of anastomotic leakage and bleeding after anterior resection for rectal cancer. *World J Surg* 2013; **37**: 227-232 [PMID: 23052807 DOI: 10.1007/s00268-012-1812-9]
- 97 **Nishigori H**, Ito M, Nishizawa Y, Nishizawa Y, Kobayashi A, Sugito M, Saito N. Effectiveness of a transanal tube for the prevention of anastomotic leakage after rectal cancer surgery.

- World J Surg* 2014; **38**: 1843-1851 [PMID: 24378550 DOI: 10.1007/s00268-013-2428-4]
- 98 **Cong ZJ**, Fu CG, Wang HT, Liu LJ, Zhang W, Wang H. Influencing factors of symptomatic anastomotic leakage after anterior resection of the rectum for cancer. *World J Surg* 2009; **33**: 1292-1297 [PMID: 19363687 DOI: 10.1007/s00268-009-0008-4]
- 99 **Hidaka E**, Ishida F, Mukai S, Nakahara K, Takayanagi D, Maeda C, Takehara Y, Tanaka J, Kudo SE. Efficacy of transanal tube for prevention of anastomotic leakage following laparoscopic low anterior resection for rectal cancers: a retrospective cohort study in a single institution. *Surg Endosc* 2015; **29**: 863-867 [PMID: 25052128 DOI: 10.1007/s00464-014-3740-2]
- 100 **Sirois-Giguère E**, Boulanger-Gobeil C, Bouchard A, Gagné JP, Grégoire RC, Thibault C, Bouchard P. Transanal drainage to treat anastomotic leaks after low anterior resection for rectal cancer: a valuable option. *Dis Colon Rectum* 2013; **56**: 586-592 [PMID: 23575397 DOI: 10.1097/DCR.0b013e31827687a4]
- 101 **Okoshi K**, Masano Y, Hasegawa S, Hida K, Kawada K, Nomura A, Kawamura J, Nagayama S, Yoshimura T, Sakai Y. Efficacy of transanal drainage for anastomotic leakage after laparoscopic low anterior resection of the rectum. *Asian J Endosc Surg* 2013; **6**: 90-95 [PMID: 23228055 DOI: 10.1111/ases.12010]
- 102 **Karliczek A**, Jesus EC, Matos D, Castro AA, Atallah AN, Wiggers T. Drainage or nondrainage in elective colorectal anastomosis: a systematic review and meta-analysis. *Colorectal Dis* 2006; **8**: 259-265 [PMID: 16630227 DOI: 10.1111/j.1463-1318.2006.00999.x]
- 103 **Rondelli F**, Bugiantella W, Vedovati MC, Balzarotti R, Avenia N, Mariani E, Agnelli G, Becattini C. To drain or not to drain extraperitoneal colorectal anastomosis? A systematic review and meta-analysis. *Colorectal Dis* 2014; **16**: O35-O42 [PMID: 24245821 DOI: 10.1111/codi.12491]
- 104 **Morowitz MJ**, Babrowski T, Carlisle EM, Olivas A, Romanowski KS, Seal JB, Liu DC, Alverdy JC. The human microbiome and surgical disease. *Ann Surg* 2011; **253**: 1094-1101 [PMID: 21422915 DOI: 10.1097/sla.0b013e31821175d7]
- 105 **Manichanh C**, Varela E, Martinez C, Antolin M, Llopis M, Doré J, Giralt J, Guamer F, Malagelada JR. The gut microbiota predispose to the pathophysiology of acute postradiotherapy diarrhea. *Am J Gastroenterol* 2008; **103**: 1754-1761 [PMID: 18564125 DOI: 10.1111/j.1572-0241.2008.01868.x]
- 106 **Shogan BD**, An GC, Schardey HM, Matthews JB, Umanskiy K, Fleshman JW, Hoepfner J, Fry DE, Garcia-Granereo E, Jeekel H, van Goor H, Dellinger EP, Konda V, Gilbert JA, Auner GW, Alverdy JC. Proceedings of the first international summit on intestinal anastomotic leak, Chicago, Illinois, October 4-5, 2012. *Surg Infect (Larchmt)* 2014; **15**: 479-489 [PMID: 25215465 DOI: 10.1089/sur.2013.114]
- 107 **Olivas AD**, Shogan BD, Valuckaite V, Zaborin A, Belogortseva N, Musch M, Meyer F, Trimble WL, An G, Gilbert J, Zaborina O, Alverdy JC. Intestinal tissues induce an SNP mutation in *Pseudomonas aeruginosa* that enhances its virulence: possible role in anastomotic leak. *PLoS One* 2012; **7**: e44326 [PMID: 22952955 DOI: 10.1371/journal.pone.0044326]
- 108 **Shogan BD**, Belogortseva N, Luong PM, Zaborin A, Lax S, Bethel C, Ward M, Muldoon JP, Singer M, An G, Umanskiy K, Konda V, Shakhsheer B, Luo J, Klabbbers R, Hancock LE, Gilbert J, Zaborina O, Alverdy JC. Collagen degradation and MMP9 activation by *Enterococcus faecalis* contribute to intestinal anastomotic leak. *Sci Transl Med* 2015; **7**: 286ra68 [PMID: 25947163 DOI: 10.1126/scitranslmed.3010658]
- 109 **Damle RN**, Cherng NB, Flahive JM, Davids JS, Maykel JA, Sturrock PR, Sweeney WB, Alavi K. Clostridium difficile infection after colorectal surgery: a rare but costly complication. *J Gastrointest Surg* 2014; **18**: 1804-1811 [PMID: 25091840 DOI: 10.1007/s11605-014-2600-7]
- 110 **Gaertner WB**, Madoff RD, Mellgren A, Kwaan MR, Melton GB. Postoperative diarrhea and high ostomy output impact postoperative outcomes after elective colon and rectal operations regardless of Clostridium difficile infection. *Am J Surg* 2015; **210**: 759-765 [PMID: 26117432 DOI: 10.1016/j.amjsurg.2015.03.032]

P- Reviewer: Konishi T, Morris DLL **S- Editor:** Qi Y
L- Editor: A **E- Editor:** Wang CH





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045