

RESEARCH

Open Access



Medicinal plants sold at traditional markets in southern Ecuador

Fani Tinitana^{1*}, Montserrat Rios^{2,3}, Juan Carlos Romero-Benavides², Marcelino de la Cruz Rot⁴ and Manuel Pardo-de-Santayana⁵

Abstract

Background: The traditional markets in southern Ecuador and within the Andean region are especially important for plant resource trading among local people, even since before Spanish colonization; therefore, ethnobotanical studies are currently necessary and important. These strategic spaces persist for the traditional medicine cultural value reflected in the higher consumption of medicinal plants, which span all socioeconomic levels of rural and urban people. The purpose of this study includes the following: 1) to create a novel list of medicinal plants sold at 33 traditional markets; 2) to establish medicinal plant use agreement amongst vendors with the Factor of Informant Consensus (FIC); and 3) to determine the most sold medicinal plant species using the Fidelity Level (FL).

Methods: This study focus on traditional markets ethnobotany utilizes the largest sample of medicinal plants market vendors up to date in Ecuador, interviewing them at 33 traditional markets, located within the Loja province. In order to determine the most sold medicinal plants and their ethnobotanical information, structured questionnaires and personal conversations were conducted with 196 medicinal plant vendors, and voucher specimens were created. Agreement among vendors about the therapeutic use of medicinal plants was measured using the FIC, and the most sold medicinal plant species were assessed with the FL.

Results and discussion: This research registered 160 medicinal plant species, grouped in 126 genera and 57 families that were sold in 33 traditional markets. The uses of medicinal plants in southern Ecuador are related to a long history of traditional medicine health practices that has persisted until today as well as high plant diversity. The 53 therapeutic uses recorded were grouped into 12 medical categories that were adapted from the World Health Organization. Three medical categories shared the highest value for FIC = 0.92, which showed a high level of agreement of market vendors for 57 medicinal plant species sold to treat ailments related with digestive, dermatological, and sensorial systems. The FL index determined 11 culturally important medicinal plant species based on the reported uses by 40 or more market vendors. Two medicinal plant species had an FL = 100 %, *Matricaria recutita* and *Gaiadendrum punctatum*, used to treat digestive and respiratory systems ailments.

(Continued on next page)

* Correspondence: ftinitana@utpl.edu.ec

¹Departamento de Ciencias Naturales, Universidad Técnica Particular de Loja, San Cayetano Alto, CP: 1101608 Loja, Ecuador

Full list of author information is available at the end of the article



(Continued from previous page)

Conclusions: In the Loja province, people continue to consume medicinal plant species sold at local markets to treat somatic and/or psychosomatic health ailments because sociocultural customs are strongly expressed in ancestral practices of wellbeing. When the largest values of FL (60.5 %–100 %) and FIC (0.81–0.92) indexes are combined, they demonstrated agreement among 196 market vendors in the use of seven medicinal plant species that were most sold for the 12 medical categories. This study stresses how important public policies are for the trade and quality of medicinal plant resources, particularly for local people practicing auto-medication. Reasons for the maintenance of traditional markets in southern Ecuador include lower cost of medicinal plants, confidence in traditional medicine, and/or sociocultural environment. In Ecuador, the sustainable management of wild medicinal plants diversity, particularly the most sold, is crucial for its conservation in nature.

Keywords: Traditional markets, Medicinal plants, Factor Informant Consensus, Fidelity Level, Loja province, Southern Ecuador

Background

Traditional markets around the world have been recognized as places for the trade of plants and their derivative products and have become exchange posts where cultures are expressed through regional trade [1–11]. Additionally, markets are a meeting place to display a diverse array of minerals, animals, and plants sold locally, which come from neighboring communities that are culturally and ecologically diverse [4, 12]. In this way, literature on traditional markets and traded medicinal plant species with their value chain flows requires more attention from scientists, because ethnobotanical information is rather scarce.

Current ethnobotanical research at traditional markets across continents, considering Asia, Africa, Oceania, and Latin America, contributes to the understanding of plant diversity through the trade of medicinal plant species and their cultural value [13–45]. In this way, market surveys can help to understand regional networks of producers, sellers, healers, and consumers by the supply and demand of medicinal plants and their derivative products [4]. The total number of inventoried medicinal plant species at a particular traditional market is important, but they do not necessarily represent all species used in the traditional medicine of a specific human group [5, 8].

According to the World Health Organization (WHO), the wellbeing of 80 % of the population in developing countries depends mostly on the use of medicinal plants through traditional medicine, spiritual therapies, and ancestral healing practices [46]. This fact is particularly evident in the ancestral practices of traditional communities living in rural areas. In Latin America, ethnobotanical studies of traditional markets and their history are needed, because the trade of medicinal plants and their derived products has local, national, regional, and international importance, especially given their growing demand [15, 47]. In the Andean-Amazonian region including Ecuador, traditional markets existed before Spanish

colonization [48]. Throughout the Spanish conquest, a new kind of market appeared within public areas named “tiánguez” for the exchange of goods; they were also strategic points for bartering, conversation, and the sharing of life experiences [47, 48].

For the Andean region, published studies of traditional markets that emphasize ethnobotanical aspects have been conducted in Venezuela [11], Colombia [18], Bolivia [19, 43], Peru [49–79], and Ecuador [80]. Within Ecuador, it is estimated that 273 medicinal plants species were sold in the herb stalls (“puestos de hierbas”) of corresponding traditional markets, which were located at six provincial capitals in the Andean and Amazonian regions [80]. These capital cities, represented by Ambato, Quito, Riobamba, Nueva Loja, Puyo, and Tena, are the main points of trade for medicinal plants and their derivatives; from these places, commerce routes begin to spread throughout the country [81, 82].

More studies are needed to investigate the medicinal plants sold in Ecuadorian markets [80–85] to determine which medicinal plant species are most sold and how these are related to local health disorders. This is particularly true for the southern region of the country and specifically for the Loja province, because although it is a region rich in plant diversity, it is a region that is deficient in traditional market studies. In this area, only a few ethnobotanical surveys have been conducted, particularly on how the “mestizo” population and indigenous communities use medicinal plant resources from wild collection and/or homegardens [9, 52, 86–89].

Nowadays, even basic inventoried information accounting for the origins of medicinal plant resources and quantities of fresh and/or dry material sold is lacking as well as consumers’ usage of these products. Studies of traditional markets are necessary in Ecuador, because large gaps in knowledge on flora trade persist. This research at Loja province includes the following objectives: 1) to create a novel list of medicinal plants sold at 33 traditional markets; 2) to establish medicinal plants

use agreement amongst vendors with the Factor of Informant Consensus (FIC) [90]; and 3) to determine the most sold medicinal plant species using the Fidelity Level (FL) [91].

Methods

Study area

The study was carried out in 33 traditional markets within the Loja province, situated in southern Ecuador, between 3°19'56"S to 4°44'36"S latitude and 79°04'28"W to 80°29'03"W longitude (Fig. 1). This region occupies 11,042 km², which is 4 % of the national territory, and borders to the south with Peru [92]. The total population of the province in 2010 was 448,966 inhabitants, consisting of 96.3 % "mestizo" Spanish speakers, and 3.7 % Saraguro indigenous people, who speak the Spanish and Kichwa languages [93].

The Loja province has abundant hydrographic features, like rivers that flow into the Pacific catchments basin. The province is dominated by the Andean mountain range, which gives rise to a very irregular topography, and altitudes between 120 and 3800 m. This region shows considerable variation in local climate,

with conditions represented by tropical dry to the west, subtropical humid in the central area, and cold humid at the east [92].

Markets

Medicinal plants were sold at 15 established markets and 18 open markets located in 13 of the 16 cities in Loja province (Fig. 1). The established market, known locally as "mercado", includes from five to ten subsectors; the most frequent of these include the following: personal use items, electronic equipment, groceries, cooked foods, dairy products, meat products, legumes, fruits, and fresh or dry raw medicinal plants. Besides the main established markets, open markets occur weekly and are known locally as "feria libre". Vendors at open markets are rural harvesters and/or small retailers who sell medicinal plants, fresh local products such as cheese, legumes, fruits, and vegetables; they commerce hens and guinea pigs.

A variety of actors were involved in the sale of medicinal plants at the studied markets. The majority of these actors included rural harvesters, small retailers, formal, and informal vendors (Table 1). The criteria applied to




Table 1 Kinds of vendors of medicinal plants and their role in traditional markets at Loja province

Vendor	Definition
Rural harvesters	Individuals who come from the rural areas surrounding the main cities of Loja province, bringing fresh medicinal plants produced and/or collected by them in nature and/or their homegardens. They always trade plant bunches in large quantities to formal vendors or small quantities to customers. They operate at open markets and/or established markets.
Small retailers	Individuals who come from rural areas surrounding the main cities of Loja province, bringing fresh medicinal plants harvested from nature or gathered from their homegardens. They occasionally go to the cities to trade plant bunches to the customers and/or to formal vendors in small quantities at open markets.
Formal vendors	Individuals who legally hold an operating license from the province government to rent a stall in the established market for trading legumes, fruits, vegetables and bunches of dry or fresh medicinal plants.
Informal vendors	Individuals who come from rural or metropolitan areas of Loja province, and are market vendors on foot. They are public resellers of fresh specific medicinal plant bunches in small quantities at established and/or open markets.

determine the types of each vendor in a market was based on how they auto-recognized their own role. The medicinal plant vendor in each market was essentially a person who used a specific know-how to trade bunches of medicinal plants, because each one is well familiarized with the therapeutic applications of every plant species sold. It is important to clarify that all interviewed vendors were just sellers and not healers.

Structured ethnobotanical questionnaires

Surveys of medicinal plants sold at 15 established markets and 18 open markets were conducted in the selected 13 cities within the Loja province between 2007 and 2013. During the visits, the first author carried out interviews with a total of 196 market vendors. After explaining the aim of the study, all vendors of medicinal plants from the 33 traditional markets were asked to participate in the research. The interviewed vendors were eighteen years or older; also, they were “mestizos” (95 %) and Saraguro indigenous people (5 %), and the large majority consisted of women (97 %).

Medicinal plants were bought from each vendor, and interviews were structured as ethnobotanical questionnaires in Spanish, being conducted by the main author with the 196 market vendors. In the field research, the first author respected the vendors who preferred to remain anonymous. The questionnaires aimed to record the specimens' information on the following: vernacular names, medicinal uses, plant morphological structures sold, and therapeutic prescriptions. All the vendors who decided to collaborate were interviewed according to mutually agreed conditions and under Ecuador's rights, especially with regards to the Convention on Biological Diversity (CDB) [94].

This research was conducted according to the code of ethics of the International Society for Ethnobiology (ISE) [95], which is also endorsed by the Society for Economic Botany (SEB). The Principle of Respect, numbered 9 in the code, recognizes the necessity for researchers to respect the integrity, morality, and spirituality of the culture, traditions, and relationships of indigenous people,

traditional societies, and local communities within their worlds.

Voucher collection and nomenclature

The nomenclature of plant families, genera, and species follows the Catalogue of Vascular Plants of Ecuador [96]. It was also compared to the TROPICOS database [97]. The 160 species were identified using the available volumes of the Flora of Ecuador [98–101] and reference material in the herbaria of the “Universidad Técnica Particular de Loja” (HUTPL) and “Universidad Nacional de Loja” (LOJA). The specimens were registered under the collection series FT (Fani Tinitana), and vouchers were deposited at HUTPL. The collection of botanical specimens sold in the 33 traditional markets authorized the Ecuadorian Ministry of the Environment (Ministerio del Ambiente del Ecuador N° 001-2013-IC-FLO-DPAP-MAE).

Quantitative analysis

All the local therapeutic uses of medicinal plants were grouped in 12 medical categories (Table 2), which were adapted from the catalogue of International Classification of Diseases made by the WHO [102]. In this research, each category proposed by the WHO allows grouping and systematizing the data related to ‘illness’ and ‘disease’ as well as to compare the results among other regional and international studies related to the markets' ethnobotany [102, 103]. Additionally, WHO recognizes in each medical category the health practice systems of traditional populations [46]. In this study, ‘illness’ refers to being ill as conceived from a sociocultural personal perception, while ‘disease’ was considered from the biomedical perspective [104].

Information recorded in the structured ethnobotanical questionnaires related to the collected 160 taxa and their medicinal uses were recorded into a data matrix for quantitative analysis. The FIC index was used to measure consensus among vendors regarding the therapeutic use of each medicinal plant [106–111]; it shows the level of homogeneity among information provided by different vendors. The FIC was calculated according to the

Table 2 Therapeutic uses of medicinal plants to treat local ailments at Loja province

Medical category	Local illnesses and diseases recognized by market vendors
Circulatory system	Anemia, bad blood circulation, high cholesterol, and high or low blood pressure
Culture-bound syndromes	"Calor encerrado", evil air, evil eye, fright, "espanto de cerro", and "pena de ausencia" or "tirisia" (see definition of syndromes in Tene et al. 2007) [88] and Rios et al. 2007 [104]
Dermatological system	Acne, fungus infection, gangrene, rash, wounds, nosebleed, hair loss, and dandruff
Digestive system	Diarrhea, constipation, sickness, hangover, flatulence, liver disorder (included inflammation), stomach infection and pain, and tooth pain
General disorders	Inflammation, cancer, fever, headache, and sunstroke
Genitourinary system	Kidney ailments (included inflammation and infection), and prostate and urinary tract disorders
Gynecological system	Vaginal disorders, abdominal pain, menstrual cramps and related disorders, ovary inflammation, and promoting labor and childbirth recovery
Hormonal system	Diabetes and galactagogue
Musculoskeletal system	Bone fracture, bruise, rheumatism, sprain, and pains
Nervous system	Nervousness
Respiratory system	Cold, cough, flu, sore throat, and measles
Sensorial system	Ear pain and eye infection

following formula: $FIC = (N_{ur} - N_t) / (N_{ur} - 1)$, where N_{ur} refers to the number of therapeutic use reports, grouped in a medical category, from market vendors for a particular medicinal plant, and N_t refers to the total number of medicinal plant species used in a particular medical category [90, 106, 107]. The FIC values range between 0 and 1, where 1 indicates the highest level of market vendor consensus.

The relative healing potential of each reported medicinal plant sold at 33 traditional markets was evaluated with the FL index [91–112]. This indicates the percentage of vendors claiming the use of a certain medicinal plant for the same therapeutic use, which was grouped in a specific medical category [113, 114]. The FL was calculated according to the following formula: $FL (\%) = (I_p \times 100 / I_u)$, where I_p is the number of market vendors who independently claim a therapeutic use of a medicinal plant species to treat a specific illness or disease, and I_u is the total number of market vendors that sold the same medicinal plant to treat any given illness or disease.

Results and discussion

Medicinal plants sold at traditional markets

This research registered 160 medicinal plant species traded in 33 traditional markets within the Loja province, which were grouped in 123 genera and 57 vascular plant families (Table 5.). In traditional markets at La Paz (Bolivia) and Cusco (Peru), a total of 129 [19] and 152 [117] medicinal plant species were respectively reported; in contrast, 400 plant species were recorded in Trujillo and Chiclayo (Peru) [22]. When compared to these previous studies, the number of medicinal plant species sold in markets within the Loja province represents an intermediate value.

The dominant plant family was Asteraceae with 19 species that represented 11.8 % of the total species, followed by 16 species of Lamiaceae (10 %), 8 species of Piperaceae (5 %) and Pteridaceae (5 %), 7 species of Amaranthaceae (4.4 %) and Solanaceae (4.4 %), 6 species of Onagraceae (3.8 %), and 5 species of Rosaceae (3.1 %) (Fig. 2). Other studies of Andean highland traditional markets also recorded Asteraceae as the family with the highest number of medicinal plant species, and Solanaceae and Lamiaceae were consistently among the most frequent families [11, 43, 117].

The most frequent medicinal plant life forms were herbs (61.1 %) and shrubs (32.2 %), followed by trees (5.5 %) and lianas (1.2 %). This data was similar to results from other studies of highland markets in Bolivia [19] and Peru [117], where the herb habit represents a large percentage due to its random occurrence, high diversity, and endemism. Herbs such as weeds, which were abundantly available in relation to other plant life forms, are an important source of food and remedies [118]. This is because they contain one or more bioactive principles and a wide variety of highly active secondary metabolic compounds [119], making these plants potentially more effective for medicinal applications [120].

Geographic status of medicinal plant species sold at traditional markets

The medicinal plants native to Ecuador belonged to 92 species (57.5 %) [82, 86, 88] and were brought to traditional markets from mountain forests, cloud forests, scrub vegetation, and the Andean paramo [121, 122]. Of the 160 species, 6 (3.8 %) were endemic to Ecuador highlands [123], corresponding to the families Asclepiadaceae (*Orthosia ellemanniae*), Asteraceae (*Achyroclinehallii*,




Fig. 2 Dominant medicinal plant families recorded in 33 traditional markets within the Loja province, southern Ecuador

Aequatorium jamesonii and *Aristeguietia persicifolia*), and Onagraceae (*Fuchsia harlingii* and *Fuchsia loxensis*), whereas 62 species (38.7 %) were introduced from different regions of the world. In relation to the role of humans in the management of medicinal plant species, the material sold in the studied traditional markets belonged to homegardens (cultivation) and natural vegetation (wild).

The five most sold medicinal plant species were very well known by vendors for their therapeutic uses and health properties. These were *Aerva saguinolenta*, *Equisetum bogotense*, *Matricaria recutita*, *Oreocallis grandiflora*, and *Ruta graveolens*. The trade of these five taxa was linked to the treatment of the most common illnesses and diseases present within the Loja province, related to the respiratory system, genitourinary system, digestive system, and culture-bound syndromes (e.g. “evil eye”, “evil air”, “frights”, and “calor encerrado”). It is important to stress that *Matricaria recutita* and *Ruta graveolens* are widely traded and used throughout South America and the Old World [19, 62, 124], because of their magical and medicinal qualities for the preparation of remedies used in therapies of soul and body.

The five plant species most commonly used in medicinal beverages, locally known as “horchata” (herbal mixture tea) and “agua aromática” (herbal tea), were native and cultivated [96]. These were *Aloysia triphylla*, *Amaranthus hybridus*, *Pelargonium graveolens*, *Equisetum bogotense*, and *Oreocallis grandiflora*. The availability of these plant species was firstly revealed by their spatial accessibility, explained by their wide spread distribution and cultivation; secondly, it was explicated by their

seasonal stock and the quantities of plant material sold throughout the year. Similarly to Venezuela, Peru, and Bolivia, these plant species are used individually or in herbal mixtures infusions [11, 19, 65], especially to treat different kinds of afflictions related to the nervous, digestive, and genitourinary systems.

Vernacular names of medicinal plant species traded in Loja province

A total of 204 vernacular names were recorded for the 160 medicinal plant species; 59.4 % of the species had at least one name, 36.1 % had two names and 4.5 % had three names. Most of the vernacular names used to identify each plant were in Spanish, followed by a few in Kichwa language from the Andean highlands. The name given to the whole plant is the same name given to the plant’s morphological structure. In the case of gathering wild plant species belonging to the *Malva* genus or cultivated hybrids of the *Fuchsia* genus, the vendors recognized all the species in the genus with the same vernacular name.

The vernacular names of the medicinal plants compiled in this study were compared to others from previous studies conducted at other traditional markets in Bolivia [19], Peru [22, 77], and Ecuador [83] as well as ethnobotanical surveys in the Andean highlands [125] and Spain [124]. The most common five medicinal plant species that share the same cosmopolitan vernacular names were *Dysphania ambrosioides* (“paico”), *Equisetum giganteum* (“cola de caballo”), *Matricaria recutita* (“manzanilla”), *Melissa officinalis* (“toronjil”), and *Rosmarinus officinalis* (“romero”).

Medicinal plants: morphological structures and therapeutic administration

The therapeutic administration recommended by the market vendors revealed the 13 kinds of medicinal plant morphological structures sold, where each one was used to treat a human organ and/or fluid. The most frequently traded morphological structures were branches for 44 medicinal plant species (27.5 %), followed by leaves (25.6 %), flowers (16.9 %), and plants without roots (16.9 %). The less frequently sold morphological structures, available only at 18 open markets, were bark, fruit, inflorescence, latex, seed, stem, style, root, and wood.

The research identified 20 modes of therapeutic administration (Table 5): 14 were prepared using fresh plant material sold at 18 open markets, and six used dried plant material sold at 15 established markets. Oral was the most frequent mode of therapeutic administration (83.8 %), prepared with fresh and/or dry plants, especially in a drink locally known as “bajeada”. Compared to other Andean markets, oral infusions have similar preparations in Bolivia [19] and Cuzco [117].

Other therapeutic administrations were rubbing (16.9 %), topical applications (8.7 %), hot baths after childbirth (5.6 %), and cleaning wounds (5 %). The majority of the prescribed medicinal plants given to patients by vendors were applied without any standardized doses. Only a few elder vendors made warnings about adverse side effects of some medicinal plants, but they never mentioned antidotes. The six most common orally administered medicinal plant species were *Aerva sanguinolenta*, *Amaranthus hybridus*, *Equisetum bogotense*, *Matricaria recutita*, *Melissa officinalis*, and *Oreocallis grandiflora*.

The therapeutic administration, locally known with the term “zumo”, generally refers to the extract of a plant morphological structure or fruit pure juice, thus differentiating “zumo” from “jugo” (juice); the latter is associated with the fruit diluted in an amount of water [104]. The five most popular medicinal plant species sold for “zumo” were *Aerva sanguinolenta* (plant without root), *Cardamine bonariensis* (plant without root), *Peperomia blanda* (plant without root), *Tradescantia zebrina* (leave), and *Verbena litoralis* (branch).

Factor of Informant Consensus (FIC)

In studies related to medicinal plants, the FIC index provides a measure of reliability for the specified statement of evidence regarding the agreement amongst a specific human group [90, 126]. The 160 medicinal plants sold in 33 traditional markets to treat different human ailments were classified into 12 medical categories, with a FIC value assigned to each (Table 3). Three medical categories shared the highest value for FIC = 0.92, which showed a high level of agreement amongst the 196 vendors for 57

medicinal plant species sold to treat the digestive, dermatological, and sensorial systems.

The digestive system has the highest value of used citations by 437 market vendors, who report 37 medicinal plant species. This is related to a high incidence of gastrointestinal ailments in southern Ecuador [86, 127], which may also be associated to stomach cancer, as this is the second most important cause of mortality in the country [128]. When comparing the large FIC values obtained in this study for the digestive system to the most important medical categories reported in previous surveys worldwide [113, 115, 116, 129], the results reveal the importance of medicinal plant species in treating ailments of the digestive system, such as gastric complaints and abdominal pains.

Additionally, therapeutic uses related with the other two medical categories, culture-bound syndromes and general disorders, had relatively high values of use citations by market vendors. The medicinal plant species responsible for this particular status was *Ruta graveolens*, used to treat seven psychosomatic complaints, and *Equisetum bogotense*, used in five common ailments. As it was confirmed by participative observation, local people believe that these two medicinal plant species were efficient in the treatment of 12 particular local ailments, revealing the persistence of cultural beliefs syndromes and the necessity of preventive medicine to avoid common ailments.

Even the lowest FIC values registered in this study, which included hormonal system (0.81), musculoskeletal system (0.83), and gynecological system (0.85), are large when compared to other studies that use this same index [106, 107, 110, 111, 130]. These low FIC values show a low agreement amongst medicinal plant market vendors, specifically in the trade of medicinal plant species associated to the treatment of the symptoms related to these ailments. This data contrasts with the results of a market study in Venezuela [30], where a FIC = 0.91 was registered for the gynecological system, as obtained with 25 interviewed market vendors.

The FIC values between 0.81 and 0.92 demonstrate the strong levels of consensus amongst 196 vendors in the multiple uses of the 160 medicinal plant species sold. Incidentally, even when the taxa number prescribed for a specific illness or disease varied, the majority of the 33 traditional markets had a common pool of regional flora sold to clients within the 12 medical categories. As in Bolivia [19] and Peru [125], the most common medicinal plant species sold were for symptoms related to the digestive system, nervous system, respiratory system, and genitourinary system.

Fidelity Level (FL)

The FL index determined 11 culturally important medicinal plant species in the local population of the Loja

Table 3 Medical categories and Factor of Informant Consensus among vendors of traditional markets at Loja province

N ^o	Medical category ^a	Number of medicinal plant species	Percentage of all medicinal plant species	Use citations bymarket vendor	Percentage of all use citations	FIC ^b
1	Digestive system	37	21.89	437	12.31	0.92
2	Dermatological system	16	9.47	186	5.24	0.92
3	Sensorial system	4	2.37	39	1.10	0.92
4	Culture-bound syndromes	34	20.12	359	10.11	0.91
5	Nervous system	20	11.83	215	6.05	0.91
6	Respiratory system	29	17.16	321	9.04	0.91
7	Genitourinary system	29	17.16	313	8.81	0.91
8	Circulatory system	16	9.47	123	3.46	0.88
9	General disorders	46	27.22	352	9.91	0.87
10	Gynecological system	28	16.57	180	5.07	0.85
11	Musculoskeletal system	13	7.69	71	2.00	0.83
12	Hormonal system	7	4.14	33	0.93	0.81

^aThe medical categories were adapted from International Classification of Diseases catalogue provided by World Health Organization and applied to group 160 medicinal plant species sold at 33 traditional markets within the Loja province, southern Ecuador

^bFIC Factor of Informant Consensus

province (Table 4), as based on the reported uses by 40 or more market vendors to treat 53 illnesses and diseases grouped in 12 medical categories. It was also useful for highlighting the most important species sold in each medical category. In this analysis, the 160 medicinal plant species mentioned by market vendors were considered, and the FL was calculated for each one. A FL of 100 % for a specific medicinal plant species indicates that all of the plant use reports mentioned the same therapeutic administrations to treat an illness or disease.

Two medicinal plant species had a FL = 100 %, *Matricaria recutita* and *Gaiadendron punctatum*, because they were used consistently for ailments in the digestive system and respiratory system, respectively. This may be

due to their greater efficacy in alleviating symptoms and the persistence of ancestral wisdom beliefs in the local population. These two species have considerable agreement amongst market vendors on their particular use and credibility and therefore could be further analyzed for developing pharmafood or pharmaceutical products.

The market vendors had a tendency to rely on 11 medicinal plant species to treat ailments related to nine medical categories. The most important seven had an FL > 60 % and represented *Matricaria recutita*, *Gaiadendron punctatum*, *Ruta graveolens*, *Melissa officinalis*, *Equisetum bogotense*, *Amaranthus hybridus*, and *Viola tricolor*. All these medicinal plant species should be studied to determine the efficacy and safety of all local

Table 4 Most used medicinal plants species for medical categories based on highest fidelity level at Loja province

N ^o	Medicinal plant species	Medical category ^a	I _p	I _t	FL value (%) ^b
1	<i>Matricaria recutita</i> L.	Digestive system	57	57	100
2	<i>Gaiadendron punctatum</i> (Ruiz & Pav.) G. Don	Respiratory system	44	44	100
3	<i>Ruta graveolens</i> L.	Culture-bound syndromes	41	46	89.1
4	<i>Melissa officinalis</i> L.	Nervous system	58	85	68.2
5	<i>Equisetum bogotense</i> Kunth	General disorders	78	116	67.2
6	<i>Amaranthus hybridus</i> L.	Circulatory system	47	73	64.4
7	<i>Viola tricolor</i> L.	Dermatological system	46	76	60.5
8	<i>Borago officinalis</i> L.	Respiratory system	49	120	40.8
9	<i>Oreocallis grandiflora</i> (Lam.) R. Br.	Genitourinary system	40	107	37.4
10	<i>Sambucus nigra</i> L.	Respiratory system	65	186	34.9
11	<i>Aerva sanguinolenta</i> (L.) Blume	Gynecological system	78	238	32.8

^aThe medical categories were adapted from International Classification of Diseases catalogue provided by World Health Organization and applied to 160 medicinal plant species sold at 33 traditional markets within the Loja province, southern Ecuador

I_p = Number of market vendors who independently cited the importance of a specific illness or disease

I_t = Total number of market vendors

^bFL value % = Fidelity Level value percentage (0 = the least, 100 = the highest efficiency)

Table 5 Medicinal plants sold in 33 traditional markets at Loja province, Southern Ecuador

Family and scientific name	Vernacular name	Medica category ^a	Morphological structure used	Therapeutic administration	Geographic status	Voucher number
Acanthaceae						
<i>Dicliptera</i> sp.	Chinche maní, chinche manilla	US	Whole plant	Oral	Native	FT1020
<i>Justicia pectoralis</i> Jacq.	Saucillo, tigresillo	CBS, DERS, GD, NS	Branch	Oral	Native	FT007
Amaranthaceae						
<i>Aerva sanguinolenta</i> (L.) Blume	Escancel	CBS, DS, DERS, GD, GS, NS, US	Plant without root	Oral, douching, topical application, poultice	Introduced	FTMAL008
<i>Alternanthera porrigens</i> (Jacq.) Kuntze	Moradilla	GS, GD	Branch, flower	Oral, bath	Native	FT0010
<i>Amaranthus caudatus</i> L.	Ataco, sangorache	CS, GS, RS	Inflorescence	Oral, bath	Native	FT0278
<i>Amaranthus hybridus</i> L.	Ataco, sangorache	CS, GD, GS, RS, US	Inflorescence	Oral	Native	FTMAL006
<i>Dysphania ambrosioides</i> (L.) Mosyakin & Clemants	Paico	CBS, DS	Branch	Rubbing	Introduced	FT0282
<i>Iresine diffusa</i> Humb. & Bonpl. ex Willd.	Escancel	DERS, DS, GS, US, NS, RS	Branch	Topical application, oral, poultice, wash	Native	FT0280
<i>Iresine herbstii</i> Hook.	Escancel	CBS, DERS, DS, GD, GS NS, US, RS	Branch	Topical application, oral, poultice, wash	Native	FT0486
Anacardiaceae						
<i>Schinus molle</i> L.	Molle	MS	Branch	Cleaning wounds	Introduced	FT194SAR
Apiaceae						
<i>Cyclospermum leptophyllum</i> (Pers.) Sprague ex Britton & P. Wilson	Culantrillo, cominillo	DS	Branch	Oral	Introduced	FT1012
<i>Eryngium</i> sp.	Pomas	DERS, GD	Flower	Cleaning wounds, oral	Native	FT0283
<i>Foeniculum vulgare</i> Mill.	Hinojo	DS	Leaf	Oral	Introduced	FT0025t
<i>Niphogeton dissecta</i> (Benth.) J.F. Macbr.	Culantrillo	DS	Plant without root	Oral	Native	FT0024t
Apocynaceae						
<i>Marsdenia cundurango</i> Rchb. f.	Condurango	GD, DS	Bark	Oral	Native	FT0196
Aquifoliaceae						
<i>Ilex guayusa</i> Loes.	Wayusa	MS	Leaf	Oral	Native	FT0285
Asclepiadaceae						
<i>Orthosia ellemanniae</i> (Morillo) Liede & Meve	Cola de caballo	US, GD	Branch	Oral	Endemic	FT037t
Asphodelaceae						
<i>Aloe vera</i> (L.) Burm. f.	Sábila	DS, DERS, US	Leaf mesophyll	Oral, topical application, bath as gel	Introduced	FT0380
Asteraceae						
<i>Achyrocline hallii</i> Hieron.	Lechugilla	DS	Whole plant	Oral	Endemic	FT0320
<i>Ambrosia arborescens</i> Mill.	Marco, altamiso	CBS, DERS	Branch	Topical application, rubbing	Native	FT045MCAT
<i>Aequatorium jamesonii</i> (S.F. Blake) C. Jeffrey	Guangalo	CBS	Branch	Rubbing	Endemic	FT003ML
<i>Ageratum conyzoides</i> L.	Pedorrera	DS	Branch	Oral	Introduced	FT0286
<i>Ambrosia peruviana</i> Willd.	Marco, altamiso	CBS	Branch	Rubbing	Native	ML005
<i>Aristeguietia persicifolia</i> (Kunth) R.M. King & H. Rob.	Ishpingo, monte de la culebra	CBS	Branch	Rubbing	Endemic	FTG164
<i>Artemisia sodiroi</i> Hieron.	Ajenjo, alcanfor	CBS, RS	Branch	Gargles, rubbing	Native	FT002MZ

Table 5 Medicinal plants sold in 33 traditional markets at Loja province, Southern Ecuador (*Continued*)

<i>Baccharis genistelloides</i> (Lam.) Pers.	Tres filos	CS, HS	Branch	Oral	Native	FT1013
<i>Baccharis latifolia</i> (Ruiz & Pav.) Pers.	Chilca larga	CBS	Branch	Rubbing	Native	FT0288
<i>Baccharis obtusifolia</i> Kunth	Chilca redonda	CBS	Branch	Rubbing	Native	FT0208
<i>Bidens triplinervia</i> Kunth	Ñachig	GS	Plant without root	Oral	Native	FT1008
<i>Chuquiraga jussieui</i> J.F. Gmel.	Chuquiragua	CS, MS	Inflorescence	Oral	Native	FT0318
<i>Cynara cardunculus</i> L.	Alcachofa	HS	Fruit	Oral	Introduced	FT0289
<i>Matricaria recutita</i> L.	Manzanilla	DERS, DS, GD, MS, RS	Plant without root	Cleaning wounds, gargles	Introduced	FT0014t
<i>Tagetes erecta</i> L.	Arrayosa	CBS	Branch, flower	Rubbing	Introduced	FT043MCAT
<i>Tagetes filifolia</i> Lag.	Anís	DS	Whole plant	Oral	Native	FT0987
<i>Tagetes</i> sp.	Chil chil	CBS	Branch	Rubbing	Native	FT0092Q
<i>Tanacetum parthenium</i> (L.) Sch. Bip.	Santa María	CBS	Branch	Rubbing, warm bath	Introduced	FT1195
<i>Taraxacum officinale</i> F.H. Wigg.	Diente de león	DS, US	Whole plant	Oral	Introduced	FT0029t
Boraginaceae						
<i>Borago officinalis</i> L.	Borrajá	RS	Flower, leaf	Oral	Introduced	FT011MAL
<i>Symphytum officinale</i> L.	Consuelda, suelda suelda	MS	Leaf	Hot bath, oral, poultice	Introduced	FT1248
Brassicaceae						
<i>Cardamine bonariensis</i> Pers.	Berro	CS	Plant without root	Oral	Native	FT0003N
<i>Lepidium thurberi</i> Wootton	Chichira	GS	Plant without root	Oral	Native	FT1249
<i>Matthiola incana</i> (L.) W.T. Aiton	Alelí	NS	Flower	Oral	Introduced and cultivated	FT1250
<i>Rorippa nasturtium-aquaticum</i> (L.) Hayek	Berro	CS	Plant without root	Oral	Introduced	FT22t9
Bromeliaceae						
<i>Guzmania</i> sp.	Clavelito de aire	NS	Leaf	Oral	Native	FT001MALA
<i>Tillandsia straminea</i> Kunth.	Flor de Cristo	NS	Flower	Oral	Native	FT1066
Burseraceae						
<i>Bursera graveolens</i> (Kunth) Triana & Planch.	Palo santo	DERS, MS, GS	Wood	Bath, incense	Native	FT961
Cactaceae						
<i>Trichocereus macrogonus</i> (Salm-Dyck) Riccob.	San Pedrillo	CBS	Vascular tissue	Topical application, Oral	Native	FT984
Campanulaceae						
<i>Lobelia</i> cf. <i>decurrens</i> Cav.	Cholo valiente	CBS	Branch	Topical application	Introduced and cultivated	FT1186
Cannaceae						
<i>Canna indica</i> L.	Achira	DG, NS, RS	Leaf	Topical application	Native	FT1251
Caprifoliaceae						
<i>Sambucus nigra</i> L.	Tilo	NS, RS	Flower	Oral	Introduced	FT1252
Caryophyllaceae						
<i>Dianthus caryophyllus</i> L.	Clavel	GS, NS	Flower	Oral	Introduced	FT1253
Chenopodiaceae						

Table 5 Medicinal plants sold in 33 traditional markets at Loja province, Southern Ecuador (*Continued*)

<i>Chenopodium album</i> L.	Palitaria	MS	Leaf	Rubbing	Introduced	FT1254
Commelinaceae						
<i>Callisia gracilis</i> (Kunth) D.R. Hunt	Cachorillo	GD	Leaf	Oral	Native	FT1255
<i>Callisia repens</i> (Jacq.) L.	Calcha	CS, RS	Leaf	Oral	Native	FT51MPAT
<i>Tradescantia zebrina</i> Heynh. ex Bosse	Calcha	CS, NS, RS	Leaf	Topical application, oral	Introduced	FT30t
Crassulaceae						
<i>Kalanchoe gastonis-bonnierei</i> Raym.-Hamet & H. Perrier.	Dulcamara, mala madre	DS, GD	Leaf	Oral	Introduced	FT1024
Cucurbitaceae						
<i>Cucurbita pepo</i> L.	Sambo	DERS	Latex	Topical application	Introduced	FT1256
Cyperaceae						
<i>Cyperus</i> sp.	Díctamo real	CS, MS	Root	Oral	Native	FT1062
Equisetaceae						
<i>Equisetum bogotense</i> Kunth	Cola de caballo, caballo chupa	DS, GD, US	Plant without root	Oral	Native	FT031t
<i>Equisetum giganteum</i> L.	Cola de caballo, caballo chupa	US	Plant without root	Oral	Native	FT1009
Ericaceae						
<i>Bejaria aestuans</i> Mutis ex L.	Payama	GD	Flower	Oral	Native	FT1257
<i>Bejaria resinosa</i> Mutis ex L. f.	Payama	GD	Flower	Oral	Native	FTE012
Euphorbiaceae						
<i>Cnidoscolus aconitifolius</i> (Mill.) I.M. Johnst.	Chaya	CS, DS, GD	Leaf	Oral	Native	FT221
Fabaceae						
<i>Amicia glandulosa</i> Kunth	Nona, urusus	RS	Flower	Oral	Native	HUTPL1975
<i>Desmodium molliculum</i> (Kunth) DC.	San Antonio	GS	Plant without root	Oral	Native	FT38MPAT
<i>Myroxylon balsamum</i> (L.) Harms	Chaquino	DS, GD	Bark	Oral	Native	FT998
<i>Otholobium mexicanum</i> (L. f.) J.W. Grimes	Guallua	DS	Branch	Oral	Native	FT1022
Gentianaceae						
<i>Centaurium erythraea</i> Rafn	Canchalagua	CS	Whole plant	Oral	Introduced	FT1016
Geraniaceae						
<i>Erodium</i> cf. <i>cicutarium</i> (L.) L'Hér. ex Aiton	Agujilla, aujilla	CBS, GD	Branch	Oral	Introduced	FTE001MC
<i>Pelargonium graveolens</i> L'Hér. ex. Aiton	Esencia de rosa	DS, GD, US	Leaf	Oral	Introduced	FT1258
<i>Pelargonium odoratissimum</i> (L.) L'Hér.	Malva olorosa	DS, GD	Branch	Oral	Introduced	FT016t
Juglandaceae						
<i>Juglans neotropica</i> Diels	Nogal	CS, MS	Leaf	Hot bath, oral	Native	FT015MPA
Lamiaceae						
<i>Clinopodium nubigenum</i> (Kunth) Kuntze	Tipo de llano	DS, RS	Plant without root	Oral	Native	FT1014
<i>Clinopodium taxifolium</i> (Kunth) Govaerts	Tipo de cerro	DS, RS	Branch	Oral	Native	FT1259
<i>Hyptis purdiei</i> Benth.	Poleo cerro, poleo negro	CBS	Branch	Rubbing	Native	FT048 MCAT

Table 5 Medicinal plants sold in 33 traditional markets at Loja province, Southern Ecuador (Continued)

<i>Melissa officinalis</i> L.	Toronjil	CBS, NS, RS	Leaf	Oral	Introduced	FT45MPAT
<i>Mentha spicata</i> L.	Hierba buena, menta, menta negra	DS, RS	Leaf	Oral	Introduced	FT1260
<i>Mentha x piperita</i> L.	Hierba buena, menta, menta negra	DS, RS	Leaf	Oral	Introduced	FT1261
<i>Minthostachys mollis</i> (Kunth) Griseb.	Poleo blanco, poleo chiquito	CBS, RS	Branch	Inhalation, oral, rubbing	Native	HUTPL1076
<i>Ocimum basilicum</i> L.	Albahaca, albahaca blanca	GS, US	Leaf	Oral	Introduced	FT46TMPA
<i>Ocimum campechianum</i> Mill.	Albahaca, albahaca negra	GS, US	Leaf	Oral	Native	FT46aMPA
<i>Origanum x majoricum</i> Camb.	Orégano, orégano de castilla	DS	Plant without root	Oral	Introduced	HUTPL5302
<i>Plectranthus unguentarius</i> Codd	Oreganón, orégano grande	DS	Leaf	Oral	Introduced	FT0185
<i>Rosmarinus officinalis</i> L.	Romero	DERS, MS, SS	Branch	Bath after childbirth, cleaning wounds, oral	Introduced	HUTPL3892
<i>Salvia scutellarioides</i> Kunth	Matico	CBS	Branch	Rubbing	Native	HUTPL931
<i>Salvia tiliifolia</i> Vahl	Santa María	CBS	Plant without root	Rubbing	Introduced	FT044MCAT
<i>Scutellaria</i> sp.	Morado	NS	Leaf	Oral	Native	FT003
<i>Thymus vulgaris</i> L.	Tomillo	DS	Plant without root	Oral	Introduced	FT1067
Linaceae						
<i>Linum usitatissimum</i> L.	Linaza	DS, GD, US	Seed	Oral	Introduced	FT47TMPA
Loranthaceae						
<i>Gaiadendron punctatum</i> (Ruiz & Pav.) G. Don	Violeta de cerro, violeta de campo	RS	Flower	Oral	Native	FT0032
Malvaceae						
<i>Alcea rosea</i> L.	Malva	GD	Flower	Oral	Introduced	HUTPL891
<i>Malva arborea</i> (L.) Webb & Berthel.	Malva rosada, malva altea	GD, US	Branch	Oral	Introduced and cultivated	FT042MC
<i>Malva parviflora</i> L.	Malva arbórea, malva blanca	GD, GS, US	Branch, flower	Oral, bath after childbirth, cleaning wounds	Introduced	FT015MCE
Moraceae						
<i>Ficus carica</i> L.	Higo	GS	Leaf	Oral	Introduced	HUTPL1192
Myricaceae						
<i>Morella parvifolia</i> (Benth.) Parra-Os.	Laurel, laurel de cera	CBS, GS	Branch	Oral	Native	FT1207
<i>Morella pubescens</i> (Humb. & Bonpl. ex Willd.) Wilbur	Laurel, laurel de cera	GS	Branch	Oral	Native	FT1023
Myrtaceae						
<i>Corymbia citriodora</i> (Hook.) K.D. Hill & L.A.S. Johnson	Eucalipto oloroso, eucalipto aromático	CBS, RS	Branch	Inhalation, oral, rubbing	Introduced	HUTPL1761
<i>Eucalyptus globulus</i> Labill.	Eucalipto blanco, eucalipto grande	CBS, RS	Branch	Inhalation, oral, rubbing	Introduced	FT1185
<i>Myrcianthes hallii</i> (O. Berg) McVaugh	Arrayán	RS	Leaf	Oral	Native	FT215G
<i>Psidium guajava</i> L.	Guayaba	DS	Fruit	Oral	Native	FT171C
Onagraceae						

Table 5 Medicinal plants sold in 33 traditional markets at Loja province, Southern Ecuador (*Continued*)

<i>Fuchsia harlingii</i> Munz	Pena pena	NS	Flower	Oral	Endemic	HUTPL5784
<i>Fuchsia hybrida</i> hort. T. ex Siebert & Voss	Pena pena	DERS, GD, NS	Flower	Cleaning wounds, oral	Introduced	FT1262
<i>Fuchsia loxensis</i> Kunth	Pena pena	NS	Flower	Oral	Endemic	FT1158
<i>Fuchsia magellanica</i> Lam.	Pena pena	NS	Flower	Oral	Introduced	FT0147
<i>Ludwigia nervosa</i> (Poir.) H. Hara	Flor de reina, mejorana de huerta	DERS, GD, NS	Flower	Oral, bath after childbirth	Native	FT194
<i>Oenothera rosea</i> L'Her. ex Aiton	Shullo	DS, US	Plant without root	Oral	Native	FT53MPAT
Orchidaceae						
<i>Epidendrum jamiesonis</i> Rchb. f.	Flor de Cristo	NS	Flower	Oral	Native	FT1063
<i>Epidendrum</i> sp.	Flor de Cristo	NS	Flower	Oral	Native	FT031t
Piperaceae						
<i>Peperomia blanda</i> (Jacq.) Kunth	Sacha congona	NS	Plant without root	Oral	Native	FT008
<i>Peperomia galioides</i> Kunth	Congona de cerro	NS	Plant without root	Oral	Native	FT124SAR
<i>Peperomia ilaloensis</i> Sodiro	Congona de castilla, congona negra	NS, SS	Plant without root	Topical application, oral	Native	FT01t
<i>Peperomia inaequalifolia</i> Ruiz & Pav.	Congona, congona grande	NS, SS	Plant without root	Topical application, oral	Native	FT1197
<i>Peperomia</i> sp.	Congona de cerro	CBS	Plant without root	Rubbing	Native	FT49MPAT
<i>Piper aduncum</i> L.	Matico	DERS, DS, GS, GD	Branch	Oral	Native	FT1019
<i>Piper carpunya</i> Ruiz & Pav.	Guaviduca de sal	HS, RS, GD	Leaf	Oral	Native	FT500
<i>Piper crassinervium</i> Kunth	Guaviduca de dulce	HS, RS	Leaf	Oral	Native	FT237
Plantaginaceae						
<i>Plantago major</i> L.	Llantén	US, GD	Whole plant	Oral	Introduced	FT13t
Poaceae						
<i>Cymbopogon citratus</i> (DC.) Stapf	Hierba luisa, paja luisa	CS, NS, US	Plant without root	Oral	Introduced	FT011t
<i>Cynodon dactylon</i> (L.) Pers.	Gramma dulce	CBS, US, GD	Plant without root	Oral	Introduced	FT008MCE
<i>Zea mays</i> L.	Pelo de choclo	US	Style	Oral	Introduced	FT1065
Polygalaceae						
<i>Polygala paniculata</i> L.	Mentol	MS	Whole plant	Poultice	Native	FT20t
Polypodiaceae						
<i>Niphidium crassifolium</i> (L.) Lellinger	Calaguala	DS, US	Root	Oral	Native	FT40T
Proteaceae						
<i>Oreocallis grandiflora</i> (Lam.) R. Br.	Cucharillo	CBS, HS, US, GD	Flower	Oral	Native	FT04t
Pteridaceae						
<i>Adiantum poiretii</i> Wikstr.	Culantrillo pata negra	GS	Leaf	Oral	Native	FT1018
<i>Adiantum raddianum</i> C. Presl	Culantrillo	GS, US, GD	Leaf	Oral	Native	MCELI35-T
<i>Cheilanthes bonariensis</i> (Willd.) Proctor.	Helecho congona	GS	Leaf	Oral, bath after childbirth	Native	MPA52T
<i>Jamesonia</i> sp.	Nido de abeja	GS	Leaf	Oral	Native	FT010
<i>Notholaena sulphurea</i> (Cav.) J. Sm.	Grano de oro	GS	Leaf	Bath after childbirth, oral	Native	FT009

Table 5 Medicinal plants sold in 33 traditional markets at Loja province, Southern Ecuador (*Continued*)

<i>Pityrogramma calomelanos</i> (L.) Link	Doradilla del sol	GS	Leaf	Bath after childbirth, oral	Native	FT013
<i>Pityrogramma ebenea</i> (L.) Proctor	Doradilla plateada, luna plateada	GS	Leaf	Bath after childbirth, oral	Native	FT014
<i>Trachypteris induta</i> (Maxon) R.M. Tryon & A.F. Tryon	Pata de gallina	GS	Leaf	Oral, bath after childbirth	Native	FT012
Rosaceae						
<i>Alchemilla aphanoides</i> Mutis ex L. f.	Saucillo	NS	Branch	Topical application, oral	Native	FT007
<i>Eriobotrya japonica</i> (Thunb.) Lindl.	Níspero, míspero	US	Leaf	Oral	Introduced	FT1063
<i>Margyricarpus pinnatus</i> (Lam.) Kuntze	Perilla	RS, DERS	Plant without root	Oral	Native	FT463
<i>Rosa cymosa</i> Tratt.	Rosa	DERS, GD, SS, US	Flower	Cleaning wounds, oral	Introduced	FT274
<i>Sanguisorba minor</i> subsp. <i>muricata</i> (Bonnier & Layens) Briq.	Pimpinela	NS	Leaf	Oral	Introduced	FT014MZ
Rubiaceae						
<i>Cinchona pubescens</i> Vahl	Cascarilla	RS	Bark	Oral	Native	FT118
Rutaceae						
<i>Citrus x junos</i> Siebold ex Tanaka	Naranja agria	DERS	Fruit	Oral	Introduced	FT007t
<i>Ruta graveolens</i> L.	Ruda	CBS, DS, GS	Branch	Bath after childbirth, oral, rubbing	Introduced	FT002
Smilacaceae						
<i>Smilax</i> sp.	Zarzaparrilla	GD, US	Root	Oral	Native	FT1263
Solanaceae						
<i>Brugmansia sanguinea</i> (Ruiz & Pav.) D. Don	Guando	CBS	Flower, leaf	Rubbing	Native	FT028ML
<i>Brugmansia x candida</i> Pers.	Guando blanco	CBS	Flower, leaf	Rubbing	Native	FT210
<i>Cestrum mariquitense</i> Kunth	Sauco negro	CS, GD	Branch	Oral, topical application	Native	FT210
<i>Cestrum racemosum</i> Ruiz & Pav.	Sauco blanco	CBS, GD	Branch	Rubbing	Native	FTS177
<i>Solanum americanum</i> Mill.	Mortño	DS, GD, NS, RS	Branch, bud	Oral, warm bath	Native	FT36t
<i>Solanum nigrescens</i> M. Martens & Galeotti	Mortño	DS, NS, RS	Bud	Oral	Native	FT1264
<i>Solanum pimpinellifolium</i> L.	Monte del gallinazo	CBS	Branch	Rubbing	Native	FT207
Tiliaceae						
<i>Triumfetta semitriloba</i> Jacq.	Abrojo, cadillo, mostrante	GD, US	Leaf	Oral	Native	FT39T
Urticaceae						
<i>Urtica dioica</i> L.	Chine	CS, MS, US,	Plant without root	Oral, rubbing	Introduced	FT1064
<i>Urtica urens</i> L.	Chine	CS, MS, US	Plant without root	Oral, rubbing	Introduced	FT275
Valerianaceae						
<i>Valeriana microphylla</i> Kunth	Valeriana, valeriana de cerro	NS	Plant without root	Oral	Native	FT1015
<i>Valeriana pyramidalis</i> Kunth	Valeriana	NS	Root	Oral	Native	FT991
Verbenaceae						
<i>Aloysia triphylla</i> Royle	Cedrón	NS	Branch	Oral	Native	FT1265
<i>Phyla scaberrima</i> (A. Juss. ex Pers.) Moldenke	Buscapina, novalgina	DS	Plant without root	Oral	Introduced	FT1240

Table 5 Medicinal plants sold in 33 traditional markets at Loja province, Southern Ecuador (*Continued*)

<i>Verbena litoralis</i> Kunth.	Verbena	CBS, GD	Branch	Oral, rubbing	Native	FT1268
Violaceae						
<i>Viola odorata</i> L.	Violeta, violeta de jardín	RS	Flower, leaf	Oral	Introduced	FT1266
<i>Viola tricolor</i> L.	Pensamiento	DERS, GD, RS	Flower	Cleaning wounds, oral	Introduced	FT1267
Zingiberaceae						
<i>Hedychium coronarium</i> J. Koenig	Caña agria	US	Stem	Oral	Introduced	FT002t

^aMedical category: CBS Culture-bound syndromes, CS Circulatory system, DERS Dermatological system, DS Digestive system, GD General disorders, GS Gynecological system, HS: Hormonal system, MS Musculoskeletal system, NS Nervous system, RS Respiratory system, SS Sensorial system, US: Urinary system

reported medical uses and also evaluated by phytochemical and pharmacological tests as well as bioactivity essays and toxicity studies.

Conclusions

Within the Loja province, people continue to use traditional medicine by consuming particular medicinal plant species sold at local markets. This also shows that sociocultural customs are strongly expressed in ancestral practices of wellbeing. Proof of the former was found by analyzing responses from 196 vendors who sold 160 fresh or dried medicinal plant species material to treat a wide spectrum of illnesses and diseases. The plant resources functioned as palliatives or, in some cases, curatives to both somatic and psychosomatic health afflictions. Nowadays in Ecuador, all stake-holders related with dynamics of traditional markets networks (e.g., all kinds of vendors and local people), who are frequently using medicinal plants require detailed research for a safety and serious use.

In the case of culture-bound syndromes plus complications with a diversity of systems such as digestive, sensorial, dermatological, respiratory, genitourinary and/or nervous, there are usually no precise therapeutical prescriptions. The separation between illnesses and diseases is very small, especially with regards to what the therapeutic administration and medical treatment should be. An example is the *Ruta graveolens* based-remedy, which is used to relieve “evil eye”, “evil air”, “frights”, and “calor encerrado” plus menstrual disorders.

The agreement among 196 market vendors in the use of seven specific most sold medicinal plant species for the 12 medical categories is fairly high, especially when the largest values of FL (60.5 %–100 %) and FIC (0.81–0.92) indexes are combined. A total of seven plant species with a high FL value for treating gastro-intestinal diseases are under investigation for their pharmacological properties by the Applied Chemical Department of the UTPL research team.

For future efforts, it should be important to focus on correlating the values of FL and FIC with the incidence of local ailments, as this will be useful to establish public health

policies related with the trade of medicinal plant species. This initiative will be effective to support traditional medicine and its therapeutic repertoire. The first step will be to choose the medicinal plant species with widespread and consistent medicinal use in southern Ecuador and to study their therapeutical applications with physicians and scientists, primarily to identify bioactive compounds.

The evidence presented in this study reaffirms the relationship between ancestral wisdom and traditional medicine, particularly in local markets within the Loja province. In fact, it is important to stress how medicinal plant resources are crucial for local people in 13 cities within the Loja province; also, it is important to understand why a high percentage of them practice auto-medication. Reasons for the maintenance of traditional markets include lower cost of plant products, confidence in traditional medicine, and/or sociocultural environment.

This research is the first contribution to understanding from the ethnobotanical point of view the human-plant dynamics of traditional markets within the Loja province, where medicinal plants have a substantial role in the lives of local people. The trade demand of medicinal plants and their derivatives over the next few years could increase, leading to the over-harvesting of wild plant species and could perhaps even endanger natural populations, (e.g., *Oreocallis grandiflora*). Sustainable management of wild medicinal plants is important for their diversity conservation and in order to avoid their extinction, particularly in the case of highly used species in traditional medicine.

Acknowledgments

The first author was supported by a doctoral fellowship with the “Universidad Técnica Particular de Loja” and the “Secretaría de Educación Superior Ciencia, Tecnología e Innovación”, and a grant from the “Universidad Politécnica de Madrid” (UPM-BSCH 2007–2008).

We extend our gratitude to the 196 vendors of the 33 traditional markets within the Loja province for sharing their wisdom.

With regard to identifying a fern species, the authors especially want to thank Dr. Benjamin Ollgaard (Biologisk Institut, Herbariet, Aarhus Universitet) for his taxonomic collaboration.

We also would like to provide a sincere recognition for the invaluable professional support from the authors to Dr. Pablo Jarrín (Universidad Regional Amazónica IKIAM), who provided useful comments to the analysis and interpretation of the measured indexes.

We would like to thank the staff at the Herbarium of the “Universidad Técnica Particular de Loja” (HUTPL) and “Universidad Nacional de Loja” (LOJA) for their taxonomical advice, particularly to Omar Cabrera and Bolívar Merino; David Duncan for his insightful comments to this manuscript; David Donoso and Daniel Horlacher who provided valuable editorial contributions; Natalia Donoso who helped to prepare the former version of the manuscript; and Raúl Sinche, Pablo Jaramillo, Ángel Cuenca, Nixon Cumbicus, Irene Jiménez, and Welinton Agreda for their cooperation during the field trips.

Funding

The first author was supported by a doctoral fellowship with the “Universidad Técnica Particular de Loja” and the “Secretaría de Educación Superior Ciencia, Tecnología e Innovación”, and a grant from the “Universidad Politécnica de Madrid” (UPM-BSCH 2007–2008).

Availability of data and materials

The dataset supporting the conclusions of this article is included within the article and its Table 5.

Authors' contributions

The first author carried out the fieldwork research for this study. All authors reviewed literature, analyzed the data, prepared the manuscript, provided revisions, and approved the final manuscript.

Competing interest

The authors declare that they have no competing interest.

Consent for publication

Not applicable.

Ethics approval and consent to participate

All the vendors who decided to collaborate were interviewed according to mutually agreed conditions and under Ecuador's rights, especially with regards to the Convention on Biological Diversity (CDB).

Author details

¹Departamento de Ciencias Naturales, Universidad Técnica Particular de Loja, San Cayetano Alto, CP: 1101608 Loja, Ecuador. ²Departamento de Química, Universidad Técnica Particular de Loja, San Cayetano Alto, Loja, Ecuador. ³Institute for Tropical Ecology and Conservation, University of Florida, Gainesville, FL 32611-0430, USA. ⁴Área de Biodiversidad y Conservación, Departamento de Biología y Geología, Universidad Rey Juan Carlos, C/ Tulipán s/n, 28933 Mostoles, Spain. ⁵Departamento de Biología (Botánica), Universidad Autónoma de Madrid, Campus de Canto Blanco, E-28049 Madrid, Spain.

Received: 28 January 2016 Accepted: 28 June 2016

Published online: 05 July 2016

References

- Hooper D, Field H, Dahlgren BE. *Useful Plants and Drugs of Iran and Iraq*. Volume 9. Chicago: 7 Field Museum of Natural History; 1937.
- Linares E, Bye RA. A study of four medicinal plant complexes of Mexico and adjacent United States. *J Ethnopharmacol*. 1987;19:153–83.
- Nicholson MS, Arzeni CB. The market medicinal plants of Monterrey, Nuevo León México. *Econ Bot*. 1993;47:184–92.
- Martin GJ. *Ethnobotany: A Methods Manual*, vol. 1. London: Earthscan Publications; 1995.
- Cunningham AB. *Applied Ethnobotany: People Wild Plant Use and Conservation*. People and Plants Conservation. London: Earthscan Publications; 2001.
- Parente CET, da Rosa MMT. Plantas comercializadas como medicinais no Município de Barra do Piraí, RJ. *Rodriguésia*. 2001;52(80):47–59.
- Hanlidou E, Karousou R, Kleftoyanni V, Kokkini S. The herbal market of Thessaloniki (N Greece) and its relation to the ethnobotanical tradition. *J Ethnopharmacol*. 2004;91:281–99.
- Williams VL, Witkowski ETF, Balkwill K. The use of incidence-based species richness estimators, species accumulation curves and similarity measure to appraise ethnobotanical inventories from South Africa. *Biodivers Conserv*. 2007;16:2495–513.
- Williams VL, Balkwill K, Witkowski ETF. Size-class prevalence of bulbous and perennial herbs sold in the Johannesburg medicinal plant markets between 1995 and 2001. *South African J Bot*. 2007;73:144–55.
- Williams VL, Witkowski ETF, Balkwill K. Volume and financial value of species traded in the medicinal plant market of Gauteng South Africa. *Int J Sustain Dev World Ecol*. 2009;14:584–603.
- Giraldo D, Baquero E, Bermúdez A, Oliveira-Miranda MA. Medicinal plant trade characterization in popular markets of Caracas. *Venezuela Acta Bot Venez*. 2009;32(2):267–301.
- Silalahi M, Walujo EB, Supriatna J, Mangunwardoyo W. The local knowledge of medicinal plants trader and diversity of medicinal plants in the Kabanjahe traditional market, North Sumatra, Indonesia. *J Ethnopharmacol*. 2015;175:432–43.
- Whitaker TW, Cutler HC. Food plants in a Mexican market. *Econ Bot*. 1966;20(1):6–16.
- White A. *Hierbas del Ecuador*. Quito: Libri Mundi; 1982. p. 379.
- Bye R, Linares E. The role of plants found in the Mexican markets and their importance in ethnobotanical studies. *J Ethnobiol*. 1983;3(1):1–13.
- Van den Berg MA. Ver-o-peso: the ethnobotany of an Amazonian market. *Adv Econ Bot*. 1984;1:140–9.
- Bye RA. Medicinal plants of the Sierra Madre: comparative study of Tarahumara and Mexican market plants. *Econ Bot*. 1986;40(1):103–24.
- Pulido MT, Cavalier J. Comercialización de productos vegetales no maderables en los mercados de Leticia y Florencia, Amazonía colombiana. In: Duivenvoorden JF, Balslev H, Cavalier J, Grandez C, Tuomisoto H, Valencia R, editors. *Evaluación de recursos vegetales no maderables en la Amazonía noroccidental*. Amsterdam: IBED, Universidad de Amsterdam; 2001. p. 265–310.
- Macía MJ, García E, Vidaurre PJ. An ethnobotanical survey of medicinal plants commercialized in the markets of La Paz and El Alto, Bolivia. *J Ethnopharmacol*. 2005;97:337–50.
- de Azevedo SKS, Silva IM. Plantas medicinais e de uso religioso comercializadas em mercados e feiras livres no Rio de Janeiro, RJ, Brasil. *Acta Bot Bras*. 2006;20(1):185–94.
- van Andel T, Behari-Ramdas J, Havinga R, Groenendijk S. The medicinal plant trade in Suriname. *Ethnobot Res Appl*. 2007;5:351–72.
- Bussmann RW, Sharon D, Vandebroek I, Jones A, Revene Z. Health for sale: the medicinal plant markets in Trujillo and Chiclayo. *Northern Peru J Ethnobiol Ethnomed*. 2007;3(37):1–9.
- Albuquerque UP, Monteiro JM, Ramos MA, Cavalcanti de Amorim EL. A pesquisa etnobotânica em mercados e feiras livres. In: Albuquerque UP, Paiva Lucena RF, da Cunha LV FC, editors. *Métodos e técnicas na pesquisa etnobotânica*. Brasil: Comunigraf Ed Recife; 2008. p. 145–60.
- Lee S, Xiao C, Pei S. Ethnobotanical survey of medicinal plants at periodic markets of Honghe Prefecture in Yunnan Province, SW China. *J Ethnopharmacol*. 2008;2008(117):362–77.
- Kool A, de Boer HJ, Krüger Å, Rydberg A, Abbad A, Björk L, Martin G. Molecular identification of commercialized medicinal plants in Southern Morocco. *PLoS One*. 2012;7:e39459.
- Leitão F, Leitão SG, Fonseca-Kruel VS, Silva IM, Martins K. Medicinal plants traded in the open-air markets in the State of Rio de Janeiro, Brazil: An overview on their botanical diversity and toxicological potential. *Rev Bras*. 2014;24(2):225–47.
- Coelho Ferreira M. Lê marché des plantes médicinales à Manaus. In *La Foret en Jeu: l'extractivisme en Amazonie centrale*. Edited by Laure E. Paris, Francia: UNESCO; 1996:173–177.
- Puentes JP, Hurrell JA. Plantas andinas y sus productos comercializados con fines medicinales y alimentarios en el Área Metropolitana Buenos Aires-La Plata, Argentina. *Bol Latinoam y del Caribe Plantas Med y Aromat*. 2015;14:206–236.
- Albuquerque UP, Monteiro JM, Ramos MA, Amorim EL. Medicinal and magic plants from a public market in northeastern Brazil. *J Ethnopharmacol*. 2007;110:76–91.
- Ledezma J, Dávila M, Mondragón A, Castillo MA, Ramírez LA. Registro y conocimiento etnobotánico de plantas medicinales por expendedores de Barquisimeto. *Venezuela Boletín del Cent Investig Biológicas*. 2007;41(4):531–44.
- Shrestha S. Marketplace plants used in ceremonial cleansing among Andean Qechuans of Ecuador. Virginia, USA: Marshall University; 2007.
- Bussmann RW, Sharon D, Ly J. From garden to market? The cultivation of native and introduced medicinal plant species in Cajamarca, Peru and implications for habitat conservation. *Ethnobot Res Appl*. 2008;6:351–61.

33. Amiri MS, Joharchi MR. Ethnobotanical investigation of traditional medicinal plants commercialized in the markets of Mashhad, Iran. *Avicena J Phytomedicine*. 2015;3:254–271.
34. Monteiro JM, de Lima AE, Amorim ELC, De Albuquerque UP. Local markets and medicinal plant commerce: a review with emphasis on Brazil. *Econ Bot*. 2010;64:352–66.
35. Lima PGC, Coelho-Ferreira M, Oliveira R. Plantas medicinais em feiras e mercados públicos do Distrito Florestal Sustentável da BR-163, estado do Pará, Brasil. *Acta Bot Bras Brasília*. 2011;25(2):422–34.
36. Van Andel T, Myren B, Van Onselen S. Ghana's herbal market. *J Ethnopharmacol*. 2012;140:368–78.
37. Meléndez M, Alvarado S, de Rojas LC. Identificación y conocimiento de las plantas medicinales expandidas en los mercados principal y libre de Maracay, estado Aragua, Venezuela. *Rev Fac Agron (Universidad Cent Venez)*. 2012;38:64–70.
38. Ladio AH, Molares S, Ochoa J, Cardoso B. Etnobotánica aplicada en Patagonia: la comercialización de malezas de uso comestible y medicinal en una feria urbana de San Carlos de Bariloche (Río Negro, Argentina). *Bol Latinoam Caribe Plantas Med Aromát*. 2013;12(1):24–37.
39. Albuquerque UP, Monteiro JM, Ramos MA, de Amorim ELC, Alves RRN. Ethnobiological Research in Public Markets. *Methods Tech Ethnobiol Ethnoecology*. 2014;110:367–378.
40. Quiroz D, Towns A, Legba SI, Swier J, Brière S, Sosef M, van Andel T. Quantifying the domestic market in herbal medicine in Benin, West Africa. *J Ethnopharmacol*. 2014;151:1100–8.
41. Towns AM, Quiroz D, Guinée L, de Boer H, van Andel T. Volume, value and floristic diversity of Gabon's medicinal plant markets. *J Ethnopharmacol*. 2014;155:1184–93.
42. Van Andel TR, Croft S, van Loon EE, Quiroz D, Towns AM, Raes N. Prioritizing West African medicinal plants for conservation and sustainable extraction studies based on market surveys and species distribution models. *Biol Conserv*. 2015;181:173–81.
43. Bussmann RW, Paniagua-Zambrana NY, Moya Huanca AL. Dangerous confusion- "Cola de Caballo"-Horsetail, in the Markets of La Paz. *Bolivia Econ Bot*. 2015;20(10):1–5.
44. Randriamiharisoa MN, Kuhlman AR, Jeannoda V, Rabarison H, Rakotoarivelo N, Randrianarivony T, Rakotoarivony F, Randrianasolo A, Bussmann RW. Medicinal plants sold in the markets of Antananarivo, Madagascar. *J Ethnobiol Ethnomed*. 2015;11:1–12.
45. Lima PGC, Coelho-Ferreira M, da Silva Santos R. Perspectives on Medicinal Plants in Public Markets across the Amazon: A Review. *Econ Bot*. 2016;70: 64–78.
46. WHO - World Health Organization: Traditional Medicine Strategy 2002–2005. 2002. [http://www.wpro.who.int/health_technology/book_who_traditional_medicine_strategy_2002_2005.pdf]. Accessed 15 Oct 2015.
47. Ocampo RA. Situación actual del comercio de plantas medicinales en América Latina. *Boletín Latinoam y del Caribe Plantas Med y Aromáticas*. 2002;1(4):35–40.
48. Salomon F. Los señores étnicos de Quito en la época de los Incas. *Pendoneros*. 1980;10:1–370.
49. De Feo V. Medicinal and magical plants in the northern Peruvian Andes. *Fitoterapia*. 1991;LXIII(5):417–40.
50. Bussmann RW, Sharon D. Traditional medicinal plant use in Northern Peru: tracking two thousand years of healing culture. *J Ethnobiol Ethnomed*. 2006;2(47):1–18.
51. Bussmann R, Sharon D. *Plantas de los cuatro vientos, flora mágica y medicinal del Perú*. Trujillo-Perú: Graficart srl; 2007.
52. Bussmann RW, Sharon D. Shadows of the colonial past-diverging plant use in Northern Peru and Southern Ecuador. *J Ethnobiol Ethnomed*. 2009;5(4):1–17.
53. Bussmann RW, Kuhlman A, Glenn A, Meyer K, Rothrock A, Townesmith A, Chait G, Malca G. Phyto-chemical analysis of peruvian medicinal plants - Análisis fitoquímico de plantas medicinales peruanas. *Arnaldoa*. 2009;16(1):105–10.
54. Bussmann RW, Sharon D. Markets, healers, vendors, collectors: The sustainability of medicinal plant use in Northern Peru. *Mt Res Dev*. 2009; 29(2):128–34.
55. Bussmann RW, Sharon D. Naming a phantom – the quest to find the identity of Ulluchu, an unidentified ceremonial plant of the Moche culture in Northern Peru. *J Ethnobiol Ethnomed*. 2009;5:8.
56. Bussmann RW, Sharon D, Díaz PD, Barocio Y. Peruvian plants canchalagua (*Schkuhria pinnata* (Lam.) Kuntze), hercampuri (*Gentianella alborosea* (Gilg.) Fabris), and corpus way (*Gentianella bicolor* (Wedd.) J. Pringle) prove to be effective in the treatment of acne. *Arnaldoa*. 2008;15(1):149–52.
57. Bussmann RW, Sharon D: From collection to market and cure—an interdisciplinary study of traditional plant use in Northern Peru. In de Albuquerque UP and Hanazaki N (eds) *Recent Developments and Case Studies in Ethnobotany*. Brazilian Society of Ethnobiology and Ethnoecology and Publication Group of Ecology and Applied Ethnobotany. Recife; 2010
58. Bussmann RW, William L Brown Center's Global Discovery Program. *Markets, Healers, Vendors, Collectors and Modern Applications – Medicinal Plant Use in Northern Peru*. *Planta Med*. 2010;76:53.
59. Bussmann RW, Glenn A, Meyer K, Kuhlman A, Townesmith A. Herbal mixtures in traditional medicine in Northern Peru. *J Ethnobiol Ethnomed*. 2010;6(10):2–11.
60. Bussmann RW, Glenn A. Cooling the heat - traditional remedies for malaria and fever in Northern Peru. *Ethnobot Res Appl*. 2010;8:125–34.
61. Bussmann RW, Glenn A, Sharon D. Antibacterial activity of medicinal plants of Northern Peru - Can traditional applications provide leads for modern science? *Indian J Tradit Knowl*. 2010;9(4):742–53.
62. Bussmann RW, Glenn A, Sharon D. Healing the body and soul: Traditional remedies for "magical" ailments, nervous system and psychosomatic disorders in Northern Peru. *African J Pharm Pharmacol*. 2010;4(9):580–629.
63. Bussmann RW, Glenn A. Medicinal plants used in northern Peru for reproductive problems and female health. *J Ethnobiol Ethnomed*. 2010;6:30.
64. Bussmann RW, Malca-García G, Glenn A, Sharon D, Chait G, Díaz D, Pourmand B, Jonat B, Somogy S, Guardado G, Aguirre C, Chan R, Meyer K, Rothrock A, Townesmith A, Effio-Carbajal J, Frías-Fernandez F, Benito M. Minimum inhibitory concentration of medicinal plants used in Northern Peru as antibacterial remedies. *J Ethnopharmacol*. 2010;132(1):101–8.
65. Bussmann RW, Glenn A. Plants used for the treatment of gastro-intestinal ailments in northern peruvian ethnomedicine. *Arnaldoa*. 2010;17(2):255–70.
66. Bussmann R, Glenn A. Peruvian medicinal plants for the treatment of liver and gallbladder ailments. *Arnaldoa*. 2010;17(2):243–53.
67. Bussmann RW, Glenn A. Medicinal plants used in Peru for the treatment of respiratory disorders. *Rev Peru Biol*. 2011;17(2):331–46.
68. Bussmann RW, Glenn A, Sharon D, Chait G, Díaz D, Pourmand K, Jonat B, Somogy S, Guardado G, Aguirre C, Chan R, Meyer K, Rothrock A, Townesmith A. Proving that traditional knowledge works: The antibacterial activity of Northern Peruvian medicinal plants. *Ethnobot Res Appl*. 2011;9:67–96.
69. Bussmann RW, Glenn A. Fighting pain: Traditional peruvian remedies for the treatment of asthma, rheumatism, arthritis and sore bones. *Indian J Tradit Knowl*. 2011;10(3):397–412.
70. Bussmann RW, Malca G, Glenn A, Sharon D, Nilsen B, Parris B, Dubose D, Ruiz D, Saleda J, Martínez M, Carillo L, Walker K, Kuhlman A, Townesmith A. Toxicity of medicinal plants used in traditional medicine in Northern Peru. *J Ethnopharmacol*. 2011;137:121–40.
71. Bussmann RW, Glenn A. Medicinal plants used in northern Peru for the treatment of bacterial and fungal infections and inflammation symptoms. *J Med Plants Res*. 2011;5(8):1297–304.
72. Bussmann RW. Traditional knowledge for modern ailments – plants used for the treatment of diabetes and cancer in northern Peru. *J Med Plants Res*. 2011;5(31):6916–30.
73. Bussmann RW, Glenn A. Traditional medicinal plants used in northern Peru for kidney problems and urinary infections. *Arnaldoa*. 2011;18(1):77–94.
74. Bussmann RW, Glenn A. Mending the heart. Plants used in peruvian ethnomedicine for heart disease, blood pressure irregularities and cholesterol control. *Arnaldoa*. 2011;18(2):167–78.
75. Bussmann RW, Paniagua-Zambrana N, Rivas Chamorro M, Molina Moreira N, Cuadros Negri MLR, Olivera J. Peril in the market-classification and dosage of species used as anti-diabetics in Lima, Peru. *J Ethnobiol Ethnomed*. 2013;9(1):37.
76. Bussmann RW. The globalization of traditional medicine in northern Peru: From shamanism to molecules. *Evidence-Based Complement Altern Med*. 2013;2013:1–46.
77. Bussmann R, Sharon D. Two decades of ethnobotanical research in southern Ecuador and northern Peru. *Ethnobiol Conserv*. 2014;3:1–50.
78. Bussmann RW, Paniagua-Zambrana N, Castañeda Sifuentes RY, Prado Velazco YA, Mandujano J. Health in a pot - The ethnobotany of emollientes and emollienteros in Peru. *Econ Bot*. 2015;69(1):83–8.
79. Azahuanche FP, Rodríguez Avalos F, León Aponte G, Sharon D, Bussmann RW, Willsky GR, Guerrero G, Willner K, Castro Dionicio I. Phytochemical and antibacterial study of medicinal plant mixtures. In search of new components. *Pueblo Cont*. 2012;23(2):339–44.
80. Cerón Martínez CE. Plantas medicinales de los Andes ecuatorianos. In: Moraes RM, Øllgaard B, Kvist LP, Borchsenius F, Balslev H, editors. *Botánica Económica de los Andes Centrales*. La Paz: Universidad Mayor de San Andrés; 2006. p. 285–93.

81. Buitrón X. Ecuador: uso y comercio de plantas medicinales, situación actual y aspectos importantes para su conservación. Cambridge: TRAFFIC International; 1999.
82. de la Torre L, Navarrete H, Muriel P, Macía MJ, Balslev H. Enciclopedia de las plantas útiles del Ecuador. Quito: Herbario QCA and Herbario AAU; 2008.
83. Pustscher J, Vogl C. An ethnobotanical survey on herbal medicine at Quito markets (Ecuador). In Proceedings of the 4th International Congress of Ethnobotany. Edited by Füsün E. Istanbul, Turkey: Ege Yayinlari; 2005:233–239.
84. Naranjo P, Coba JL. Etnomedicina en el Ecuador, vol. 3. Corporación Editora Nacional: Quito; 2003.
85. de la Torre L, Muriel P, Balslev H. Etnobotánica en los Andes del Ecuador. In: Moraes RM, Øllgaard B, Kvist LP, Borchsenius F, Balslev H, editors. Botánica Económica de los Andes Centrales. La Paz: Universidad Mayor de San Andrés; 2006. p. 246–67.
86. Bussmann RW, Sharon D. Traditional medicinal plant use in Loja province, southern Ecuador. *J Ethnobiol Ethnomed*. 2006;2:44.
87. Van den Eynden V. Use and management of edible non-crop plants in southern Ecuador. Ghent: Thesis, Ghent University; 2004.
88. Elleman L. Saraguroerne og deres planter—en gruppe højlandsindianeres anvendelse af den naturlige vegetation. Aarhus: Thesis, Aarhus University; 1990.
89. Tene V, Malagón O, Finzi PV, Vidari G, Armijos C, Zaragoza T. An ethnobotanical survey of medicinal plants used in Loja and Zamora-Chinchipe, Ecuador. *J Ethnopharmacol*. 2007;111:63–81.
90. Trotter RT, Logan MH. Informant consensus: A new approach for identifying potentially effective medicinal plants. In: Etkin NL, editor. *Plants in Indigenous Medicine and Diet*. New York: Redgrave Publishers; 1986. p. 91–112.
91. Friedman J, Yaniv Z, Dafni A, Palewitch D. A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among Bedouins in the Negev desert. *Israel J Ethnopharmacol*. 1986;16(2–3):275–87.
92. Cueva Ortiz J, Chalan LA. Cobertura vegetal y uso actual del suelo de la provincia de Loja. Informe Técnico, Naturaleza y Cultura Internacional: Loja; 2010.
93. Instituto Nacional de Estadística: Censo de Población y Vivienda del Ecuador. 2010
94. Tapia Paredes JE. Perspectivas para la protección normativa de los conocimientos tradicionales frente al régimen de propiedad intelectual en el comercio internacional. Sede Ecuador, Quito: Thesis, Universidad Andina Simón Bolívar; 2014.
95. International Society of Ethnobiology (2006). International Society of Ethnobiology Code of Ethics (with 2008 additions). [http://www.ethnobiology.net/wp-content/uploads/ISECodeofEthics_Spanish.pdf]
96. Jørgensen PM, León-Yáñez S, editors. *Catalogue of Vascular Plants of Ecuador*. Saint Louis: Monographs in Systematic Botany from the Missouri Botanical Garden; 1999.
97. Tropicos. Missouri Botanical Garden plant data base [<http://www.tropicos.org>]. Accessed from 05 Jan 2012 to 20 Oct 2012.
98. Harling G, Sparre B (eds.). *Flora of Ecuador*. Opera Botanica Series B (1–4, 1973–1975; 5–25, 1976–1986). Department of Systematic Botany, University of Göteborg and Botany Section, Riksmuseum, Stockholm; 1973–1986
99. Harling G, Anderson L (eds.). *Flora of Ecuador (26–77)*. Department of Systematic Botany, University of Göteborg, Göteborg; 1987–2006
100. Harling G, Persson C (eds.). *Flora of Ecuador (78–87)*. Department of Systematic Botany, University of Göteborg, Göteborg; 2007–2010
101. Persson C, Ståhl B. *Flora of Ecuador (91, 22–26)*. Göteborg: Botanical Institute, University of Göteborg; 2014.
102. International Classification of Diseases. World Health Organization [<http://apps.who.int/classifications/icd10/browse/2015/en>]. Accessed from 20 to 25 Feb 2015.
103. World Health Organization: International Statistical Classification of Diseases and Related Health Problems. In WHO Library Cataloguing. Volume 2. 10th Revision; 2010:39
104. Alexiades MN. Collecting ethnobotanical data: An introduction to basic concepts and techniques. In: Alexiades MN, editor. *Selected Guidelines for Ethnobotanical Research*. New York: The New York Botanical Garden; 1996. p. 1–57.
105. Rios M, Koziol MJ, Pedersen HB, Granda G. *Plantas útiles del Ecuador: aplicaciones, retos y perspectivas/Useful plants of Ecuador: Applications, challenges, and perspectives*. Quito: Ediciones Abya-Yala; 2007.
106. Heinrich M, Ankli A, Frei B, Weimann C, Sticher O. Medicinal plants in Mexico: Healers' consensus and cultural importance. *Soc Sci Med*. 1998;47(11):1859–71.
107. Gradé JT, Tabuti JRS, Van Damme P. Four footed pharmacists: Indications of self-medicating livestock in Karamoja, Uganda. *Econ Bot*. 2009;63(1):29–42.
108. Ghorbani A, Langenberger G, Feng L, Sauerborn J. Ethnobotanical study of medicinal plants utilized by Hani ethnicity in Naban River Watershed National Nature Reserve, Yunnan, China. *J Ethnopharmacol*. 2011;134(3):651–67.
109. Kaval I, Behçet L, Caklioglu U. Ethnobotanical study on medicinal plants in Geçitli and its surrounding (Hakkari-Turkey). *J Ethnopharmacol*. 2014;155(1):171–84.
110. Mahomoodally MF, Muthoorah LD. An ethnopharmacological survey of natural remedies used by the chinese community in Mauritius. *Asian Pac J Trop Biomed*. 2014;4 Suppl 1:S387–99.
111. Abbet C, Mayor R, Roguet D, Spichiger R, Hamburger M, Potterat O. Ethnobotanical survey on wild alpine food plants in Lower and Central Valais (Switzerland). *J Ethnopharmacol*. 2014;151(1):624–34.
112. Dembélé U, Lykke AM, Koné Y, Témé B, Kouyaté AM. Use-value and importance of socio-cultural knowledge on *Carapa procera* trees in the Sudanian zone in Mali. *J Ethnobiol Ethnomed*. 2015;11:14.
113. Kal E, Asfaw Z, Kelbessa E, Van Damme P. Ethnomedicinal study of plants used for human ailments in Ankober District, north Shewa zone, Amhara Region, Ethiopia. *J Ethnobiol Ethnomed*. 2013;9:63.
114. Song M-J, Kim H, Heldenbrand B, Jeon J, Lee S. Ethnopharmacological survey of medicinal plants in Jeju Island, Korea. *J Ethnobiol Ethnomed*. 2013;9:48.
115. Quave CL, Pieroni A. A reservoir of ethnobotanical knowledge informs resilient food security and health strategies in the Balkans. *Nat Plants*. 2015;1:1–6.
116. Mall B, Gauchan DP, Chhetri RB. An ethnobotanical study of medicinal plants used by ethnic people in Parbat district of western Nepal. *J Ethnopharmacol*. 2015;165:103–17.
117. Huamantupa I, Cuba M, Urrunaga R, Paz E, Ananya N, Callalli M, Pallqui N, Coasaca H. Riqueza, uso y origen de plantas medicinales expendidas en los mercados de la ciudad del Cusco. *Rev Peru Biol*. 2011;18(3):283–91.
118. Bye R. The role of humans in the diversification of plants in Mexico. In: Ramamoorthy TP, Bye RA, Lot A, Fa JE, editors. *Biological Diversity of Mexico: Origins and Distribution*. México: Jardín Botánico del Instituto de Biología, Universidad Nacional Autónoma de México; 1993. p. 707–31.
119. Stepp JR. The role of weeds as sources of pharmaceuticals. *J Ethnopharmacol*. 2004;92(2–3):163–6.
120. de Albuquerque UP, de Medeiros PM, de Almeida ALS, Monteiro JM, de Freitas Lins Neto EM, de Melo JG, dos Santos JP ALS. Medicinal plants of the caatinga (semi-arid) vegetation of NE Brazil: A quantitative approach. *J Ethnopharmacol*. 2007;114(3):325–54.
121. Aguirre Z, Madsen JE, Cotton E, Balslev H. Botánica austroecuatoriana: estudios sobre los recursos vegetales en las provincias de El Oro, Loja y Zamora-Chinchipe. Quito: Ediciones Abya Yala; 2002.
122. Aguirre Z, Yaguana C, Merino B. *Plantas medicinales de la zona andina de la provincia de Loja, Ecuador*. Loja: Herbario y Jardín Botánico "Reinaldo Espinosa"; 2014
123. León-Yáñez S, Valencia R, Pitman N, Endara C, Navarrete H. *Libro rojo de las plantas endémicas del Ecuador*. 2th ed. Quito: Publicaciones del Herbario QCA, Pontificia Universidad Católica del Ecuador; 2011.
124. Pardo-de-Santayana M. *Estudios etnobotánicos en Campoo (Cantabria): conocimiento y uso tradicional de plantas*, vol. 33. Madrid: Editorial CSIC-CSIC Press; 2008.
125. Monigatti M, Bussmann RW, Weckerle CS. Medicinal plant use in two Andean communities located at different altitudes in the Bolívar Province, Peru. *J Ethnopharmacol*. 2013;145(2):450–64.
126. Malla B, Chhetri RB. Indigenous knowledge on medicinal non-timber forest products (NTFP) in Parbat district of Nepal. *Indo Glob J Pharm Sci*. 2012;2(2): 213–25.
127. Jerves-Andrade L, Cuzco N, Tobar V, Ansaloni R, Maes L, Wilches I. Medicinal plants used in south Ecuador for gastrointestinal problems: An evaluation of their antibacterial potential. *J Med Plants Res*. 2014;8:1310–20.
128. GLOBOCAN: Estimated Cancer Incidence, Mortality, and Prevalence Worldwide in 2012. IARC; 2014
129. Tariq A, Mussarat S, Adnan M, Abd-Allah EF, Hashem A, Alqarawi AA, Ullah R. Ethnomedicinal evaluation of medicinal plants used against gastrointestinal complaints. *Biomed Res Int*. 2015;2015:1–14.
130. Uddin MZ, Hassan MA. Determination of informant consensus factor of ethnomedicinal plants used in Kalenga forest, Bangladesh. *Bangladesh J Plant Taxon*. 2014;21:83–91.