

Oxidative Stress in Neurodegenerative Diseases

Ewa Niedzielska¹ · Irena Smaga¹ · Maciej Gawlik¹ · Andrzej Moniczewski¹ ·
Piotr Stankowicz¹ · Joanna Pera² · Małgorzata Filip^{1,3} 

Received: 5 March 2015 / Accepted: 1 July 2015 / Published online: 22 July 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract The pathophysiologies of neurodegenerative diseases, including amyotrophic lateral sclerosis (ALS), Parkinson's disease (PD), and Alzheimer's disease (AD), are far from being fully explained. Oxidative stress (OS) has been proposed as one factor that plays a potential role in the pathogenesis of neurodegenerative disorders. Clinical and preclinical studies indicate that neurodegenerative diseases are characterized by higher levels of OS biomarkers and by lower levels of antioxidant defense biomarkers in the brain and peripheral tissues. In this article, we review the current knowledge regarding the involvement of OS in neurodegenerative diseases, based on clinical trials and animal studies. In addition, we analyze the effects of the drug-induced modulation of oxidative balance, and we explore pharmacotherapeutic strategies for OS reduction.

Keywords Alzheimer's disease · Amyotrophic lateral sclerosis · Antioxidant defense molecule · Non-enzymatic and enzymatic antioxidant · Oxidative stress biomarker · Parkinson's disease · ESR—electron spin resonance · IST—immuno-spin trapping technique

Abbreviations

3-NT	3-Nitrotyrosine
4-HDA	4-Hydroxyalkenal
5-OHC	5-Hydroxycytosine
5-OHU	5-Hydroxyuracil
6-OHDA	6-Hydroxydopamine
⁶² Cu-ATSM	Copper-diacetyl-bis(N4-methylthiosemicarbazone)
8-OHA	2,8-Hydroxyadenine
8-OHdG	8-Hydroxy-2'-deoxyguanosine
8-OHG	8-Hydroxyguanine
AD	Alzheimer's disease
AGE	Advanced glycation end product
ALE	Advanced lipid peroxidation end product
ALS	Amyotrophic lateral sclerosis
AOPP	Advanced oxidation protein products
APOE	Apolipoprotein E
APP	Amyloid precursor protein
ATP	Adenosine triphosphate
Aβ	Beta amyloid
CAT	Catalase
COMT	Catechol-O-methyltransferase
COX	Cyclooxygenase
CSF	Cerebrospinal fluid
CTL	Creatol (5-hydroxycreatinine)
CTX	Cortex
CYP 450	Cytochrome P450
DHE	Docosahexaenoic acid
eNOS	Endothelial nitric oxide synthetase
EPA	Eicosapentaenoic acid
ESR	Electron spin resonance
ETC	Electron transport chain
F2-isoPs	F2-isoprostanes
F4-NP	F4-neuroprostane
FALS	Familial amyotrophic lateral sclerosis

✉ Małgorzata Filip
mal.fil@if-pan.krakow.pl

¹ Department of Toxicology, Chair of Toxicology, Faculty of Pharmacy, Jagiellonian University, Medical College, Medyczna 9, 30-688 Kraków, Poland

² Department of Neurology, Faculty of Medicine, Jagiellonian University, Medical College, Botaniczna 3, 31-503 Krakow, Poland

³ Laboratory of Drug Addiction Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Smętna 12, 31-343 Kraków, Poland

FDA	Food and Drug Administration	Prx	Peroxiredoxin
GPx	Glutathione peroxidase	Prx2	Peroxiredoxin-2
GPx1	Glutathione peroxidase-1	PSEN 1	Gene encoding presenilin 1
GR	Glutathione reductase	RIA	Radioimmunoassay
GSH	Glutathione	RNS	Reactive nitrogen species
GSH/	Reduced/oxidized glutathione ratio	ROS	Reactive oxygen species
GSSG		s.c.	Subcutaneous
GSSG	Oxidized glutathione	SAG	Superoxide anion generation
GSSG/	Oxidized/reduced glutathione ratio	SALS	Sporadic amyotrophic lateral sclerosis
GSH		SN	Substantia nigra
GST	Glutathione <i>S</i> -transferase	SNpc	Substantia nigra pars compacta
HD	Huntington disease	SOD	Superoxide dismutase
HETE	Hydroxyeicosatetraenoic acid	SOD1	Copper/zinc superoxide dismutase
HIF-1a	Hypoxia-inducible factor-1a	STR	Striatum
HIP	Hippocampus	TAC	Total antioxidant capacity
HNE	4-Hydroxynonenal	TBARS	Thiobarbituric acid reactive substances
HODE	Hydroxyoctadecadienoic acid	TQ/TQH2	Tocopherylquinone/tocopheryl hydroquinone
iNOS	Inducible nitric oxide synthetase	Trx	Thioredoxin
i.c.v.	Intracerebroventricular	TSE	Aqueous extract of tomato seeds
i.p.	Intraperitoneal	TT	Total thiol
i.v.	Intravenous	Tyr	Tyrosine
Intra-SNc	Intra-substantia nigra injection	UQ/UQH2	Ubiquinone/ubiquinol
inj.		XO	Xanthine oxidase
Intra-STR	Intrastriatal injection	ω -3 FA	Omega-3 fatty acid
inj.			
IsoF	Isofuran		
IST	Immuno-spin trapping technique		
LF	Lipofuscin		
LOX	Lipoxygenase		
LPO	Lipid hydroperoxide		
Lyso PC	Lysophosphatidylcholine		
MAO	Monoaminoxidase		
MAO-B	Monoamine oxidase B		
MDA	Malondialdehyde		
MPO	Myeloperoxidase		
MPTP	1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine		
NADH	Reduced form of nicotinamide adenine dinucleotide		
MRI	Magnetic resonance imaging		
NADPH	Reduced form of nicotinamide adenine dinucleotide phosphate		
NF	Neurofuran		
nNOS	Inducible nitric oxide synthetase		
NOX	NADPH oxidase		
ONOO ⁻	Peroxynitrite		
NS	Nitrosative stress		
Opht A	Ophthalmic acid		
ox-LDL	Oxidized LDL		
OS	Oxidative stress		
p.o.	Per os		
PC	Protein carbonyl		
PD	Parkinson's disease		
PET	Positron emission tomography		

Introduction

Identifying factors that contribute to neurodegenerative processes in the brain is one of the major goals of modern medicine. Currently, there are several hypotheses concerning the mechanisms that lead to the damage and death of brain cells in neurodegenerative diseases, such as excitotoxic effects by excitatory amino acids, disturbed cellular energy metabolism, and oxidative stress (OS), which is caused by free radicals or other reactive molecules.

The excessive production of reactive species and insufficient activity of antioxidant defense mechanisms have been implicated in the pathogenesis of many neurodegenerative diseases, including amyotrophic lateral sclerosis (ALS), Parkinson's disease (PD), Alzheimer's disease (AD), and Huntington disease (HD) [1].

Reactive Species, Sources, and Cellular Pro-/Antioxidant Balance

It is thought that the biological oxidants that cause oxidative damage comprise the products of endogenous and exogenous processes that involve oxygen and nitrogen. Reactive species that contain oxygen are produced during aerobic respiration, cellular metabolism, and defense against pathogens [2]. The chemical potential of the oxygen molecule relies on its

electron structure (two unpaired electrons in its basic triplet state). It promotes one-electron reactions that form the basis for respiration (reduction of oxygen molecules in four single-electron reactions), microsomal electron transport chains (ETC) (via cytochrome P-450 (CYP 450)), and oxidative burst activity in macrophages [3].

The high dynamics of the chemical processes that are achieved in elementary single-electron reactions are desirable and are the source of reactive molecules, which are either undesirable side products (respiration and metabolism) or in excess of the established requirements (defense process). These reactive molecules are known as reactive oxygen species (ROS) and reactive nitrogen species (RNS). Among them, the best known are singlet oxygen ($^1\text{O}_2$), superoxide anion radicals ($\text{O}_2^{\cdot-}$), hydroxyl radicals (HO^\bullet), hydrogen peroxide (H_2O_2), nitric oxide (NO), and peroxy nitrite anions (ONOO^-) [4, 5].

At physiological concentrations, ROS/RNS play important regulatory and mediator functions, but an uncontrolled increase in ROS/RNS concentrations leads to a chain of radical reactions that increases the risk of damage to biological molecules in a living organism. This is caused by the high reactivity of ROS and RNS with lipids, proteins, carbohydrates, and nucleic acids. Hence, establishing an antioxidant barrier is required to limit the amount of ROS/RNS to a level that is not threatening to the integrity of biological systems. Excessive formation of ROS/RNS that exceeds the maximum capacity of the antioxidant barrier leads to a disturbance in the pro-/antioxidant equilibrium and, finally, to the development of the state known as OS.

OS can be triggered by radicals produced by either exogenous processes (e.g., xenobiotics, cold, viral and bacterial infections, ionizing radiation, ultrasound or photo-oxidation, poor diet, alcohol consumption, and smoking) or endogenous processes, which are the basic biochemical reactions in the body mentioned earlier (Fig. 1).

ROS production as a side effect of aerobic respiration occurs on the inner membrane of the mitochondrion [6] (Fig. 2). The respiratory chain (mitochondrial ETC) consists of a series of membrane-bound complexes, such as complex I (reduced form of nicotinamide adenine dinucleotide (NADH)/ubiquinone reductase), complex II (succinate ubiquinone reductase), complex III (ubiquinol cytochrome c reductase), complex IV (cytochrome c oxidase), and complex V (adenosine triphosphate (ATP) synthase) [7]. The final acceptor of electrons and protons, an oxygen molecule, undergoes four-electron reduction, which can lead to the production of water molecules. During ETC, single electrons leak to reduce molecular oxygen and to form $\text{O}_2^{\cdot-}$ and, later, H_2O_2 and HO^\bullet [8] (Fig. 1).

Because of its high consumption of oxygen and its high lipid content, the brain is particularly vulnerable to damage caused by ROS and RNS. The extent of the damage varies, depending on, among other factors, the source and type of the

reactive species. More active molecules, such as HO^\bullet and ONOO^- , interfere with other molecules at the site of their production, while less active ones, such as $^1\text{O}_2$ and $\text{O}_2^{\cdot-}$, can diffuse over longer distances and produce effects in more specific locations [9]. They can also react with each other and form more active forms, as in the case of the reaction of $\text{O}_2^{\cdot-}$ with NO, which creates ONOO^- [10].


In mitochondria, NO is produced from L-arginine and L-citrulline in a reaction that is catalyzed by nitric oxide synthase (NOS), which has three isoforms with different tissue localizations. Cellular Ca^{2+} -dependent neuronal NOS (nNOS) is expressed in astrocytes, microglia, and macrophages, and endothelial NOS (eNOS) is expressed in the vascular endothelium and Ca^{2+} -independent inducible NOS (iNOS). NO is involved in many important processes within the central nervous system, such as the regulation of cerebral blood flow and memory. In addition, it plays a significant role in the regulation of the immune system, including the modulation of cytokine production. The released NO acts on neighboring cells, leading to somatic mutations and affecting cell cycle regulatory proteins, apoptosis, and DNA repair [11].

RNS are important for the generation of OS. ONOO^- is rapidly decomposed into HO^\bullet , nitrogen dioxide radical (NO_2^\bullet), and nitryl cation (NO_2^+). All of these can damage nerve cells [12].

These highly reactive compounds induce changes in the structure and function of cell membranes, proteins, lipoproteins, enzymes, hormones, and genetic material. In particular, membranes are a primary target for ROS. Conversion products of lipid peroxidation lead to the decomposition of polyunsaturated fatty acids and the formation of the final products, i.e., the reactive aldehydes, such as malondialdehyde (MDA) and 4-hydroxynonenal (HNE). These compounds react with DNA or protein molecules and modify their structure and functions [13, 14].

There are several mechanisms designed to protect the organism from the harmful effects of ROS and RNS. The ultimate amount of ROS/RNS is under strict control in the body as a result of enzymatic and non-enzymatic defense mechanisms. The production of ROS- and RNS-induced damage (the final effect of OS) in tissue can be confirmed by the presence of tissue-specific and non-specific biomarkers [15–20]. Several markers of OS and antioxidant activity are presented in Fig. 2. Recent technical advances used to detect and identify ROS/RNS biomarkers and free radical metabolism are electron spin resonance (ESR), the immuno-spin trapping technique (IST), and radioimmunoassay (RIA) [21, 22].

The cellular antioxidant system, designed to prevent damage to tissue, is composed of antioxidant enzymes and other non-enzymatic compounds that have the ability to reduce different chemical structures [21]. These compounds are responsible for maintaining the balance between pro- and antioxidant

Fig. 1 Generation of reactive species (based on [280–283])

agents and alleviating OS (see Table 1). The essential components of the enzymatic antioxidant defense are superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and glutathione reductase (GR), while the non-enzymatic antioxidants include glutathione (GSH), thioredoxin (Trx), vitamins A, E, and C, flavonoids, trace elements, and proteins, e.g., albumin, ceruloplasmin, and metallothionein.

A large body of evidence confirms a relationship between OS and the development of neurodegenerative diseases. The increased neuronal ROS production and accumulation of oxidative damage that occurs with age correlate well with the extent of neurodegeneration. In the following sections of this article, we present the current knowledge on the relationships between the intensity of OS and the initiation and progression of the major neurodegenerative diseases AD, PD, and ALS. The assessment of these relationships is based on biomarkers of OS and indicators of the pro-/antioxidant balance in nervous tissue that are correlated with the typical characteristics of neurodegenerative processes.

Amyotrophic lateral sclerosis (ICD-10 G12.21)

ALS (also known also as Charcot's or Lou Gehrig's disease) is a severe neurodegenerative disease that is characterized by progressive upper motor neuron loss in the cerebral cortex and lower motor neuron loss in the brainstem and spinal cord [23, 24]. This results in spasticity, muscle wasting, and weakness, leading finally to paralysis and difficulties with speech, swallowing, and breathing. ALS may have either a limb onset (80 % cases) or a bulbar onset (20 % cases) [24, 25]. There is

currently no cure for ALS and only riluzole, which acts on glutamate signaling, has been registered for the treatment of the disease. Riluzole was shown to slow disease progression and to improve limb function; however, the survival of patients was prolonged by only 2–3 months and death due to respiratory failure occurred in most cases within 3–5 years of the diagnosis [26].

The main pathological hallmark of ALS is the formation of cytoplasmic aggregates in degenerating motor neurons and surrounding oligodendrocytes, but those inclusions are also present in the frontal and temporal cortices, hippocampus, and cerebellum [27].

Only 5–10 % of ALS cases are inherited (familial ALS (FALS)); the remaining cases have no clear genetic background (sporadic ALS (SALS)) [24], and their pathogenesis is still poorly understood. Studies concerning the mechanisms of ALS development indicate that many factors, including excitotoxicity, mitochondrial dysfunction, endoplasmic reticulum stress, neuroinflammation, and OS, can be involved in this process. The two forms of the disease are clinically indistinguishable because the symptoms and pathological changes in SALS and FALS are similar. FALS is caused by mutations in some genes, such as those coding for SOD1, FUS RNA binding protein, TAR DNA binding protein, vesicle-associated membrane protein B, valosin-containing protein, optineurin, alsin, senataxin, spatascin, angiogenin, or ubiquilin-2 [24, 26]. Some of these gene mutations have also been found in SALS patients [28, 29].

The most common known genetic mutation for ALS is the recently described expanded GGGGCC hexanucleotide repeat in the non-coding region of the *C9orf72* gene, on




Fig. 2 Source of radical and OS biomarkers. *8-OHA* 2,8-hydroxyadenine, *HNE* 4-hydroxynonenal, *5-OHC* 5-hydroxycytosine, *5-OHU* 5-hydroxyuracil, *8-OHdG* 8-hydroxy-2'-deoxyguanosine, *8-OH-Gua* 8-hydroxyguanine, *AGEs* advanced glycation end products, *ALES* advanced lipid peroxidation end products, *AOPP* advanced oxidation products, *CTL* creatol, *COX* cyclooxygenases, *CYP* cytochrome P450, *ETC* electron transport chain, *F2-isoPs* F2-isoprostanes, *F4-NPs* F4-neuroprostanes, *HETE* hydroxyeicosatetraenoic acids, *HODE* hydroxyoctadecadienoic acid,

HIF-1a hypoxia-inducible factor-1a, *IsoF* isofuran, *LF* lipofuscin, *LOX* lipoxygenases, *Lyso PC* lysophosphatidylcholines, *MPO* myeloperoxidase, *NOX* NADPH oxidases, *NFS* neurofuran, *Opht A* ophthalmic acid, *GSSG* oxidized glutathione, *ox-LDL* oxidized LDL, *GSSG/GSH* oxidized/reduced glutathione, *Prxs* peroxiredoxins, *PC* protein carbonyl, *TBARS* thiobarbituric acid reactive substances, *TQ/TQH2* tocopherylquinone/tocopheryl hydroquinone, *TAC* total antioxidant capacity, *Tyr* tyrosines, *UQ/UQH2* ubiquinone/ubiquinol, *XO* xanthine oxidase

chromosome 9p21 [30]. Another common mutation is localized in *SOD1*. *SOD1* in its native state metabolizes $O_2^{\cdot-}$ to molecular misfolding [31]. Pathogenic mutations of *SOD1* enzyme can be present in different regions of the enzyme, e.g., G93A (glycine 93 changed to alanine), H46R (histidine at codon 46 changed to arginine), or A4V (alanine at codon 4 changed to valine). Mutated *SOD1* can form cytotoxic protein aggregates alone or with other proteins what possibly leads to loss of the enzymatic function or to acquiring the toxic properties [32, 33]. As *SOD1* plays a crucial role in $O_2^{\cdot-}$ clearance, its functional loss can lead to increased levels of OS. Then, the remaining wild-type *SOD1* may become itself a target of oxidative modification after which it dissociates from dimers to monomers and further forms aggregates with toxic properties of mutant forms of *SOD1*. That was shown in in vitro studies

[34, 35]. Accumulation of abnormal *SOD1* was also confirmed in the spinal cord [34] in animal studies [36, 37] as well as in ALS patients [38].

Clinical Studies

OS Biomarkers

Post-mortem studies on tissue samples from SALS and FALS patients support the hypothesis of oxidative damage of proteins, lipids, and DNA. For instance, HNE and 3-nitrotyrosine (3-NT) levels were increased in CSF in SALS patients [39, 40], while a rise in 8-hydroxy-2'-deoxyguanosine (8-OHdG) has been described in both

Table 1 Enzymatic and non-enzymatic antioxidants against OS

Agents	Mechanism
Enzymatic	
Superoxide dismutases (SOD) CuZn-SOD, located in the cytoplasm Mn-SOD, located in the mitochondria EC-SOD extracellular	Scavenges superoxide anion to form H ₂ O ₂
Catalases (CATs)	Peroxisomes remove the hydrogen peroxide
Peroxiredoxins (Prxs)	Reduce free hydrogen peroxide to water
Glutathione peroxidases (GPx)	Catalyzes the reduction of H ₂ O ₂ and alkyl hydroperoxides to H ₂ O
Glutathione reductases (GRs)	Catalyzes the reduction of oxidized glutathione (GSSG) to reduced glutathione (GSH)
Glutathione S-transferases (GSTs)	Catalyze the conjugation of xenobiotics and endogenous or exogenous electrophilic compounds to reduced glutathione
Thioredoxin reductases (TrxRs)	Repair oxidative damages of proteins
Heme oxygenase-1 (HO-1)	Creation of products with antioxidant properties
Metalloproteins	
Metallothionein (MT)	Physiological metal chelation (Zn, Cu, Se) and xenobiotic metals (Cd, Hg, Ag, As)
Albumin	Chelation of metal ions (Fe, Cu)
Ceruloplasmin (CP)	Chelation of metal ions (Cu)
Ferritin	Chelation of metal ions (Fe)
Transferrin	Chelation of metal ions (Fe)
Non-enzymatic	
Vitamin C (ascorbic acid)	ROS scavenger
Vitamin E (α -tocopherol)	ROS scavenger
Vitamin A (retinol)	ROS scavenger
Carotenoids	ROS scavenger
Melatonin	ROS scavenger
Reduced nicotinamide adenine dinucleotide phosphate (NADPH)	Coenzyme used in anabolic reactions
Adenosine (ADO)	Endogenous activator of the cellular antioxidant defense system
Uric acid	ROS scavenger
Ubiquinol (the reduced form of coenzyme Q-10)	ROS scavenger
Polyphenols	ROS scavenger, metal chelation, induction of antioxidant enzymes
Phenolic acids	
Flavonoids	
Stilbenes	
GSH reduced glutathione	ROS scavenger
Amino acids (cysteine, homocysteine, methionine, taurine)	ROS scavenger
α -Lipoic acid	ROS scavenger
Lycopene	ROS scavenger
Carotenoids	ROS scavenger
S-adenosyl-L-methionine	ROS scavenger
Selenium	ROS scavenger

Data from [274–279]

SALS and FALS patients [41, 42]. However, no correlation between these markers and the severity or duration of the disease has been found [43].

Raised levels of thiobarbituric acid reactive substances (TBARS) and advanced oxidation protein products (AOPP) and decreased ferric-reducing ability of plasma (FRAP) were detected in the plasma or erythrocytes of SALS patients

[44–46], but plasma protein carbonyl (PC) levels surprisingly did not differ between SALS patients and control subjects [16]. More conclusive results came from a study where urine from SALS patients contained a higher level of isoprostanoids (IsoPs) and 8-OHdG compared to a control group [16, 47], suggesting that IsoPs and 8-OHdG could be considered markers of OS in ALS. These studies included only 50

participants with SALS compared to 46 control subjects. Thus, these results should be confirmed in larger cohorts.

The presence of OS biomarkers in regions of the CNS that are critical for ALS suggests that they are implicated in motor neuron degeneration. This fact is supported by very recent positron emission tomography (PET) imaging data in humans, which confirmed that OS were enhanced in the motor cortex in ALS patients compared with controls. Moreover, the observed OS increase in the mild stage of the disease led to the conclusion that OS may be an important factor associated with the development of neurodegeneration in ALS patients [48].

Antioxidant Defense Biomarkers

Most studies concerning antioxidant defense biomarkers in ALS patients have shown changes in peripheral tissues or in CSF but rarely in the brain. For example, GSH levels were reduced in erythrocytes in ALS patients [45]. In contrast, an earlier study demonstrated a lower GSSG/GSH ratio with a concomitant rise in NO in the CSF of SALS patients [49]. According to the latter study, the lower GSSG/GSH level was caused by the decreased production of O_2^- -oxidizing GSH because of the shift of oxidation to NO production or oxidation [49]. Very recent in vivo imaging studies have shown decreased GSH levels in the motor cortex of ALS patients by using the J-editing technique or 62 copper-diacyl-bis(N4-methylthiosemicarbazone (^{62}Cu -ATSM)) PET technique [25, 48].

The activity of SOD decreased in red blood cells and the CSF of FALS- and SALS-diagnosed patients [42, 50–52]. Interestingly, the reduction in the SOD1 protein concentration in erythrocytes of FALS patients correlated with *SOD1* gene mutations [42]. However, SALS patients displayed enhanced activity of SOD1 in CSF [53], indicating that this different outcome may depend on either the ALS type, the disease duration, or the sampling time [54].

Apart from SOD, the activity of CAT, another antioxidant defense enzyme, was also found to be diminished in red blood cells in FALS [51] and SALS [51, 52] patients. Another study revealed that CAT activity decreased in erythrocytes with disease progression, which may suggest a link between this parameter and ALS duration [45].

Results regarding GPx or GR activity in ALS patients are controversial. GPx type 3 (also known as plasma GPx) and GR levels were lower in the serum or in red blood cells, respectively, in ALS patients [45, 55]. In contrast, another study showed enhanced GPx in the serum and CSF of ALS patients [56], while GR activity was elevated in the erythrocytes in both SALS and FALS patients [51, 52]. One study reported no change in GR activity in red blood cells in SALS patients [57].

In ALS patients, the plasma and/or CSF levels of other antioxidants (i.e., vitamin E, β -carotene, vitamin C, uric acid,

and thiol group-containing molecules, such as ubiquinol-10) were either elevated [53] or not changed [44, 58, 59].

These divergent results could be explained by the heterogeneity of ALS [60]. In fact, ALS-diagnosed patients vary in their rate of disease progression and may differ in the number of years of survival. The median survival from onset to death in ALS varies from 20 to 48 months; however, many studies have reported a survival time of longer than 10 years in 10–20 % of patients [60]. Moreover, a study from Cova et al. [57] showed that the activity of GPx, GR, and CuZn SOD in erythrocytes was decreased in patients who had a faster disease progression rate and that SOD1 activity significantly decreased along the disease course in subjects with a more impaired functional status. All of these results support the hypothesis that ALS has many variants or mimic syndromes that differ in their pathogenic mechanisms and in their profile of enzymatic and non-enzymatic antioxidant responses [57].

Pharmacological Strategies to Reduce OS

Several pharmacotherapeutic agents with antioxidant properties have been attempted to slow ALS progression; however, most of them failed to do so (Table 2). Vitamin E, when combined with riluzole, diminished TBARS levels, enhanced GPx activity in the plasma, and prolonged the milder stages of the ALS Health State scale, but it did not affect survival and motor function [61].

Another study investigated a very high dose of vitamin E as an add-on therapy to riluzole for ALS treatment; however, this treatment with vitamin E, at a dose of 5,000 mg/day for 18 months, failed to slow disease progression [62]. On the other hand, another study showed that intake of a combination of polyunsaturated fatty acids and vitamin E reduced the risk of developing ALS [63].

Edavarone (MCI-186) is another free radical scavenger that is already approved to treat cerebral infarction and to investigate ALS. It eliminates lipid peroxide and hydroxyl radicals by transferring an electron to the radical and thereby exerting a protective effect on neurons. In ALS patients, edavarone was shown to diminish nitrosative stress (NS) in CSF [64] and is now being tested in a phase III clinical trial in Japan for ALS treatment [65]. Coenzyme Q10 was administered for 9 months in a dose of 2,700 mg/day to ALS patients in a multicenter trial that unfortunately showed a lack of compound efficacy for ALS treatment [66]. This study also determined 8-OHdG levels in the plasma; however, according to the authors, the results for this measure will only be available later [66].

Other antioxidant medications, i.e., acetylcysteine, creatine, or selegiline, had no significant effect on survival or the rate of clinical progression of ALS [67–69]. In these studies, the influence on antioxidant defense biomarkers and oxidant

Table 2 Clinical trials in ALS patients with agents/drugs showing antioxidant properties

Antioxidant	Time point of antioxidant status determination from the beginning of therapy	Oxidative damage biomarker change	Reference
Vitamin E (500 mg twice a day) with riluzole (100 mg/day)	3 months	↑GPx in plasma, ↓TBARS in plasma	[61]
Selegiline hydrochloride (5 mg twice a day p.o.)	6 months	nd	[69]
Acetylcysteine (50 mg/kg/day s.c.)	12 months	nd	[68]
Creatine (5 g twice a day p.o.)	16 months	nd	[67]
Coenzyme Q10 (1,800 and 2,700 mg/day p.o.)	9 months	nd	[66]
Edaravone (60 mg/day i.v.)	2 weeks administration every 2 weeks, 6 replications	↓3-NT in CSF	[64]
Melatonin (300 mg/day) per rectum	4 months	↓PC in serum	[70]

↓ decrease, ↑ increase, *nd* not determined, *3-NT* 3-nitrotyrosine, *8-OHG* 8-hydroxyguanosine, *CSF* cerebrospinal fluid, *GPx* glutathione peroxidase, *i.v.* intravenously, *p.o.* per os, *PC* protein carbonyl, *s.c.* subcutaneous, *TBARS* thiobarbituric acid reactive substances

damage was not evaluated. In another study, melatonin was used as a potential neuroprotective compound and, when normalized to control values, was found to elevate the level of PC in the serum of 31 SALS patients [70].

The novel antioxidant AEOL 10150, which is a small molecule that catalytically consumes ROS and RNS, is considered to be the most promising compound under evaluation in a clinical trial [71]. In a small, open-label study in ALS patients, AEOL 10150 was shown to be a safe and well-tolerated drug [71]. However, neither efficacy nor measurement of oxidative biomarkers in ALS patients who were on AEOL 10150 has been published.

The antioxidant medications used in ALS clinical trials have so far been unable to slow the progression of the disease. Moreover, a limited number of clinical studies have investigated oxidative damage or changes in the antioxidant defense status after antioxidant therapy.

Anti-ALS Drugs that Modulate the Oxidative Balance

No data.

Animal Studies

OS Biomarkers

The most popular animal models of ALS are based on SOD1 mutant transgenic mice (for example: SOD1 G93A, SOD1 H46R, SOD1 H46R/H48Q, SOD1 A4V, etc. (for more details, see [72])). Other ALS models include Wobbler mice and rodent models with genetic mutations in TAR DNA binding protein of 43 kDa (TDP-43), VAMP-associated protein B, dynactin or FUS/TLS, or C9orf72 [73].

Changes in oxidative biomarkers have been detected in ALS models in rodents (Table 3). For example, free radical levels were elevated in the spinal cord of SOD1 G93A mice

post-mortem [74, 75], and trapped radical adducts were also recently detected in the spinal cord of these mice *in vivo* [76]. In line with this latter observation, increased levels of PC in the spinal cord and in the motor cortex [77, 78], MDA and 4-HDA in the brain and CSF [79], and HNE, HNE-adducts, and 8-OHG in the spinal cord [80, 81] have been observed in several ALS animal models (see Table 3).

It has been established that enhanced nitration processes are also present in animal models of ALS. For example, 3-NT was found in the spinal cords of aged SOD1 G93A mice together with the presence of overnitrated proteins (actin or ATPase) in the spinal cord [82], in the motor and sensory cortices [78], and in the peripheral blood mononuclear cells [83] in transgenic mouse models of ALS and that these were observed even before the onset of the disease.

OS and NS are widely present in ALS, and *in vitro* studies suggest that this mechanism can facilitate the formation of protein aggregates [84]. In support of this conclusion, recent studies have revealed that a selective nNOS inhibitor increased survival in SOD1 transgenic mice [85].

Antioxidant Defense Biomarkers

In general, transgenic mouse models overexpressing mutant human *SOD1* showed increased activity by the enzyme [32]. However, in the end stage of the disease, SOD1 activity in SOD1 G93A mice remained at the same level as wild-type SOD in non-transgenic mice [86]. What is interesting is that SOD1 knockout mice did not show an ALS-like phenotype [32].

Other antioxidant biomarkers, including GSH, GPx, and GR, are also changed during the course of ALS. The levels of GSH differed in mice carrying different SOD1 mutations. For example, in SOD1 G93A mutant mice, the GSH level decreased and the level of GSSG rose in the lumbar spinal cord [87], while in the same tissue in SOD1 mice with the

Table 3 OS biomarkers in ALS animal model

Animal model	OS biomarkers	Reference
Mice SOD1 G93A-2Gur	↑PC in spinal cord	[77]
Mice SOD1 G93A	↑MDA, ↑4-HDA in brain and CSF ∅ MDA, ∅ 4-HDA in skeletal muscles and heart ∅ PC in brain, CSF, skeletal muscles, liver, heart	[79]
Mice SOD1 G93A	↑MDA, ↑HNE in spinal cord	[81]
Mice SOD1 G93A	↑In vivo trapped radical adducts	[76]
Mice SOD1 G93A	↑8-OHdG in spinal cord	[80]
Mice SOD1 G93A	↑PC and ↑3-NT in motor and sensory CTX	[78]
Rats SOD1 G93A	↑Overnitrated proteins in mononuclear cells from peripheral blood	[83]
Mice SOD1 G93A	↑3-NT in spinal cord in presymptomatic stage	[82]

↓ decrease, ↑ increase, ∅ no changes, *nd* not determined, *3-NT* 3-nitrotyrosine, *4-HDA* 4-hydroxyalkenal, *8-OHdG* 8-hydroxy-2'-deoxyguanosine, *CSF* cerebrospinal fluid, *CTX* cortex, *MDA* malondialdehyde, *PC* protein carbonyl

H46R/H48Q mutations, the GSH level was not changed [88]. A recent study by Vargas et al. [88] showed that decreased GSH content accelerated neurological deficits in the SOD1 G93A mouse model. The mechanism of GSH reduction is linked to the nuclear transcription factor Nrf2 because its transfection into the SOD G93A mouse brain led to the up-regulation of GSH synthesis in astrocytes and reduced the most apparent neurological and biochemical symptoms of the disease [89].

In animal models of ALS, motor neurons have displayed overexpression of Prx2 and glutathione peroxidase-1 (GPx1). The number of neurons containing Prx2 and GPx1 decreased in the terminal stage of ALS [90], suggesting a breakdown of this redox system at the advanced stages of the disease. As discussed by Kato et al. [91], a possible reason for Prx2 and GPx1 breakdown might be related to the co-aggregation of these enzymes with mutant SOD1 and the exacerbation of mutant SOD1-mediated toxicity in neuronal Lewy body-like hyaline inclusions (LBHIs), which was shown in the spinal cords of transgenic rats [91]. Similarly, GPx3 protein levels in the serum of SOD1 H46R rats were increased in the pre-symptomatic stage and decreased gradually with disease progression [55]. However, another study did not reveal significance for the role of GPx in ALS [92], and further investigation is therefore necessary to clarify this problem.

Because enhanced SOD activity in ALS animal models remains enhanced in most of the disease stages and decreases only to the control level of non-transgenic mice in the end stage of the disease, this raises the question of whether these ALS animal models are suitable to study ALS antioxidant defense biomarkers because an ALS key enzyme, SOD, is decreased in ALS patients. These contradictory results, obtained in animals and humans, do not allow researchers to draw conclusions regarding the significance of these biomarkers in animal ALS models.

Pharmacotherapeutic Strategy to Reduce OS

Many substances possessing antioxidant properties have been proposed as ALS treatment strategies (Table 4). For example, coenzyme Q10 was shown recently to be unable to prolong survival when given after ALS onset [93]. As coenzyme Q10 is characterized by rather poor CNS availability (which possibly explains its small pharmacological effects), its reduced form, ubiquinol-10, has better bioavailability and antioxidant properties and was also investigated. However, similar to its parent drug, ubiquinol-10 did not prolong lifespan. Nevertheless, it was noted that poor CNS availability after oral dosing was observed in this study, which possibly explains the lack of pharmacological effects, similar to the case with its parent drug [93].

Creatine was the next drug that extended survival, but not age of onset, in ALS animals [94–96]. It has neuroprotective properties and buffers against ATP depletion in mitochondria. Its dysfunction can lead to ATP decreases, which may contribute to cell death [97]. Because mitochondrial swelling and vacuolization are among the earliest pathological features in ALS mice with the SOD mutations [98], creatine could be useful for protecting mitochondria and for disease treatment in such a preclinical model. In most studies, creatine was delivered in the diet as a 1 or 2 % food supplementation. In addition to its effect of prolonging the lifespan in a mouse ALS model, creatine also improved motor performance [96] and provided protection from motor neuron loss at 120 days of age in these mice [98]. Moreover, creatine was shown to prevent the rise of 3-NT in the spinal cord and ROS production in the dialysate from microdialysis in ALS animals [98], and it also had a positive effect on weight retention [99]. However, one study found that creatine administration had no effect on the disturbed muscle function [100]. Moreover, co-administration of creatine with the cyclooxygenase-2 inhibitors rofecoxib and celecoxib or of creatine with minocycline

Table 4 Therapeutic trials with agents/drugs with antioxidant properties in ALS animal models

Drug, dose, route of administration	Animal model	Therapy start after	Changes in oxidative defense and damage biomarkers	Onset change	Survival change	Reference
Ubiquinol-10 800 mg/kg/day p.o.	Mice SOD1 G93A	Post-onset	nd	∅	∅	[93]
Coenzyme Q10 800 mg/kg/day p.o.	Mice SOD1 G93A	Post-onset	nd	∅	∅	
Creatine (2 % in diet)	Mice SOD1 G93A	Pre-onset	nd	∅	↑	[95]
Creatine (2 % in diet) + celecoxib (12 % in diet)	Mice SOD1 G93A	Pre-onset	nd	∅	↑	[94]
Creatine (2 % in diet) + rofecoxib (5 % in diet)	Mice SOD1 G93A	Pre-onset	nd	∅	↑	
Creatine (2 % in diet) + minocycline (22 mg/kg/day i.p.)	Mice SOD1 G93A	Pre-onset	nd	∅	↑	[96]
Vitamin E 200 IU/day, 5 days/week for 30 days	Mice SOD1 G93A	Pre-onset	↓8-OHG in spinal cord	↑	∅	[103]
EUK-8 and EUK-134 33 mg/kg i.p. 3 times a week	Mice SOD1 G93A	Pre-onset	↓MDA, ↓PC in spinal cord	∅	↑	[102]
Edaravone 15 mg/kg i.p.	Mice SOD1 G93A	At symptoms onset	↓abnormal SOD1 deposition in spinal cord	na	∅	[36]
AEOL 10150 2.5 mg/kg i.p.	Mice SOD1 G93A	At symptoms onset	↓MDA, ↓3-NT in spinal cord	na	↑	[108]
Ammonium tetrathiomolybdate 5 mg/kg i.p.	Mice SOD1 G93A	Pre-onset	↓LPO, ↓SOD in spinal cord	↑	↑	[86]
		Post-onset	nd	↑	↑	[105]
Resveratrol 25 mg/kg i.p.	Mice SOD1 G93A	Pre-onset	↓MDA in spinal cord	↑	↑	[106]
Melatonin 30 mg/kg, 10 μl/g body weight i.p.	Mice SOD1 G93A	Pre-onset	nd	↑	↑	[104]
Melatonin 5 mg/kg, 2.5 mg/kg or 50 mg/kg i.p.	Mice SOD1 G93A	Pre-onset	↑4-HNE, ↑SOD1 in spinal cord	↓	↓	[107]

↑ increase, ↓ decrease, ∅ no change, na not applicable, nd not determined, 3-NT- 3 nitrotyrosine, i.p. intraperitoneally, LPO lipid hydroperoxide, MDA malondialdehyde, PC protein carbonyls, ROS reactive oxygen species, s.c. subcutaneously, SOD superoxide dismutase

(a tetracycline antibiotic with neuroprotective and anti-inflammatory effects) [101] may be even more beneficial for extending survival (even by 30 %) [94, 96].

Other antioxidants that extended survival in ALS mice include EUK-8 and EUK-134. In preclinical studies, these two synthetic SOD/CAT mimetics reduced the levels of OS, as measured by MDA, PC, and prolonged survival, but not disease onset, in ALS mice [102]. On the other hand, treatment with vitamin E significantly delayed ALS onset with no effect on survival but with a diminishing effect on 8-hydroxyguanine (8-OHG) levels in the spinal cord [103]. Drugs that shared both effects (ALS onset delay and lifespan extension) were melatonin, ammonium tetrathiomolybdate (a copper-chelating drug), and resveratrol (a substance that originates in plants and is found in highest amounts in red wine and the skin of red grapes) [86, 104–106]. However, in case of melatonin, the data were not clear as it was given in a dose range of 2.5–50 mg/kg intraperitoneal (i.p.) (in the same animal model) and also produced a surprisingly shortened survival, accelerated disease onset, enhanced lipid peroxidation in the spinal cord, and increased spinal motoneuron loss [107]. A possible reason for these results might be due, according to authors, to melatonin's effect on upregulating the gene expression of human toxic SOD1, which presumably

overrides any of the anti-oxidant properties of melatonin. Such a hypothesis needs to be tested in future studies.

Other substances used in preclinical ALS models, i.e., resveratrol, which, when given before disease onset, decreased MDA levels [106], and ammonium tetrathiomolybdate, which decreased spinal SOD activity [86]. The latter compound also ameliorated ALS-like symptoms in SOD1 G93A mice, probably by chelating the copper ions from the Cys111 site in the SOD-1 enzyme, which is important because various SOD-1 mutations affect Cu and Zn metal-binding, thereby promoting toxic protein aggregation [86]. Moreover, successful effects were achieved from treatment with ammonium tetrathiomolybdate when it was given after disease onset, which is satisfactory because this regime of administration closely reflects clinical practice.

Edaravone and AEOL 10150 are recently studied drugs for ALS that were administered at ALS onset. Although edaravone did not prolong survival in an ALS mouse model, it diminished SOD1 deposition in the anterior horn of the spinal cord and slowed disease progression and motor neuron degeneration [36]. Another substance that gave more promising results is AEOL 10150, which, when administered to ALS mice, decreased 3-NT and MDA levels in the spinal cord, extended animal survival [108], provided better preservation

of motor neuron architecture, and diminished the level of astrogliosis [108].

In conclusion, according to animal studies, antioxidants may become putative ALS therapies because many of them extend the lifespan and diminish OS in ALS animals. However, it must be remembered that many of them, when tested in humans, have not yet shown benefits for survival time and motor function amelioration.

Anti-ALS Drugs that Modulate Oxidative Balance

Riluzole was shown to extend the lifespan in mice in the SOD1 G93A ALS animal model, but it did not change the disease onset [109] or have a satisfactory effect on the latter parameter [110]. No studies concerning oxidative defense or oxidative damage biomarkers were conducted; however, *in vitro* studies showed that riluzole had antioxidant properties in cultured cortical neurons [111].

Parkinson's Disease (ICD-10 G20)

PD is a progressive degenerative disorder that is characterized by the loss of dopamine-producing neurons in the substantia nigra (SN) and by the presence of Lewy bodies in the SN and locus coeruleus. Clinical manifestations of PD include resting tremor, muscle rigidity, slowing of voluntary movements (bradykinesia), a tendency to fall (postural instability), and a mask-like facial expression [112]. The pathological hallmarks of PD, Lewy bodies, contain various proteins, including α -synuclein, ubiquitin, Parkin, and neurofilaments.

PD occurs in sporadic (non-inherited; 90–95 % cases) and familial (inherited; 5–10 % cases) forms. Mutations in the α -synuclein gene cause one of the familial forms of PD via autosomal dominant inheritance [113]. Recently, it was suggested that OS and imbalances between ROS/RNS and antioxidant defense mechanisms are linked to the etiology of PD.

Clinical Studies

Oxidative Biomarkers

Many studies have demonstrated the presence of OS and its markers in the brain and CSF in PD patients. Cholesterol lipid hydroperoxide and MDA were found to be up to 10-fold higher in SN in *post-mortem* brains of PD patients compared with other brain regions and age-matched controls [114]. In contrast, a recent paper demonstrated lower levels of MDA in the caudate nucleus and putamen and increased MDA levels in the frontal cortex in the *post-mortem* analyses of PD brains compared to healthy age-matched controls. Those results suggested that the non-SN regions, such as the caudate nucleus or the putamen, may have different compensatory mechanisms

against OS could protect them from oxidative damage [115]. Moreover, HNE and acrolein-modified proteins were found in the neocortex and brainstem and in the SN, respectively, of PD patients [116–118]. According to *in vitro* studies, HNE modification of α -synuclein has been shown to trigger oligomerization and fibrillization of unmodified α -synuclein in the nervous system, which might lead to dopaminergic neuron injury. A recent observation suggests that HNE-modified proteins should be considered to be important players in PD pathophysiology [119].

Despite a number of studies supporting lipid peroxidation in the brains of PD patients, the levels of F2-isoprostanes were not elevated in the SN of PD patients [120], and the reason for this observed difference needs to be explained.

PCs were also found in PD patients' brains compared to age-matched controls [121]. PCs are present not only in the brain regions specific for PD (the SN, caudate nucleus, and putamen) but also in other brain areas not directly linked with PD. Because most PD patients are treated with L-DOPA, this observation may suggest that L-DOPA, the "gold standard" for PD therapy, may contribute to carbonyl formation because it was shown to have pro-oxidant properties *in vitro* [122]. Interestingly, brain regions from individuals with putative pre-symptomatic PD (incidental Lewy body disease) showed no PC rise. This information suggests that in PD, oxidative protein damage occurs late during disease progression and/or that L-DOPA treatment contributes to carbonyl formation [121]. Oxidative damage to proteins in PD also occurred through nitration, and 3-NT was found to be increased within Lewy bodies in the SN pars compacta (SNpc) of PD patients [123].

Another pathology found in PD brains is DNA and RNA damage [124, 125]. The factors 8-OHG and 8-OHdG were elevated in various parts of the PD brain compared to controls; however, the most striking rise was detected in the SN [124, 125]. Similarly, increased levels of 8-OHG and 8-OHdG were observed in CSF [126, 127]. However, these studies came to opposite conclusions with respect for the correlation between 8-OHG levels and disease duration.

In the CSF of living PD patients, enhanced levels of HNE and MDA have been shown as well [128–130], but different results were obtained by Shukla et al. [131]. Moreover, markers of oxidative damage in PD patients were also detected in the serum and urine [132–135], but their use as indicators of the course of the disease is far from being useful for clinical practice because the existing data are contradictory [136–138]. As has been suggested [136–138], these differences may be due to the variability in methods used to measure OS markers.

The results of many studies have demonstrated the presence of OS in the brain, CSF, serum, and urine of PD patients; however, none of the OS markers has been established as a specific biomarker for PD disease or as a marker for PD disease progression.

Antioxidant Defense Biomarkers

PD is characterized by a selective loss of GSH in the SN (40 % compared to control subjects) but not in other parts of the brain [139]. This decrease is one of the earliest biochemical changes that has been observed in the disease [140–143], and it results in a selective drop in mitochondrial complex I activity, another hallmark of PD [144]. On the other hand, a substantial rise in SOD levels has been observed in the SN and basal ganglia in PD patients [145], while no change in activities of CAT, GPx, and GR was found compared to age-matched controls [145]. Another study showed some deficiency in GPx in the SN in Parkinsonian patients [146], but the weak (ca. 19 %) reduction in such a small number of patients ($n=11$) cannot be considered to be conclusive. Another small pilot study in PD patients ($n=8$) indicated a peripheral (in blood) increase in GSH/GSSG [147] when the patients went off of PD medications (dopamine receptor agonists) for 12 h in comparison to GSH/GSSG levels when they were on medications. This suggests that PD medications may play a disadvantageous role that leads to enhanced peripheral oxidative stress; however, the small sample size excludes a final conclusion [147].

Pharmacological Strategies to Reduce OS

Because there is an overwhelming amount of evidence showing that OS and NS in PD patients leads to an increase in pathological damage in the SN, many approaches have been attempted to reduce ROS/RNS production. One of the possible strategies was to supplement GSH. As shown by Sechi et al. [148], in nine patients in the early stage of the disease [148], GSH (600 mg twice daily) administered intravenously for 30 days reduced (by 42 %) the clinical disability of PD patients, and this effect lasted for 2–4 months. Furthermore, a recent study of *N*-acetylcysteine (150 mg/kg), which is a GSH precursor, revealed an elevation in brain GSH levels and the blood GSH/GSSG ratio after 1-h of intravenous (i.v.) infusion [149]. Unfortunately, no results concerning the clinical status of PD patients have been described. However, *N*-acetylcysteine, in a smaller dose (70 mg/kg p.o., every 12 h over 2 days), produced negligible changes in GSH levels in the CSF and no immediate improvement to symptoms (according to the Unified Parkinson's Disease Rating Scale and the Montreal Cognitive Assessment) [150].

Magnetic resonance imaging (MRI) studies showed a rise in iron concentrations in the SN in PD patients [151]. Because iron can lead to ROS production in PD patients, an iron-binding compound, deferiprone, has been tested in a pilot study in PD patients (FAIRPARK trial, registered as ClinicalTrials.gov NCT00943748). Patients in early-stage PD who were on a stabilized dopamine regimen received

deferiprone (30 mg/kg/day) for 12 months (early-start paradigm, $n=19$), while the other group received the drug for only 6 months, with the latter, shorter therapy starting 6 months later than the early-start paradigm (delayed-start paradigm, $n=18$) [152]. The earlier therapy start diminished SN iron deposits to a greater extent than the delayed-start paradigm and improved motor performance vs. placebo and vs. the delayed-start paradigm, according to the Unified Parkinson's Disease Rating Scale [152]. Moreover, in deferiprone-treated patients, GPx and SOD activity in the CSF increased, which supports the connection between the chelator treatment and the antioxidant response.

Vitamin E (α -tocopherol) was also suggested as a way to diminish the OS and to reduce clinical symptoms in PD. However, the clinical trial The Deprenyl and Tocopherol Antioxidant Therapy of Parkinsonism study (DATATOP study) revealed no evidence of any beneficial effect for α -tocopherol (2,000 IU/day) in either slowing functional declines or ameliorating the clinical features of Parkinson's disease. It is of note that no analysis of OS biomarkers was performed in that trial [153, 154]. Since DATATOP, no clinical trials using vitamin E as a potential PD medication have been conducted. In fact, vitamin E was only used in PD clinical trials as a supplement for coenzyme Q10 or as a placebo [155] or a control [156].

Another potent antioxidant, coenzyme Q10 (1200 mg a day), in the first reported multicenter, randomized, placebo-controlled, and double-blind trial slowed functional declines compared to placebo [157]. Lower doses or different formulations of coenzyme Q10 displayed no symptomatic effects on midstage PD [158]. Recent clinical trial did not show a benefit for coenzyme Q10 (1,200 or 2,400 mg/day) in 600 patients in early-stage PD [155].

An early clinical study on creatine monohydrate (10 g/day for 12 months) in 67 subjects who were within 5 years of a PD diagnosis showed some positive results for treating behavioral difficulties [159], and an additional 18-month follow-up study confirmed this finding [160]. On the other hand, a smaller ($n=31$ creatine PD group, $n=17$ placebo PD group), but longer, 2-year clinical trial demonstrated that creatine had no effect on PD scores or dopamine transporter imaging [161]. In September 2013, the NINDS discontinued the NET-PD LS-1 study (phase III clinical trial with a total of 1,720 planned participants, ClinicalTrials.gov identifier: NCT00449865) that started in 2007 because the results obtained from a study of creatine used for the treatment of early stage PD did not demonstrate a statistically significant difference between the active substance and placebo [162].

In conclusion, although evidence for the link between OS and damage in PD is overwhelming, suggesting the potential efficacy of antioxidant drugs, most clinical trials have so far failed to support this statement.

Anti-Parkinsonian Strategies to Restore Oxidative Balance

Administration of zonisamide, an anticonvulsant drug prescribed to treat resting tremor in PD, inhibited the rise of 8-OHdG levels in the urine of PD patients. As the 8-OHdG rise correlates with disease progression and aging, it can be presumed that zonisamide could be helpful in defending against OS-evoked DNA modifications in PD patients. Other drugs used for treatment of PD (i.e., L-DOPA, monoamine oxidase B (MAO-B) inhibitors, catechol-*O*-methyltransferase (COMT) inhibitors, and the dopamine receptor agonists ropinirole and pramipexole) have not shown such properties [163].

Interesting findings were reported in a study that measured GSH levels in venous blood in PD subjects who were on- and off-medication while performing acute physical exercises, because we know that this type of physical activity leads to GSH depletion and GSSG rise [164]. Surprisingly, the off-medication patients had a lower drop in GSH level than the on-medication group. This finding suggests that patients in the off-medication state handled acute stress better than those in the on-medication state, indicating that medication may impede the ability to tolerate acute OS [165]. Similar conclusions were obtained in a very recent study by Nikolova et al. [166], who observed a greater rise in PC and 8-OHdG levels in the blood of PD patients who were treated with L-DOPA in comparison to untreated PD patients, demonstrating that administration of L-DOPA may cause greater OS than PD itself [166].

Animal Studies

Oxidative Biomarkers

The most popular animal models of PD include pharmacological (6-hydroxydopamine (6-OHDA), 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), rotenone, and paraquat) as well as several genetic (with mutations in the α -synuclein, PINK1, Parkin, or LRRK2 genes) models [167].

The 6-OHDA model (Table 5), wherein the toxin is injected directly into the SNpc, medial forebrain bundle, or striatum, was the first animal model of PD associated with dopaminergic neuronal death within the SNpc [168].

Another PD model utilizes MPTP, a highly lipophilic molecule that rapidly crosses the blood–brain barrier, leading to an irreversible and selective loss of dopaminergic neurons in the SN in non-human primates [169, 170] and in rodents [162, 171], although the latter species was less sensitive to MPTP than primates [172]. Other chemical models are based on an insecticide, rotenone, or paraquat, an herbicide. Rotenone, when given i.v. in a low dose to rats, produces selective degeneration of SN dopaminergic neurons that is accompanied by α synuclein-positive LB-like inclusions [173]. Paraquat is

used less widely than MPTP, rotenone, or 6-OHDA models and is used instead as an addition to other toxic agents, such as the fungicide maneb [174]. It was reported to cause selective degeneration of nigrostriatal dopaminergic neurons in mice [175]. The mechanism of action of MPTP (a precursor form of the reactive metabolite MPP⁺), rotenone, or paraquat is based on the inhibition of mitochondrial complex I, which leads to ROS production [167, 174, 175].

As has been demonstrated in numerous studies, OS is widely present in all of these toxin-based models (see Table 5). 6-OHDA, when injected into the striatum, increased MDA, HNE, PC, and 3-NT levels in this structure and increased TBARS in SN [176–179]. Similarly, in the MPTP and in the rotenone models, elevated levels of lipid peroxidation products [180–185] and oxidatively modified proteins [185, 186] were observed in various parts of the brain (striatum, cortex, SN, hippocampus, cerebellum, and midbrain). In addition to lipid damage, increased 3-NT levels were also detected following the use of MPTP in the SN, striatum, and ventral midbrain [181, 187]. MPTP or rotenone-treated animals also showed oxidatively modified RNA or DNA in the SN or striatum [185, 188, 189]. In the paraquat and maneb PD models, enhanced lipid peroxidation in the nigrostriatal areas of animal brains was also shown [190] (Table 5). A very recent report from Kumar et al. [175] demonstrated for the first time that α -synuclein radical is formed in the midbrain in paraquat- and maneb-treated mice, indicating that radical formation is mediated by peroxynitrite and depends on both NADPH oxidase and iNOS. This interesting result regarding α -synuclein radical formation was obtained by using the immuno-spin trapping method in combination with immunoprecipitation [175]. Moreover, it was noted that protein radicals such as α -synuclein radical may trigger protein aggregation, which plays a causal role in dopaminergic neuronal death [175].

For review of genetic models and OS, see the excellent paper [167].

All toxin-based models share common characteristics, including the ability to produce ROS and further oxidative damage, which causes death in dopaminergic neurons and reflects part of the pathology observed in PD. Although all of those models have drawbacks, they are useful for testing neuroprotective therapies.

Antioxidant Defense Biomarkers

A characteristic shared feature observed in all toxin-based models is a drop in GSH level in key PD structures [177, 183–187, 191–196] (Table 5). Importantly, lower GSH levels make nigrostriatal neurons more susceptible to oxidative damage and further degeneration.

Studies using 6-OHDA also showed a reduction in activity by SOD, CAT, and glutathione *S*-transferase (GST) in striatum

Table 5 Changes in OS and anti-OS defense biomarkers in toxin-based model of PD

Animal model	Route of administration, dosage	Animal species	OS biomarkers	Oxidative defense biomarkers	Reference
6-OHDA	Unilateral intra-STR inj. 10 µg/2 µl	Rats (Wistar)	↑TBARS in SN	↓GSH, ↓CAT ↓SOD in SN	[191]
	Bilateral intra-STR inj. 10 µg/2 µl	Rats (Fischer)	↑HNE, ↑PC in STR	nd	[176]
	Unilateral intra-STR inj. 300 µg/10 µl	Rats (Sprague-Dawley)	↑PC in STR	↓GSH and ↓SOD in STR	[177]
	Intra-STR inj. 8 mg/2 ml	Rats (Wistar)	↑MDA in STR	↓SOD, GST in STR	[178]
	Intra-STR inj. 8 µg/4 µl	Rats (Wistar)	↑3-NT, ↑HNE in STR	nd	[179]
MPTP	20 µl/kg intra-SN inj.	Rats (Wistar)	↑MDA in SN	↓SOD in SN	[180]
	25 mg/kg/day for 5 days	Mice (C57BL/6)	↑iNOS, ↑HNE, ↑3-NT in SN	nd	[181]
	Intra-SN inj.	Rats (Wistar)	↑MDA, ↑nitrate in STR and CTX	↓GSSH, ↓CAT in STR and CTX	[182]
	24 mg/kg sc. for 6 days	Mice (C57BL/6)	↑MDA in STR	↓GSH, ↓SOD, ↓GPx, ↓CAT in STR	[197]
	100 µg/1 µl, bilateral infusions	Rats (Wistar)	↑LPO in STR	↓GSH, ↑SOD in STR	[183]
	15 mg/kg s.c. for 3 days	Mice (C57BL/6)	↑3-NT, ↑HNE in ventral midbrain	↓GSH, ↑GSSG in ventral midbrain	[187]
	24 mg/kg s.c. (2 times/day at 12-h interval) for 2 days	Mice (C57BL/6)	↑8-OHG in SN	nd	[188]
	20 mg/kg i.p. 3 times at 2-h interval	Mice (C57BL/6J)	↑8-OHG in STR	nd	[189]
	30 µg/kg i.p. 2 times at 16-h interval	Mice (Balb/c)	nd	↓GSH, ↑SOD in SN, nucleus caudate, and putamen	[192]
	Rotenone	2.5 mg/kg, i.p., for 10 days	Rats (Wistar)	nd	↓GSH in HIP, ↓CAT in HIP and STR
One-sided intra-SN inj. 6 µg		Rats (Sprague-Dawley)	nd	↓GSH, ↑Cu-Zn SOD, ↑CAT in SN, Ø GSH, Ø SOD in STR	[194]
3 mg/kg i.p. for 30 days		Rats (Wistar)	nd	↓GSH, ↓SOD in STR	[195]
2 mg/kg s.c. for 35 days		Rats (Sprague-Dawley)	↑TBARS, ↑SAG in the midbrain regions and cerebellum and CTX	↓GSH, ↓CAT in midbrain	[196]
2 mg/kg s.c. for 35 days		Rats (Sprague-Dawley)	↑MDA in midbrain regions and cerebellum	↓GSH, ↓SOD, and ↑CAT in midbrain and CTX ↑NO in midbrain	[184]
1 mg/kg i.p. for 3 weeks		Mice (CFT-Swiss)	↑PC in HIP, CTX, STR	↓GSH, ↓TT in HIP, ↓GPx, ↓CAT in CTX and HIP, ↓SOD in CTX and STR	[186]
Paraquat + maneb	1.5 mg/kg/48 h s.c., 6 doses	Rats (Albino)	↑LPO, ↑PC, ↓mtDNA in STR	↓GSH, ↓SOD in STR	[185]
	Paraquat (10 mg/kg) + maneb (30 mg/kg), twice a week, for 9 weeks	Mice (Swiss albino)	↑MDA, ↑NO in nigrostriatal tissues	↑GST in nigrostriatal regions	[190]
	Paraquat (10 mg/kg, i.p.), and maneb (30 mg/kg, i.p.), twice a week, for 6 weeks	Mice (C57BL/6J)	α-Synuclein radical in the midbrain	nd	[175]

↑ increase, ↓ decrease, Ø no change, *nd* not determined, *6-OHDA* 6-hydroxydopamine, *CAT* catalase, *CTX* cortex, *GPx* glutathione peroxidase, *GSH* glutathione, *GST* glutathione *S*-transferase, *HIP* hippocampus, *HNE* 4-hydroxynonenal, *i.c.v.* intracerebroventricular, *intra-SN inj.* intra-substantia nigra injection, *LPO* lipid hydroperoxide, *MDA* malonyldialdehyde, *MPTP* 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, *NO* nitric oxide, *NOS* nitric oxide synthetase, *PC* protein carbonyl, *SAG* superoxide anion generation, *s.c.* subcutaneous, *SN* substantia nigra, *SOD* superoxide dismutase, *STR* striatum, *TBARS* thiobarbituric acid reactive substances

and SN [177, 178, 191]. On the other hand, results from the MPTP model are inconclusive regarding SOD and CAT activity. Moreover, some of the MPTP studies showed increased SOD activity in the SN [192] and striatum [183], while others reported diminished SOD activity in these regions [180, 197]. These differences may have resulted from the use of different doses of the toxin, varied routes of drug administration (intra-cranial versus i.p. or s.c.), and different use of rodent species in the studies (see Table 5).

Similar to SOD activity, CAT activity cannot be considered a biomarker of OS in rodent PD models as its activity was both diminished [182, 191, 193, 194, 196, 197] and enhanced [184]. Moreover, GPx activity was diminished in striatum in an MPTP model [197], while GST activity was found to be elevated in a maneb and paraquat PD animal model [190] (Table 5).

All of these reports on the enhanced activities of SOD, CAT, and GST suggest the presence of mechanisms in brain

areas that defend against exposure to PD toxin models. On the other hand, diminished activities or levels of antioxidant enzymes may indicate that these defense mechanisms were overcome and that the degeneration process had begun.

Pharmacotherapeutic Strategy to Reduce OS

Several agents, such as valproic acid [178] and melatonin [198], effectively reversed changes in antioxidant defense biomarkers and oxidative damage in the 6-OHDA rat model of PD (Table 6).

There are also data in the literature showing that other agents and drugs have antioxidant activity (i.e., ceftriaxone [182] and *N*-acetylcysteine [199], which decreased oxidative damage and/or enhanced antioxidant defense biomarkers in the striatum, cortex or SN in rodent MPTP models).

Ibuprofen (a non-steroidal anti-inflammatory drug [193]), acetyl-L-carnitine (a natural compound reported to prevent mitochondrial injury deriving from oxidative damage in vivo), α -lipoic acid (given alone or in combination with acetyl-L-carnitine [185]), and centrophenoxine (a potent nootropic agent that acts as an antioxidant) [184] were demonstrated to enhance GSH levels and CAT and SOD activity and to

Table 6 Therapeutic trials with substances possessing antioxidant properties in PD animal models and their influence on changes of biomarkers of OS and anti-oxidative defense

Model	Biomarkers of OS and anti-oxidative defense	Drug and route of administration	Changes in biomarkers of OS and anti-oxidative defense	Reference
6-OHDA, rats (Wistar)	↑MDA, ↓SOD, ↓GPx, ↓CAT in STR	Melatonin (10 mg/kg i.p. for 30 days)	↓MDA, ↑SOD, ↑GPx, ↑CAT in STR	[198]
6-OHDA, 8 mg/2 ml intra-STR inj., rats	↑MDA, ↓SOD, ↓GST in STR	Valproic acid (300 mg/kg i.p. for 10 days)	↓MDA, ↑SOD in STR	[178]
MPTP intra-SN inj., rats	↓GSH, ↓CAT, ↑nitrate, ↑LPO in STR and CTX	Ceftriaxone (100 mg/kg or 200 mg/kg i.p.)	↑GSH, ↑CAT, ↓nitrate, ↓LPO in STR and CTX	[182]
		Ceftriaxone (100 mg/kg i.p. and ropinirol 1, 5 or 3 mg/kg i.p.)	↑GSH, ↑CAT, ↓nitrate, ↓LPO in STR and	
MPTP mice (C57BL/6J)	↑LPO, ↑SOD, ↓GPx, ↓GSH in SNpc	<i>N</i> -acetylcysteine	↓LPO, ↓SOD, ↑GPx, ↑GSH in SNpc	[199]
Rotenone, 2.5 mg/kg, i.p. for 10 days, rats	↓GSH in HIP, ↓CAT in HIP and STR	Ibuprofen (15 mg/kg, p.o. 22 days post 10-day rotenone treatment)	↑GSH in HIP vs. rotenone group, ↑CAT in HIP and STR vs. rotenone group	[193]
Rotenone, 3 mg/kg i.p. for 30 days, rats	↓GSH, ↓SOD, ↑MDA in STR	Lycopene (10 mg/kg, p.o. for 30 days, a combined treatment with rotenone)	↑GSH, ↓SOD, ↑CAT in HIP, STR	[195]
Rotenone, 1 mg/kg/day i.p. for 3 weeks, mice (CFT-Swiss)	↓GSH in HIP, ↓TT in HIP, ↓GPx, ↓CAT in CTX and HIP, ↓SOD in CTX and STR, ↑PC in HIP, CTX, STR	TSE—aqueous extract of tomato seeds (100 mg/kg p.o. for 3 weeks, 1 h before rotenone injection)	↑GSH, ↑CAT, ↑TT in HIP, ↓PC in STR, ↑SOD in CTX and STR, ↑GPx in CTX and HIP ↓PC in STR, ↑SOD in CTX and STR	[186]
Rotenone, 1.5 mg/kg/48 h/6 doses, s.c., rats (Albino)	↑LPO, ↑PC, ↓GSH, SOD, ↓mtDNA in STR	Acetyl-L-carnitine (100 mg/kg/day, p.o.), α -lipoic acid (50 mg/kg/day, p.o.) or their combination	↓LPO, ↑GSH, ↑SOD, and ↑mtDNA in STR after each drug alone or after combination, ↓PC, ↑CAT in STR only after combination	[185]
Rotenone, 2 mg/kg s.c. for 35 days, rats (Sprague-Dawley)	↑MDA in midbrain and cerebellum, ↓GSH, ↓SOD, and ↑CAT in midbrain and CTX, ↑NO in midbrain	Centrophenoxine (100 mg/kg i.p., co-administration with rotenone for 35 days)	↓MDA in midbrain and cerebellum, ↑GSH, ↑SOD, ↑CAT in midbrain and CTX, ↓NO in midbrain	[184]
Maneb (30 mg/kg) and paraquat (10 mg/kg) i.p. twice a week for 9 weeks, mice (Swiss albino)	↑MDA, ↑NO, ↑GST in nigrostriatal regions	Silymarin (40 mg/kg i.p. for 9 weeks), maneb, and paraquat were administered 2 h after this injection Melatonin (30 mg/kg i.p. for 9 weeks, maneb and paraquat were administered 2 h after this injection)	↓MDA, ↓NO, and ↓GST in nigrostriatal tissues	[190]

↑ increase, ↓ decrease, *nd* not determined, *CAT* catalase, *CTX* cortex, *GPx* glutathione peroxidase, *GSH* glutathione, *GST* glutathione transferase, *GST* glutathione *S*-transferase, *HIP* hippocampus, *HNE* 4-hydroxynonenal, *i.c.v.* intracerebroventricular, *i.p.* intraperitoneal, *intra-SNc inj.* intra-substantia nigra injection, *intra-STR inj.* intrastriatal injection, *LPO* lipid hydroperoxide, *MDA* malonyldialdehyde, *NO* nitric oxide, *NOS* nitric oxide synthetase, *PC* protein carbonyl, *s.c.* subcutaneous, *SN* substantia nigra, *SNpc* substantia nigra pars compacta, *SOD* superoxide dismutase, *STR* striatum, *TBARS* thiobarbituric acid reactive substances, *TT* total thiol

decrease lipid peroxidation in investigated brain regions in a rat rotenone model (Table 6).

Prevention of oxidative damage and the presence of antioxidant defense biomarkers have been documented following treatment with natural compounds, such as lycopene [195], aqueous extract of tomato seeds (TSE) [186], and melatonin [190].

Many different agents may improve antioxidant brain status in different PD models. However, it should be noted that most of these agents were given before or concomitantly with rotenone, MPTP, or other PD-causing toxins. To definitively answer whether these agents can also show efficacy in reducing the consequences of exposure to prior administration of PD-inducing toxins, further studies are required. This is especially true because the latter type of drug administration would be a better model for evaluating any pharmacological strategy for reducing OS in PD patients.

Anti-Parkinsonian Drugs that Modulate Oxidative Balance

Most anti-parkinsonian drugs may improve brain antioxidant status in PD preclinical tests (Table 7). Ropinirole, a second-generation, non-ergoline dopamine receptor agonist with D2-like receptor selectivity and a chemical structure similar to that of dopamine was found to enhance GSH levels and CAT [182] activity and to diminish nitrate levels [182] in the striatum in MPTP-lesioned animals.

Other anti-parkinsonian drugs, such as selegiline (a selective irreversible MAO-B inhibitor) [196], deferoxamine [177], and pramipexole (a non-ergoline dopamine agonist) [200],

increased GSH levels in the striatum, SN, or cortex. Deferoxamine also decreased a protein oxidative damage biomarker [177] and enhanced SOD activity in the striatum, while selegiline reduced superoxide anion generation (SAG) and increased CAT activity in midbrain regions and the cortex [196]. Interestingly, L-DOPA, the most commonly used drug in PD treatment, did not restore the reduced GSH levels in the SN in the MPTP mouse model [200].

The above studies suggest that antiparkinsonian drugs, with the exception of L-DOPA, display some antioxidant properties, which may be considered as part of their mode of action and efficacy in PD treatment.

Alzheimer's Disease (ICD-10 G30)

AD is the most common neurodegenerative disease and is characterized by memory loss, dysfunctions in cognitive abilities (e.g., executive function, attention, language, and visuospatial skills), confusion, aggression, and mood swings [113, 201] and leads to death within 5 to 9 years after the diagnosis [202].

The pathogenesis of AD is not yet clearly understood. The aggregation of extracellular insoluble protein plaques (composed of beta amyloid (A β)) and intracellular neurofibrillary tangles (NFTs, composed of tau protein) are critical hallmarks of AD [113, 201]. However, many ongoing pathological processes lead to regional neuron loss, beginning in the medial temporal lobe [201] and following in other brain regions, such as the hippocampus and cerebral cortex [113].

Many clinical trials and animal studies have recognized free radicals as mediators of injury in AD patients and AD models.

Table 7 Therapeutic trials with anti-parkinsonian drugs in PD animal models and their influence on biomarkers of OS and of anti-oxidative defense

Model	Oxidative defense biomarkers	Drug	Changes in oxidative defense biomarkers	Reference
MPTP 25 mg/kg i.p. for 5 days administered to 6C57BL/6J mouse	↓GSH in SN	L-DOPA (200 mg/kg i.p. 2 injections/day for 4 weeks, co-administration with MPTP)	∅ GSH in SN	[200]
MPTP 1 μ mol/2 μ l intra-SN administered to Wistar rat	↓GSH, ↓CAT, ↑nitrate, and ↑LPO in STR and CTX	Ropinirole 1, 5, or 3 mg/kg i.p. for 14 days, after MPTP treatment	↑GSH, ↑CAT, ↓nitrate (only 1.5 mg/kg) in STR and CTX	[182]
MPTP 25 mg/kg i.p. for 5 days administered to C57BL/6J mouse	↓GSH in SN	Pramipexole 1 mg/kg i.p. 2 injections/day for 4 weeks, co-administration with MPTP	↑GSH in SN	[200]
Rotenone 2 mg/kg s.c. 35 days administered to Sprague-Dawley rats	↑TBARS, ↑SAG in midbrain regions and cerebellum, ↓GSH and ↓CAT in midbrain regions and CTX	Deprenyl 10 mg/kg p.o. co-administration with rotenone for 35 days	↓TBARS, ↓SAG in midbrain; ↑GSH, ↑CAT in midbrain and CTX	[196]
6-OHDA 300 μ g/10 μ l unilateral intra-STR inj. administered to Sprague-Dawley rats	↑PC, ↓GSH, and ↓SOD in STR	Deferoxamine 50 mg/kg p.o. for 14 days, co-administration with 6-OHDA	↓PC, ↑GSH, and ↑SOD in STR	[177]

↑ increase, ↓ decrease, ∅ no change, *na* not applicable, 6-OHDA 6-hydroxydopamine, CAT catalase, CTX cortex, GSH glutathione, *i.p.* intraperitoneal, LPO lipid hydroperoxide, MPTP 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, PC protein carbonyl, *s.c.* subcutaneous, SAG superoxide anion generation, SOD superoxide dismutase, STR striatum, TBARS thiobarbituric acid reactive substances

Clinical Studies

Oxidative Biomarkers

The first report of the involvement of OS in AD pathology came from a paper by Martins et al. [203], in which elevated activities of glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase were documented in *post-mortem* studies of brains of AD patients compared to age-matched controls [203]. The latter increase was proposed to be a response to enhanced brain peroxide metabolism.

Other *post-mortem* studies on brains and CSF from AD patients showed ROS-mediated injuries. For instance, AD patients had increased levels of MDA and HNE, iso- and neuroprostanes, and acrolein compared to controls [204]. It was suggested that these peroxidated lipids formed adducts with proteins and that they might thereby play a role in AD pathogenesis [201].

In addition to lipids, protein damage due to OS has also been reported in AD. In fact, increased PC levels in the frontal and parietal cortices and the hippocampus were found in *post-mortem* studies of the brains of AD patients, while PC was absent in the cerebellum, where no AD pathology was present [205]. Furthermore, evidence of oxidative DNA modification was found in AD patients as an increase in 8-OHG in human brain homogenates [201].

In AD patients, ROS production seems to be enhanced; furthermore, increases in RNS were also detected. Such evidence of RNS modification was identified both in astrocytes and in neurons in AD patient brains examined *post-mortem* [206]. The changes in astrocytes were found to co-localize with an increase in iNOS, eNOS, and nNOS expression. The latter increases were noted specifically in cortical pyramidal cells [206]. In another study, increased expression of iNOS and eNOS was observed to be directly associated with A β deposits, showing that beta amyloid might induce NOS to produce NO, which might lead to 3-NT formation [207].

The presence of 3-NT was also reported in the cerebral blood vessels of AD patients *post-mortem* [206]. These findings were associated with reduced NO bioavailability in plasma and further hypoperfusion in AD patients because NO promotes vascular smooth muscle relaxation and thereby regulates blood flow. As was reported in recent studies, enhanced NO production is a likely cause of production of ONOO⁻ [208, 209].

Another set of oxidative damage biomarkers, 8-OHdG and 8-OHG, were elevated in AD ventricular CSF [210] and in brains in both mitochondrial and nuclear DNA compared with age-matched controls [211].

Consistent data showing enhanced levels of MDA, HNE, iso- and neuroprostanes, acrolein, PC, 8-OHG, 8-OHdG, and 3-NT in the CNS of AD patients can be considered to be proof that OS and NOS are significant contributors to brain damage.

Antioxidant Defense Biomarkers

Pivotal antioxidant enzymes, including GPx, CAT, and SOD, display changed levels in the brains of AD patients [212, 213]. However, the data are not consistent. For instance, elevated levels of antioxidant enzymes (mainly SOD) in the hippocampus and amygdala of AD patients have been reported [208]. On the other hand, in AD patients, decreased levels of SOD, GPx, and CAT were found in the frontal and temporal cortex [214], while decreases in GSH were observed in the brain and erythrocytes of AD patients [212, 213]. Evidence in support of changes in antioxidant enzymes comes from a recent study that identified genetic polymorphisms in the *GPx-1* and *GST* genes that were positive risk factors for AD [215, 216].

The GSH levels were reduced not only in AD but also in mild cognitive impairment (MCI), which is considered to be a preclinical stage of AD [217]. MCI patients also showed a decreased GSH/GSSG ratio and a reduction in SOD and GST activity in the hippocampus compared to age-matched controls [218], which suggests that alterations in GSH metabolism may be considered as an early biomarker of AD onset.

The plasma levels of antioxidants, such as albumin, bilirubin, uric acid, lycopene, vitamin A, vitamin C, and vitamin E, are decreased in AD patients [219, 220], although there are some reports indicating the opposite direction of these changes [221]. Differences in results might be caused by measurement of antioxidants at different disease stages (fully developed disease vs. subclinical stage of the disease) [219–221].

Pharmacological Strategy to Reduce OS

As OS is present in AD patients, some clinical studies have aimed to test the ability of antioxidant substances to diminish ROS production and to alleviate or to slow the course of the disorder (Table 8).

Most studies on the effects of the administration of vitamins that possess antioxidant activity have provided inconclusive information showing that they diminished lipid peroxidation in CSF but had no positive effects on cognitive or functional aspects. For example, AD patients taking cholinesterase inhibitors and vitamin C (1,000 mg/day) or vitamin E (400 IU/day) supplements for 1 year showed decreased CSF oxidation, but no difference in cognition was observed [222]. Similarly, 16-week treatment with a combination of vitamin E (800 IU/day), vitamin C (500 mg/day) and α -lipoic acid (900 mg/day) decreased CSF F2-isoprostane levels, which suggested a reduction of OS in the brain, but surprisingly, this therapy accelerated cognitive decline (according to Mini-Mental State Examination scores), leading to the conclusion that a combination of antioxidants should not be used for AD therapy [223]. On the other hand, very recent results from a large, clinical, double-blind, randomized trial (TEAM-AD VA, NCT00235716) showed that a much larger dose of vitamin

Table 8 Clinical trials with substances/drugs with antioxidant properties in AD patients

Antioxidant	Duration of therapy	Oxidative damage biomarker change	Reference
Vitamin E (α -tocopherol, 800 IU/day) + vitamin C (500 mg/day) + α -lipoic acid (900 mg/day)	16 weeks	↓F2-isoprostane in CSF	[223]
Coenzyme Q10 (400 mg \times 3 times/day)	16 weeks	Ø F2-isoprostane in CSF	[223]
ω -3 (3 g/day contained 675 mg DHA and 975 mg EPA)	12 months	Ø F2-isoprostane in urine, Ø PC in plasma	[228]
ω -3 + α -lipoic acid (ω -3, 3 g/day contained 675 mg DHA and 975 mg EPA + α -lipoic acid, 600 mg/day in one tablet)	12 months	Ø F2-isoprostane in urine Ø PC in plasma	[228]
Vitamin C (1,000 mg/day) + vitamin E (400 IU/day)	12 months	↓oxidation of CSF	[222]
Curcumin (1 or 4 g/day)	6 months	Ø F2-isoprostane in plasma	[230]
Curcuminoids (2 or 4 g/day)	24 weeks	Ø F2-isoprostane in CSF	[229]
Idebenone (120, 240, or 360 mg/day)	12 months	nd	[225]

↓ decrease, Ø no change, nd not determined, CSF cerebrospinal fluid, DHE docosahexaenoic acid, EPA eicosapentaenoic acid, ω -3 omega-3 fatty acids

E (2,000 IU/day) than was used in previous trials resulted in a slower functional decline compared with placebo in mild to moderate AD [224]. Although the latter study suggests that vitamin E can have a positive influence on AD, no OS biomarkers have been measured in parallel in the AD patients who participated in that trial, which limits the final conclusion.

Administration of other antioxidants, including coenzyme Q10 as well as its synthetic analogue, idebenone (which possesses a better ability to pass the blood–brain barrier), in AD patients did not provide any positive results with regards to the volume of ROS-dependent tissue damage or cognitive function improvements [223, 225]. Similarly, administration of omega-3 (ω -3) fatty acids also did not yield a positive outcome for slowing the rate of decline of cognitive (Mental State Examination (MMSE); Alzheimer’s Disease Assessment Scale—cognitive subscale) or functional (Activities of Daily Living/Instrumental Activities of Daily Living) abilities [226, 227]. Different results were reported in a recent study, where 12-month ω -3 fatty acid supplementation caused a delay in progression of functional impairment in AD patients, while combined supplementation of ω -3 and α -lipoic acid resulted in slowing global cognitive declines (MMSE) [228]. Although positive cognitive outcomes were obtained, no changes after ω -3 or ω -3 plus α -lipoic acid supplementation were observed in OS biomarkers, suggesting a different mechanism for their actions that lead to improved cognitive and functional measures [228].

Curcumin, which is a natural polyphenolic compound and an in vitro blocker of A β aggregation, did not diminish the enhancement of F2-isoprostane levels in the CSF [229] or plasma [230], or the A β _{1–40} level in plasma [230], and it did not ameliorate neuropsychological test results in AD patients [229, 230]. As suggested by Ringman et al. [229], low bioavailability of the drug and low plasma levels due to poor uptake from the gastrointestinal tract might be the reasons for its lack of efficacy in the latter studies. There is some hope that curcumin efficacy can be improved through the use of its

lipidated forms, which are predicted to have better uptake compared to the nonlipidated form [231]. In a study in healthy, middle-aged volunteers, the lipidated form of curcumin (80 mg/day) decreased A β _{1–40} levels in plasma [231], suggesting that further trials using lipidated curcumin should be considered in AD patients.

More promising results came from a study using resveratrol. The Copenhagen City Heart Study reported that monthly or weekly consumption of red wine was associated with a lower risk of dementia [232]. The clinical study NCT01504854, also called the “Phase II Study to Evaluate the Impact on Biomarkers of Resveratrol Treatment in Patients with Mild to Moderate AD,” was undertaken in 2011 to evaluate the effectiveness of resveratrol for changing AD CSF biomarkers (total tau, A β ₄₂, A β ₄₀, and phospho-tau181), the effect of resveratrol treatment on hippocampal atrophy and regional cortical thinning, and the influence of resveratrol administration on the outcomes of several clinical scales used to assess the severity of dementia [233]. According to the report at <http://clinicaltrials.gov> record (accessed 15 May 2015), the study has been completed, but no results have yet been published.

Anti-Alzheimer’s Disease Therapy and Oxidative Balance

Acetylcholinesterase (AChE) inhibitors (donepezil, rivastigmine, galantamine, and tacrine) and the NMDA receptor antagonist memantine are the most commonly used drugs in AD pharmacotherapy. Only some clinical studies that have investigated the influence of these drugs on oxidative balance in AD patients are currently available (see Table 9). One of them showed no positive effects of AChE inhibitors on OS parameters (CAT and GR levels) in the blood of AD patients compared with AD drug-naïve patients [234]. In another study, donepezil enhanced GSH levels, while rivastigmine diminished advanced glycation end products (AGEs) in the plasma of AD patients. However, other examined parameters, namely total

Table 9 Clinical trials of anti-Alzheimer drugs and their influence on OS biomarkers

Anti-Alzheimer medication (dose)	Duration of therapy	Oxidative damage biomarker change	Reference
Donepezil (10 mg/day)	≥24 months	↑GSH, Ø AGEs, TAC, PC in plasma	[235]
Tivastigmine (9.5 mg/day)	≥24 months	↓AGEs, Ø GSH, TAC, PC in plasma	
Donepezil (10 mg/day) + memantine (20 mg/kg)	≥24 months	Ø GSH, TAC, PC, AGEs in plasma	
Memantine (20 mg/day)	For 6 months	↓oxidation rates of lipids in plasma	[236]

↑ increase, ↓ decrease, Ø no change, AGEs advanced glycation end products, GSH glutathione, PC protein carbonyl, TAC total antioxidant capacity

antioxidant capacity (TAC) and PC, have not been improved by those drugs [235]. Combined therapy with memantine and donepezil failed to improve GSH, TAC, PC, or AGEs [235]. A very recent study revealed that 6-month treatment with memantine decreased the oxidation rate of plasma lipids in AD patients compared with untreated patients [236]. The above clinical trials included small sample sizes and should initiate future examinations evaluating the effect of different types of AD medications on OS markers in AD patients.

Animal Studies

Oxidative Biomarkers

AD can be modeled by several procedures in animal. Injection with scopolamine (i.p.), streptozotocin (intracerebroventricular (i.c.v.)), A β (i.c.v.), or apolipoprotein E (APOE) in transgenic models is used to study sporadic AD, while amyloid precursor protein (APP) and presenilin 1 (PSEN1) and PSEN2 transgenic models are used to examine familial AD [237–240]. For detailed descriptions of AD animal models, see [241–246].

In both pharmacological and genetic models of AD, disordered OS biomarkers are present in animal brains (Table 10). MDA, HNE, or TBARS were enhanced in all of those models in the cerebral cortex or/and hippocampus or/and the whole brain [237–241, 247–255]. Oxidative modification of proteins has also been demonstrated in the cortex and whole brain homogenate of transgenic AD mice [253, 255] and in the cerebral cortex and hippocampus of an A β ₁₋₄₂ mouse model [247]. In addition to OS due to oxygen, there is also proof of the presence of NS in whole brain lysates from the APP23 transgenic AD mouse model [251].

Antioxidant Defense Biomarkers

Antioxidant defense biomarkers have been found to be changed in AD models (see Table 10). Diminished levels of GSH in the cerebral cortex or hippocampus or in whole brain lysates have been demonstrated in pharmacologically induced AD animal models [237, 238, 240, 241, 247, 250, 254, 256].

Furthermore, the activities of enzymes connected with GSH metabolism, such as GPx and GR, and the enzymes involved in antioxidant defense (SOD and CAT) were reduced in the hippocampus and cerebral cortex in pharmacological and genetic models [237, 238, 241, 248, 249, 254–256]. It should be noted that some studies demonstrated no change in CAT and SOD activity in whole brain lysates in Wistar rats in the streptozotocin model [240], while enhanced SOD, GPx, and GR were observed in the mouse cerebral cortex and hippocampus following i.c.v. A β ₁₋₄₂ injection [247]. It is also important to mention that in transgenic models, the changes depend on animal age. For example, APP^{sw}/PS1^{de9} mice at 2.5 months of age did not exhibit any significant changes in measures of OS and cognitive function, whereas 3.5-month-old mice showed diminished SOD and GPx activity and increased MDA and PC levels that were accompanied by spatial memory impairments [255].

Pharmacotherapeutic Strategy to Reduce OS

Several preclinical studies on AD have shown that many antioxidants can both diminish OS and improve cognitive impairments (Table 11). Among different compounds of special interest are vitamin E, vitamin C, and α -lipoic acid. Vitamin E given 7 days before A β ₁₋₄₂ decreased MDA and protein carbonyls in the mouse hippocampus and cortex [247]. Similarly, α -lipoic acid enrichment decreased HNE levels in AbPP Tg2576 mouse brains but did not decrease 3-NT levels [251]. In AbPP Tg2576 mice that overexpress a mutant form of APP (beta amyloid (β A), an (A4) precursor protein) and show impaired learning, an R- α -lipoic acid-enriched diet, administered for 10 months, decreased HNE levels in total brain homogenates and also attenuated HNE protein adducts that accumulated around amyloid deposits in the hippocampal and cortical region, but it had little effect on cognitive performance and brain A β load. This latter study seems to suggest that a long-term antioxidant therapy that reduced oxidative modifications provided a limited benefit [251].

In contrast to the study that used α -lipoic acid, vitamin C, when administered to other transgenic lines, such as APP/PSEN1 mice, did not decrease enhanced MDA levels in the cortex or A β plaque deposits in the cortex and

Table 10 OS biomarkers and OS defense biomarkers in pharmacologically developed and in transgenic AD animal models

Animal model (dosage)	Animal species	OS biomarkers	OS defense biomarkers	Reference
A β ₁₋₄₂ (i.c.v. injection)	Mice	↑MDA in cerebral CTX and HIP	↓SOD, ↓GPx, ↓GSH, ↑GSSG in cerebral CTX and HIP	[237]
A β ₁₋₄₂ injection (i.c.v. injection)	Mice (Chinese Kun Ming)	↑MDA in cerebral CTX and HIP	↓SOD, ↓GPx, ↓GSH in HIP and cerebral CTX	[241]
A β ₁₋₄₂ (i.c.v. injection 400 pmol)	Mice (C57BL/6)	↑MDA, ↑PC in cerebral CTX and HIP	↑Mn-SOD, ↑Zn, Cu-SOD, ↑GPx (only up to 2 days after A β ₁₋₄₂ injection), ↑GR (only 2 h after A β ₁₋₄₂ injection) in cerebral CTX and HIP	[247]
Scopolamine (2 mg/kg i.p. once per day for 2 weeks)	Mice (Kun Ming)	↑MDA in HIP	↓SOD, ↓GSH in HIP	[238]
Scopolamine (1 mg/kg, i.p. single injection)	Mice (Swiss)	↑MDA in CTX and HIP	↓SOD, ↓GPx, ↓GSH-Rx in CTX and HIP	[248]
Scopolamine (1.4 mg/kg, i.p. single injection)	Mice (Swiss)	↑MDA in whole brain lysate	↓CAT in whole brain lysate	[249]
AF64A, a cholinotoxin (2 nmol/2 μ l, bilaterally single i.c.v. injection)	Rat (Wistar)	↑MDA in HIP	nd	[239]
Streptozotocin (3 mg/kg bilaterally i.c.v. injection on days 1 and 3)	Rat (Wistar)	↑MDA in whole brain lysate	↓GSH, \emptyset CAT, \emptyset SOD in whole brain lysate	[240]
Streptozotocin (2.57 mg/kg bilaterally single i.c.v. injection)	Mice (Swiss albino)	↑TBARS in HIP	↓GSH, ↓GPx, ↓GR in HIP	[250]
Streptozotocin (3.0 mg/kg i.c.v. single injection, 5 μ l/injection per site)	Rat (Wistar)	↑HNE, ↑MDA, ↑TBARS, ↑PC in HIP	↓GSH, ↓GPx, ↓GR, ↓CAT, ↓SOD in HIP	[256]
AbPP Tg2576 transgenic mice		↑HNE, ↑3-NT in whole brain	nd	[251]
APP/PSEN1 transgenic mice		↑MDA in HIP	nd	[252]
APP23 transgenic mice		↑PC in CTX	nd	[253]
Heterozygote APP transgenic mice		↑TBARS in the brain homogenate	↓GSH, ↓SOD in the brain homogenate	[254]
APP ^{swe} /PS1 ^{dE9} transgenic mice		↑MDA, ↑PC in the brain homogenate	↓SOD, ↓GPx in the brain homogenate	[255]

↑ increase, ↓ decrease, *nd* not determined, 3-NT 3-nitrotyrosine, CTX cortex, GSH glutathione, CAT catalase, GPx glutathione peroxidase, GR glutathione reductase, GSSG oxidized glutathione, HIP hippocampus, HNE 4-hydroxynonenal, i.c.v. intracerebroventricular, i.p. intraperitoneal, MDA malonyldialdehyde, Mn-SOD manganese superoxide dismutase (located in mitochondria), PC protein carbonyl, SOD superoxide dismutase, TBARS thiobarbituric acid reactive substances, Zn, Cu-SOD copper/zinc superoxide dismutase (located in cytoplasm)

hippocampal regions in either middle-aged or aged animals [252], although the drug administration improved memory, according to tests that suggested that cognitive rescue was achieved, to some degree, even in animals that suffered from severe neuropathology. The lack of effect of vitamin C on A β plaque deposits seems to result from the late introduction of medication in this test because A β plaques, considered an end point in the disease process, are detectable in these mice at 4–5 months, which was before the beginning of the test [252]. It is also possible that ascorbate had an effect on soluble A β [252]. Reduced vitamin C supplementation has been shown to cause enhanced OS in the form of MDA in APP/PSEN1 mice, which are not able to endogenously synthesize vitamin C [257]. This latter observation led to the conclusion that vitamin C may not be an anti-OS medication per se, but its deficiency in AD patients may lead to oxidative damage. Interestingly,

another study showed that the long-lasting incretin hormone analogue D-Ala²GIP (glucose-dependent insulinotropic polypeptide) was able to decrease OS biomarkers (i.e., 8-OHG) and amyloid plaque load in 12- and 19-month-old APP^{swe}/PS1 mice [258]. Many natural compounds that possess antioxidant properties have been tested in animal models as AD treatments. Imperatorin and hesperidin diminished brain damage due to OS, and most of them enhanced the power of oxidative defenses [238–241, 259]. Moreover, meloxicam (an anti-inflammatory drug) and selegiline, given alone or in combination, inhibited lipid peroxidation, prevented a decrease in CAT activity, and showed memory-enhancing capacity in a scopolamine AD model [249]. Another compound, S-allyl cysteine, which is a sulfur-containing amino acid that was reported to have antioxidant and neurotrophic activity, prevented cognitive and neurobehavioral impairments,

Table 11 Studies with substances/drugs with antioxidant properties in different AD animal models and their influence on oxidative damage and antioxidant defense and biomarkers

Model	Oxidative damage and defense biomarkers	Drug and route of administration	Changes in oxidative defense biomarkers	Reference
A β_{1-42} i.c.v. to mice (Chinese Kun Ming)	↑MDA, ↓SOD, ↓GPx, ↓GSH in HIP and cerebral CTX	Schisantherin A 0.1 mg/kg for 5 days i.c.v., injection started after 3 days from A β_{1-42} injection	↓MDA in cerebral CTX, ↑SOD, ↑GPx, ↑GSH in HIP and cerebral CTX	[241]
A β_{1-42} i.c.v. to mice (C57BL/6)	↑MDA, ↑PC, ↑Mn-SOD, ↑Zn, Cu-SOD, ↑GPx, ↑GR in cerebral CTX and HIP	Vitamin E 150 mg/kg, p.o. for 27 days, administration began 7 days before A β_{1-42} i.c.v.	↓MDA, ↓PC, ↓Mn-SOD, ↓Zn, Cu-SOD, ↑GPx, Ø GR in cerebral CTX and HIP	[247]
AF64A (a cholinotoxin) 2 nmol/2 μ l, bilaterally i.c.v. to rats (Wistar)	↑MDA in HIP	Piperine 5 or 10 mg/kg p.o. 2 weeks before and 1 week after AF64A	↓MDA in HIP	[239]
Streptozotocin 2.57 mg/kg i.c.v. to mice (Swiss albino)	↑TBARS, ↓GSH, ↓GPx, ↓GR in HIP	S-allyl cysteine 30 mg/kg i.p. for 15 days pre-treatment before streptozocin	↓TBARS, ↑GSH, ↑GPx, ↑GR in HIP	[250]
Scopolamine 1 mg/kg i.p. single injection to mice (Swiss) 20 min before pretest (memory acquisition) or 15 min after pretest (memory consolidation)	↑MDA, ↓SOD, ↓GPx, ↓GR in CTX and HIP	Imperatorin 1, 5, or 10 mg/kg i.p. 2 \times /day for 7 days (on 7th day (pretest): 10 min before scopolamine injection or 15 min after scopolamine injection)	↓MDA, ↑SOD in CTX and HIP, ↑GPx in CTX and HIP, ↑GR in CTX	[248]
Mice (AbPP Tg2576)	↑HNE, ↑3-NT in brain homogenates	α -lipoic acid 30 mg/kg/day enriched diet for 10 months	↓HNE, Ø 3-NT in brain homogenates	[251]
Mice (APP/PSEN1 transgenic)	↑MDA in HIP	Vitamin C 125 mg/kg i.p. for 12 days	Ø MDA in HIP	[252]
Mice (Gulo $^{-/-}$ APP/PSEN1)	MDA level in CTX on vitamin C content standard diet (0.33 g/L of drinking water) not changed	Vitamin C- low diet content 0.099 g/L of drinking water	↑MDA in CTX	[257]
Mice (APPswe/PS1)	nd	Melatonin 5 mg/kg p.o. for 5.5 months	↓MDA, ↓PC in HIP	[260]
Mice (heterozygote APP)	↑TBARS, ↓GSH, ↓SOD in the brain homogenate	Melatonin 10 mg/kg/day for 4 months intragastrically	↓TBARS, ↑GSH, ↑SOD in the brain homogenate	[254]
Mice (APPswe/PS1)	nd	Long-lasting incretin hormone analogue D-Ala ² GIP 35 days at 25 nmol/kg i.p. once daily	↓8-OHG in CA1 HIP region (in 12 and 19 months old mice)	[258]
Mice (APPswe/PS1)	↑H ₂ O ₂ , ↑MDA, ↓GSH, ↓TAC in whole brain lysate	Hesperidin 100 mg/kg per day for 16 weeks in chow	↓H ₂ O ₂ , ↓MDA, ↑GSH, ↑TAC in whole brain lysate	[259]

↑ increase, ↓ decrease, Ø no changes, 3-NT 3-nitrotyrosine, CAT catalase, CTX cortex, D-Ala²GIP glucose-dependent insulinotropic polypeptide, GPx glutathione peroxidase, GR glutathione reductase, GSH glutathione, GSSG oxidized glutathione, HIP hippocampus, HNE 4-hydroxynonenal, i.c.v. intracerebroventricular, i.p. intraperitoneal, MDA malonyldialdehyde, Mn-SOD manganese superoxide dismutase (located in mitochondria), PC protein carbonyl, SOD superoxide dismutase, TAC total antioxidant capacity, TBARS thiobarbituric acid reactive substances, Zn, Cu-SOD copper/zinc superoxide dismutase (located in cytoplasm)

prevented ROS damage in the hippocampus, and augmented endogenous antioxidant enzymes in a streptozocin AD model [250]. Similar results were obtained when melatonin was given chronically to a genetic AD mouse model, as the drug alleviated OS and enhanced GSH levels [254, 260]. Moreover, results from Feng et al. [254] showed that OS is an early event in AD pathogenesis and that antioxidant therapies may be beneficial if given at this stage of the disease [254].

As shown above, results from animal AD models that have used various pharmacological compounds to reduce OS and to alleviate memory deficits in AD are promising but do not yet parallel the results obtained in clinical trials.

Anti-Alzheimer's Disease Drugs that Modulate Oxidative Balance

Medications used to treat Alzheimer's disease have been shown to have antioxidant properties that depend on the dose and AD model (Table 12). For example, tacrine, the first anticholinesterase inhibitor approved by the Food and Drug Administration (FDA), was shown to suppress OS in an animal AD model [261]. In fact, tacrine (50–800 μ g/kg i.m.) increased the FRAP value, which serves as a measure of "antioxidant power" [262], without elevating any marker of OS-associated damage in brain tissue. The effect of tacrine may therefore be considered to be positive when this drug is

Table 12 Trials with anti-Alzheimer drugs in different AD animal models and their influence on oxidative damage and anti-oxidative defense biomarkers

Model	Oxidative defense biomarkers	Drug and route of administration	Changes in oxidative defense biomarkers	Reference
Scopolamine 2 mg/kg i.p. once per day for 2 weeks to mice (Kun Ming)	↑MDA, ↓SOD, ↓GSH in HIP	Donepezil (3 mg/kg/day p.o. once per day for 2 weeks)	↓MDA, ↓SOD, ↑GSH in HIP	[238]
Aβ ₁₋₄₂ 3 μl of 1 mg/ml solution. i.c.v. to mice (Chinese Kun Ming)	↑MDA, ↓SOD, ↓GPx, ↓GSH in HIP and cerebral CTX	Donepezil (0.01 mg/kg/day i.c.v. for 14 days)	↓MDA, ↑GSH, ↑GPx, Ø SOD in HIP and cerebral CTX	[241]
Mice APPswe/PS1 (transgenic model)	↑H ₂ O ₂ , ↑MDA, ↓GSH, ↓TAC	Donepezil (2.5 mg/kg/day for 16 weeks)	↓H ₂ O ₂ , ↓MDA, ↓GSH, Ø TAC	[259]
Scopolamine 1.4 mg/kg i.p. once a day for 9 days to mice (Swiss)	↑MDA, ↓CAT in whole brain lysate	Donepezil (5 mg/kg p.o. once a day for 9 days before scopolamine administration)	↓MDA, ↑CAT in whole brain lysate	[249]
Streptozotocin 0.5 mg/kg i.c. on 1st and 3rd day to mice (Swiss albino)	↑MDA, ↓GSH in whole brain lysate	Tacrine (5 mg/kg/day p.o.) or donepezil (5 mg/kg/day p.o.) for 7 days	↓MDA, ↑GSH (not significant) in whole brain lysate	[261]
Colchicine 5 μg/5 μl i.c.v. injection to rats (Wistar)	↑MDA, ↓GSH in brain	Rivastigmine (2.5 mg/kg p.o. for 28 days started 7 days before colchicine injection)	Ø MDA, Ø GSH in brain	[264]
Kainic acid (KA) 0.4 μg/2 μl single unilateral intrahippocampal injection to rats (Wistar)	↑MDA, ↑nitrate, ↓GSH, ↑GSSG in HIP	Galantamine (2.5 mg or 5 mg/kg for 14 days starting from the day of KA injection)	↑MDA, ↑nitrate, ↓GSH, ↑GSSG in HIP	[266]
Aβ ₂₅₋₃₅ i.c.v. injection once a day for 3 consecutive days to rats (Wistar)	↑iNOS in HIP	Memantine (5 mg/kg/day i.p. for 8 days, 3 times after Aβ ₂₅₋₃₅ injection and for 5 following post-Aβ ₂₅₋₃₅ days)	↓iNOS in HIP	[268]
Age-induced memory impairment in rats (Wistar) (24 months old)	↑PC in HIP and CTX	Memantine (20 mg/kg i.p. for 21 days)	↓PC in HIP and CTX	[267]
Streptozocin 10 μl injection of 3 mg/kg bilaterally on 1st and 3rd day to rats (Sprague-Dawley)	↑ROS, ↑nitrate in CTX and HIP	Memantine (10 mg/kg p.o. for 13 days starting from STZ injection)	↓ROS, ↓nitrate in CTX and HIP	[269]
Kainic acid 200 ng/10 μl bilaterally i.c.v. to rats (Sprague-Dawley)	↑ROS, ↑MDA, in STR, CTX, cerebellum, HIP, ↑nitrite in CTX, HIP	Memantine (10 mg/kg p.o. for 13 days starting from the day of OKA injection)	↓ROS, ↓MDA in STR, CTX, cerebellum, HIP, ↓nitrite in CTX, HIP	[270]

↑ increase, ↓ decrease, Ø no changes, *nd* not determined, 3-NT 3-nitrotyrosine, *AlCl*₃ aluminum chloride, *CAT* catalase, *CTX* cortex, *GPx* glutathione peroxidase, *GR* glutathione reductase, *GSH* glutathione, *GSSG* oxidized glutathione, *HIP* hippocampus, *HNE* 4-hydroxynonenal, *i.c.* intracerebral, *i.c.v.* intracerebroventricular, *i.p.* intraperitoneal, *KA* kainic acid, *MDA* malonyldialdehyde, *PC* protein carbonyl, *SOD* superoxide dismutase, *STR* striatum, *TAC* total antioxidant capacity, *TBARS* thiobarbituric acid reactive substances

used in doses that stimulate the antioxidant system without inducing oxidative damage in brain tissue [263].

Donepezil is another cholinesterase inhibitor used in AD patients that, when given in doses as low as 3 mg/kg [259], 5 mg/kg [238, 249], or even 0.01 mg/kg [241] in a mouse AD model, both increased antioxidant power (CAT, SOD, GSH, or GPx, depending on the dose; see Table 12) and diminished lipid peroxidation [238, 241, 249]. However, donepezil, when given in a similar dose of 2.5 mg/kg, failed to combat OS biomarkers and to stimulate antioxidant defenses in the APPswe/PS1 transgenic mouse AD model [259]. Those contradictory results come from studies using non-transgenic and transgenic animal AD models, which means that the multiple adaptations developed for use in these transgenic animals could be the reason for the observed difference in outcomes.

Another medication used in AD treatment is rivastigmine. This drug neither attenuated lipid peroxidation nor restored GSH depletion in the brains of rats in an AD model [264], although an older study indicated antioxidant properties for rivastigmine when AD was induced in rats by aluminum chloride administration [265]. Such differences in the effects of rivastigmine might be caused either by differences in the AD model used in the study (aluminum chlorate p.o. vs. colchicine i.c.v. models) or by differences in the rivastigmine dose regimen (0.3 mg/kg for 3 months vs. 2.5 mg/kg p.o. for 28 days). Based on the above scant reports, it is too soon to either confirm or exclude rivastigmine as an effective OS scavenger in AD.

A single report showed the ability of another AChE inhibitor, galantamine, to reduce OS. In a cognitive impairment

animal model, galantamine decreased lipid peroxidation, nitrate, and GSSG levels, enhanced SOD activity, and impaired GSH levels following kainic acid intrahippocampal injection, and it restored cognitive deficits as well [266].

Memantine has also been widely studied in preclinical AD models. For example, it was shown that memantine reduced oxidative damage to proteins in the cortex and hippocampus but not in the striatum, resulting in the reversal of concomitant age-induced recognition memory deficits in aged rats [267]. Other studies found that memantine diminished the level of inducible forms of NOS in an A β _{25–35} AD model [268] and ROS and nitrate levels in the hippocampus and cortex in a streptozotocin AD model [269] and in a kainic acid-induced model of dementia [270]. However, memantine was shown to have neuroprotective properties not only in AD models but also in 3-nitropropionic acid [271], rotenone [272], and diisopropylphosphorofluoridate (DFP) toxicity models [273].

There is a wide range of evidence showing that several drugs used to treat AD have antioxidant properties, suggesting that at least part of their efficacy in animal models may come from that action.

Summary and Conclusions

In general, the presence of OS in the pathophysiology of many neurodegenerative disorders, including ALS, PD, and AD, is a well-recognized phenomenon. The results of many in vitro and in vivo preclinical and clinical studies have consistently demonstrated that OS is one of the crucial players in the degeneration that occurs in the nervous system. The imbalance between OS and antioxidant defense systems seems to be a universal condition in neurodegeneration. However, what can be surprising is that the results of many studies often provide different results when trying to determine the exact mechanisms that underlie OS and to determine which of the markers of OS could be clinically useful. What has been shown to be elevated in one study does not necessarily have to rise in another. In preclinical studies, these divergent results could be explained by the use of different models, different species, or different methodologies. As for the clinical setting, it must be stressed that the number of patients available for study is usually small because they are in different stages of their diseases, there are often coexisting comorbidities, and, last but not least, they often take many other medications with different pro- or antioxidant properties. The analysis of potential biomarkers under these conditions is extremely difficult. Therefore, assessing the real efficacy of potential antioxidant drugs is a challenge. However, there are some data, if even modest, that some of the existing drugs possess anti-oxidant properties and that they could slow down neurodegenerative processes and improve our understanding of the significance of OS in the pathobiology of these untreatable conditions.

The results of clinical and preclinical studies have demonstrated the presence of elevated levels of OS biomarkers as well as impairments to antioxidant defenses in the brain and peripheral tissues in PD, AD, and ALS. As the currently available therapies for these neurodegenerative diseases are not sufficiently effective for treating disease symptoms, novel substances are searched for. Among these, drugs with antioxidant activity, which are widely studied as a possible anti-neurodegenerative PD, AD, or ALS agents, can efficiently normalize biomarkers of the oxidant/antioxidant balance in animal models. Most such drugs have so far failed to slow down the progression of the disease or to prolong the lives of patients. Some exceptions within these anti-neurodegenerative drugs exist, and they give hope and inspire further research.

Acknowledgments This work has been supported by the statutory funds of the Jagiellonian University (K/ZDS/004127) and by the grant from the National Science Centre no. UMO-2011/01/B/NZ4/00708.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Li J, O W, Li W, Jiang Z, Ghanbari HA (2013) Oxidative stress and neurodegenerative disorders. *Int J Mol Sci* 14:24438–24475
- Falkowski PG, Godfrey LV (2008) Electrons, life and the evolution of earth's oxygen cycle. *Philosophical Transactions of the Royal Society of London Ser B, Biol Sci* 363:2705–2716
- Paiva CN, Bozza MT (2014) Are reactive oxygen species always detrimental to pathogens? *Antioxid Redox Signal* 20:1000–1037
- Popa-Wagner A, Mitran S, Sivanesan S, Chang E, Buga A (2013) ROS and brain diseases: the good, the bad, and the ugly. *Oxidative Med Cell Longevity* 2013
- Metodiewa D, KoÅvska C (2000) Reactive oxygen species and reactive nitrogen species: relevance to cyto(neuro)toxic events and neurologic disorders. An overview. *Neurotox Res* 1:197–233
- Venditti P, Di Stefano L, Di Meo S (2013) Mitochondrial metabolism of reactive oxygen species. *Mitochondrion* 13:71–82
- Li X, Fang P, Mai J, Choi ET, Wang H, Yang X (2013) Targeting mitochondrial reactive oxygen species as novel therapy for inflammatory diseases and cancers. *J Hematol Oncol* 6
- Mailloux RJ, McBride SL, Harper M (2013) Unearthing the secrets of mitochondrial ROS and glutathione in bioenergetics. *Trends Biochem Sci* 38:592–602
- Cordeiro RM (2014) Reactive oxygen species at phospholipid bilayers: distribution, mobility and permeation. *Biochim Biophys Acta* 1838:438–444
- Blough NV, Zafiriou OC (1985) Reaction of superoxide with nitric oxide to form peroxonitrite in alkaline aqueous solution. *Inorg Chem* 24:3502–3504
- Forstermann U, Sessa WC (2012) Nitric oxide synthases: regulation and function. *Eur Heart J* 33:829–837

12. Bedard K, Krause K (2007) The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 87:245–313
13. Fritz KS, Petersen DR (2013) An overview of the chemistry and biology of reactive aldehydes. *Free Radic Biol Med* 59:85–91
14. Albarracin SL, Stab B, Casas Z, Sutachan JJ, Samudio I, Gonzalez J, Gonzalo L, Capani F, Morales L, Barreto GE (2012) Effects of natural antioxidants in neurodegenerative disease. *Nutr Neurosci* 15:1–9
15. Pinchuk I, Shoval H, Dotan Y, Lichtenberg D (2012) Evaluation of antioxidants: scope, limitations and relevance of assays. *Chem Phys Lipids* 165:638–647
16. Miller E, Bieta, Morel A, Saso L, Saluk J (2014) Isoprostanes and neuroprostanes as biomarkers of oxidative stress in neurodegenerative diseases. *Oxidat Med Cell Longevity* 2014
17. Ho E, Karimi Galoughi K, Liu C, Bhindi R, Figtree GA (2013) Biological markers of oxidative stress: applications to cardiovascular research and practice. *Redox Biol* 1:483–491
18. Zitka O, Krizkova S, Skalickova S, Kopel P, Babula P, Adam V, Kizek R (2013) Electrochemical study of DNA damaged by oxidation stress. *Comb Chem High Throughput Screen* 16:130–141
19. Li J, Liu D, Sun L, Lu Y, Zhang Z (2012) Advanced glycation end products and neurodegenerative diseases: mechanisms and perspective. *J Neurol Sci* 317:1–5
20. Yavuz BB, Yavuz B, Halil M, Cankurtaran M, Ulger Z, Cankurtaran ES, Aytemir K, Ariogul S (2008) Serum elevated gamma glutamyltransferase levels may be a marker for oxidative stress in Alzheimer's disease. *Int Psychogeriatr* 20:815–823
21. Kadiiska MB, Basu S, Brot N, Cooper C, Saari Csallany A, Davies MJ, George MM, Murray DM, Jackson Roberts L III, Shigenaga MK, Sohal RS, Stocker R, Van Thiel DH, Wiswedel I, Hatch GE, Mason RP (2013) Biomarkers of oxidative stress study V: ozone exposure of rats and its effect on lipids, proteins, and DNA in plasma and urine. *Free Radic Biol Med* 61:408–415
22. Gomez-Mejiba SE, Zhai Z, Della-Vedova MC, Muñoz MD, Chatterjee S, Towner RA, Hensley K, Floyd RA, Mason RP, Ramirez DC (2014) Immuno-spin trapping from biochemistry to medicine: advances, challenges, and pitfalls. focus on protein-centered radicals. *Biochim Biophys Acta - Gen Subj* 1840:722–729
23. Cozzolino M, Carri MT (2012) Mitochondrial dysfunction in ALS. *Prog Neurobiol* 97:54–66
24. Chen S, Sayana P, Zhang X, Le W (2013) Genetics of amyotrophic lateral sclerosis: an update. *Molec Neurodegen* 8
25. Weiduschat N, Mao X, Hupf J, Armstrong N, Kang G, Lange DJ, Mitsumoto H, Shungu DC (2014) Motor cortex glutathione deficit in ALS measured in vivo with the J-editing technique. *Neurosci Lett* 570:102–107
26. Parakh S, Spencer DM, Halloran MA, Soo KY, Atkin JD (2013) Redox regulation in amyotrophic lateral sclerosis. *Oxidat Med Cell Longevity*
27. Blokhuis AM, Groen EJM, Koppers M, Van Den Berg LH, Pasterkamp RJ (2013) Protein aggregation in amyotrophic lateral sclerosis. *Acta Neuropathol* 125:777–794
28. Schymick JC, Talbot K, Traynor BJ (2007) Genetics of sporadic amyotrophic lateral sclerosis. *Hum Mol Genet* 16:R233–R242
29. Corcia P, Blasco H, Camu W (2014) Genetics of amyotrophic lateral sclerosis. *Presse Med* 43:555–562
30. Mendez EF, Sattler R (2015) Biomarker development for C9orf72 repeat expansion in ALS. *Brain Res* 1607:26–35
31. Beckman G, Lundgren E, Tamvik A (1973) Superoxide dismutase isozymes in different human tissues, their genetic control and intracellular localization. *Hum Hered* 23:338–345
32. Saccon RA, Bunton-Stasyshyn RKA, Fisher EMC, Fratta P (2013) Is SOD1 loss of function involved in amyotrophic lateral sclerosis? *Brain* 136:2342–2358
33. Pasinelli P, Belford ME, Lennon N, Bacskai BJ, Hyman BT, Trotti D, Brown RH Jr (2004) Amyotrophic lateral sclerosis-associated SOD1 mutant proteins bind and aggregate with Bcl-2 in spinal cord mitochondria. *Neuron* 43:19–30
34. Rakhit R, Crow JP, Lepock JR, Kondejewski LH, Cashman NR, Chakrabarty A (2004) Monomeric Cu, Zn-superoxide dismutase is a common misfolding intermediate in the oxidation models of sporadic and familial amyotrophic lateral sclerosis. *J Biol Chem* 279:15499–15504
35. Ezzi SA, Urushitani M, Julien J (2007) Wild-type superoxide dismutase acquires binding and toxic properties of ALS-linked mutant forms through oxidation. *J Neurochem* 102:170–178
36. Ito H, Wate R, Zhang J, Ohnishi S, Kaneko S, Ito H, Nakano S, Kusaka H (2008) Treatment with edaravone, initiated at symptom onset, slows motor decline and decreases SOD1 deposition in ALS mice. *Exp Neurol* 213:448–455
37. Watanabe M, Dykes-Hoberg M, Cizewski Culotta V, Price DL, Wong PC, Rothstein JD (2001) Histological evidence of protein aggregation in mutant SOD1 transgenic mice and in amyotrophic lateral sclerosis neural tissues. *Neurobiol Dis* 8:933–941
38. Forsberg K, Jonsson PA, Andersen PM, Bergemalm D, Graffmo KS, Hultdin M, Jacobsson J, Rosquist R, Marklund SL, Brännström T (2010) Novel antibodies reveal inclusions containing non-native SOD1 in sporadic ALS patients. *PLoS ONE* 5
39. Tohgi H, Abe T, Yamizaki K, Murata T, Ishizaki E, Isobe C (1999) Remarkable increase in cerebrospinal fluid 3-nitrotyrosine in patients with sporadic amyotrophic lateral sclerosis. *Ann Neurol* 46:129–131
40. Smith RG, Henry YK, Mattson MP, Appel SH (1998) Presence of 4-hydroxynonenal in cerebrospinal fluid of patients with sporadic amyotrophic lateral sclerosis. *Ann Neurol* 44:696–699
41. Bogdanov M, Brown RH Jr, Matson W, Smart R, Hayden D, O'Donnell H, Flint Beal M, Cudkowicz M (2000) Increased oxidative damage to DNA in ALS patients. *Free Radic Biol Med* 29:652–658
42. Ihara Y, Nobukuni K, Takata H, Hayabara T (2005) Oxidative stress and metal content in blood and cerebrospinal fluid of amyotrophic lateral sclerosis patients with and without a Cu, Zn-superoxide dismutase mutation. *Neurol Res* 27:105–108
43. Barber SC, Shaw PJ (2010) Oxidative stress in ALS: key role in motor neuron injury and therapeutic target. *Free Radic Biol Med* 48:629–641
44. Oteiza PI, Uchitel OD, Carrasquedo F, Dubrovski AL, Roma JC, Fraga CG (1997) Evaluation of antioxidants, protein, and lipid oxidation products in blood from sporadic amyotrophic lateral sclerosis patients. *Neurochem Res* 22:535–539
45. Babu GN, Kumar A, Chandra R, Puri SK, Singh RL, Kalita J, Misra UK (2008) Oxidant-antioxidant imbalance in the erythrocytes of sporadic amyotrophic lateral sclerosis patients correlates with the progression of disease. *Neurochem Int* 52:1284–1289
46. LoGerfo A, Chico L, Borgia L, et al. (2014) Lack of association between nuclear factor erythroid-derived 2-like 2 promoter gene polymorphisms and oxidative stress biomarkers in amyotrophic lateral sclerosis patients. *Oxidat Med Cell Longevity*, vol. 2014, Article ID 432626, 9 pages, 2014. doi:10.1155/2014/432626
47. Mitsumoto H, Santella R, Liu X, Bogdanov M, Zipprich J, Wu H, Mahata J, Kilty M, Bednarz K, Bell D, Gordon PH, Hornig M, Mehrazin M, Naini A, Flint Beal M, Factor-Litvak P (2008) Oxidative stress biomarkers in sporadic ALS. *Amyotrophic Lat Scler* 9:177–183
48. Ikawa M, Okazawa H, Tsujikawa T, Muramatsu T, Kishitani T, Kamisawa T, Matsunaga A, Yamamura O, Mori T, Hamano T, Kiyono Y, Nakamoto Y, Yoneda M (2014) Increased cerebral oxidative stress in amyotrophic lateral sclerosis: a 62CU-ATSM pet study. *Neurology* 82

49. Tohgi H, Abe T, Yamazaki K, Murata T, Ishizaki E, Isobe C (1999) Increase in oxidized NO products and reduction in oxidized glutathione in cerebrospinal fluid from patients with sporadic form of amyotrophic lateral sclerosis. *Neurosci Lett* 260:204–206
50. Boll M, Alcaraz-Zubeldia M, Montes S, Murillo-Bonilla L, Rios C (2003) Raised nitrate concentration and low SOD activity in the CSF of sporadic ALS patients. *Neurochem Res* 28:699–703
51. Nikolic-Kokic A, Stevic Z, Blagojevic D, Davidovic B, Jones DR, Spasic MB (2006) Alterations in anti-oxidative defence enzymes in erythrocytes from sporadic amyotrophic lateral sclerosis (SALS) and familial ALS patients. *Clin Chem Lab Med* 44:589–593
52. Apostolski S, Marinkovic Z, Nikolic A, Blagojevic D, Spasic MB, Michael Michelson A (1998) Glutathione peroxidase in amyotrophic lateral sclerosis: the effects of selenium supplementation. *J Environ Pathol Toxicol Oncol* 17:325–329
53. Kocic AN, Stevic Z, Stojanovic S, Blagojevic DP, Jones DR, Pavlovic S, Niketic V, Apostolski S, Spasic MB (2005) Biotransformation of nitric oxide in the cerebrospinal fluid of amyotrophic lateral sclerosis patients. *Redox Rep* 10:265–270
54. Süßmuth SD, Brettschneider J, Ludolph AC, Tumani H (2008) Biochemical markers in CSF of ALS patients. *Curr Med Chem* 15:1788–1801
55. Tanaka H, Shimazawa M, Takata M, Kaneko H, Tsuruma K, Ikeda T, Warita H, Aoki M, Yamada M, Takahashi H, Hozumi I, Minatsu H, Inuzuka T, Hara H (2013) ITIH4 and Gpx3 are potential biomarkers for amyotrophic lateral sclerosis. *J Neurol* 260:1782–1797
56. Kuzma M, Jamrozik Z, Baranczyk-Kuzma A (2006) Activity and expression of glutathione S-transferase pi in patients with amyotrophic lateral sclerosis. *Clin Chim Acta* 364:217–221
57. Cova E, Bongioanni P, Cereda C, Metelli MR, Salvaneschi L, Bernuzzi S, Guareschi S, Rossi B, Ceroni M (2010) Time course of oxidant markers and antioxidant defenses in subgroups of amyotrophic lateral sclerosis patients. *Neurochem Int* 56:687–693
58. De Bustos F, Jiménez-Jiménez FJ, Molina JA, Esteban J, Guerrero-Sola A, Zurdo M, Orti-Pareja M, Tallón-Barranco A, Gómez-Escalonilla C, Ramírez-Ramos C, Arenas J, Enriquez De Salamanca R (1998) Cerebrospinal fluid levels of alpha-tocopherol in amyotrophic lateral sclerosis. *J Neural Transm* 105:703–708
59. Paraskevas GP, Kapaki E, Libitaki G, Zournas C, Segditsa I, Papageorgiou C (1997) Ascorbate in healthy subjects, amyotrophic lateral sclerosis and Alzheimer's disease. *Acta Neurol Scand* 96:88–90
60. Johnson WM, Wilson-Delfosse AL, Mieryl JJ (2012) Dysregulation of glutathione homeostasis in neurodegenerative diseases. *Nutrients* 4:1399–1440
61. Desnuelle C, Dib M, Garrel C, Favier A (2001) A double-blind, placebo-controlled randomized clinical trial of alpha-tocopherol (vitamin E) in the treatment of amyotrophic lateral sclerosis. *Amyotroph Lateral Sclerosis Motor Neuron Disord* 2:9–18
62. Graf M, Ecker D, Horowski R, Kramer B, Riederer P, Gerlach M, Hager C, Ludolph AC (2005) High dose vitamin E therapy in amyotrophic lateral sclerosis as add-on therapy to riluzole: results of a placebo-controlled double-blind study. *J Neural Transm* 112:649–660
63. Veldink JH, Kalmijn S, Groeneveld G, Wunderink W, Koster A, De Vries JHM, Van Der Luyt J, Wokke JHJ, Van Den Berg LH (2007) Intake of polyunsaturated fatty acids and vitamin E reduces the risk of developing amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* 78:367–371
64. Yoshino H, Kimura A (2006) Investigation of the therapeutic effects of edaravone, a free radical scavenger, on amyotrophic lateral sclerosis (phase II study). *Amyotroph Later Sclerosis: Off Public World Fed Neurol Res Group Motor Neuron Dis* 7:241–245
65. <https://clinicaltrials.gov/show/NCT01492686>. Accessed 15 May 2015
66. Kaufmann P, Thompson JLP, Levy G, Buchsbaum R, Shefner J, Krivickas LS, Katz J, Rollins Y, Barohn RJ, Jackson CE, Tiryaki E, Lomen-Hoerth C, Armon C, Tandan R, Rudnicki SA, Reznia K, Sufit R, Pestronk A, Novella SP, Heiman-Patterson T, Kasarskis EJ, Pioro EP, Montes J, Arbing R, Vecchio D, Barsdorf A, Mitsumoto H, Levin B (2009) Phase II trial of CoQ10 for ALS finds insufficient evidence to justify phase III. *Ann Neurol* 66:235–244
67. Groeneveld GJ, Veldink JH, Van der Tweel I, Kalmijn S, Beijer C, De Visser M, Wokke JHJ, Franssen H, Van den Berg LH (2003) A randomized sequential trial of creatine in amyotrophic lateral sclerosis. *Ann Neurol* 53:437–445
68. Louwese ES, Weverling GJ, Bossuyt PMM, Meyjes FEP, De Jong JMBV (1995) Randomized, double-blind, controlled trial of acetylcysteine in amyotrophic lateral sclerosis. *Arch Neurol* 52:559–564
69. Lange DJ, Murphy PL, Diamond B, Appel V, Lai EC, Younger DS, Appel SH (1998) Selegiline is ineffective in a collaborative double-blind, placebo-controlled trial for treatment of amyotrophic lateral sclerosis. *Arch Neurol* 55:93–96
70. Weishaupt JH, Bartels C, Pölking E, Dietrich J, Rohde G, Poeggeler B, Mertens N, Sperling S, Bohn M, Hüther G, Schneider A, Bach A, Sirén A, Hardeland R, Bähr M, Nave K, Ehrenreich H (2006) Reduced oxidative damage in ALS by high-dose enteral melatonin treatment. *J Pineal Res* 41:313–323
71. Zoccollella S, Santamato A, Lamberti P (2009) Current and emerging treatments for amyotrophic lateral sclerosis. *Neuropsychiat Dis Treatment* Vol 5, Issue 1, 2009, Pages 577–595 5:577–595
72. Turner BJ, Talbot K (2008) Transgenics, toxicity and therapeutics in rodent models of mutant SOD1-mediated familial ALS. *Prog Neurobiol* 85:94–134
73. McGoldrick P, Joyce PI, Fisher EMC, Greensmith L (2013) Rodent models of amyotrophic lateral sclerosis. *Biochim Biophys Acta (BBA) - Mol Basis Dis* 1832:1421–1436
74. Bogdanov MB, Ramos LE, Xu Z, Beal MF (1998) Elevated 'hydroxyl radical' generation in vivo in an animal model of amyotrophic lateral sclerosis. *J Neurochem* 71:1321–1324
75. Liu D, Wen J, Liu J, Li L (1999) The roles of free radicals in amyotrophic lateral sclerosis: reactive oxygen species and elevated oxidation of protein, DNA, and membrane phospholipids. *FASEB J* 13:2318–2328
76. Towner RA, Smith N, Saunders D, Lupu F, Silasi-Mansat R, West M, Ramirez DC, Gomez-Mejiba SE, Bonini MG, Mason RP, Ehrenshaft M, Hensley K (2013) In vivo detection of free radicals using molecular MRI and immuno-spin trapping in a mouse model for amyotrophic lateral sclerosis. *Free Radic Biol Med* 63:351–360
77. Poon HF, Hensley K, Thongboonkerd V, Merchant ML, Lynn BC, Pierce WM, Klein JB, Calabrese V, Butterfield DA (2005) Redox proteomics analysis of oxidatively modified proteins in G93A-SOD1 transgenic mice—a model of familial amyotrophic lateral sclerosis. *Free Radic Biol Med* 39:453–462
78. Liu D, Bao F, Wen J, Liu J (2007) Mutation of superoxide dismutase elevates reactive species: comparison of nitration and oxidation of proteins in different brain regions of transgenic mice with amyotrophic lateral sclerosis. *Neuroscience* 146:255–264
79. Miana-Mena FJ, González-Mingot C, Larrodé P, Muñoz MJ, Oliván S, Fuentes-Broto L, Martínez-Ballarín E, Reiter RJ, Osta R, García JJ (2011) Monitoring systemic oxidative stress in an animal model of amyotrophic lateral sclerosis. *J Neurol* 258:762–769

80. Morimoto N, Miyazaki K, Kurata T, Ikeda Y, Matsuura T, Kang D, Ide T, Abe K (2012) Effect of mitochondrial transcription factor a overexpression on motor neurons in amyotrophic lateral sclerosis model mice. *J Neurosci Res* 90:1200–1208
81. Seo J, Baek I, Leem Y, Kim T, Cho Y, Lee SM, Park YH, Han P (2011) SK-PC-B70M alleviates neurologic symptoms in G93A-SOD1 amyotrophic lateral sclerosis mice. *Brain Res* 1368:299–307
82. Casoni F, Basso M, Massignan T, Gianazzail E, Cheroni C, Salmons M, Bendotti C, Bonetto V (2005) Protein nitration in a mouse model of familial amyotrophic lateral sclerosis: possible multifunctional role in the pathogenesis. *J Biol Chem* 280:16295–16304
83. Nardo G, Pozzi S, Mantovani S, Garbelli S, Marinou K, Basso M, Mora G, Bendotti C, Bonetto V (2009) Nitroproteomics of peripheral blood mononuclear cells from patients and a rat model of ALS. *Antioxid Redox Signal* 11:1559–1567
84. Basso M, Samengo G, Nardo G, Massignan T, D'Alessandro G, Tartari S, Cantoni L, Marino M, Cheroni C, de Biasi S, Giordana MT, Strong MJ, Estevez AG, Salmons M, Bendotti C, Bonetto V (2009) Characterization of detergent-insoluble proteins in ALS indicates a causal link between nitrative stress and aggregation in pathogenesis. *PLoS ONE* 4
85. Cha CI, Chung YH, Shin C, Shin DH, Kim YS, Gurney ME, Lee KW (2000) Immunocytochemical study on the distribution of nitrotyrosine in the brain of the transgenic mice expressing a human Cu/Zn SOD mutation. *Brain Res* 853:156–161
86. Tokuda E, Ono S, Ishige K, Watanabe S, Okawa E, Ito Y, Suzuki T (2008) Ammonium tetrathiomolybdate delays onset, prolongs survival, and slows progression of disease in a mouse model for amyotrophic lateral sclerosis. *Exp Neurol* 213:122–128
87. Chi L, Ke Y, Luo C, Gozal D, Liu R (2007) Depletion of reduced glutathione enhances motor neuron degeneration in vitro and in vivo. *Neuroscience* 144:991–1003
88. Vargas MR, Johnson DA, Johnson JA (2011) Decreased glutathione accelerates neurological deficit and mitochondrial pathology in familial ALS-linked hSOD1G93A mice model. *Neurobiol Dis* 43:543–551
89. Vargas MR, Johnson DA, Sirkis DW, Messing A, Johnson JA (2008) Nrf2 activation in astrocytes protects against neurodegeneration in mouse models of familial amyotrophic lateral sclerosis. *J Neurosci* 28:13574–13581
90. Kato S, Kato M, Abe Y, Matsumura T, Nishino T, Aoki M, Itoyama Y, Asayama K, Awaya A, Hirano A, Ohama E (2005) Redox system expression in the motor neurons in amyotrophic lateral sclerosis (ALS): immunohistochemical studies on sporadic ALS, superoxide dismutase 1 (SOD1)-mutated familial ALS, and SOD1-mutated ALS animal models. *Acta Neuropathol* 110:101–112
91. Kato S, Saeki Y, Aoki M, Nagai M, Ishigaki A, Itoyama Y, Kato M, Asayama K, Awaya A, Hirano A, Ohama E (2004) Histological evidence of redox system breakdown caused by superoxide dismutase 1 (SOD1) aggregation is common to SOD1-mutated neurons in humans and animal models. *Acta Neuropathol* 107:149–158
92. Cudkovic ME, Pastuszka KA, Sapp PC, Mathews RK, Leahy J, Pasinelli P, Francis JW, Jiang D, Andersen JK, Brown RH Jr (2002) Survival in transgenic ALS mice does not vary with CNS glutathione peroxidase activity. *Neurology* 59:729–734
93. Lucchetti J, Marino M, Papa S, Tortarolo M, Guiso G, Pozzi S, Bonetto V, Caccia S, Beghi E, Bendotti C, Gobbi M (2013) A mouse model of familial ALS has increased CNS levels of endogenous Ubiquinol9/10 and does not benefit from exogenous administration of Ubiquinol10. *PLoS ONE* 8
94. Klivenyi P, Kiaei M, Gardian G, Calingasan NY, Beal MF (2004) Additive neuroprotective effects of creatine and cyclooxygenase 2 inhibitors in a transgenic mouse model of amyotrophic lateral sclerosis. *J Neurochem* 88:576–582
95. Peña-Altamira E, Crochemore C, Virgili M, Contestabile A (2005) Neurochemical correlates of differential neuroprotection by long-term dietary creatine supplementation. *Brain Res* 1058:183–188
96. Zhang W, Narayanan M, Friedlander RM (2003) Additive neuroprotective effects of minocycline with creatine in a mouse model of ALS. *Ann Neurol* 53:267–270
97. Matthews RT, Ferrante RJ, Klivenyi P, Yang L, Klein AM, Mueller G, Kaddurah-Daouk R, Beal MF (1999) Creatine and cyclocreatine attenuate MPTP neurotoxicity. *Exp Neurol* 157:142–149
98. Klivenyi P, Ferrante RJ, Matthews RT, Bogdanov MB, Klein AM, Andreassen OA, Mueller G, Wermer M, Kaddurah-Daouk R, Beal MF (1999) Neuroprotective effects of creatine in a transgenic animal model of amyotrophic lateral sclerosis. *Nat Med* 5:347–350
99. Choi J, Küstermann E, Dedeoglu A, Jenkins BG (2009) Magnetic resonance spectroscopy of regional brain metabolite markers in FALS mice and the effects of dietary creatine supplementation. *Eur J Neurosci* 30:2143–2150
100. Derave W, Van Den Bosch L, Lemmens G, Eijnde BO, Robberecht W, Hespel P (2003) Skeletal muscle properties in a transgenic mouse model for amyotrophic lateral sclerosis: effects of creatine treatment. *Neurobiol Dis* 13:264–272
101. Levkovitch-Verbin H, Waserzoog Y, Vander S, Makarovsky D, Piven I (2014) Minocycline upregulates pro-survival genes and downregulates pro-apoptotic genes in experimental glaucoma. *Graefes Arch Clin Exp Ophthalmol* 252:761–772
102. Jung C, Rong Y, Doctrow S, Baudry M, Malfroy B, Xu Z (2001) Synthetic superoxide dismutase/catalase mimetics reduce oxidative stress and prolong survival in a mouse amyotrophic lateral sclerosis model. *Neurosci Lett* 304:157–160
103. Chang Y, Kong Q, Shan X, Tian G, Ilieva H, Cleveland DW, Rothstein JD, Borchelt DR, Wong PC, Lin C-G (2008) Messenger RNA oxidation occurs early in disease pathogenesis and promotes motor neuron degeneration in ALS. *PLoS ONE* 3
104. Zhang Y, Cook A, Kim J, Baranov SV, Jiang J, Smith K, Cormier K, Bennett E, Browner RP, Day AL, Carlisle DL, Ferrante RJ, Wang X, Friedlander RM (2013) Melatonin inhibits the caspase-1/cytochrome c/caspase-3 cell death pathway, inhibits MT1 receptor loss and delays disease progression in a mouse model of amyotrophic lateral sclerosis. *Neurobiol Dis* 55:26–35
105. Tokuda E, Okawa E, Watanabe S, Ono S, Marklund SL (2013) Dysregulation of intracellular copper homeostasis is common to transgenic mice expressing human mutant superoxide dismutase-1s regardless of their copper-binding abilities. *Neurobiol Dis* 54:308–319
106. Song L, Chen L, Zhang X, Li J, Le W (2014) Resveratrol ameliorates motor neuron degeneration and improves survival in SOD1G93A mouse model of amyotrophic lateral sclerosis. *BioMed Res Int* 2014
107. Dardiotis E, Panayiotou E, Feldman ML, Hadjisavvas A, Malas S, Vonta I, Hadjigeorgiou G, Kyriakou K, Kyriakides T (2013) Intraperitoneal melatonin is not neuroprotective in the G93ASOD1 transgenic mouse model of familial ALS and may exacerbate neurodegeneration. *Neurosci Lett* 548:170–175
108. Crow JP, Calingasan NY, Chen J, Hill JL, Beal MF (2005) Manganese porphyrin given at symptom onset markedly extends survival of ALS mice. *Ann Neurol* 58:258–265
109. Gurney ME, Cutting FB, Zhai P, Doble A, Taylor CP, Andrus PK, Hall ED (1996) Benefit of vitamin E, riluzole, and gabapentin in a transgenic model of familial amyotrophic lateral sclerosis. *Ann Neurol* 39:147–157
110. Snow RJ, Turnbull J, Da Silva S, Jiang F, Tarnopolsky MA (2003) Creatine supplementation and riluzole treatment provide similar

- beneficial effects in copper, zinc superoxide dismutase (G93A) transgenic mice. *Neuroscience* 119:661–667
111. Koh J, Kim D, Hwang JY, Kim YH, Seo JH (1999) Antioxidative and proapoptotic effects of riluzole on cultured cortical neurons. *J Neurochem* 72:716–723
 112. <http://www.icd10data.com/ICD10CM/Codes/G00-G99/G20-G26/G20-G20>. Accessed 15 May 2015
 113. Gandhi SAA (2012) Mechanism of oxidative stress in neurodegeneration. *Oxid Med Cell Longev* 2012:428010
 114. Dexter DT, Carter CJ, Wells FR, Javoy-Agid F, Agid Y, Lees A, Jenner P, Marsden CD (1989) Basal lipid peroxidation in substantia nigra is increased in Parkinson's disease. *J Neurochem* 52:381–389
 115. Mythri RB, Venkateshappa C, Harish G, Mahadevan A, Muthane UB, Yasha TC, Srinivas Bharath MM, Shankar SK (2011) Evaluation of markers of oxidative stress, antioxidant function and astrocytic proliferation in the striatum and frontal cortex of Parkinson's disease brains. *Neurochem Res* 36:1452–1463
 116. Castellani RJ, Perry G, Siedlak SL, Nunomura A, Shimohama S, Zhang J, Montine T, Sayre LM, Smith MA (2002) Hydroxynonenal adducts indicate a role for lipid peroxidation in neocortical and brainstem Lewy bodies in humans. *Neurosci Lett* 319:25–28
 117. Shamoto-Nagai M, Maruyama W, Hashizume Y, Yoshida M, Osawa T, Riederer P, Naoi M (2007) In parkinsonian substantia nigra, α -synuclein is modified by acrolein, a lipid-peroxidation product, and accumulates in the dopamine neurons with inhibition of proteasome activity. *J Neural Transm* 114:1559–1567
 118. Yoritaka A, Hattori N, Uchida K, Tanaka M, Stadtman ER, Mizuno Y (1996) Immunohistochemical detection of 4-hydroxynonenal protein adducts in Parkinson disease. *Proc Natl Acad Sci U S A* 93:2696–2701
 119. Xiang W, Schlachetzki JCM, Helling S, Bussmann JC, Berlinghof M, Schäffer TE, Marcus K, Winkler J, Klucken J, Becker C (2013) Oxidative stress-induced posttranslational modifications of α -synuclein: specific modification of α -synuclein by 4-hydroxy-2-nonenal increases dopaminergic toxicity. *Mol Cell Neurosci* 54:71–83
 120. Fessel JP, Hulette C, Powell S, Roberts LJ II, Zhang J (2003) Isofurans, but not F2-isoprostanes, are increased in the substantia nigra of patients with Parkinson's disease and with dementia with Lewy body disease. *J Neurochem* 85:645–650
 121. Alam ZI, Daniel SE, Lees AJ, Marsden DC, Jenner P, Halliwell B (1997) A generalised increase in protein carbonyls in the brain in Parkinson's but not incidental Lewy body disease. *J Neurochem* 69:1326–1329
 122. Ogata M, Kaneya D, Shin-Ya K, Li L, Abe Y, Katoh H, Seki S, Seki Y, Gonda R, Urano S, Endo T (2005) Trapping effect of eugenol on hydroxyl radicals induced by L-DOPA in vitro. *Chem Pharm Bull* 53:1167–1170
 123. Good PF, Hsu A, Werner P, Perl DP, Warren Olanow C (1998) Protein nitration in Parkinson's disease. *J Neuropathol Exp Neurol* 57:338–342
 124. Zhang J, Perry G, Smith MA, Robertson D, Olson SJ, Graham DG, Montine TJ (1999) Parkinson's disease is associated with oxidative damage to cytoplasmic DNA and RNA in substantia nigra neurons. *Am J Pathol* 154:1423–1429
 125. Alam ZI, Jenner A, Daniel SE, Lees AJ, Cairns N, Marsden CD, Jenner P, Halliwell B (1997) Oxidative DNA damage in the Parkinsonian brain: an apparent selective increase in 8-hydroxyguanine levels in substantia nigra. *J Neurochem* 69:1196–1203
 126. Kikuchi A, Takeda A, Onodera H, Kimpara T, Hisanaga K, Sato N, Nunomura A, Castellani RJ, Perry G, Smith MA, Itoyama Y (2002) Systemic increase of oxidative nucleic acid damage in Parkinson's disease and multiple system atrophy. *Neurobiol Dis* 9:244–248
 127. Isobe C, Abe T, Terayama Y (2010) Levels of reduced and oxidized coenzymeQ-10 and 8-hydroxy-2'-deoxyguanosine in the cerebrospinal fluid of patients with living Parkinson's disease demonstrate that mitochondrial oxidative damage and/or oxidative DNA damage contributes to the neurodegenerative process. *Neurosci Lett* 469:159–163
 128. Boll M, Alcaraz-Zubeldia M, Montes S, Rios C (2008) Free copper, ferroxidase and SOD1 activities, lipid peroxidation and NOx content in the CSF. A different marker profile in four neurodegenerative diseases. *Neurochem Res* 33:1717–1723
 129. Selley ML (1998) (E)-4-Hydroxy-2-nonenal may be involved in the pathogenesis of Parkinson's disease. *Free Radic Biol Med* 25:169–174
 130. Ilic TV, Jovanovic M, Jovicic A, Tomovic M (1999) Oxidative stress indicators are elevated in de novo Parkinson's disease patients. *Funct Neurol* 14:141–147
 131. Shukla R, Rajani M, Srivastava N, Barthwal MK, Dikshit M (2006) Nitrite and malondialdehyde content in cerebrospinal fluid of patients with Parkinson's disease. *Int J Neurosci* 116:1391–1402
 132. Sanders LH, Timothy Greenamyre J (2013) Oxidative damage to macromolecules in human Parkinson disease and the rotenone model. *Free Radic Biol Med* 62:111–120
 133. Sato S, Mizuno Y, Hattori N (2005) Urinary 8-hydroxydeoxyguanosine levels as a biomarker for progression of Parkinson disease. *Neurology* 64:1081–1083
 134. Seet RCS, Lee CJ, Lim ECH, Tan JH, Quek AML, Chong W, Looi W, Huang S, Wang H, Chan Y, Halliwell B (2010) Oxidative damage in Parkinson disease: measurement using accurate biomarkers. *Free Radic Biol Med* 48:560–566
 135. Lee C-J, Seet RCS, Huang SH, Long LH, Halliwell B (2009) Different patterns of oxidized lipid products in plasma and urine of dengue fever, stroke, and Parkinson's disease patients: cautions in the use of biomarkers of oxidative stress. *Antioxid Redox Signal* 11:407–420
 136. Ahlskog JE, Uitti RJ, Low PA, Tyce GM, Nickander KK, Petersen RC, Kokmen E (1995) No evidence for systemic oxidant stress in Parkinson's or Alzheimer's disease. *Mov Disord* 10:566–573
 137. Molina J, Jiménez-Jiménez FJ, Fernández-Calle P, Lalinde L, Tenias J, Ponal M, Vazquez A, Codoceo R (1992) Serum lipid peroxides in patients with Parkinson's disease. *Neurosci Lett* 136:137–140
 138. Connolly J, Siderowf A, Clark CM, Mu D, Pratico D (2008) F2 isoprostane levels in plasma and urine do not support increased lipid peroxidation in cognitively impaired parkinson disease patients. *Cogn Behav Neurol* 21:83–86
 139. Sian J, Dexter DT, Lees AJ, Daniel S, Agid Y, Javoy-Agid F, Jenner P, Marsden CD (1994) Alterations in glutathione levels in Parkinson's disease and other neurodegenerative disorders affecting basal ganglia. *Ann Neurol* 36:348–355
 140. Perry TL, Yong VW (1986) Idiopathic Parkinson's disease, progressive supranuclear palsy and glutathione metabolism in the substantia nigra of patients. *Neurosci Lett* 67:269–274
 141. Perry TL, Godin DV, Hansen S (1982) Parkinson's disease: a disorder due to nigral glutathione deficiency? *Neurosci Lett* 33:305–310
 142. Danielson SR, Andersen JK (2008) Oxidative and nitrative protein modifications in Parkinson's disease. *Free Radic Biol Med* 44:1787–1794
 143. Riederer P, Sofic E, Rausch W, Schmidt B, Reynolds GP, Jellinger K, Youdim MBH (1989) Transition metals, ferritin, glutathione, and ascorbic acid in parkinsonian brains. *J Neurochem* 52:515–520

144. Chinta SJ, Andersen JK (2008) Redox imbalance in Parkinson's disease. *Biochim Biophys Acta Gen Subj* 1780:1362–1367
145. Marttila RJ, Lorentz H, Rinne UK (1988) Oxygen toxicity protecting enzymes in Parkinson's disease: increase of superoxide dismutase-like activity in the substantia nigra and basal nucleus. *J Neurol Sci* 86:321–331
146. Kish SJ, Morito C, Hornykiewicz O (1985) Glutathione peroxidase activity in Parkinson's disease brain. *Neurosci Lett* 58:343–346
147. Mbangata MB, Kartha RV, Mishra U, Coles LD, Tuite PJ, Cloyd JC (2014) Oxidative stress status in patients with Parkinson's disease on and off medication. *Mov Disord* 29:S143
148. Sechi G, Deledda MG, Bua G, Satta WM, Deiana GA, Pes GM, Rosati G (1996) Reduced intravenous glutathione in the treatment of early Parkinson's disease. *Prog Neuro-Psychopharmacol Biol Psychiatry* 20:1159–1170
149. Holmay MJ, Terpstra M, Coles LD, Mishra U, Ahlskog M, Öz G, Cloyd JC, Tuite PJ (2013) N-acetylcysteine boosts brain and blood glutathione in gaucher and Parkinson diseases. *Clin Neuropharmacol* 36:103–106
150. Katz M, Swanson RA, Glass GA (2014) Cerebrospinal fluid concentrations of N-acetylcysteine after oral administration: phase I trial in Parkinson's disease. *Mov Disord* 29:S247
151. Gorell JM, Ordidge RJ, Brown GG, Deniau J, Buderer NM, Helpert JA (1995) Increased iron-related MRI contrast in the substantia nigra in Parkinson's disease. *Neurology* 45:1138–1143
152. Devos D, Moreau C, Devedjian JC, Kluza J, Petraut M, Laloux C, Jonneaux A, Ryckewaert G, Garçon G, Rouaix N, Duhamel A, Jissendi P, Dujardin K, Auger F, Ravasi L, Hopes L, Grolez G, Firdaus W, Sablonniere B, Strubi-Vuillaume I, Zahr N, Destée A, Corvol J, Pörtl D, Leist M, Rose C, Defebvre L, Marchetti P, Cabantchik ZI, Bordet R (2014) Targeting chelatable iron as a therapeutic modality in Parkinson's disease. *Antioxid Redox Signal* 21:195–210
153. Shoulson I (1998) DATATOP: a decade of neuroprotective inquiry. *Ann Neurol* 44:S160–S166
154. Shoulson I, Fahn S, Oakes D, Kieburtz K, Lang A, Langston JW, LeWitt P, Olanow CW, Penney JB, Tanner C, Rudolph A, Pelusio RM (1993) Effects of tocopherol and deprenyl on the progression of disability in early Parkinson's disease. *N Engl J Med* 328:176–183
155. Beal MF (2014) A randomized clinical trial of high-dosage coenzyme q10 in early parkinson disease no evidence of benefit. *JAMA Neurol* 71:543–552
156. Su W, Chen H, Zhang Z, Chen B, Wang L, Sun X, Shen Y, Li Y, Geng T, Zhao W, Zhang X (2004) A multi-center, randomized, vitamin E controlled and opening clinical trial of selegiline in patients with Parkinson's disease. *Chin J Neurol* 37:413–416
157. Shults CW, Oakes D, Kieburtz K, Flint Beal M, Haas R, Plumb S, Juncos JL, Nutt J, Shoulson I, Carter J, Kompoliti K, Perlmutter JS, Reich S, Stern M, Watts RL, Kurlan R, Molho E, Harrison M, Lew M (2002) Effects of coenzyme Q 10 in early Parkinson disease: evidence of slowing of the functional decline. *Arch Neurol* 59:1541–1550
158. Storch A, Jost WH, Vieregge P, Spiegel J, Greulich W, Durner J, Müller T, Kupsch A, Henningsen H, Oertel WH, Fuchs G, Kuhn W, Niklowitz P, Koch R, Herting B, Reichmann H (2007) Randomized, double-blind, placebo-controlled trial on symptomatic effects of coenzyme Q10 in Parkinson disease. *Arch Neurol* 64:938–944
159. NINDS (2006) A randomized, double-blind, futility clinical trial of creatine and minocycline in early Parkinson disease. *Neurology* 66:664–671
160. Kieburtz K, Tilley B, Ravina B, Galpern W, Shannon K, Tanner C, Wooten GF (2008) A pilot clinical trial of creatine and minocycline in early Parkinson disease: 18-month results. *Clin Neuropharmacol* 31:141–150
161. Bender A, Koch W, Elstner M, Schombacher Y, Bender J, Moeschl M, Gekeler F, Muller-Myhsok B, Gasser T, Tatsch K, Klopstock T (2006) Creatine supplementation in Parkinson disease: a placebo-controlled randomized pilot trial. *Neurology* 67:1262–1264
162. <http://parkinsontrial.ninds.nih.gov/netpd-LS1-study-termination.htm>. Accessed 15 May 2015
163. Hirayama M, Goto S, Nakamura T, Mizutani Y, Suzuki J, Watanabe H, Sobue G (2014) Oxidative stress marker (urinary 8-hydroxy-2'-deoxyguanosine) was modified by treatment of various Parkinson's drug. *Mov Disord* 29:S140
164. Sen CK (1999) Glutathione homeostasis in response to exercise training and nutritional supplements. *Mol Cell Biochem* 196:31–42
165. Elokda A, Di Francisco-Donoghue J, Lamberg EM, Werner WG (2010) Effects of exercise induced oxidative stress on glutathione levels in Parkinson's disease on and off medication. *J Neurol* 257:1648–1653
166. Nikolova GD, Grigorov BG, Zheleva AM, Gadjeva VG (2014) Influence of therapy on some important final products of oxidation of lipids, proteins and nucleic acids in patients with Parkinson's diseases. *Adv Biol Chem* 4:253–260
167. Blesa J, Phani S, Jackson-Lewis V, Przedborski S (2012) Classic and new animal models of Parkinson's disease. *J Biomed Biotechnol* 2012
168. Ungerstedt U (1971) Postsynaptic supersensitivity after 6-hydroxy-dopamine induced degeneration of the nigro-striatal dopamine system. *Acta Physiol Scandinav, Suppl* 367:69–93
169. Langston JW, Forno LS, Rebert CS, Irwin I (1984) Selective nigral toxicity after systemic administration of 1-methyl-4-phenyl-1,2,5,6-tetrahydropyridine (MPTP) in the squirrel monkey. *Brain Res* 292:390–394
170. Blesa J, Juri C, Collantes M, Peñuelas I, Prieto E, Iglesias E, Martí-Climent J, Arbizu J, Zubieta JL, Rodríguez-Oroz MC, García-García D, Richter JA, Cavada C, Obeso JA (2010) Progression of dopaminergic depletion in a model of MPTP-induced parkinsonism in non-human primates. An 18F-DOPA and 11C-DTBZ PET study. *Neurobiol Dis* 38:456–463
171. Ohashi S1, Mori A, Kurihara N, Mitsumoto Y, Nakai M (2006) Age-related severity of dopaminergic neurodegeneration to MPTP neurotoxicity causes motor dysfunction in C57BL/6 mice. *Neurosci Lett* 401(1–2):183–187 (2006 Jun 19;401(1–2):183–7. **Epub 2006 Apr 3.**)
172. Shimohama S, Hisahara S (2011) Toxin-induced and genetic animal models of Parkinson's disease. *Parkinson's Dis*
173. Betarbet R, Sherer TB, MacKenzie G, Garcia-Osuna M, Panov AV, Greenamyre JT (2000) Chronic systemic pesticide exposure reproduces features of Parkinson's disease. *Nat Neurosci* 3:1301–1306
174. McCormack AL, Atienza JG, Johnston LC, Andersen JK, Vu S, Di Monte DA (2005) Role of oxidative stress in paraquat-induced dopaminergic cell degeneration. *J Neurochem* 93:1030–1037
175. Kumar A, Leinisch F, Kadiiska MB, Corbett J, Mason RP (2015) Formation and implications of alpha-synuclein radical in maneb and paraquat-induced models of Parkinson's disease. *Mol Neurobiol*
176. Smith MP, Cass WA (2007) Oxidative stress and dopamine depletion in an intrastriatal 6-hydroxydopamine model of Parkinson's disease. *Neuroscience* 144:1057–1066
177. Haleagrahara N, Siew CJ, Ponnusamy K (2013) Effect of quercetin and desferrioxamine on 6-hydroxydopamine (6-OHDA) induced neurotoxicity in striatum of rats. *J Toxicol Sci* 38:25–33
178. Dagdelen M, Akkaya H, Genç E (2013) The role of valproic acid and levodopa on oxidative stress in a 6-hydroxydopamine

- lesioned rat model of Parkinson's disease. *Eur Neuropsychopharmacol* 23:S197
179. Inden M, Kitamura Y, Kondo J, Hayashi K, Yanagida T, Takata K, Tsuchiya D, Yanagisawa D, Nishimura K, Taniguchi T, Shimohama S, Sugimoto H, Akaike A (2005) Serofendic acid prevents 6-hydroxydopamine-induced nigral neurodegeneration and drug-induced rotational asymmetry in hemi-parkinsonian rats. *J Neurochem* 95:950–961
 180. Su-Fen Y, Zheng-Qin Y, Qin W, An-Sheng S, Xie-Nan H, Jing-Shan S (2001) Protective effect and mechanism of Ginkgo biloba leaf extracts for Parkinson disease induced by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *Acta Pharmacol Sin* 22:1089–1093
 181. Ghosh A, Kanthasamy A, Joseph J, Anantharam V, Srivastava P, Dranka BP, Kalyanaraman B, Kanthasamy AG (2012) Anti-inflammatory and neuroprotective effects of an orally active apocynin derivative in pre-clinical models of Parkinson's disease. *J Neuroinflammation* 9
 182. Bisht R, Kaur B, Gupta H, Prakash A (2014) Ceftriaxone mediated rescue of nigral oxidative damage and motor deficits in MPTP model of Parkinson's disease in rats. *Neurotoxicology* 44:71–79
 183. Barbiero JK, Santiago R, Tonin FS, Boschen S, da Silva LM, de Paula Werner MF, da Cunha C, Lima MMS, Vital MABF (2014) PPAR- α agonist fenofibrate protects against the damaging effects of MPTP in a rat model of Parkinson's disease. *Prog Neuro-Psychopharmacol Biol Psychiatry* 53:35–44
 184. Verma R, Nehru B (2009) Effect of centropheoxine against rotenone-induced oxidative stress in an animal model of Parkinson's disease. *Neurochem Int* 55:369–375
 185. Zaitone SA, Abo-Elmatty DM, Shaalan AA (2012) Acetyl-l-carnitine and (alpha)-lipoic acid affect rotenone-induced damage in nigral dopaminergic neurons of rat brain, implication for Parkinson's disease therapy. *Pharmacol Biochem Behav* 100:347–360
 186. Gokul K, Muralidhara (2014) Oral supplements of aqueous extract of tomato seeds alleviate motor abnormality, oxidative impairments and neurotoxicity induced by rotenone in mice: relevance to Parkinson's disease. *Neurochem Res* 39:1382–1394
 187. Liang L, Huang J, Fulton R, Day BJ, Patel M (2007) An orally active catalytic metalloporphyrin protects against 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine neurotoxicity in vivo. *J Neurosci* 27:4326–4333
 188. Chen L, Gao Y, Li X, Shen D, Sun F (2005) Melatonin protects against MPTP/MPP⁺-induced mitochondrial DNA oxidative damage in vivo and in vitro. *J Pineal Res* 39:34–42
 189. Fujita K, Seike T, Yutsudo N, Ohno M, Yamada H, Yamaguchi H, Sakumi K, Yamakawa Y, Kido MA, Takaki A, Katafuchi T, Tanaka Y, Nakabeppu Y, Noda M (2009) Hydrogen in drinking water reduces dopaminergic neuronal loss in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson's disease. *PLoS ONE* 4
 190. Singhal NK, Srivastava G, Patel DK, Jain SK, Singh MP (2011) Melatonin or silymarin reduces maneb- and paraquat-induced Parkinson's disease phenotype in the mouse. *J Pineal Res* 50:97–109
 191. Ahmad M, Saleem S, Ahmad AS, Yousuf S, Ansari MA, Khan MB, Ishrat T, Chaturvedi RK, Agrawal AK, Islam F (2005) Ginkgo biloba affords dose-dependent protection against 6-hydroxydopamine-induced parkinsonism in rats: neurobehavioural, neurochemical and immunohistochemical evidences. *J Neurochem* 93:94–104
 192. Thomas B, Mohanakumar KP (2004) Melatonin protects against oxidative stress caused by 1-methyl-4-phenyl-1, 2,3,6-tetrahydropyridine in the mouse nigrostriatum. *J Pineal Res* 36:25–32
 193. Zaminelli T, Gradowski RW, Bassani TB, Barbiero JK, Santiago RM, Maria-Ferreira D, Baggio CH, Vital MABF (2014) Antidepressant and antioxidative effect of ibuprofen in the rotenone model of Parkinson's disease. *Neurotoxicity Res*
 194. Karuppagounder SS, Madathil SK, Pandey M, Haobam R, Rajamma U, Mohanakumar KP (2013) Quercetin up-regulates mitochondrial complex-I activity to protect against programmed cell death in rotenone model of Parkinson's disease in rats. *Neuroscience* 236:136–148
 195. Kaur H, Chauhan S, Sandhir R (2011) Protective effect of lycopene on oxidative stress and cognitive decline in rotenone induced model of Parkinson's disease. *Neurochem Res* 36:1435–1443
 196. Khurana N, Gajbhiye A (2013) Ameliorative effect of *Sida cordifolia* in rotenone induced oxidative stress model of Parkinson's disease. *Neurotoxicology* 39:57–64
 197. Chen C, Yin M, Hsu C, Liu T (2007) Antioxidative and anti-inflammatory effects of four cysteine-containing agents in striatum of MPTP-treated mice. *Nutrition* 23:589–597
 198. Kaya Y, Ozsoy O, Onullut E, Aslan M, Yildirim FB, Anullar A (2013) Beneficial effect of melatonin treatment on substantia nigra in an experimental model of Parkinson's disease. *J Neurol Sci* 30:142
 199. Sharma A, Kaur P, Kumar V, Gill KD (2007) Attenuation of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine induced nigrostriatal toxicity in mice by N-acetyl cysteine. *Cell Mol Biol* 53:48–55
 200. Shin JY, Park H, Ahn YH, Lee PH (2009) Neuroprotective effect of l-dopa on dopaminergic neurons is comparable to pramipexol in MPTP-treated animal model of Parkinson's disease: a direct comparison study. *J Neurochem* 111:1042–1050
 201. Sutherland GT, Chami B, Youssef P, Witting PK (2013) Oxidative stress in Alzheimer's disease: primary villain or physiological by-product? *Redox Rep* 18:134–141
 202. Wolfson C, Wolfson DB, Asgharian M, M'Lan CE, Østbye T, Rockwood K, Hogan DB (2001) A reevaluation of the duration of survival after the onset of dementia. *N Engl J Med* 344:1111–1116
 203. Martins RN, Harper CG, Stokes GB, Masters CL (1986) Increased cerebral glucose-6-phosphate dehydrogenase activity in Alzheimer's disease may reflect oxidative stress. *J Neurochem* 46:1042–1045
 204. Sultana R, Butterfield DA (2010) Role of oxidative stress in the progression of Alzheimer's disease. *J Alzheimer's Dis* 19:341–353
 205. Ansari MA, Scheff SW (2010) Oxidative stress in the progression of Alzheimer disease in the frontal cortex. *J Neuropathol Exp Neurol* 69:155–167
 206. Lüth H, Münch G, Arendt T (2002) Aberrant expression of NOS isoforms in Alzheimer's disease is structurally related to nitrotyrosine formation. *Brain Res* 953:135–143
 207. Lüth H, Holzer M, Gärtner U, Staufenbiel M, Arendt T (2001) Expression of endothelial and inducible NOS-isoforms is increased in Alzheimer's disease, in APP23 transgenic mice and after experimental brain lesion in rat: evidence for an induction by amyloid pathology. *Brain Res* 913:57–67
 208. Massaad CA (2011) Neuronal and vascular oxidative stress in Alzheimer's disease. *Curr Neuropharmacol* 9:662–673
 209. Toda N, Ayajiki K, Okamura T (2009) Cerebral blood flow regulation by nitric oxide in neurological disorders. *Can J Physiol Pharmacol* 87:581–594
 210. Lovell MA, Gabbita SP, Markesbery WR (1999) Increased DNA oxidation and decreased levels of repair products in Alzheimer's disease ventricular CSF. *J Neurochem* 72:771–776
 211. Mecocci P, MacGarvey U, Beal MF (1994) Oxidative damage to mitochondrial DNA is increased in Alzheimer's disease. *Ann Neurol* 36:747–751

212. Lovell MA, Xie C, Markesbery WR (1998) Decreased glutathione transferase activity in brain and ventricular fluid in Alzheimer's disease. *Neurology* 51:1562–1566
213. Casado Á, Encarnación López-Fernández M, Concepción Casado M, De La Torre R (2008) Lipid peroxidation and antioxidant enzyme activities in vascular and Alzheimer dementias. *Neurochem Res* 33:450–458
214. Marcus DL, Thomas C, Rodriguez C, Simberkoff K, Tsai JS, Strafaci JA, Freedman ML (1998) Increased peroxidation and reduced antioxidant enzyme activity in Alzheimer's disease. *Exp Neurol* 150:40–44
215. Spalletta G, Bernardini S, Bellincampi L, Federici G, Trequatrini A, Ciappi F, Bria P, Caltagirone C, Bossù P (2007) Glutathione S-transferase p1 and t1 gene polymorphisms predict longitudinal course and age at onset of Alzheimer disease. *Am J Geriatr Psychiatr* 15:879–887
216. Paz-Y-Miño C, Carrera C, López-Cortés A, Muñoz MJ, Cumbal N, Castro B, Cabrera A, Sánchez ME (2010) Genetic polymorphisms in apolipoprotein e and glutathione peroxidase 1 genes in the Ecuadorian population affected with Alzheimer's disease. *Am J Med Sci* 340:373–377
217. Aoyama K, Nakaki T (2013) Impaired glutathione synthesis in neurodegeneration. *Int J Mol Sci* 14:21021–21044
218. Sultana R, Piroddi M, Galli F, Butterfield DA (2008) Protein levels and activity of some antioxidant enzymes in hippocampus of subjects with amnesic mild cognitive impairment. *Neurochem Res* 33:2540–2546
219. Foy CJ, Passmore AP, Vahidassr MD, Young IS, Lawson JT (1999) Plasma chain-breaking antioxidants in Alzheimer's disease, vascular dementia and Parkinson's disease. *QJM - Month J Assoc Phys* 92:39–45
220. Kim T, Pae C, Yoon S, Jang W, Lee NJ, Kim J, Lee S, Lee C, Paik I, Lee C (2006) Decreased plasma antioxidants in patients with Alzheimer's disease. *Int J Geriatr Psychiatr* 21:344–348
221. Giavarotti L, Simon KA, Azzalis LA, Fonseca FLA, Lima AF, Freitas MCV, Brunialti MKC, Salomão R, Moscardi AAVS, M. Montañó MBM, Ramos LR, Junqueira VBC (2013) Mild systemic oxidative stress in the subclinical stage of Alzheimer's disease. *Oxidat Med Cell Longevity*
222. Arlt S, Müller-Thomsen T, Beisiegel U, Kontush A (2012) Effect of one-year vitamin C- and E-supplementation on cerebrospinal fluid oxidation parameters and clinical course in Alzheimer's disease. *Neurochem Res* 37:2706–2714
223. Galasko DR, Peskind E, Clark CM, Quinn JF, Ringman JM, Jicha GA, Cotman C, Cottrell B, Montine TJ, Thomas RG, Aisen P (2012) Antioxidants for Alzheimer disease: a randomized clinical trial with cerebrospinal fluid biomarker measures. *Arch Neurol* 69:836–841
224. Dysken MW, Sano M, Asthana S, Vertrees JE, Pallaki M, Llorente M, Love S, Schellenberg GD, McCarten JR, Malphurs J, Prieto S, Chen P, Loreck DJ, Trapp G, Bakshi RS, Mintzer JE, Heidebrink JL, Vidal-Cardona A, Arroyo LM, Cruz AR, Zachariah S, Kowall NW, Chopra MP, Craft S, Thielke S, Turvey CL, Woodman C, Monnell KA, Gordon K, Tomaska J, Segal Y, Peduzzi PN, Guarino PD (2014) Effect of vitamin E and memantine on functional decline in Alzheimer disease: the TEAM-AD VA cooperative randomized trial. *JAMA - J Am Med Assoc* 311:33–44
225. Thal LJ, Grundman M, Berg J, Erstrom K, Margolin R, Pfeiffer E, Weiner MF, Zamrini E, Thomas RG (2003) Idebenone treatment fails to slow cognitive decline in Alzheimer's disease. *Neurology* 61:1498–1502
226. Quinn JF, Raman R, Thomas RG, Yurko-Mauro K, Nelson EB, Van Dyck C, Galvin JE, Emond J, Jack CR Jr, Weiner M, Shinto L, Aisen PS (2010) Docosahexaenoic acid supplementation and cognitive decline in Alzheimer disease: a randomized trial. *JAMA - J Am Med Assoc* 304:1903–1911
227. Freund-Levi Y, Eriksdotter-Jönghagen M, Cederholm T, Basun H, Faxén-Irving G, Garlind A, Vedin I, Vessby B, Wahlund L, Palmblad J (2006) ω -3 fatty acid treatment in 174 patients with mild to moderate Alzheimer disease: OmegAD study—a randomized double-blind trial. *Arch Neurol* 63:1402–1408
228. Shinto L, Quinn J, Montine T, Dodge HH, Woodward W, Baldauf-Wagner S, Waichunas D, Bumgarner L, Bourdette D, Silbert L, Kaye J (2014) A randomized placebo-controlled pilot trial of omega-3 fatty acids and alpha lipoic acid in Alzheimer's disease. *J Alzheimer's Dis* 38:111–120
229. Ringman JM, Frautschy SA, Teng E, Begum AN, Bardens J, Beigi M, Gyls KH, Badmaev V, Heath DD, Apostolova LG, Porter V, Vanek Z, Marshall GA, Hellemann G, Sugar C, Masterman DL, Montine TJ, Cummings JL, Cole GM (2012) Oral curcumin for Alzheimer's disease: tolerability and efficacy in a 24-week randomized, double blind, placebo-controlled study. *Alzheimer's Res Therapy* 4
230. Baum L, Lam CWK, Cheung SK, Kwok T, Lui V, Tsoh J, Lam L, Leung V, Hui E, Ng C, Woo J, Chiu HFK, Goggins WB, Zee BC, Cheng KF, Fong CYS, Wong A, Mok H, Chow MSS, Ho PC, Ip SP, Ho CS, Yu XW, Lai CYL, Chan M, Szeto S, Chan IHS, Mok V (2008) Six-month randomized, placebo-controlled, double-blind, pilot clinical trial of curcumin in patients with Alzheimer disease. *J Clin Psychopharmacol* 28:110–113
231. Disilvestro RA, Joseph E, Zhao S, Bomser J (2012) Diverse effects of a low dose supplement of lipidated curcumin in healthy middle aged people. *Nutrition* J 11
232. Truelsen T, Thudium D, Grønbaek M (2002) Amount and type of alcohol and risk of dementia: the Copenhagen City Heart Study. *Neurology* 59:1313–1319
233. <https://clinicaltrials.gov/ct2/show/NCT01504854>. Accessed 15 May 2015
234. Klugman A, Naughton DP, Isaac M, Shah I, Petroczi A, Tabet N (2012) Antioxidant enzymatic activities in Alzheimer's disease: the relationship to acetylcholinesterase inhibitors. *J Alzheimer's Dis* 30:467–474
235. Gubandru M, Margina D, Tsitsimpikou C, Goutzourelas N, Tsarouhas K, Ilie M, Tsatsakis AM, Kouretas D (2013) Alzheimer's disease treated patients showed different patterns for oxidative stress and inflammation markers. *Food Chem Toxicol* 61:209–214
236. Snalina N, Alessenko A, Gavrilova S, Gurianova S, Prochorov A, Kononova E, Fedorova Y (2014) Memantine changes lipids spectrum and lipid peroxidation in animal brain and plasma of patients with Alzheimer's disease. *FEBS J* 281:86
237. Hu D, Li C, Han N, Miao L, Wang D, Liu Z, Wang H, Yin J (2012) Deoxyschizandrin isolated from the fruits of *Schisandra chinensis* ameliorates a β induced memory impairment in mice. *Planta Med* 78:1332–1336
238. Hou X, Wu D, Zhang C, Yan R, Yang C, Rong C, Zhang L, Chang X, Su R, Zhang S, He W, Qu Z, Li S, Su Z, Chen Y, Wang Q, Fang S (2014) Bushen-Yizhi formula ameliorates cognition deficits and attenuates oxidative stress-related neuronal apoptosis in scopolamine-induced senescence in mice. *Int J Mol Med* 34:429–439
239. Chonpathompikunlert P, Wattanathorn J, Muchimapura S (2010) Piperine, the main alkaloid of Thai black pepper, protects against neurodegeneration and cognitive impairment in animal model of cognitive deficit like condition of Alzheimer's disease. *Food Chem Toxicol* 48:798–802
240. Veerendra Kumar MH, Gupta YK (2003) Effect of *Centella asiatica* on cognition and oxidative stress in an intracerebroventricular streptozotocin model of Alzheimer's disease in rats. *Clin Exp Pharmacol Physiol* 30:336–342

241. Li X, Zhao X, Xu X, Mao X, Liu Z, Li H, Guo L, Bi K, Jia Y (2014) Schisantherin A recovers A(beta)-induced neurodegeneration with cognitive decline in mice. *Physiol Behav* 132:10–16
242. Fan Y, Hu J, Li J, Yang Z, Xin X, Wang J, Ding J, Geng M (2005) Effect of acidic oligosaccharide sugar chain on scopolamine-induced memory impairment in rats and its related mechanisms. *Neurosci Lett* 374:222–226
243. Richards JG, Higgins GA, Ouagazzal A, Ozmen L, Kew JNC, Bohrmann B, Malherbe P, Brockhaus M, Loetscher H, Czech C, Huber G, Bluethmann H, Jacobsen H, Kemp JA (2003) PS2APP transgenic mice, coexpressing hPS2mut and hAPPswe, show age-related cognitive deficits associated with discrete brain amyloid deposition and inflammation. *J Neurosci* 23:8989–9003
244. Cavanaugh SE, Pippin JJ, Barnard ND (2014) Animal models of Alzheimer disease: historical pitfalls and a path forward. *ALTEX* 31:279–302
245. Götz J, Ittner LM (2008) Animal models of Alzheimer's disease and frontotemporal dementia. *Nat Rev Neurosci* 9:532–544
246. Lannert H, Hoyer S (1998) Intracerebroventricular administration of streptozotocin causes long-term diminutions in learning and memory abilities and in cerebral energy metabolism in adult rats. *Behav Neurosci* 112:1199–1208
247. Jhoo JH, Kim H, Nabeshima T, Yamada K, Shin E, Jhoo W, Kim W, Kang K, Jo SA, Woo JI (2004) β -Amyloid (1–42)-induced learning and memory deficits in mice: involvement of oxidative burdens in the hippocampus and cerebral cortex. *Behav Brain Res* 155:185–196
248. Budzynska B, Boguszewska-Czubara A, Kruk-Slomka M, Skalicka-Wozniak K, Michalak A, Musik I, Biala G (2015) Effects of imperatorin on scopolamine-induced cognitive impairment and oxidative stress in mice. *Psychopharmacology (Berl)* 232:931–942
249. Goverdhan P, Sravanthi A, Mamatha T (2012) Neuroprotective effects of meloxicam and selegiline in scopolamine-induced cognitive impairment and oxidative stress. *Int J Alzheimer's Dis*
250. Javed H, Khan MM, Khan A, Vaibhav K, Ahmad A, Khuwaja G, Ahmed ME, Raza SS, Ashafaq M, Tabassum R, Siddiqui MS, El-Agnaf OM, Safhi MM, Islam F (2011) S-allyl cysteine attenuates oxidative stress associated cognitive impairment and neurodegeneration in mouse model of streptozotocin-induced experimental dementia of Alzheimer's type. *Brain Res* 1389:133–142
251. Siedlak SL, Casadesus G, Webber KM, Pappolla MA, Atwood CS, Smith MA, Perry G (2009) Chronic antioxidant therapy reduces oxidative stress in a mouse model of Alzheimer's disease. *Free Radic Res* 43:156–164
252. Harrison FE, Hosseini AH, McDonald MP, May JM (2009) Vitamin C reduces spatial learning deficits in middle-aged and very old APP/PSEN1 transgenic and wild-type mice. *Pharmacol Biochem Behav* 93:443–450
253. Hartl D, Schuldt V, Forler S, Zabel C, Klose J, Rohe M (2012) Presymptomatic alterations in energy metabolism and oxidative stress in the APP23 mouse model of Alzheimer disease. *J Proteome Res* 11:3295–3304
254. Feng Z, Qin C, Chang Y, Zhang J (2006) Early melatonin supplementation alleviates oxidative stress in a transgenic mouse model of Alzheimer's disease. *Free Radic Biol Med* 40:101–109
255. Zhang W, Bai M, Xi Y, Hao J, Liu L, Mao N, Su C, Miao J, Li Z (2012) Early memory deficits precede plaque deposition in APPswe/PS1dE9 mice: involvement of oxidative stress and cholinergic dysfunction. *Free Radic Biol Med* 52:1443–1452
256. Khan MB, Khan MM, Khan A, Ahmed ME, Ishrat T, Tabassum R, Vaibhav K, Ahmad A, Islam F (2012) Naringenin ameliorates Alzheimer's disease (AD)-type neurodegeneration with cognitive impairment (AD-TNDCI) caused by the intracerebroventricular-streptozotocin in rat model. *Neurochem Int* 61:1081–1093
257. Harrison FE, May JM, McDonald MP (2010) Vitamin C deficiency increases basal exploratory activity but decreases scopolamine-induced activity in APP/PSEN1 transgenic mice. *Pharmacol Biochem Behav* 94:543–552
258. Duffy AM, Hölscher C (2013) The incretin analogue D-Ala2GIP reduces plaque load, astrogliosis and oxidative stress in an APP/PS1 mouse model of Alzheimer's disease. *Neuroscience* 228:294–300
259. Wang D, Liu L, Zhu X, Wu W, Wang Y (2014) Hesperidin alleviates cognitive impairment, mitochondrial dysfunction and oxidative stress in a mouse model of Alzheimer's disease. *Cell Mol Neurobiol*
260. Otalora BB, Popovic N, Gambini J, Popovic M, Viña J, Bonet-Costa V, Reiter RJ, Camello PJ, Rol MÁ, Madrid JA (2012) Circadian system functionality, hippocampal oxidative stress, and spatial memory in the APPswePS1dE9 transgenic model of Alzheimer disease: effects of melatonin or ramelteon. *Chronobiol Int* 29:822–834
261. Saxena G, Singh SP, Agrawal R, Nath C (2008) Effect of donepezil and tacrine on oxidative stress in intracerebral streptozotocin-induced model of dementia in mice. *Eur J Pharmacol* 581:283–289
262. Benzie IFF, Strain JJ (1996) The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': the FRAP assay. *Anal Biochem* 239:70–76
263. Kracmarova A, Bandouchova H, Pikula J, Pohanka M (2012) Tacrine is implicated in oxidative stress in the laboratory guinea pig model. *Neuroendocrinol Lett* 33:136–144
264. Khurana S, Jain S, Mediratta PK, Banerjee BD, Sharma KK (2012) Protective role of curcumin on colchicine-induced cognitive dysfunction and oxidative stress in rats. *Hum Exp Toxicol* 31:686–697
265. Mahdy K, Shaker O, Wafay H, Nassar Y, Hassan H, Hussein A (2012) Effect of some medicinal plant extracts on the oxidative stress status in Alzheimer's disease induced in rats. *Eur Rev Med Pharmacol Sci* 16:31–42
266. Kumar A, Prakash A, Pahwa D (2011) Galantamine potentiates the protective effect of rofecoxib and caffeic acid against intrahippocampal kainic acid-induced cognitive dysfunction in rat. *Brain Res Bull* 85:158–168
267. Pietá Dias C, Martins de Lima MN, Presti-Torres J, Dornelles A, Garcia VA, Siciliani Scalco F, Rewsaat Guimarães M, Constantino L, Budni P, Dal-Pizzol F, Schröder N (2007) Memantine reduces oxidative damage and enhances long-term recognition memory in aged rats. *Neuroscience* 146:1719–1725
268. Arif M, Chikuma T, Ahmed MM, Nakazato M, Smith MA, Kato T (2009) Effects of memantine on soluble A β 25–35-induced changes in peptidergic and glial cells in Alzheimer's disease model rat brain regions. *Neuroscience* 164:1199–1209
269. Rai S, Kamat PK, Nath C, Shukla R (2014) Glial activation and post-synaptic neurotoxicity: the key events in streptozotocin (ICV) induced memory impairment in rats. *Pharmacol Biochem Behav* 117:104–117
270. Kamat PK, Tota S, Saxena G, Shukla R, Nath C (2010) Okadaic acid (ICV) induced memory impairment in rats: a suitable experimental model to test anti-dementia activity. *Brain Res* 1309:66–74
271. Nasr P, Carbery T, Geddes JW (2009) N-Methyl-D-aspartate receptor antagonists have variable affect in 3-nitropropionic acid toxicity. *Neurochem Res* 34:490–498
272. Rojas JC, Saavedra JA, Gonzalez-Lima F (2008) Neuroprotective effects of memantine in a mouse model of retinal degeneration induced by rotenone. *Brain Res* 1215:208–217
273. Zaja-Milatovic S, Gupta RC, Aschner M, Milatovic D (2009) Protection of DFP-induced oxidative damage and

- neurodegeneration by antioxidants and NMDA receptor antagonist. *Toxicol Appl Pharmacol* 240:124–131
274. LÄ¼ J, Lin PH, Yao Q, Chen C (2010) Chemical and molecular mechanisms of antioxidants: experimental approaches and model systems. *J Cell Mol Med* 14:840–860
275. Rahal A, Kumar A, Singh V, Yadav B, Tiwari R, Chakraborty S, Dhama K (2014) Oxidative stress, prooxidants, and antioxidants: the interplay. *BioMed Res Int* 2014
276. Townsend DM, Tew KD (2003) The role of glutathione-S-transferase in anti-cancer drug resistance. *Oncogene* 22:7369–7375
277. Tabart J, Kevers C, Pincemail J, Defraigne J, Dommes J (2009) Comparative antioxidant capacities of phenolic compounds measured by various tests. *Food Chem* 113:1226–1233
278. Siasos G, Tousoulis D, Tsigkou V, Kokkou E, Oikonomou E, Vavuranakis M, Basdra EK, Papavassiliou AG, Stefanadis C (2013) Flavonoids in atherosclerosis: an overview of their mechanisms of action. *Curr Med Chem* 20:2641–2660
279. Jomova K, Valko M (2011) Advances in metal-induced oxidative stress and human disease. *Toxicology* 283:65–87
280. Lushchak VI (2012) Glutathione homeostasis and functions: potential targets for medical interventions. *J Amino Acids* 2012
281. Djamali A (2007) Oxidative stress as a common pathway to chronic tubulointerstitial injury in kidney allografts. *Am J Physiol - Renal Physiol*
282. Bartosz G (2009) Reactive oxygen species: destroyers or messengers? *Biochem Pharmacol* 77:1303–1315
283. Brieger K, Schiavone S, Miller FJ, Krause K- (2012) Reactive oxygen species: from health to disease. *Swiss Medical Weekly* 142