

Matrix metalloproteinases in exercise and obesity

Jonathan Jaoude¹
Yunsuk Koh²

¹Department of Biology, ²Department of Health, Human Performance, and Recreation, Baylor University, Waco, TX, USA

Abstract: Matrix metalloproteinases (MMPs) are zinc- and calcium-dependent endoproteinases that have the ability to break down extracellular matrix. The large range of MMPs' functions widens their spectrum of potential role as activators or inhibitors in tissue remodeling, cardiovascular diseases, and obesity. In particular, MMP-1, -2, and -9 may be associated with exercise and obesity. Thus, the current study reviewed the effects of different types of exercise (resistance and aerobic) on MMP-1, -2, and -9. Previous studies report that the response of MMP-2 and -9 to resistance exercise is dependent upon the length of exercise training, since long-term resistance exercise training increased both MMP-2 and -9, whereas acute bout of resistance exercise decreased these MMPs. Aerobic exercise produces an inconsistent result on MMPs, although some studies showed a decrease in MMP-1. Obesity is related to a relatively lower level of MMP-9, indicating that an exercise-induced increase in MMP-9 may positively influence obesity. A comprehensive understanding of the relationship between exercise, obesity, and MMPs does not exist yet. Future studies examining the acute and chronic responses of these MMPs using different subject models may provide a better understanding of the molecular mechanisms that are associated with exercise, obesity, and cardiovascular disease.

Keywords: cardiovascular disease, gelatinases, collagenases, TIMP

Introduction

The property of matrix metalloproteinases

Matrix metalloproteinases (MMPs) were first observed in 1962 by Jerome Gross and Charles Lapiere in tadpole tissue that exhibited collagenolytic activity.¹ Eisen et al² were able to isolate human MMPs 6 years after its first discovery. MMPs are zinc- and calcium-dependent endoproteinases that play a crucial role in the remodeling of extracellular matrix (ECM) by breaking down its protein components.³ MMPs can be categorized, on the basis of substrate specificity and homology, into the following six family groups: collagenases, gelatinases, stromelysins, matrilysins, membrane-type MMPs, and other MMPs (Figure 1; Table 1). All MMPs share common domain structures that degrade various ECM and nonmatrix.^{4,5} Specific propeptide and catalytic domains exist (ie, MMP-7 and -26) along with a hemopexin-like, four-bladed, β -propeller domain located on the C-terminus, which is connected to a linker or hinge region (MMP-1, -3, -8, -11, -12, -13, -18, -19, -20, -21, -27, and -28).⁶ These are the domains and regions that are involved in substrate recognition and inhibitor binding.⁷⁻⁹ MMP-2 and -9 also have a fibronectin-like domain of three type II repeats.⁶

Correspondence: Yunsuk Koh
Department of Health, Human Performance, and Recreation, Baylor University, One Bear Place #97313, Waco, TX 76798, USA
Tel +1 254 710 4002
Fax +1 254 710 3527
Email yunsuk_koh@baylor.edu




Figure 1 General structure of MMP groups with SP, propeptide, catalytic, and hemopexin domains.

Notes: The active site of zinc bound to the catalytic domain and the CS can be found in all MMPs. Gelatinase FN repeats are found in the catalytic domain of gelatinases. A furin-cleavage site (F) between the prodomain and catalytic domain can be found in MT-MMPs. Some MMPs have a TM domain attached to the hemopexin domain (MT-MMP). Other MT-MMPs have GPI-anchor domain.

Abbreviations: CS, cysteine switch; FN, fibronectin; GPI, glycosylphosphatidylinositol; MMP, matrix metalloproteinase; MT-MMP, membrane type-matrix metalloproteinase; SP, signal peptide; TM, transmembrane.

Activation of MMPs

There are three general activation mechanisms of MMPs, including the pro-MMP cleavage, phosphorylation, and oxidative stressors. The cleavage of pro-MMP is known as “cysteine switch” that exists in all known MMPs.¹⁰ The cleavage targets an intramolecular complex between a single cysteine residue in the prodomain and the zinc ion in the catalytic domain (Figure 1). This cleavage of MMPs is a required activation before any other forms of catalytic activation may occur. MMPs can be activated by phosphorylation. One study examined ubiquitous MMP-2 in human connective tissue and found that five of 29 potential phosphorylation sites are targets for protein kinase C to alter MMP-2 activity. Enzymatic property changes were confirmed through zymograph, gelatin dequenching assays, and analysis of kinetic parameters.¹¹ MMPs can also be activated by oxidative stressors such as homocysteine (Hcy), nitric oxide (NO), and hydrogen sulfide (H₂S). Hcy is a metabolite of the amino acids cysteine and methionine that activates MMPs by the extracellular signal-regulated kinase pathway.^{12,13} In mice, high levels of Hcy showed increased aortic MMP-2 and -9 along with

increased aortic blood pressure, resistance, pulse rate, wall thickness, and extracellular collagen accumulation.¹⁴ On the other hand, H₂S, a metabolite of Hcy, has been clinically used to treat atherosclerosis, since it deactivates MMP-9 and activates MMP-2 in heart tissue. Thus, H₂S may promote angiogenesis and inhibit antiangiogenic factors.¹⁵ H₂S is an important factor for an anaerobic pathway conversion process of Hcy, and plays a major role in vasodilation and antioxidant health by normalizing the levels of redox stress and MMPs, as seen in vascular remodeling of damaged carotid artery.¹⁶ Injected H₂S into the carotid artery showed increased levels of MMP-9 and decreased levels of MMP-2.¹⁷ One study reported that H₂S-knockout mice showed more outgrowth in the vessels, as compared to donor H₂S that increased the levels of neointima formation in the carotid artery. This result supports evidence on how H₂S can be seen as a clinical route to treat atherosclerosis.¹⁷ Although NO plays an important role in MMP regulation, its exact mechanism has not been completely understood. Moreover, the results of previous studies investigating the MMP regulation by NO or NO-dependent pathways are equivocal. One study

Table 1 Classification of MMPs

MMP	Category	Enzyme	Target(s)	Inhibitor(s)
MMP-1	Collagenases	Collagenase-1	Collagens (I–III, VII, VIII, and X), gelatin, aggrecan, L-selectin, IL-1 β , proteoglycans, entactin, ovostatin, MMP-2, MMP-9	Batimastat (BB-94), BB-1101, MMI270B, Metastat (CMT-3), Doxycycline, FN-439, Ilomastat, Marimastat (BB-2516), Minocycline
MMP-2	Gelatinases	Gelatinase-A	Gelatin, collagen IV–VI, X, elastin, fibronectin	TIMP-4, Batimastat (BB-94), BB-1101, MMI270B, Doxycycline, Ilomastat, Marimastat (BB-2516), Minocycline
MMP-3	Stromelysins	Stromelysin-1	Collagens (III–V, and IX), gelatin, aggrecan, perlecan, decorin, laminin, elastin, casein, osteonectin, ovostatin, entactin, plasminogen, MBP, IL-1 β , MMP-2/TIMP-2, MMP-7, MMP-8, MMP-9, MMP-13	Batimastat (BB-94), BB-1101, MMI270B, Doxycycline, FN-439, Ilomastat, Marimastat (BB-2516), Minocycline
MMP-7	Matrilysins	Matrilysin (PUMP)	Collagens (IV, X), gelatin, aggrecan, decorin, fibronectin, laminin, elastin, casein, transferrin, plasminogen, β 4-integrin, MMP-1, MMP-2, MMP-9, MMP-9/TIMP-1	Batimastat (BB-94), BB-1101, Doxycycline, Marimastat (BB-2516), Minocycline
MMP-8	Collagenases	Collagenase-2/neutrophil	Collagens (I–III, V, VII, VIII, and X), gelatin, aggrecan, fibronectin	TIMP-1, Batimastat (BB-94), BB-1101, MMI270B, Metastat (CMT-3), Doxycycline, FN-439, Ilomastat, Marimastat (BB-2516)
MMP-9	Gelatinases	Gelatinase-A	Collagens (IV, V, VII, X, and XIV), gelatin, entactin, aggrecan, elastin, fibronectin, osteonectin, plasminogen, MBP, IL-1 β	TIMP-1, Batimastat (BB-94), BB-1101, MMI270B, FN-439, Ilomastat, Marimastat (BB-2516), Minocycline
MMP-10	Stromelysins	Stromelysin-2	Collagens (III–V), gelatin, casein, aggrecan, elastin, MMP-1, MMP-8	
MMP-11	Stromelysins	Stromelysin-3	Unknown (casein)	
MMP-12	Other enzymes	Macrophage metalloelastase	Collagen IV, gelatin, elastin, casein, fibronectin, vitronectin, laminin, entactin, fibrinogen, fibrin, plasminogen	BB-1101
MMP-13	Collagenases	Collagenase-3	Collagens (I–IV, IX, X, and XIV), gelatin, plasminogen, aggrecan, perlecan, fibronectin, osteonectin, MMP-9	BB-1101, Metastat (CMT-3), MMI270B, Doxycycline
MMP-14	MT-MMP	MT1-MMP	Collagens (I–III), gelatin, casein, fibronectin, laminin, vitronectin, entactin, proteoglycans, MMP-2, MMP-13	TIMP-1, TIMP-2, BB-1101, Ilomastat, Marimastat (BB-2516)
MMP-15	MT-MMP	MT2-MMP	Fibronectin, entactin, laminin, aggrecan, perlecan; MMP-2	
MMP-16	MT-MMP	MT3-MMP	Collagen III, gelatin, casein, fibronectin, MMP-2	
MMP-17	Stromelysins	Homology tostromelysin-2 (51.6%)		
MMP-17	MT-MMP	MT4-MMP		TIMP-1, TIMP-2
MMP-18	Collagenases	Collagenase-4	Type I collagen	
MMP-19	Other enzymes	RASI 1	Type I collagen	
MMP-20	Other enzymes	Enamelysin	Amelogenin, aggrecan	
MMP-21	Other enzymes	MMP identified on chromosome 1		
MMP-22	Other enzymes	MMP identified on chromosome 1		
MMP-23	Other enzymes	From human ovary cDNA		
MMP-24	MT-MMP	MT5-MMP	Fibronectin, but not collagen type I or laminin	
MMP-25	MT-MMP	MT6-MMP	Progelatinase A	TIMP-1, TIMP-4
MMP-26	Matrilysins	Matrilysin-2	Collagen IV, fibronectin, fibrinogen, gelatin, α (1)-proteinase inhibitor	
MMP-28	Other enzymes	Epilysin		
MMP-29	Other enzymes	Unnamed		

Abbreviations: MMP, matrix metalloproteinase; IL, interleukin; MT-MMP, membrane type-matrix metalloproteinase; RASI, rice-amylose/subtilisin inhibitor; cDNA, complementary DNA; CMT, chemically modified tetracycline; MMI, matrix metalloproteinase inhibitor; TIMP, tissue inhibitor of metalloproteinase.

reported that NO inhibited MMP-9 expression in activated astrocytes,¹⁸ while another study showed that NO inhibitor rather decreased MMP-9 in rat cardiac allografts.¹⁹ The three MMP regulators mentioned above can give more insight on how MMP levels can be regulated in ways that can benefit the human body against diseases. However, more knowledge of the exact mechanisms related to each regulator is necessary for a full understanding of the MMP processes.

Inhibition of MMPs

Two types of MMP inhibitors exist: endogenous and exogenous inhibitors. Tissue inhibitors of metalloproteinase (TIMPs) are endogenous inhibitors that can be secreted (TIMP-1, TIMP-2, TIMP-4) or bound to ECM components (TIMP-3).²⁰ They inactivate MMPs by forming bonds with catalytic zinc in 1:1 ratios within the MMP structure.²¹ They do so by creating noncovalent interactions between the N-terminal domain of the TIMP and the active site of MMPs. Recent studies have shown TIMPs' therapeutic potential in cardiovascular diseases (CVDs). The MMP-inhibitory effects of TIMP-1 involve binding of C-terminal domain to pro-MMP-2 and pro-MMP-9. TIMP-1 may specifically help ECM manipulation in ischemia in such a way that it may be a surrogate marker for increased ECM turnover.²² TIMP-2 reverses ECM remodeling in human cardiac tissue in a dose-dependent manner.²³ TIMP-3 prevented degradation of cardiac tissue matrix after a myocardial infarction by reducing MMPs in vascular smooth muscle cells.²⁴ Changes in the ECM of atrial fibroblast in rheumatic heart disease have been associated with TIMP-4 expression.²⁵

Exogenous inhibitors include hydroxamic acid derivative and thiirane gelatinase inhibitor SB-3CT on several MMPs.^{26–29} Batimastat (BB-94), a hydroxamic acid derivative, has been recently found to treat aneurysms on using nanoparticle technology to directly administer anti-MMP target to abdominal aorta aneurysm.³⁰ In rat aorta cell culture, administration of BB-94 decreased 90% of MMP-9 activity and 10% of MMP-2 activity, while showing no effect on TIMP-2 activity. The effect of change in MMP activity was tested by injecting BB-94 directly into the abdominal aorta. The blank control showed a 269% increase in the aneurysm, while the treatment group showed only a 40% increase in expansion. Another derivative, marimastat, has shown success as a therapeutic drug in repressing non-small-cell lung cancer in a Phase I trial.³¹ Oral administration of marimastat was added to the accepted treatment with carboplatin and paclitaxel in order to test whether or not marimastat would affect the kinetics of the treatment. TIMPs and hydroxamic

acid derivatives play significant roles in general physiology and pathology, and thus can develop as a therapeutic target in the future.

The roles of MMPs in CVD and obesity

Breakdown of ECM by MMPs includes physiological processes such as embryonic development, reproduction, and tissue remodeling, as well as disease processes. MMPs are involved in the remodeling process of cell membrane and in cell behaviors like proliferation, migration, and apoptosis.³² CVD is the number one leading cause of death in the world and includes any disease associated with the heart and blood vessels, such as myocardial infarction, heart failure, atherosclerosis, stroke, aortic aneurysms, and so on.³³ MMPs are expressed and activated in many different types of CVDs including atherosclerosis, myocardial infarction, and cardiac dysfunction.^{34,35} In particular, MMP-2 and -9 degrade a major element in the basement membrane, collagen IV, which helps in cellular arrangement of skeletal muscle.³⁶ MMPs also play a role in disease diagnoses. For instance, MMP-2 and -9 are found to be independent predictors for kidney disease progression and its associated mortality.³⁷ Obesity is strongly associated with CVDs and occurs when pre-existing, fully differentiated adipocytes are enlarged by excess energy input and accumulate to a point where the pathological expansion becomes a concern.³⁸ Higher levels of MMPs are associated with obesity and CVDs. For instance, MMP-1, -2, -3, -7, -9, -10, -11, and -12 are found at higher levels in atherosclerotic arteries.³⁹ Among these MMPs, the current review focused on MMP-1, -2, and -9 due to their strong association with obesity and CVDs.

MMP-1

MMP-1 may be involved in plaque burden, although plaque morphology was not tested.⁴⁰ A strong correlation between MMP-1 mRNA and lupus erythematosus atherosclerosis has been reported.⁴¹ Elevated MMP-1 was also associated with myocardial infarction and angiographic coronary artery disease, although the mechanistic pathway was not examined.⁴² One study promoted inflammation and atherosclerosis using C-reactive protein to examine the effects of MMP-1 in human mammary arteries and carotid arteries. Both MMP-1 and its mRNA expression had increased significantly, suggesting an association with inflammation and plaque vulnerability.⁴³ Higher levels of MMP-1 have also been observed in several types of human carotid atherosclerosis, and also, histological associations with plaque instability have been

found.⁴⁴ However, controversial results were obtained on the association of MMP-1 with obesity, since the expression of MMP-1 may be different between obese and nonobese people. According to a recent study, MMP-1 may stimulate tissue remodeling during adipose tissue expansion in obesity. Certain MMP-1 alleles showed increased frequency with high body mass index, potentially suggesting a defensive role.⁴⁵ In contrast, MMP-1 was reported to have a strong association with nonobese individuals.⁴⁶ In spite of the contradictory effect of MMP-1 on obesity, the majority of evidence leans toward MMP-1 increasing with obesity.

MMP-2

MMP-2 has been reported to promote atherosclerosis. Elevated MMP-2 may negatively affect vascular permeability and play an important role in the progression of heart failure.⁴⁷ One study reported that MMP-2 knockout mice showed decrease in atherogenesis.⁴⁸ Increased duration of ischemia and delayed functional recovery have been linked to elevated MMP-2 levels,⁴⁹ while decreased MMP-2 levels provided protection from cardiac dysfunction.⁵⁰ The role of MMP-2 in voluntary exercise and its effects on infarct size have been investigated in rats.⁵¹ A 6-week voluntary wheel-running exercise, where the rats self-selected the time, duration, and intensity in a nonstressful environment, was used as the exercise protocol. Induced ischemia by left anterior descending coronary artery occlusion *ex vivo* and angina provoked by epinephrine plus phentolamine protocols were used on the rats. Serum MMP-2, coronary effluent MMP-2 activity, and infarct size all showed a significant decrease after exercise. These results show that MMP-2 may be soon viewed as a cardioprotective molecule for myocardial infarctions and perfusions.

MMP-9

It has been reported that MMP-9 may be involved in elastase action related to aortic stiffening and development of isolated systolic hypertension in healthy and younger individuals.⁵² However, MMP-9 may attenuate atherosclerotic development and prevent plaque development, since MMP-9 knockout mice showed plaque development.⁵³ Both MMP-2 and -9 have been linked to increased inflammation under high coronary risk events and high plaque instability.^{54,55} An early study showed that genetically obese rats had high levels of MMP-2 and low levels of MMP-9,⁵⁶ suggesting that MMP-2 may be involved in adipose ECM degradation. In addition, high-fat diet-induced obese (HFDIO) mice had lower MMP-9 mRNA, and an antigrowth myostatin (MSTN), which is known to

be suppressed in HFDIO condition, showed resistance to HFDIO.⁵⁷ This indicates that decreased levels of MMP-2 and increased MMP-9 levels can prevent or regulate obesity development.

The effects of exercise on MMPs

It is evident that exercise favorably affects CVDs and obesity. According to the recent studies examining the effects of exercise on the regulation of MMPs, the responses of MMPs to exercise may be more dependent upon the mode and length of exercise performed in a variety of subject models (Table 2). Thus, understanding the relationship between MMPs and exercise is particularly important in obese population, since obesity is strongly associated with CVDs and other types of metabolic diseases.⁵⁸ In the current review, we examined how different types of exercise (resistance and aerobic) influence MMP-1, -2, and -9.

Resistance training and MMPs

Based on the previously published studies, the responses of MMPs to resistance training are more likely related to duration of exercise training. Resistance training lasting from 5 to 12 weeks may increase MMP-2 and -9 in both animal and human subjects,⁵⁹⁻⁶² whereas acute bout of resistance training may decrease these MMPs.⁶³ In an animal study, HFDIO rats that performed resistance training had significantly increased MMP-2 levels in bicep and gastrocnemius muscles. A comparable difference in MMP-2 levels between obese and nonobese rats indicated that it was related to the levels of obesity and also suggested that this exercise-induced increase in MMP-2 in rats may prevent obesity. In this regard, MMP-2 could potentially be a negative regulator of obesity.⁶² In another study, rats fed with high-fat diet performed a 12-week strength training program that consisted of vertical ladder exercise three times a week with weights attached to their tails. Following 12 weeks of training, rats showed increase in muscle MMP-2, suggesting that high levels of MMP-2 may be inversely related to obesity.⁵⁹ One study examined the effects of oxidative stress on MMP-2 and -9 in skeletal muscle using the exercise protocol composed of leg presses for 45 minutes, four times a week for 5 weeks. The levels of MMP-2 increased by day 10, and MMP-2 mRNA in myofibril increased with training as well. The responses of MMP-9 and its mRNA activity at day 10 were relatively lower as compared to those of MMP-2, but held the same level of activity after the first exercise.⁶⁰ One study examining the relationship between diabetes and MMPs reported that patients with type 2 diabetes who performed a rowing

Table 2 Effects of different types of exercise on MMPs

Study	Samples	Training type	Length of study	Tissue	Outcome	P-value
Souza et al ⁶²	Wistar rats (N=32; C=8, C-OB=8, EG=8, EG-OB=8)	RT	12 weeks	Skeletal muscle	MMP-2 ↑	P<0.05
Leite et al ⁵⁹	Wistar rats (N=32; C=8, C-OB=8, EG=8, EG-OB=8)	RT	12 weeks	Left ventricle	MMP-2 ↑	P<0.01
Nascimento et al ⁶³	Obese elderly women (N=10)	RT	Acute	Plasma	MMP-2 ↓ MMP-9 ↓	P<0.05
Rullman et al ⁶⁰	Healthy males (N=10)	RT	5 weeks	Skeletal muscle	MMP-2 ↑ MMP-9 ↑	P<0.001 P<0.05
Scheede-Bergdahl et al ⁶¹	Type 2 diabetic males (N=22; C=10, T2D=12)	RT	8 weeks	Skeletal muscle	MMP-2 ↑	P<0.05
Donley et al ⁶⁹	MetS females (C=22, MetS=22)	AT	8 weeks	Plasma	MMP-1 ↓	P<0.05
Rullman et al ⁶⁸	Healthy males (N=10)	AT	Acute	Skeletal muscle	MMP-2 ↔ MMP-9 ↑	P<0.05 P<0.05
Shon et al ⁶⁵	ApoE ^{-/-} mice (N=68; C=15, C-OB=26, EG=9, EG-OB=18)	AT	10 weeks	Atheroma	MMP-2 ↓ MMP-9 ↓	P<0.05 P<0.05
Posa et al ⁵¹	Wistar rats (N=70; C=35, EG=35)	AT	6 weeks	Plasma	MMP-2 ↓	P<0.001
Kwak et al ⁶⁴	Aging rat model (N=40; C=10, EG=10, O=10, O-EG=10)	AT	12 weeks	Left ventricle	MMP-1 ↓ MMP-2 ↓ MMP-9 ↔	P<0.05 P<0.05 NS
De Aro et al ⁶⁶	Wistar rats (N=77; C=11, 1d1h=11, 1d3h=11, 3d1h=11, 3d3h=11, 6d1h=11, 6d3h=11)	AT	1, 3, 6 days	Calcaneal tendon	MMP-2 ↔, ↔, ↑ MMP-9 ↔, ↔, ↔	P<0.05 NS
Nishijima et al ⁶⁷	Wistar rats (N=4, EG=16)	AT	1 week	Hippocampus	MMP-2 ↔ MMP-9	NS P<0.05

Note: “↑” indicates an increased level; “↓” indicates a decreased level; and “↔” indicates no change.

Abbreviations: ApoE^{-/-}, apolipoprotein E knockout; AT, aerobic exercise; C, control; C-OB, control obese; d, days; EG, exercise group; EG-OB, obese exercise group; h, hours; MetS, metabolic syndrome; MMP, matrix metalloproteinase; NS, not shown; O, old; O-EG, old exercise group; RT, resistance training; T2D, type 2 diabetes.

exercise at 65%–70% of VO₂ (volume of oxygen intake) peak significantly increased MMP-2 mRNA in skeletal muscle, suggesting that exercise-induced changes in MMP-2 may benefit type 2 diabetes.⁶¹ In contrast, obese elderly women who performed acute eccentric resistance training (ten repetitions at 110% of ten-repetition maximum) had decreased levels of both MMP-2 and -9 in the plasma after 48 hours of the training session. The authors concluded that this exercise-induced reduction in MMP-2 and -9 may be a positive sign in the transient defense of inflammatory MMPs associated with obesity and atherosclerosis.⁶³

Aerobic exercise and MMPs

The impact of aerobic exercise training on MMPs may be related to duration of exercise. In general, the long-term aerobic exercise training lasting up to 12 weeks may decrease both MMP-2 and -9,^{51,64,65} while these MMPs increase following acute bouts of exercise.^{66–68} Moreover, only limited information regarding the effects of aerobic exercise on MMP-1 is currently available. MMP-1 in serum and heart tissue decreased following aerobic exercise training (up to 12 weeks) in female subjects with metabolic syndrome⁶⁹ and in aged mice.⁶⁴ MMP-9 and its mRNA activity in skeletal muscle significantly increased after a 65-minute aerobic

cycling exercise, while MMP-2 mRNA did not change.⁶⁸ In an animal study, apoE^{-/-} mice were given a Western diet to develop plaque while participating in a 30-minute treadmill exercise program (5 days/week) for 10 weeks. Exercise intervention did not attenuate aortic plaque, but MMP-2 and -9 significantly decreased.⁶⁵ Another study examining the effects of voluntary wheel-running exercise for 6 weeks reported a reduction in serum MMP-2 as part of a cardio-protective mechanism against cardiac injury.⁵¹ Aged rats that performed a 45-minute aerobic exercise on a treadmill for up to 12 weeks (5 days/week) showed decrease in both MMP-1 and -2 in the heart tissue, while MMP-9 was not altered.⁶⁴ To examine the influence of exercise with various durations and resting periods, Wistar rats were separated into several groups that performed either 1, 3, or 6 days of exercise training (three sessions per day) with an 1-, 3-, or 6-hour resting period between each session. No significant difference was seen in either protein MMP-2 or -9 in the calcaneal tendon in any of the groups, except the group that performed the 6-day with 3-hour rest exercise protocol which showed an increase in MMP-2.⁶⁶ One study examined the responses of MMP-2 and -9 levels of hippocampi samples in rats following treadmill exercise. The rats exercised at moderate intensity on the treadmill for 30 minutes for 7 days, and hippocampi

samples were analyzed by gel zymography to examine the changes in the proteolytic activity of MMP-2 and -9 at 0, 6, 12, and 24 hours postexercise. Results showed that MMP-2 did not change significantly, whereas the 12-hour samples exhibited a significant increase in MMP-9.⁶⁷

Conclusion

The previous studies we reviewed show that the response of MMPs following resistance exercise is more related to the length of resistance exercise training. In general, long-term resistance exercise training may increase both MMP-2 and -9, while acute bouts of resistance exercise may decrease these MMPs. Furthermore, aerobic exercise training leads to an inconsistent result in MMP-2 and -9, although some studies showed a reduction in MMP-1. Also, a relatively lower level of MMP-9 has been observed in obese subjects, indicating that an exercise-induced increase in MMP-9 may play a positive role in obesity.

Elevated MMP-1 has been shown to be strongly associated with CVDs and obesity. Thus, a treatment targeting on lowering MMP-1 may benefit the most in patients with CVDs or obesity. In this regard, an exercise intervention should include a long-term aerobic training as it can reduce the MMP-1 levels and prevent obesity. Reducing MMP-2 would improve prognosis of CVDs, and long-term aerobic training would help reduce MMP-2 levels as well. However, the potential drawback of long-term aerobic training would be decreased level of MMP-9, which may hinder some beneficial effects of exercise on obesity and plaque development. Therefore, it is important to balance aerobic exercise with long-term resistance training, which may increase MMP-9, thereby offering the most positive effects of exercise training on improvement in MMPs. Future studies examining the effects of different types of aerobic and resistance exercise training or a combination of these two forms of training would give more insight on how exercise influences MMPs. Also, exercise intensity may play a role in changes in MMPs, although it has not been fully examined. Therefore, findings on how different types, intensities, and duration of exercises and when to measure MMPs will open a new spectrum of molecular mechanism that is mostly unknown. Finding which MMPs are associated with obesity, exercise, and CVDs is the first step toward understanding the mechanism of this potentially powerful group of proteins. More knowledge of MMPs could lead to more effective treatment or approach to weight loss or new drugs that safely help patients with obesity, CVDs, or other types of diseases. Not all major MMPs have been fully tested with different types and duration of exercise regimens. This calls for research on different MMPs

in exercise to get a better understanding on how the human body can naturally manipulate MMP metabolism to benefit the body as a whole.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Gross J, Lapiere CM. Collagenolytic activity in amphibian tissues: a tissue culture assay. *Proc Natl Acad Sci U S A*. 1962;48:1014–1022.
2. Eisen AZ, Jeffrey JJ, Gross J. Human skin collagenase. Isolation and mechanism of attack on the collagen molecule. *Biochim Biophys Acta*. 1968;151(3):637–645.
3. Renaud S, Leppert D. Matrix metalloproteinases in neuromuscular disease. *Muscle Nerve*. 2007;36(1):1–13.
4. Nagase H, Woessner JF, Jr. Matrix metalloproteinases. *J Biol Chem*. 1999;274(31):21491–21494.
5. Puente XS, Sanchez LM, Overall CM, Lopez-Otin C. Human and mouse proteases: a comparative genomic approach. *Nat Rev Genet*. 2003; 4(7):544–558.
6. Sekhon B. Matrix metalloproteinase – an overview. *Res Rep Biol*. 2010; 1:1–20.
7. Chaudhary AK, Singh M, Bharti AC, Asotra K, Sundaram S, Mehrotra R. Genetic polymorphisms of matrix metalloproteinases and their inhibitors in potentially malignant and malignant lesions of the head and neck. *J Biomed Sci*. 2010;17:10.
8. Kadoglou NP, Liapis CD. Matrix metalloproteinases: contribution to pathogenesis, diagnosis, surveillance and treatment of abdominal aortic aneurysms. *Curr Med Res Opin*. 2004;20(4):419–432.
9. Cho C, Bunch DO, Faure JE, et al. Fertilization defects in sperm from mice lacking fertilin beta. *Science*. 1998;281(5384):1857–1859.
10. Van Wart HE, Birkedal-Hansen H. The cysteine switch: a principle of regulation of metalloproteinase activity with potential applicability to the entire matrix metalloproteinase gene family. *Proc Natl Acad Sci U S A*. 1990;87(14):5578–5582.
11. Sariahmetoglu M, Crawford BD, Leon H, et al. Regulation of matrix metalloproteinase-2 (MMP-2) activity by phosphorylation. *FASEB J*. 2007;21(10):2486–2495.
12. Moshal KS, Sen U, Tyagi N, et al. Regulation of homocysteine-induced MMP-9 by ERK1/2 pathway. *Am J Physiol Cell Physiol*. 2006;290(3): C883–891.
13. Bescond A, Augier T, Chareyre C, Garcon D, Hornebeck W, Charpiot P. Influence of homocysteine on matrix metalloproteinase-2: activation and activity. *Biochem Biophys Res Commun*. 1999;263(2):498–503.
14. Ovechkin AV, Tyagi N, Sen U, et al. 3-Deazaadenosine mitigates arterial remodeling and hypertension in hyperhomocysteinemic mice. *Am J Physiol Lung Cell Mol Physiol*. 2006;291(5):L905–911.
15. Givvimani S, Munjal C, Gargoum R, et al. Hydrogen sulfide mitigates transition from compensatory hypertrophy to heart failure. *J Appl Physiol (1985)*. 2011;110(4):1093–1100.
16. Vacek TP, Gillespie W, Tyagi N, Vacek JC, Tyagi SC. Hydrogen sulfide protects against vascular remodeling from endothelial damage. *Amino Acids*. 2010;39(5):1161–1169.
17. Yang G, Li H, Tang G, et al. Increased neointimal formation in cystathionine gamma-lyase deficient mice: role of hydrogen sulfide in alpha5beta1-integrin and matrix metalloproteinase-2 expression in smooth muscle cells. *J Mol Cell Cardiol*. 2012;52(3):677–688.
18. Shin CY, Lee WJ, Choi JW, et al. Down-regulation of matrix metalloproteinase-9 expression by nitric oxide in lipopolysaccharide-stimulated rat primary astrocytes. *Nitric Oxide*. 2007;16(4):425–432.
19. Egi K, Conrad NE, Kwan J, Schulze C, Schulz R, Wildhirt SM. Inhibition of inducible nitric oxide synthase and superoxide production reduces matrix metalloproteinase-9 activity and restores coronary vasomotor function in rat cardiac allografts. *Eur J Cardiothorac Surg*. 2004; 26(2):262–269.

20. Melendez-Zajgla J, Del Pozo L, Ceballos G, Maldonado V. Tissue inhibitor of metalloproteinases-4. The road less traveled. *Mol Cancer*. 2008;7:85.
21. Brinckerhoff CE, Matrisian LM. Matrix metalloproteinases: a tail of a frog that became a prince. *Nat Rev Mol Cell Biol*. 2002;3(3):207–214.
22. Dinh W, Futh R, Scheffold T, et al. Increased serum levels of tissue inhibitor of metalloproteinase-1 in patients with acute myocardial infarction. *Int Heart J*. 2009;50(4):421–431.
23. Ngu JM, Teng G, Meijndert HC, et al. Human cardiac fibroblast extracellular matrix remodeling: dual effects of tissue inhibitor of metalloproteinase-2. *Cardiovasc Pathol*. 2014;23(6):335–343.
24. Jia ZB, Tian H, Kang K, et al. Expression of the tissue inhibitor of metalloproteinase-3 by transplanted VSMCs modifies heart structure and function after myocardial infarction. *Transpl Immunol*. 2014;30(4):149–158.
25. Sun Y, Huang ZY, Wang ZH, et al. TGF-beta1 and TIMP-4 regulate atrial fibrosis in atrial fibrillation secondary to rheumatic heart disease. *Mol Cell Biochem*. 2015;406(1–2):131–138.
26. Kruger A, Soeltl R, Sopov I, et al. Hydroxamate-type matrix metalloproteinase inhibitor batimastat promotes liver metastasis. *Cancer Res*. 2001;61(4):1272–1275.
27. Kohno T, Hochigai H, Yamashita E, Tsukihara T, Kanaoka M. Crystal structures of the catalytic domain of human stromelysin-1 (MMP-3) and collagenase-3 (MMP-13) with a hydroxamic acid inhibitor SM-25453. *Biochem Biophys Res Commun*. 2006;344(1):315–322.
28. Skarja GA, Brown AL, Ho RK, May MH, Sefton MV. The effect of a hydroxamic acid-containing polymer on active matrix metalloproteinases. *Biomaterials*. 2009;30(10):1890–1897.
29. Gu Z, Cui J, Brown S, et al. A highly specific inhibitor of matrix metalloproteinase-9 rescues laminin from proteolysis and neurons from apoptosis in transient focal cerebral ischemia. *J Neurosci*. 2005;25(27):6401–6408.
30. Nosoudi N, Nahar-Gohad P, Sinha A, et al. Prevention of abdominal aortic aneurysm progression by targeted inhibition of matrix metalloproteinase activity with batimastat-loaded nanoparticles. *Circ Res*. 2015;117(11):e80–89.
31. Goffin JR, Anderson IC, Supko JG, et al. Phase I trial of the matrix metalloproteinase inhibitor marimastat combined with carboplatin and paclitaxel in patients with advanced non-small cell lung cancer. *Clin Cancer Res*. 2005;11(9):3417–3424.
32. Murphy G, Nagase H. Localizing matrix metalloproteinase activities in the pericellular environment. *FEBS J*. 2011;278(1):2–15.
33. Mozaffarian D, Benjamin EJ, Go AS, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics – 2015 update: a report from the American Heart Association. *Circulation*. 2015;131(4):e29–322.
34. Galis ZS, Sukhova GK, Lark MW, Libby P. Increased expression of matrix metalloproteinases and matrix degrading activity in vulnerable regions of human atherosclerotic plaques. *J Clin Invest*. 1994;94(6):2493–2503.
35. Thorp EB. Contrasting inflammation resolution during atherosclerosis and post myocardial infarction at the level of monocyte/macrophage phagocytic clearance. *Front Immunol*. 2012;3:39.
36. Hopps E, Caimi G. Matrix metalloproteinases in metabolic syndrome. *Eur J Intern Med*. 2012;23(2):99–104.
37. Hsu TW, Kuo KL, Hung SC, Huang PH, Chen JW, Tarn DC. Progression of kidney disease in non-diabetic patients with coronary artery disease: predictive role of circulating matrix metalloproteinase-2, -3, and -9. *PLoS One*. 2013;8(7):e70132.
38. Spalding KL, Arner E, Westermark PO, et al. Dynamics of fat cell turnover in humans. *Nature*. 2008;453(7196):783–787.
39. Back M, Ketelhuth DF, Agewall S. Matrix metalloproteinases in atherothrombosis. *Prog Cardiovasc Dis*. 2010;52(5):410–428.
40. Lehrke M, Greif M, Broedl UC, et al. MMP-1 serum levels predict coronary atherosclerosis in humans. *Cardiovasc Diabetol*. 2009;8:50.
41. Zhang HY, Bao SM, Shou WL, et al. Expression of matrix metalloproteinase-1 mRNA in peripheral blood mononuclear cells of systemic lupus erythematosus patients and its relationship with atherosclerosis. *Chin Med J (Engl)*. 2009;122(21):2593–2597.
42. Horne BD, Camp NJ, Carlquist JF, et al. Multiple-polymorphism associations of 7 matrix metalloproteinase and tissue inhibitor metalloproteinase genes with myocardial infarction and angiographic coronary artery disease. *Am Heart J*. 2007;154(4):751–758.
43. Montero I, Orbe J, Varo N, et al. C-reactive protein induces matrix metalloproteinase-1 and -10 in human endothelial cells: implications for clinical and subclinical atherosclerosis. *J Am Coll Cardiol*. 2006;47(7):1369–1378.
44. Nikkari ST, O'Brien KD, Ferguson M, et al. Interstitial collagenase (MMP-1) expression in human carotid atherosclerosis. *Circulation*. 1995;92(6):1393–1398.
45. Nho YK, Ha E, Yu KI, et al. Matrix metalloproteinase-1 promoter is associated with body mass index in Korean population with aged greater or equal to 50 years. *Clin Chim Acta*. 2008;396(1–2):14–17.
46. Huang HL, Wu S, Hsu LA, et al. Genetic variants associated with circulating MMP1 levels near matrix metalloproteinase genes on chromosome 11q21-22 in Taiwanese: interaction with obesity. *BMC Med Genet*. 2013;14:30.
47. Gielen S, Schuler G, Adams V. Cardiovascular effects of exercise training: molecular mechanisms. *Circulation*. 2010;122(12):1221–1238.
48. Kuzuya M, Nakamura K, Sasaki T, Cheng XW, Itohara S, Iguchi A. Effect of MMP-2 deficiency on atherosclerotic lesion formation in apoE-deficient mice. *Arterioscler Thromb Vasc Biol*. 2006;26(5):1120–1125.
49. Kandasamy AD, Chow AK, Ali MA, Schulz R. Matrix metalloproteinase-2 and myocardial oxidative stress injury: beyond the matrix. *Cardiovasc Res*. 2010;85(3):413–423.
50. Fert-Bober J, Leon H, Sawicka J, et al. Inhibiting matrix metalloproteinase-2 reduces protein release into coronary effluent from isolated rat hearts during ischemia-reperfusion. *Basic Res Cardiol*. 2008;103(5):431–443.
51. Posa A, Szabo R, Kupai K, et al. Cardioprotective effects of voluntary exercise in a rat model: role of matrix metalloproteinase-2. *Oxid Med Cell Longev*. 2015;2015:876805.
52. Yasmin, McEniery CM, Wallace S, et al. Matrix metalloproteinase-9 (MMP-9), MMP-2, and serum elastase activity are associated with systolic hypertension and arterial stiffness. *Arterioscler Thromb Vasc Biol*. 2005;25(2):372.
53. Johnson JL, George SJ, Newby AC, Jackson CL. Divergent effects of matrix metalloproteinases 3, 7, 9, and 12 on atherosclerotic plaque stability in mouse brachiocephalic arteries. *Proc Natl Acad Sci U S A*. 2005;102(43):15575–15580.
54. Nascimento Dda C, Durigan Rde C, Tibana RA, Durigan JL, Navalta JW, Prestes J. The response of matrix metalloproteinase-9 and -2 to exercise. *Sports Med*. 2015;45(2):269–278.
55. Heo SH, Cho CH, Kim HO, et al. Plaque rupture is a determinant of vascular events in carotid artery atherosclerotic disease: involvement of matrix metalloproteinases 2 and 9. *J Clin Neurol*. 2011;7(2):69–76.
56. Chavey C, Mari B, Montheuil MN, et al. Matrix metalloproteinases are differentially expressed in adipose tissue during obesity and modulate adipocyte differentiation. *J Biol Chem*. 2003;278(14):11888–11896.
57. Biga PR, Froehlich JM, Greenlee KJ, Galt NJ, Meyer BM, Christensen DJ. Gelatinases impart susceptibility to high-fat diet-induced obesity in mice. *J Nutr Biochem*. 2013;24(8):1462–1468.
58. Lavie CJ, Milani RV, Ventura HO. Obesity and cardiovascular disease: risk factor, paradox, and impact of weight loss. *J Am Coll Cardiol*. 2009;53(21):1925–1932.
59. Leite RD, Durigan Rde C, de Souza Lino AD, et al. Resistance training may concomitantly benefit body composition, blood pressure and muscle MMP-2 activity on the left ventricle of high-fat fed diet rats. *Metabolism*. 2013;62(10):1477–1484.
60. Rullman E, Norrbom J, Stromberg A, et al. Endurance exercise activates matrix metalloproteinases in human skeletal muscle. *J Appl Physiol (1985)*. 2009;106(3):804–812.
61. Scheede-Bergdahl C, Bergdahl A, Schjerling P, Qvortrup K, Koskinen SO, Dela F. Exercise-induced regulation of matrix metalloproteinases in the skeletal muscle of subjects with type 2 diabetes. *Diab Vasc Dis Res*. 2014;11(5):324–334.

62. Souza MV, Leite RD, Souza Lino AD, et al. Resistance training improves body composition and increases matrix metalloproteinase 2 activity in biceps and gastrocnemius muscles of diet-induced obese rats. *Clinics (Sao Paulo)*. 2014;69(4):265–270.
63. Nascimento DD, Navalta JW, Durigan JL, et al. Acute eccentric resistance exercise decreases matrix metalloproteinase activity in obese elderly women. *Clin Physiol Funct Imaging*. 2016;36:039–145.
64. Kwak HB, Kim JH, Joshi K, Yeh A, Martinez DA, Lawler JM. Exercise training reduces fibrosis and matrix metalloproteinase dysregulation in the aging rat heart. *FASEB J*. 2011;25(3):1106–1117.
65. Shon SM, Park JH, Nahrendorf M, et al. Exercise attenuates matrix metalloproteinase activity in preexisting atherosclerotic plaque. *Atherosclerosis*. 2011;216(1):67–73.
66. De Aro AA, Ferrucci DL, Borges FP, Stach-Machado DR, Macedo DV, Pimentel ER. Exhaustive exercise with different rest periods changes the collagen content and MMP-2 activation on the calcaneal tendon. *Anat Rec (Hoboken)*. 2014;297(2):281–288.
67. Nishijima T, Kawakami M, Kita I. A bout of treadmill exercise increases matrix metalloproteinase-9 activity in the rat hippocampus. *Neurosci Lett*. 2015;594:144–149.
68. Rullman E, Rundqvist H, Wagsater D, et al. A single bout of exercise activates matrix metalloproteinase in human skeletal muscle. *J Appl Physiol (1985)*. 2007;102(6):2346–2351.
69. Donley DA, Fournier SB, Reger BL, et al. Aerobic exercise training reduces arterial stiffness in metabolic syndrome. *J Appl Physiol (1985)*. 2014;116(11):1396–1404.

Vascular Health and Risk Management

Publish your work in this journal

Vascular Health and Risk Management is an international, peer-reviewed journal of therapeutics and risk management, focusing on concise rapid reporting of clinical studies on the processes involved in the maintenance of vascular health; the monitoring, prevention and treatment of vascular disease and its sequelae; and the involvement of

metabolic disorders, particularly diabetes. This journal is indexed on PubMed Central and MedLine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/vascular-health-and-risk-management-journal>

Dovepress