

Figure 2 Strong positivity of the intracytoplasmic eosinophilic material following immunohistochemical staining with cystatin C.

reported before.⁸⁻¹⁰ Further studies are necessary to assess the true prevalence of mucous gastroduodenal cells with eosinophilic bodies

in Japanese nationals and to clarify the functional importance of the Cystatin C material retained in these cells.

This study was supported by a grant of the Foundations of Promotions of Cancer Research, Tokyo, Japan, and by a grant of the Swedish Medical Research Council project 03X-0623.

- 1 Helander HF. The cells of the gastric mucosa. *Int Rev Cytol* 1981;70:217-89
- 2 Junqueira LC, Carneiro J. Digestive tract. In: *Basic histology* 4th ed. Los Angeles: Lange Medical, 1983:361-2.
- 3 Owen DA. Normal histology of the stomach. *Am J Surg Pathol* 1986;10:48-61
- 4 Rhodin JAG. The alimentary system. In: *Histology: A text and atlas*. London: Oxford University Press 1974:544-51.
- 5 Rubio CA. Five types of pyloric cells in the antral mucosa of the stomach. *Path Res Pract* 1992;188:157-61.
- 6 Abrahamson M, Olafsson I, Palsdottir A, et al. Structure and expression of the human cystatin C gene. *Biochem J* 1990;268:287-94.
- 7 Rubio CA, Hirota T, Itabashi M, Hirohashi S, Kato Y. Quantitation of gastric intestinal metaplasia by morphology in Japanese patients. *Jpn J Cancer Res* 1992; 83:491-4.
- 8 Rubio CA, Kato Y. Six types of intraepithelial vacuoles in the human gastric mucosa. *Path Res Pract* 1988;183: 321-25.
- 9 Rubio CA, Kato Y, Kitagawa T, Sugano H. A pitfall in the interpretation of intestinal metaplasia of the stomach. *Acta Pathol Microbiol Immunol Scand (A)* 1986; 94:165-6.
- 10 Thompson IW, Day DW, Wright NA. Subnuclear vacuolated mucous cells: a novel abnormality of simple mucin-secreting cells of non-specialised gastric mucosa and Brunner's glands. *Histopathology* 1987;11:1067-81.

J Clin Pathol 1992;45:1120-1121

Reassessment of the rate of fixative diffusion

R D Start, C M Layton, S S Cross, J H F Smith

Abstract

The diffusion of fixatives is slow. Early work using plasma gels and animal tissues showed the distance penetrated by a fixative to be a simple function of the fixation time but this relation has not been established in human tissues. The rates of diffusion into whole human spleens were measured for three primary fixatives over periods ranging from one to 25 days. A positive correlation was demonstrated between penetration distance (mm) and fixation time (hours). The diffusion rates were slower than those in previous studies. These results have possible implications for the handling of surgical specimens.

(*J Clin Pathol* 1992;45:1120-1121)

Tissues are usually fixed before microscopic examination and effective fixation is essential for accurate histological interpretation. Fixation depends not only on the rate at which a fixative reacts with tissue components, but also on the ability of the fixative to diffuse into tissues. An important rate limiting factor within the complex process of fixation is the slow speed of fixative diffusion into tissues.

Our understanding of this subject is largely based on early studies which examined the diffusion of fixatives into plasma gels and animal tissues.¹⁻³ These studies indicated that fixatives obey the laws of diffusion and that the distance penetrated by a fixative depends on a simple function of fixation time. This distance in millimetres is equal to the square root of the fixation time in hours, multiplied by a coefficient of diffusibility (K) for that fixative.^{2,3} More complicated formulae have been proposed,⁴ but no further studies have tested this relation nor examined the diffusion of fixatives into human tissues. This study was designed to examine the rate of fixative diffusion into a human tissue in the context of routine surgical specimens using necropsy material.

Department of
Histopathology,
Northern General
Hospital, Sheffield

R D Start
C M Layton
S S Cross
J H F Smith


Correspondence to
Dr Roger D Start,
Department of Pathology,
University of Sheffield
Medical School, PO Box
596, Beech Hill Road,
Sheffield S10 2UL

Accepted for publication
3 June 1992

Diffusibility coefficients of fixatives

Fixative	K*	r (standard error)
4% formaldehyde solution	0.55 (0.78)	0.88 (p < 0.001)
Saturated picric acid solution	0.40 (0.50)	0.79 (0.001 < p < 0.01)
5% acetic acid solution	0.90 (1.20)	0.82 (0.001 < p < 0.01)

*Figures in parentheses are the values of K from Medawar, 1941²



The predicted diffusion of 4% formaldehyde solution into 10 mm and 50 mm diameter specimens.

Methods

The spleen was chosen because of the relatively homogenous solid structure, convenient size, and availability. Ten human spleens were immersed whole in 4% formaldehyde solution for periods of between one and 25 days after removal at necropsy. The hilar vessels were ligated before immersion and the specimens were kept at a constant temperature of 20°C which approximates to the average constant temperature in our laboratory. Three complete cross-sections were then taken from the central portion of the long axis of each spleen and immersed in 0.02% chromic acid solution for 14 days to macerate any unfixated tissue.¹ This produced a sharp line of demarcation between the fixed outer rim and the rest of the tissue. The depth of penetration, as represented by the thickness of the outer rim, was calculated from the mean of 100 Vernier micrometer measurements on each slice after a simple correction for tissue shrinkage.¹ Each measurement was made in a horizontal plane towards the central point of the slice. Similar fixation experiments were performed using saturated picric acid solution and 5% acetic acid solution. The correlation between the depth of penetration (mm) and the square root of the fixation time (hours) was assessed by linear analysis of raw data from which the correlation coefficient (r) and its standard error based on 10 data points were calculated for each fixative. The coefficients of diffusibility were determined by regression analysis.

Results

The results are presented in the table. A positive correlation was demonstrated between the depth of penetration and the square root of the fixation time for each of the fixatives. The

predicted diffusion of 4% formaldehyde solution into hypothetical, solid 10 mm and 50 mm diameter specimens is shown in the figure.

Discussion

The diffusion of fixatives is usually measured on uniform coagulants such as gelatine or plasma gels.^{2,3} The rates obtained using animal tissue are generally lower because of barriers such as cell membranes.¹ The diffusion rates of the primary fixatives in our human tissue model are slower than in these previous studies and may result from differences in tissue structure together with less controllable factors such as impedance by the splenic capsule. The presence of fixed tissue is also believed to slow subsequent inward diffusion of fixatives.³ Within the limitations of a simple model, our results do not support this theory and are more consistent with earlier studies which showed that fixatives obey the laws of diffusion and appear to neither retard nor facilitate their own penetration.^{2,3}

Primary fixation can affect histological interpretation.⁵ Delayed fixation has been shown to influence the number of observable mitotic figures in tissues,^{6,7} and fixation may influence the immunoreactivity of tissue antigens.⁸ The commonest primary fixative is 4% formaldehyde solution which will penetrate just 2.4 mm in 24 hours. Small specimens measuring less than 10 mm in diameter will fix by simple diffusion. Larger specimens should be thinly sliced, partly dissected or perfused to facilitate rapid and even fixation. Although this study has only investigated the diffusion of a limited number of fixatives into a single human tissue, the observations emphasize the need for prompt and efficient fixation of surgical specimens.

- 1 Underhill BL. The rate of penetration of fixatives. *J R Microsc Soc* 1932;52:113-20.
- 2 Medawar PB. The rate of penetration of fixatives. *J R Microsc Soc* 1941;61:46-57.
- 3 Baker JR. *Principles of biological microtechnique*. London: Methuen, 1960:31-43.
- 4 Dempster WT. Rates of penetrating fixing fluids. *J Am Anat* 1960;107:59-72.
- 5 Hopwood D. Fixation and fixatives. In: Bancroft JD, Stevens A, *Theory and practice of histological techniques*. Edinburgh: Churchill Livingstone, 1982:21-42.
- 6 Graem N, Helweg-Larsen K. Mitotic activity and delay in fixation of tumour tissue. *Acta Pathol Microbiol Scand [A]* 1979;87:375-8.
- 7 Start RD, Flynn MS, Cross SS, Rogers K, Smith JHF. Is the grading of breast carcinoma affected by delay in fixation? *Virchows Arch (Pathol Anat)* 1991;419:475-7.
- 8 Pelstring RJ, Allred DC, Esther RJ, Lampkin SR, Banks PM. Differential antigen preservation during tissue autolysis. *Hum Pathol* 1991;22:237-41.