

RESEARCH PAPER

Cryptotanshinone inhibits oxidized LDL-induced adhesion molecule expression via ROS dependent NF- κ B pathways

Wenwen Zhao, Chuanhong Wu, and Xiuping Chen

State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Macao, China

ABSTRACT

Adhesion molecules, such as intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and E-selectin, play important roles in the initial stage of atherosclerosis. Cryptotanshinone (CPT), a natural compound isolated from *Salvia miltiorrhiza* Bunge, exhibits anti-atherosclerotic activity although the underlying mechanisms remain elusive. In this study, the protective effect of CPT against oxidized low-density lipoprotein (ox-LDL)-induced adhesion molecule expression was investigated in human umbilical vein endothelial cells. Ox-LDL significantly induced ICAM-1, VCAM-1, and E-selectin expression at the mRNA and protein levels but reduced eNOS phosphorylation and NO generation, which were reversed by CPT pretreatment. Sodium nitroprusside, a NO donor, N-acetyl-L-cysteine (NAC), a reactive oxygen species (ROS) scavenger, and BAY117082, a NF- κ B inhibitor, inhibited ox-LDL-induced ICAM-1, VCAM-1, and E-selectin expression. Ox-LDL-induced ROS production was significantly inhibited by CPT and NAC. Furthermore, ox-LDL activated the NF- κ B signaling pathway by inducing phosphorylation of IKK β and I κ B α , promoting the interaction of IKK β and I κ B α , and increasing p65 nuclear translocation, which were significantly inhibited by CPT. In addition, CPT, NAC, and BAY117082 inhibited ox-LDL-induced membrane expression of ICAM-1, VCAM-1, E-selectin, and endothelial-monocyte adhesion and restored eNOS phosphorylation and NO generation. Results suggested that CPT inhibited ox-LDL-induced adhesion molecule expression by decreasing ROS and inhibiting the NF- κ B pathways, which provides new insight into the anti-atherosclerotic mechanism of CPT.

ARTICLE HISTORY

Received 4 May 2015
Revised 31 October 2015
Accepted 7 November 2015

KEYWORDS

cryptotanshinone;
endothelial cells; endothelial
dysfunction; Ox-LDL; ROS

Introduction

Adhesion molecules, such as intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and E-selectin, are transmembrane proteins that mediate adhesion and interactions between cells or a cell and extracellular matrix.¹ Increased expression and activation of these molecules in vascular endothelial cells mediate the attachment of circulating monocytes/leukocytes to the surface of endothelial cells, which is an early step in atherosclerosis development.² P- and E-selectin expressed on the surface of activated endothelial cells bind to carbohydrate ligands on leukocytes.³ The subsequent firm attachment of monocytes to the endothelium is mediated by the interaction of ICAM-1 or VCAM-1 and integrin VLA-4 on the endothelium and monocytes, respectively.² Therefore, the important role of adhesion molecules in the initial step of atherosclerosis makes them potential drug targets for therapeutic intervention of atherosclerosis.⁴

Oxidized low-density lipoprotein (ox-LDL), the oxidized form of LDL caused by reactive oxygen species (ROS), has been implicated as the key risk factor in the pathogenesis of atherosclerosis. Up-regulation of endothelial surface adhesion molecules, such as ICAM-1, VCAM-1, E-, and P-selectin, and enhanced monocyte-endothelial adhesion by ox-LDL have been determined to play important roles in atherosclerosis.^{5–7}

Cryptotanshinone (CPT), an active ingredient isolated from traditional Chinese herb *Salvia miltiorrhiza* Bunge, demonstrates anti-tumor, anti-inflammation, and anti-Alzheimer's disease effects.^{8–10} A previous study reported that CPT decreases ox-LDL-induced secretion of soluble ICAM-1 (sICAM-1) and soluble VCAM-1 (sVCAM-1) in human umbilical vein endothelial cells (HUVECs),¹¹ but the underlying mechanisms remain unclear. In this study, the potential mechanisms of CPT inhibiting ox-LDL-induced adhesion molecule expression were investigated.

Materials and methods

Materials

CPT (>95%) was purchased from Chengdu Pufeide Biological Technology Co., LTD. (China). MTT and 5-(6)-chloromethyl-2',7'-dichlorodihydrofluorescein diacetate (CM-DCFH₂-DA) were obtained from Sigma (USA). Antibodies for NF- κ B p65, phosphorylated eNOS (p-eNOS), eNOS, phosphorylated I κ B α (p-I κ B α), I κ B α , phosphorylated IKK β (p-IKK β), IKK β , Histone H3, ICAM-1, VCAM-1, E-selectin, and GAPDH were acquired from Cell Signaling Technology (USA). Antibodies for Nrf-1 and Nrf-2 were procured from Santa Cruz (USA). Sodium nitroprusside (SNP) and 3-amino,4-aminomethyl-2',7'-difluorescein diacetate (DAF-FM) were purchased from Beyotime (China). Protein A/G PLUS-Agarose was obtained from Santa Cruz (USA). Hoechst 33342 was acquired from Invitrogen (USA). Primers and other real-time PCR-related materials were purchased from Sangon Biotech (China) and TaKaRa Bio Group (Japan). BCA protein kits were purchased from Thermo Fisher (USA). Ox-LDL was obtained from Yiyuan Biotechnology Co., LTD. (China).

Cell culture

Primary HUVECs (Gibco, Life Technologies Corp. C-003-5C) from a newborn were cultured in Ham's F-12K (Kaighn's) medium (Gibco) with endothelial cell growth supplement from bovine neural tissue (ECGS) in a humidified incubator at 37°C and 5% CO₂. Cells at passages 2–4 were used.

Human monocyte cell line (THP-1) obtained from ATCC was cultured in RPMI 1640 medium containing 10% fetal calf serum, 2 mM glutamine, 100 U/mL penicillin, and 100 μ g/mL streptomycin. Cells were cultured at 37°C in a humidified atmosphere of 5% CO₂ and 95% air.

MTT assay

Confluent HUVECs in 96-well microplates were treated with ox-LDL (1.0–20 μ g/mL) or CPT (0–1.0 μ M) for 24 h. After the removal of medium, MTT (5 mg/mL) was added to each well and incubated for another 4 h. The supernatant was then removed, and 100 μ L of DMSO was added to each well to solubilize the formazan crystals. The absorbance was measured at 540 nm using an automated microplate reader (PerkinElmer).

Measurement of ROS

Approximately 1.0×10^6 /well cells were seeded in a 6-well plate overnight. After ox-LDL (10 μ g/mL)

treatment for 2 h with or without CPT (50 nM) pretreatment, cells were incubated with CM-DCFH₂-DA (10 μ M) in the dark at 37°C for 40 min. Cells were washed with PBS and detached with trypsin/EDTA, and cellular fluorescence was analyzed by flow cytometry (Becton Dickinson FACS Canto™).

Immunoprecipitation

The immunoprecipitation assay was performed as described in our previous report¹² with minor revisions. In brief, after ox-LDL treatment with or without CPT pretreatment, cell lysate was collected, and the protein content was determined. The cell lysate was then incubated with anti-IKK β antibody (2 μ g) for 2 h at 4°C, followed by incubation with 20 μ L of protein A/G plus-agarose beads overnight with constant shaking. The beads were washed 3 times with ice-cold radio immunoprecipitation assay buffer, and bound protein was extracted by adding 40 μ L of 2 \times SDS sample buffer and boiling for 5 min. The complex was subjected to SDS-PAGE and visualized by Western blot.

Measurement of NO

Cells were treated with ox-LDL (10 μ g/mL) for 2 h with or without CPT (50 nM) pretreatment. NO production was detected by fluorescent probe DAF-FM (5 μ M) at 37°C for 30 min in the dark. Cells were rinsed with PBS, fixed in 2% paraformaldehyde (v/v) at 4°C for 5 min, and examined by fluorescent microscopy. To quantitatively determine NO formation, cells were trypsinized, and the fluorescence was detected by a flow cytometer.

Endothelial–monocyte adhesion assay

The endothelial–monocyte adhesion assay was performed as described in our previous report.¹³ In brief, THP-1 cells were labeled with Hoechst 33342 (10 μ g/mL) for 30 min in the dark. The labeled THP-1 cells were incubated with ox-LDL-treated endothelial cells (with or without CPT pretreatment) for 3 h at 37°C. The non-adherent cells were removed with PBS. Endothelial–monocyte adhesion was observed by fluorescent microscopy.

Immunofluorescence assay

Cells (5×10^4 cells/well) were seeded on glass slides in 12-well plates. After ox-LDL treatment (with or without CPT pretreatment), the slides were fixed with 4% PFA

for 30 min. The slides were then permeabilized with PBS-T (containing 0.1% Triton x-100 in PBS solution) and blocked with PBS-B (containing 4% BSA in PBS solution). After incubation with the primary (1:1000) and secondary antibodies (1:2000), the cells were stained with Hoechst 33342 in the dark for 30 min. The location and expression of adhesion molecule were observed by fluorescence microscopy.

Real-time RT-PCR

Total RNA was extracted with TRIzol Reagent. About 2 μ g RNA was reverse-transcribed into cDNA using First Strand cDNA Synthesis Kit (Toyobo, Japan). Real-time PCR was performed using SYBR Green PCR reagents (Applied Biosystems). The specific primers are as follows: for ICAM-1, 5'-GGC TGG AGC TGT TTG AGA AC-3' (forward) and 5'-ACT GTG GGG TTC AAC CTC TG' (reverse); for VCAM-1, 5'-AAA AGC GGA GAC AGG AGA CA-3' (forward) and 5'-AGC ACG AGA AGC TCA GGA GA' (reverse); for E-selectin, 5'-TCT CTC AGC TCT CAC TTG-3' (forward) and 5'-TTC TTC TTG CTG CAC CTC T' (reverse); and for GAPDH, 5'-CGA GAT CCC TCC AAA ATC AA-3' (forward) and 5'-TTC ACA CCC ATG GAC GAA CAT-3' (reverse). The expression levels of target genes were determined by normalizing to GAPDH expression.

Western blot

Total proteins were extracted from treated cells, and the protein contents were quantified by BCA Protein Assay Kit. About 30 μ g of proteins was subjected to SDS-PAGE and transferred onto a PVDF membrane. After blocking with 5% non-fat milk in TBST (20 mM Tris-HCl, 500 mM NaCl, and 0.1% Tween 20) at room temperature for 2 h, membranes were incubated with

primary antibodies (1:2000) overnight at 4°C followed by secondary antibodies (1:10,000). The protein-antibody complexes were detected by an ECL Advance Western Blot Detection Kit. The intensity of the bands was quantitated with QuantityOne software (Bio-Rad).

Statistical analysis

Data were expressed as the mean \pm SD from at least 3 separate experiments. The differences between groups were analyzed using Prism 5.0 (Graph Pad Software Inc., San Diego, CA, USA) by one-way ANOVA, followed by Student-Newman-Keuls test. $p < 0.05$ was considered statistically significant.

Results

CPT inhibited ox-LDL-induced adhesion molecule expression and restored NO production

Given that both ox-LDL and CPT were cytotoxic to a panel of cells, we first tested their effect on HUVEC viability. Both ox-LDL and CPT were cytotoxic to endothelial cells. However, ox-LDL at 10 μ g/mL and CPT at 50 nM revealed no obvious cytotoxicity (Figs. 1A and B). Ox-LDL remarkably induced ICAM-1, VCAM-1, and E-selectin expression at both protein and mRNA levels, which were significantly reversed by CPT pretreatment (Figs. 2A and B). Furthermore, ox-LDL treatment decreased the protein expression of *p*-eNOS and intracellular NO production, which were also restored by CPT pretreatment (Figs. 2C, D, and E). SNP, a NO donor, also dramatically reversed the ox-LDL-induced down-regulation of *p*-eNOS expression (Fig. 2F) and upregulation of ICAM-1, VCAM-1, and E-selectin expression (Fig. 2G). CPT alone showed no effect on the protein expression of eNOS, *p*-eNOS, ICAM-1, VCAM-1, and E-selectin (Supplementary Fig. 1).




Figure 1. Cytotoxic effect of ox-LDL and CPT on HUVECs. Confluent HUVECs (1.0×10^4 cells/well) in 96-well plates were exposed to different concentrations of ox-LDL or CPT for 24 h, and the cell viability was examined with MTT assay. * $p < 0.05$, ** $p < 0.01$ vs. Cont. Cont, control group.




Figure 2. CPT inhibited ox-LDL-induced adhesion molecule expression and restored NO production. HUVECs were treated with ox-LDL (10 $\mu\text{g}/\text{mL}$) for 24 h with or without CPT pretreatment (50 nM) for 1 h. The protein (A, C) and mRNA (B) expression levels were determined by Western blot and RT-PCR, respectively. HUVECs were incubated with ox-LDL for 2 h with or without CPT (50 nM) pretreatment for 1 h. Intracellular NO production was determined by a flow cytometer (D) and detected by fluorescence microscope (60 \times) (E) after DAF-FM staining. HUVECs were stimulated with ox-LDL (10 $\mu\text{g}/\text{mL}$) for 24 h with or without CPT (50 nM) or SNP (1 mM) pretreatment, and the protein expression was measured by Western blot (F and G). SNP, sodium nitroprusside. ** $p < 0.01$, *** $p < 0.001$ vs. Cont. Cont, control group; ## $p < 0.01$, ### $p < 0.001$ vs. ox-LDL group.

CPT inhibited ox-LDL-induced adhesion molecule expression and restored NO production via reducing ROS

Compared with the untreated group, ox-LDL treatment induced a significant right shift of ROS peaks, suggesting increased ROS formation (Fig. 3A). CPT and NAC pretreatment reversed the right shift of ROS peaks, indicating the inhibition of ROS. Statistical analysis demonstrated that ox-LDL induced more than twofold of ROS production, which was significantly suppressed by CPT or NAC pretreatment in HUVECs (Fig. 3B). Furthermore, NAC pretreatment, similar to CPT, dramatically inhibited ox-LDL-induced protein (Fig. 3C) and mRNA (Fig. 3D) expression levels of ICAM-1, VCAM-1, and E-selectin. In addition, ox-LDL-induced decreased expression of *p*-eNOS and NO production were also partly restored by CPT and NAC (Figs. 3E, F, and G).

CPT inhibited ox-LDL-induced adhesion molecule expression and restored NO production via NF- κ B signaling

Ox-LDL treatment significantly increased the nuclear expression of NF- κ B p65, which was inhibited by CPT (Fig. 4A). Ox-LDL displayed no effect on either Nrf-1 or Nrf-2 protein expression. The IKK kinase complex, which is essentially composed of 2 kinases, namely, IKK α and IKK β , is the core element of the NF- κ B cascade. IKK β plays a critical role in the phosphorylation of I κ B α and subsequent activation of NF- κ B. Thus, the effect of CPT on IKK β was determined. Ox-LDL induced the phosphorylation of IKK β and I κ B α , which was inhibited by CPT (Fig. 4B). Furthermore, immunoprecipitation assay showed that ox-LDL enhanced the binding of IKK β with I κ B α , which was also decreased by CPT (Fig. 4C). Both CPT and BAY117082 (a NF- κ B inhibitor) obviously reversed ox-LDL-induced ICAM-1, VCAM-1, and E-selectin expression (Figs. 4D and E), as well as restored *p*-eNOS expression and NO production (Figs. 4F and G). In addition, compared with SNP treated group, combination with BAY117082 further decreased the expression ICAM-1, VCAM-1, and E-selectin induced by ox-LDL (Fig. 4H).

CPT inhibited ox-LDL-induced membrane adhesion molecule expression and endothelial-monocyte adhesion

Endothelial cells expressed/secreted 2 forms of ICAM-1 and VCAM-1: membranous ICAM-1 and sICAM-1 and membranous VCAM-1 and sVCAM-1.¹⁴⁻¹⁶ Membranous

ICAM-1 and VCAM-1 and transmembrane protein E-selectin mediate adhesion and interactions between cells or a cell and extracellular matrix¹⁷; thus, their expression levels were determined. Ox-LDL treatment induced intensive membrane fluorescence, suggesting the increased membrane expression of ICAM-1, VCAM-1, and E-selectin, which was significantly inhibited by CPT, NAC, or BAY117082 pretreatment (Fig. 5A). Furthermore, ox-LDL stimulation increased the adhesion of THP-1 cells to HUVECs, which was also dramatically inhibited by CPT, NAC, or BAY117082 pretreatment (Fig. 5B).

Discussion

Ox-LDL plays a crucial role in the initiation and progression of atherosclerosis by inducing endothelial activation and dysfunction, promoting the proliferation and migration of SMCs, facilitating the formation of foam cells, and enhancing endothelial-monocyte adhesion.¹⁸⁻²⁰ Accumulated studies revealed that a panel of natural products, such as curcumin, berberine, and tanshinone IIA,²¹⁻²³ can inhibit ox-LDL-induced adhesion molecule expression. A previous study observed that CPT improves ox-LDL-induced adhesion molecule expression,¹¹ but the underlying mechanisms remain elusive. In the current study, we revealed that CPT inhibited ox-LDL-induced adhesion molecule expression via the ROS-dependent pathway mediated by NF- κ B.

MTT assay showed that CPT demonstrated significant cytotoxicity to HUVECs with a high concentration (>0.1 μ M). Thus, its effect was tested at non-cytotoxic concentrations. Endothelial dysfunction (ED) contributes to a panel of cardiovascular diseases, including hypertension, atherosclerosis, and coronary artery disease. The characteristics of ED include a lack of NO such as inhibiting and decreasing NO bioavailability²⁴ and/or overexpression of adhesion molecules, such as ICAM-1, VCAM-1, E-, and P-selectin.^{25,26} A previous study demonstrated that CPT inhibits ox-LDL-induced sICAM-1 and sVCAM-1 secretion in HUVEC culture medium.¹¹ In the present study, we determined that ox-LDL induced the up-regulation of ICAM-1, VCAM-1, and E-selectin expression at both protein and mRNA levels and decreased NO production, thereby suggesting that ox-LDL caused ED. CPT pretreatment reversed these alterations, indicating that CPT protected ox-LDL-induced ED.

ROS is a major target of oxidant stress, playing a critical role in the pathophysiology of several vascular diseases and disorders.²⁷ Ox-LDL has been established to induce intracellular ROS formation in endothelial cells.^{28,29} Consistent with these reports, elevated ROS levels were observed after ox-LDL exposure under our experimental conditions. Similar to our recent report




Figure 3. CPT inhibited ox-LDL-induced adhesion molecule expression and restored NO production via ROS. HUVECs were pretreated with CPT (50 nM) or NAC (5 mM) for 1 h, followed by ox-LDL (10 μ g/mL) treatment for 2 h. ROS generation was measured by flow cytometry (A, B). HUVECs were pretreated with CPT or NAC for 1 h and then treated with ox-LDL for 24 h. The protein expression was measured by Western blot (C, E), and mRNA expression was determined by RT-PCR (D). Intracellular NO production was determined by a flow cytometer (F) and detected with fluorescence microscope (60 \times) (G) after DAF-FM staining. ** $p < 0.01$, *** $p < 0.001$ vs Cont. Cont, control group; ## $p < 0.01$, ### $p < 0.001$ vs. ox-LDL group.




Figure 4. CPT inhibited ox-LDL-induced adhesion molecule expression and restored NO production via NF- κ B. HUVECs were treated with ox-LDL (10 μ g/mL) for 24 h with or without CPT (50 nM) SNP (1 mM) or BAY117082 (10 μ M) pretreatment. The protein expression was measured by Western blot (A–D, F, H), and mRNA expression was determined by RT-PCR (E). Intracellular NO production was detected with fluorescence microscope after DAF-FM staining (60 \times) (G). ** p < 0.01, *** p < 0.001 vs. Cont. Cont, control group; ### p < 0.01 vs. ox-LDL group; # \S < 0.05 vs. SNP + ox-LDL group.

that CPT inhibited TNF- α -induced ROS formation in HUVECs,³⁰ CPT also dramatically inhibited ox-LDL-induced ROS formation in endothelial cells. Furthermore, NAC, a ROS scavenger, exhibited a similar effect on ox-LDL-induced adhesion molecule expression and ROS production. Collectively, these data suggested that CPT inhibited ox-LDL-induced adhesion molecule expression in a ROS-dependent manner.

Several lines of evidence indicated that ox-LDL suppresses NF- κ B and incapacitates the protection from apoptosis in activated endothelial cells,^{28,31} whereas other studies oppositely highlighted that ox-LDL induces NF- κ B activation and the subsequent expression of pro-inflammatory genes. Nrf-1 and Nrf-2 are 2 important oxidative sensitive transcription factors involved in regulating the expression of adhesion molecules.^{32–35} We found that ox-LDL had no influence on either




Figure 5. CPT inhibited ox-LDL-induced membrane expression of adhesion molecules and THP-1-HUVEC adhesion. HUVECs were treated with ox-LDL (10 $\mu\text{g}/\text{mL}$) for 24 h with or without CPT (50 nM), BAY117082 (10 μM), or NAC (5 mM) pretreatment. The membrane expression of adhesion molecules was examined by immunofluorescence assay (60 \times) (A). HUVECs were treated with ox-LDL (10 $\mu\text{g}/\text{mL}$) for 24 h with or without CPT (50 nM), BAY117082 (10 μM), or NAC (5 mM) pretreatment. The cells were then incubated with labeled THP-1 cells for 3 h. The attached THP-1 cells were visualized by inverted fluorescent microscopy (20 \times) (B).

Nrf-1 or Nrf-2 protein expression, but it dramatically increased the nuclear expression of p65, suggesting that ox-LDL induced p65 nuclear translocation. Ox-LDL induced the phosphorylation of IKK β and I κ B α and enhanced their interactions, indicating that ox-LDL activated the NF- κ B pathway by promoting p65 nuclear translocation mediated by enhancing IKK β -

I κ B α interactions and their phosphorylation.³⁶ The inhibitory effect of CPT suggested that CPT regulated IKK β and I κ B α , the upper stream factors of the NF- κ B pathways. Thus, potential targets for CPT may exist in the NF- κ B system. The inhibitory effect of BAY117082, a NF- κ B inhibitor, on ox-LDL-induced adhesion molecule expression further confirmed the




Figure 6. Mechanisms underlying CPT inhibited ox-LDL-induced adhesion molecule expression in HUVECs.

involvement of NF- κ B. Collectively, these data indicated that CPT suppressed ox-LDL-induced NF- κ B activation and the subsequent expression of adhesion molecules.

Several studies suggested that NO can suppress the expression of VCAM-1, ICAM-1, and E-selectin in response to various pro-inflammatory cytokines.³⁷ In the present study, we found that SNP, a NO donor, dramatically reversed the ox-LDL-induced expression of adhesion molecules, eNOS phosphorylation, and NO generation. Furthermore, the ox-LDL-induced downregulation of eNOS phosphorylation and decrease of NO generation were reversed by NAC and BAY117082. These data proved that NO negatively regulated adhesion molecule expression mediated by NF- κ B in response to ox-LDL, which formed a branch for the regulation of adhesion molecules. The decreased intracellular NO partly resulted from the decreased expression of eNOS because eNOS was the main source of NO production in endothelial cells. However, this phenomenon might also be due to the overproduction of ROS, because certain types of ROS, especially superoxide anion, can react with NO to produce highly reactive and cytotoxic products, such as peroxynitrite (ONOO⁻) and peroxynitrous.^{29,38} Thus, the inhibitory effect of CPT on ox-LDL-induced adhesion molecule expression might be due to its direct effect on the NF- κ B system or/and indirectly mediated by eNOS/NO. To make clear whether NO was located in NF- κ B pathway or worked independent of NF- κ B pathway, combined treatment with SNP and BAY117082 was performed. Decreased expression of adhesion molecules in response to SNP was further decreased by BAY117082. This suggested that NO was located in downstream of NF- κ B. However, the exact role of eNOS/NO in the regulation of adhesion molecule expression in response to ox-LDL requires further investigation.

Monocyte adhesion to endothelial cells was an important early event in atherogenesis, which was partly controlled by expression of adhesion molecules on the endothelial cell surface.^{39,40} The immunofluorescence results indicated that ox-LDL induced the increased membrane protein expression of ICAM-1, VCAM-1, and E-selectin. Furthermore, ox-LDL induced enhanced THP-1 cell adhesion to endothelial cells, suggesting that these adhesion molecules played an important role in this process. The inhibitory effect of NAC and BAY117082 provided further evidence for the involvement of ROS and NF- κ B in ox-LDL-induced adhesion molecule expression. Thus, the inhibitory effect of CPT indicated that CPT may enhance atherosclerosis treatment in the early stage.

In conclusion, the present work provided evidence that CPT, a natural compound, inhibited adhesion

molecule expression upon ox-LDL stimulation by regulating the NF- κ B pathway mediated through decreasing ROS formation (Fig. 6). This data provides new insights into the beneficial effect of CPT in cardiovascular diseases.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

This study was supported by the National Natural Science Foundation of China (No 81160048), the Science and Technology Development Fund, Macao S.A.R (FDCT) (No. 021/2012/A1), and the Research Fund of University of Macau (MRG007/CXP/2013/ICMS).

References

- [1] Davies MJ, Gordon JL, Gearing AJH, Pigott R, Woolf N, Katz D, Kyriakopoulos A. The Expression of the Adhesion Molecules Icam-1, Vcam-1, Pecam, and E-Selectin in Human Atherosclerosis. *J Pathol* 1993; 171:223–9; PMID:7506307; <http://dx.doi.org/10.1002/path.1711710311>
- [2] Ling S, Nheu L, Komesaroff PA. Cell adhesion molecules as pharmaceutical target in atherosclerosis. *Mini Rev Med Chem* 2012; 12:175–83; PMID:22070689; <http://dx.doi.org/10.2174/138955712798995057>
- [3] Varki A. Selectin ligands. *Proc Natl Acad Sci U S A* 1994; 91:7390–7; PMID:7519775; <http://dx.doi.org/10.1073/pnas.91.16.7390>
- [4] Lutters BC, Leeuwenburgh MA, Appeldoorn CC, Moleenaar TJ, Van Berkel TJ, Biessen EA. Blocking endothelial adhesion molecules: a potential therapeutic strategy to combat atherogenesis. *Curr Opin Lipidol* 2004; 15:545–52; PMID:15361790; <http://dx.doi.org/10.1097/00041433-200410000-00008>
- [5] Frostegard J, Wu R, Haegerstrand A, Patarroyo M, Lefvert AK, Nilsson J. Mononuclear leukocytes exposed to oxidized low density lipoprotein secrete a factor that stimulates endothelial cells to express adhesion molecules. *Atherosclerosis* 1993; 103:213–9; PMID:7507327; [http://dx.doi.org/10.1016/0021-9150\(93\)90264-U](http://dx.doi.org/10.1016/0021-9150(93)90264-U)
- [6] Kim JA, Territo MC, Wayner E, Carlos TM, Parhami F, Smith CW, Haberland ME, Fogelman AM, Berliner JA. Partial characterization of leukocyte binding molecules on endothelial cells induced by minimally oxidized LDL. *Arterioscler Thromb* 1994; 14:427–33; PMID:8123647; <http://dx.doi.org/10.1161/01.ATV.14.3.427>
- [7] Jeng JR, Chang CH, Shieh SM, Chiu HC. Oxidized low-density lipoprotein enhances monocyte-endothelial cell binding against shear-stress-induced detachment. *Biochimica et Biophysica Acta* 1993; 1178:221–7; PMID:7688576; [http://dx.doi.org/10.1016/0167-4889\(93\)90013-F](http://dx.doi.org/10.1016/0167-4889(93)90013-F)
- [8] Chen W, Lu Y, Chen G, Huang S. Molecular evidence of cryptotanshinone for treatment and prevention of

- human cancer. *Anti-Cancer Agents Med Chem* 2013; 13:979–87; PMID:23272908; <http://dx.doi.org/10.2174/18715206113139990115>
- [9] Tang S, Shen XY, Huang HQ, Xu SW, Yu Y, Zhou CH, Chen SR, Le K, Wang YH, Liu PQ. Cryptotanshinone suppressed inflammatory cytokines secretion in RAW264.7 macrophages through inhibition of the NF-kappaB and MAPK signaling pathways. *Inflammation* 2011; 34:111–8; PMID:20490642; <http://dx.doi.org/10.1007/s10753-010-9214-3>
- [10] Yoo KY, Park SY. Terpenoids as potential anti-Alzheimer's disease therapeutics. *Molecules (Basel, Switzerland)* 2012; 17:3524–38; PMID:22430119; <http://dx.doi.org/10.3390/molecules17033524>
- [11] Ang KP, Tan HK, Selvaraja M, Kadir AA, Somchit MN, Akim AM, Zakaria ZA, Ahmad Z. Cryptotanshinone attenuates in vitro oxLDL-induced pre-lesional atherosclerotic events. *Planta Med* 2011; 77:1782–7; PMID:21614753; <http://dx.doi.org/10.1055/s-0030-1271119>
- [12] Ou HC, Song TY, Yeh YC, Huang CY, Yang SF, Chiu TH, Tsai KL, Chen KL, Wu YJ, Tsai CS. EGCG protects against oxidized LDL-induced endothelial dysfunction by inhibiting LOX-1-mediated signaling. *J Appl Physiol* 2010; 108:1745–56; PMID:20203069; <http://dx.doi.org/10.1152/jappphysiol.00879.2009>
- [13] Zhao W, Ma G, Chen X. Lipopolysaccharide induced LOX-1 expression via TLR4/MyD88/ROS activated p38MAPK-NF-kappaB pathway. *Vasc Pharmacol* 2014; 63:162–72; PMID:25135647; <http://dx.doi.org/10.1016/j.vph.2014.06.008>
- [14] Budnik A, Grewe M, Gyufko K, Krutmann J. Analysis of the production of soluble ICAM-1 molecules by human cells. *Exp Hematol* 1996; 24:352–9; PMID:8641365.
- [15] Calabresi PA, Tranquill LR, Dambrosia JM, Stone LA, Maloni H, Bash CN, Frank JA, McFarland HF. Increases in soluble VCAM-1 correlate with a decrease in MRI lesions in multiple sclerosis treated with interferon beta-1b. *Ann Neurol* 1997; 41:669–74; PMID:9153530; <http://dx.doi.org/10.1002/ana.410410517>
- [16] Pigott R, Dillon LP, Hemingway IH, Gearing AJH. Soluble Forms of E-Selectin, Icam-1 and Vcam-1 Are Present in the Supernatants of Cytokine Activated Cultured Endothelial-Cells. *Biochem Biophys Res Commun* 1992; 187:584–9; PMID:1382417; [http://dx.doi.org/10.1016/0006-291X\(92\)91234-H](http://dx.doi.org/10.1016/0006-291X(92)91234-H)
- [17] Ala A, Dhillon AP, Hodgson HJ. Role of cell adhesion molecules in leukocyte recruitment in the liver and gut. *Int J Exp Pathol* 2003; 84:1–16; PMID:12694483; <http://dx.doi.org/10.1046/j.1365-2613.2003.00235.x>
- [18] Li D, Mehta JL. Antisense to LOX-1 inhibits oxidized LDL-mediated upregulation of monocyte chemoattractant protein-1 and monocyte adhesion to human coronary artery endothelial cells. *Circulation* 2000; 101:2889–95; PMID:10869259; <http://dx.doi.org/10.1161/01.CIR.101.25.2889>
- [19] Kaperonis EA, Liapis CD, Kakisis JD, Dimitroulis D, Papavassiliou VG. Inflammation and atherosclerosis. *Eur J Vasc Endovasc Surg* 2006; 31:386–93; PMID:16359887; <http://dx.doi.org/10.1016/j.ejvs.2005.11.001>
- [20] Ou HC, Chou FP, Lin TM, Yang CH, Sheu WH. Protective effects of honokiol against oxidized LDL-induced cytotoxicity and adhesion molecule expression in endothelial cells. *Chem Biol Interact* 2006; 161:1–13; PMID:16580656; <http://dx.doi.org/10.1016/j.cbi.2006.02.006>
- [21] Xu S, Liu Z, Huang Y, Le K, Tang F, Huang H, Ogura S, Little PJ, Shen X, Liu P. Tanshinone II-A inhibits oxidized LDL-induced LOX-1 expression in macrophages by reducing intracellular superoxide radical generation and NF-kappaB activation. *Transl Res* 2012; 160:114–24; PMID:22677363; <http://dx.doi.org/10.1016/j.trsl.2012.01.008>
- [22] Huang Z, Cai X, Li S, Zhou H, Chu M, Shan P, Huang W. Berberine attenuated monocyte adhesion to endothelial cells induced by oxidized low density lipoprotein via inhibition of adhesion molecule expression. *Mol Med Rep* 2013; 7:461–5; PMID:23241897.
- [23] Hasan ST, Zingg JM, Kwan P, Noble T, Smith D, Meydani M. Curcumin modulation of high fat diet-induced atherosclerosis and steatohepatosis in LDL receptor deficient mice. *Atherosclerosis* 2014; 232:40–51; PMID:24401215; <http://dx.doi.org/10.1016/j.atherosclerosis.2013.10.016>
- [24] Gryglewski RJ, Palmer RM, Moncada S. Superoxide anion is involved in the breakdown of endothelium-derived vascular relaxing factor. *Nature* 1986; 320:454–6; PMID:3007998; <http://dx.doi.org/10.1038/320454a0>
- [25] Salvia Eotlif, vivo mBoaamai, inflammatory responses. *Phytotherapy Research* Kubes P, Suzuki M, Granger DN. Nitric oxide: an endogenous modulator of leukocyte adhesion. *Proc Natl Acad Sci U S A* 1991; 88:4651–5; PMID:1675786; <http://dx.doi.org/10.1073/pnas.88.11.4651>
- [26] Papaharalambus CA, Griendling KK. Basic mechanisms of oxidative stress and reactive oxygen species in cardiovascular injury. *Trends Cardiovasc Med* 2007; 17:48–54; PMID:17292046; <http://dx.doi.org/10.1016/j.tcm.2006.11.005>
- [27] Szocs K. Endothelial dysfunction and reactive oxygen species production in ischemia/reperfusion and nitrate tolerance. *Gen Physiol Biophys* 2004; 23:265–95; PMID:15638116.
- [28] Cominacini L, Pasini AF, Garbin U, Davoli A, Tosetti ML, Campagnola M, Rigoni A, Pastorino AM, Lo Cascio V, Sawamura T. Oxidized low density lipoprotein (ox-LDL) binding to ox-LDL receptor-1 in endothelial cells induces the activation of NF-kappaB through an increased production of intracellular reactive oxygen species. *J Biol Chem* 2000; 275:12633–8; PMID:10777555; <http://dx.doi.org/10.1074/jbc.275.17.12633>
- [29] Cominacini L, Rigoni A, Pasini AF, Garbin U, Davoli A, Campagnola M, Pastorino AM, Lo Cascio V, Sawamura T. The binding of oxidized low density lipoprotein (ox-LDL) to ox-LDL receptor-1 reduces the intracellular concentration of nitric oxide in endothelial cells through an increased production of superoxide. *J Biol Chem* 2001; 276:13750–5; PMID:11278710.
- [30] Potenza MA, Marasciulo FL, Chieppa DM, Brigiani GS, Formoso G, Quon MJ, Montagnani M. Insulin resistance in spontaneously hypertensive rats is associated with endothelial dysfunction characterized by imbalance between NO and ET-1 production. *Am J*

- Physiol Heart Circulat Physiol 2005; 289:H813–22; PMID:15792994; <http://dx.doi.org/10.1152/ajpheart.00092.2005>
- [31] Heermeier K, Leicht W, Palmethofer A, Ullrich M, Wanner C, Galle J. Oxidized LDL suppresses NF-kappaB and overcomes protection from apoptosis in activated endothelial cells. *J Am Soc Nephrol* 2001; 12:456–63; PMID:11181793.
- [32] Collins T, Read MA, Neish AS, Whitley MZ, Thanos D, Maniatis T. Transcriptional regulation of endothelial cell adhesion molecules: NF-kappa B and cytokine-inducible enhancers. *FASEB J* 1995; 9:899–909; PMID:7542214
- [33] Foncea R, Carvajal C, Almarza C, Leighton F. Endothelial cell oxidative stress and signal transduction. *Biol Res* 2000; 33:89–96; PMID:15693275; <http://dx.doi.org/10.4067/S0716-97602000000200008>
- [34] Kim CK, Cho DH, Lee KS, Lee DK, Park CW, Kim WG, Lee SJ, Ha KS, Goo Taeg O, Kwon YG, et al. Ginseng Berry Extract Prevents Atherogenesis via Anti-Inflammatory Action by Upregulating Phase II Gene Expression. *Evid Based Complement Alternat Med* 2012; 2012:490301; PMID:23243449.
- [35] Drabarek B, Dymkowska D, Szczepanowska J, Zablocki K. TNFalpha affects energy metabolism and stimulates biogenesis of mitochondria in EA.hy926 endothelial cells. *Int J Biochem Cell Biol* 2012; 44:1390–7; PMID:22687752; <http://dx.doi.org/10.1016/j.biocel.2012.05.022>
- [36] Oeckinghaus A, Ghosh S. The NF-kappaB family of transcription factors and its regulation. *Cold Spring Harb Perspect Biol* 2009; 1:a000034; PMID:20066092; <http://dx.doi.org/10.1101/cshperspect.a000034>
- [37] Liao JK. Linking endothelial dysfunction with endothelial cell activation. *J Clin Invest* 2013; 123:540–1; PMID:23485580; <http://dx.doi.org/10.1172/JCI66843>
- [38] Vaziri ND, Ni ZM, Oveisi F, Liang KH, Pandian R. Enhanced nitric oxide inactivation and protein nitration by reactive oxygen species in renal insufficiency. *Hypertension* 2002; 39:135–41; PMID:11799092; <http://dx.doi.org/10.1161/hy0102.100540>
- [39] Ross R. The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature* 1993; 362:801–9; PMID:8479518; <http://dx.doi.org/10.1038/362801a0>
- [40] Jang Y, Lincoff AM, Plow EF, Topol EJ. Cell adhesion molecules in coronary artery disease. *J Am Coll Cardiol* 1994; 24:1591–601; PMID:7963103; [http://dx.doi.org/10.1016/0735-1097\(94\)90162-7](http://dx.doi.org/10.1016/0735-1097(94)90162-7)