



Published in final edited form as:

Bio Protoc. 2012 November 20; 2(22): .

Mouse ESC Differentiation to Nkx2.1+ Lung and Thyroid Progenitors

Tyler A. Longmire¹, Laertis Ikonomou², and Darrell N. Kotton^{2,*}

¹Center for Regenerative Medicine, Boston University and Boston Medical Center, Boston, USA

²Center for Regenerative Medicine, Boston University and Boston Medical Center, Boston, USA

Abstract

The *de novo* derivation of lung progenitors from pluripotent stem cells provides the opportunity to model early lung development *in vitro* and allows easy access to cells for tissue engineering or basic cell biology studies. This detailed protocol allows the generation of lung and thyroid progenitors from mouse embryonic stem cell (ESC) or induced pluripotent stem cell (iPSC) lines. When used together with a published Nkx2.1-GFP knock-in ESC line, the protocol allows tracking and purification of lung and thyroid progenitors by sorting on the GFP reporter based on the induction of the earliest known marker of lung and thyroid cell fate, Nkx2.1. After sorting, a pure population of Nkx2.1+ cells can then be replated for further expansion, differentiation, and maturation in culture in serum-free conditions.

Materials and Reagents

1. Mouse ESCs or iPSCs carrying a GFP reporter knocked in to the Nkx2.1 locus (Nkx2.1-GFP ESCs) (Longmire *et al.*, 2012)
2. 1× 0.05% Trypsin-EDTA (Life Technologies, Gibco®, catalog number: 25300-054)
3. Defined Fetal Bovine Serum (Hyclone, catalog number: SH30070.03)
4. IMDM powder (Life Technologies, Invitrogen™, catalog number: 12200-036)
5. NaHCO₃ (Sigma-Aldrich, catalog number: S-5761)
6. Pen/Strep (Life Technologies, Invitrogen™, catalog number: 15140-148) (10,000 U Penicillin and 10 mg Streptomycin per ml)
7. Cellgro water (VWR, catalog number: 45000-672)
8. Ham's F-12 (Cellgro, catalog number: 10-080-CV)
9. B-27 supplement with RA (Life Technologies, Invitrogen™, catalog number: 17504-044)

*For correspondence: dkotton@bu.edu.

10. N-2 supplement (Life Technologies, Invitrogen™, catalog number: 17502-048)
11. BSA Fraction V 7.5% in PBS (Life Technologies, Invitrogen™, catalog number: 15260-037)
12. 1 -thioglycerol (MTG) (Sigma, M6145-25ml)
13. 200 mM L-Glutamine (Life Technologies, Invitrogen™, catalog number: 25030-081)
14. Ascorbic Acid (Sigma-Aldrich, catalog number: A4544-25G)
15. 1M HEPES (Gibco, 15630-080)
16. CaCl₂ (Sigma-Aldrich, catalog number: C4901)
17. BSA (Sigma-Aldrich, catalog number: A9418-10G)
18. 100× ITS supplement (BD Biosciences, catalog number: 354352)
19. mNoggin (R&D Systems, catalog number: 1967-NG-025)
20. SB431542 (Sigma-Aldrich, catalog number: S4317)
21. mWnt3a (R&D Systems, catalog number: 1324-WN-010)
22. hBMP4 (R&D Systems, catalog number: 314-BP-050)
23. hEGF (R&D Systems, catalog number: 236-EG-01M)
24. mFGF2 (R&D Systems, catalog number: catalog number: 3139-FB-025)
25. mFGF7 (R&D Systems, catalog number: 5028-KG-025)
26. hFGF10 (R&D Systems, catalog number: 345-FG-025)
27. Heparin sodium salt (Sigma-Aldrich, catalog number: H4784-250mg)
28. Dexamethasone (Sigma-Aldrich, catalog number: D4902)
29. 8 - Br - cAMP (Sigma-Aldrich, catalog number: B7880)
30. IBMX (Sigma-Aldrich, catalog number: I5879)
31. DMSO, Hybri-Max (Sigma-Aldrich, catalog number: D2650)
32. Ethanol (Sigma-Aldrich, catalog number: E7023)
33. Activin A (R&D Systems, catalog number: 338-AC)
34. PBS (Life Technologies, Gibco®, catalog number: 14190-250)
35. 0.1% Gelatin in ultrapure water (EMD Millipore, catalog number: ES-006-B)
36. Cxcr4 Antibody: APC Rat anti-mouse CD184 (Cxcr4) (BD-Pharmigen, catalog number: 558644)

37. cKit Antibody: PE Rat anti-mouse CD117 (cKit) (BD-Pharmigen, catalog number: 553355)
38. APC Isotype: APC Rat IgG_{2b}, K (BD-Pharmigen, catalog number: 553991)
39. PE Isotype: PE Rat IgG_{2b}, K (BD-Pharmigen, catalog number: 553989)
40. IMDM (see Recipes)
41. Serum free differentiation medium (SFD) (see Recipes)
42. Complete serum free differentiation medium (cSFDM) (see Recipes)
43. BASE medium for DCI+K (see Recipes)
44. Anteriorization medium (see Recipes)
45. Ventralization medium (see Recipes)
46. DCI+K medium (see Recipes)
47. Preparation of 10× cAMP+IBMX stock (see Recipes)

Equipment

1. P100 Petri dish (100 mm × 15 mm Bacteriological Petri Dish, nontreated polystyrene, BD Falcon™, catalog number: 351029)
2. P150 Petri dish (150 mm × 15 mm Bacteriological Petri Dish, nontreated polystyrene, BD Falcon™, catalog number: 351058)
3. 12 × 75 mm, 5 ml polystyrene round bottom test tube with a cell strainer cap (BD, 352235)
4. 1.5-ml Eppendorf Snap-Cap microcentrifuge tubes (Thermo Fisher Scientific, catalog number: 05-402-25)
5. Centrifuges
6. LSRII flow cytometer
7. 0.22µm filter

Procedure

- A. Timeline
 1. Timepoint: 0 h
 - a. Nkx2-1-GFP mouse ESCs or iPSCs are cultured in 2i_LIF (serum-free, feeder-free) conditions. For each experiment, a new vial of passage 23 is thawed and cells are used after two passages.

- 6.** Timepoint: Day 22
 - a.** Remove media and rinse with PBS.
 - b.** Switch to DCI+K media.
 - c.** Harvest cells on Day 25 (procedure same as on Day 15).

- B.** Gelatin coating

Apply 0.1% gelatin (dissolved in ultrapure water) for 25-30 min at room temperature (*e.g.* 1 ml in a well of a 6-well plate). Aspirate gelatin, rinse with PBS and aspirate again. The plate is now ready to use.

- C.** FACS for Cxcr4/cKit
 - 1.** Cell count
 - a.** Remove 1 ml of EBs, spin down and trypsinize (0.5 ml trypsin), count cells.
 - b.** Based on previous cell count, remove a culture volume that corresponds to $2\text{-}2.5 \times 10^6$ cells.
 - c.** Repeat procedure for undifferentiated ES cells.

 - 2.** Preparation of cells for staining (all steps on ice)
 - a.** Spin down EBs, resuspend in 1 ml trypsin (60 sec at 37 °C), monodisperse with a P1000 pipette.
 - b.** Inactivate with 1 ml serum, spin down (5 min, $300 \times g$, 4 °C) and wash once with 5 ml IMDM.
 - c.** Resuspend in 500 μl PBS⁺ (PBS+2% FBS), cell concentration should be $0.4\text{ - }0.5 \times 10^6$ cells/100 μl .
 - d.** Prepare 2×5 Eppendorf tubes (unstained, isotypes, Cxcr4, cKit, Cxcr4/cKit (double)), mark each tube series (D5 endoderm or undifferentiated ES cells) and transfer 100 μl of each culture to each tube.

 - 3.** Cell staining
 - a.** Add the appropriate antibodies per tube (*e.g.* no antibodies in the “unstained” tube, both antibodies in the “double” tube).

Antibodies used for this protocol as of 08/15/12:

Isotype: APC Rat IgG_{2b}, K

Isotype: PE Rat IgC_{2b}, K

Cxcr4: APC Rat α -mouse CD184

cKit: PE Rat α -mouse CD117

- b.** Vortex briefly and transfer on ice for 30 min (cover with aluminum foil, vortex once again at 15 min).
- c.** Add 1 ml PBS⁺ per tube, spin at $300 \times g$ for 5 min in a tabletop centrifuge, carefully aspirate supernatant.
- d.** Resuspend pellet in 500 μ l PBS⁺.
- e.** Transfer to FACS polystyrene tubes with the cell strainer cap.
- f.** Take cells to LSRII for analysis.

Recipes

- 1.** IMDM
 - 1 packet IMDM powder
 - 3.02 g NaHCO₃
 - 10 ml Pen/Strep
 - 1 L Cellgro water
 - Check pH (acceptable range 6.9 - pH 7.3)
- 2.** Serum free differentiation medium (SFD)
 - IMDM-375 ml
 - Ham's F-1-125 ml
 - B-27 supplement with RA-5 ml
 - N-2 supplement-2.5 ml
 - BSA 7.5% in PBS-3.3 ml
- 3.** Complete serum free differentiation medium (cSFDM)
 - SFD -100 ml
 - MTG 300 μ l of stock (Stock: 26 μ l MTG to 2 ml IMDM)
 - 200 mM L-Glut -1 ml

- Ascorbic Acid -1 ml of stock (Stock: 5 mg/ml distilled water, prepare fresh!)
4. BASE medium for DCI+K
Ham's F-12-243.7 ml
1.0 M HEPES (pH 7.4)-3.75 ml
1.0 M CaCl₂-200 µl
BSA-0.625 g
100× ITS supplement-2.5 ml
 5. Anteriorization medium
cSFDM-10 ml
mNoggin -100 ng/ml (Stock: 10 µg/ml)
SB431542 -10 µM (Stock: 10 mM in DMSO)
 6. Ventralization medium
cSFDM - 10 ml
mWnt3a - 100 ng/ml (Stock: 100 µg/ml)
hBMP4 - 10 ng/ml (Stock: 10 µg/ml)
hEGF - 20 ng/ml (Stock: 20 µg/ml)
mFGF2 - 250 ng/ml (Stock: 100 µg/ml)
mFGF7 - 10 ng/ml (Stock: 10 µg/ml)
hFGF10 - 10 ng/ml (Stock: 10 µg/ml)
Heparin sodium salt - 100 ng/ml (Stock: 1 mg/ml)
 7. DCI+K medium
BASE media - 25 ml
Dexamethasone - 50 nM (Stock: 250 µM in ethanol)
KGF (FGF7) - 10 ng/ml (Stock: 10 µg/ml)
cAMP+IBMX - 0.1 mM (Stock: 1 mM cAMP+1 mM IBMX)
 8. Preparation of 10× cAMP+IBMX stock
Dissolve 22.22 mg IBMX in 1 ml of DMSO (0.1 M IBMX stock, store at -20 °C)
To prepare the 1 mM cAMP+1mM IBMX (10×) stock, dissolve 21.5 mg 8BrcAMP in 49.5 ml of BASE media and add 0.5 ml of IBMX stock
0.22 µm filter
Store at 4 °C for up to 4 weeks

Acknowledgments

This protocol was originally published as part of: Longmire *et al.* (2012). The authors wish to thank all members of the Kotton laboratory for helpful discussions and editing. DNK is supported by NIH PO1 HL047049-16A1, 1RC2HL101535-01, 1R01 HL095993-01, 1R01 HL108678, a USAMRRA Award, an Alpha-1 Foundation Award, and an ARC award from the Evans Center for Interdisciplinary Research at Boston University. TAL is supported by NIH training grant T32 HL007035. LI is supported by R01 HL111574 and an ATS/ChILD Foundation Award.

References

- Gonzales LW, Guttentag SH, Wade KC, Postle AD, Ballard PL. Differentiation of human pulmonary type II cells in vitro by glucocorticoid plus cAMP. *Am J Physiol Lung Cell Mol Physiol.* 2002; 283(5):L940–951. [PubMed: 12376347]
- Longmire TA, Ikonomou L, Hawkins F, Christodoulou C, Cao Y, Jean JC, Kwok LW, Mou H, Rajagopal J, Shen SS, Downton AA, Serra M, Weiss DJ, Green MD, Snoeck HW, Ramirez MI, Kotton DN. Efficient derivation of purified lung and thyroid progenitors from embryonic stem cells. *Cell Stem Cell.* 2012; 10(4):398–411. [PubMed: 22482505]
- Wray J, Kalkan T, Smith AG. Revolutionizing Drug Discovery with Stem Cell Technology. *Biochem Soc Trans.* 2010; 38:1027–1032. [PubMed: 20658998]