



Published in final edited form as:

J Magn Reson. 2016 August ; 269: 50–54. doi:10.1016/j.jmr.2016.05.006.

A Triarylmethyl Spin Label for Long-Range Distance Measurement at Physiological Temperatures Using T_1 Relaxation Enhancement

Zhongyu Yang^{1,†}, Michael D. Bridges¹, Carlos J. López¹, Olga Yu. Rogozhnikova^{2,3}, Dmitry V. Trukhin^{2,3}, Evan K. Brooks¹, Victor Tormyshev^{2,3,*}, Howard J. Halpern^{4,*}, and Wayne L. Hubbell^{1,*}

¹Jules Stein Eye Institute and Department of Chemistry and Biochemistry, University of California, Los Angeles, Los Angeles, CA, 90095

²N.N. Vorozhtsov Novosibirsk Institute of Organic Chemistry, Novosibirsk 630090, Russia

³Novosibirsk State University, Novosibirsk 630090, Russia

⁴The Center for EPR Imaging in vivo Physiology, Department of Radiation and Cellular Oncology, University of Chicago, Chicago, IL 60637, USA

Abstract

Site-directed spin labeling (SDSL) in combination with Electron Paramagnetic Resonance (EPR) spectroscopy has become an important tool for measuring distances in proteins on the order of a few nm. For this purpose pairs of spin labels, most commonly nitroxides, are site-selectively introduced into the protein. Recent efforts to develop new spin labels are focused on tailoring the intrinsic properties of the label to either extend the upper limit of measurable distances at physiological temperature, or to provide a unique spectral lineshape so that selective pairwise distances can be measured in a protein or complex containing multiple spin label species. Triarylmethyl (TAM) radicals are the foundation for a new class of spin labels that promise to provide both capabilities. Here we report a new methanethiosulfonate derivative of a TAM radical that reacts rapidly and selectively with an engineered cysteine residue to generate a TAM containing side chain (TAM1) in high yield. With a TAM1 residue and Cu^{2+} bound to an engineered Cu^{2+} binding site, enhanced T_1 relaxation of TAM should enable measurement of interspin distances up to 50 Å at physiological temperature. To achieve favorable TAM1-labeled protein concentrations without aggregation, proteins are tethered to a solid support either site-

*Corresponding Authors. torm@nioch.nsc.ru, h-halpern@uchicago.edu, hubbellw@jsei.ucla.edu Phone: (310) 206-8830.

† Present address: Department of Chemistry and Biochemistry, North Dakota State University, Fargo, ND, 58102

The authors declare no competing financial interests.


Supplementary Material

Details of synthesis of the TAM-MTS, preparation of protein samples, EPR experiments, discussion of modeling procedure, and additional data analysis are provided in the Supplementary Material.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

selectively using an unnatural amino acid or via native lysine residues. The methodology is general and readily extendable to complex systems, including membrane proteins.

Graphical abstract



Keywords

Triarylmethyl radical; spin labeling; saturation recovery; distance measurement at physiological temperature; relaxation enhancement

Distance constraints in proteins provide direct information on structure and structural transitions therein. Therefore, methodological development and optimization of distance measurements in proteins is an important endeavor in protein science and structural biology. Typical distances of interest in proteins are on the order of tens of Angstroms (\AA). Among the available experimental methods that can measure distances on this scale, site-directed spin labeling (SDSL) combined with electron paramagnetic resonance (EPR) spectroscopy stands out as a unique and powerful approach, applicable to proteins and protein complexes of any degree of complexity or size, and with relatively high sensitivity (μ g of proteins required).

In the SDSL-EPR strategy, spin label pairs are introduced site-selectively [1,2], and the magnetic interactions between the pairs are measured as a metric for the interspin distance. Nitroxides are commonly used as spin labels, and the nitroxide side chain R1 [3,4] is the most popular (Figure 1). Conformationally constrained nitroxide side chains R1p [5] and RX [6] that limit the contribution of side chain flexibility to the determined distances have also been developed (Fig. 1).

Magnetic dipolar interactions are directly measured by double electron-electron resonance (DEER) [7] and double quantum coherence (DQC) [8]. In either case, the probability distribution of interspin distances in the range of 15–70 \AA is obtained. However, in most cases both DEER and DQC are limited to cryogenic temperatures to slow electron spin relaxation rates and rotational tumbling of the entire protein, which would average out the orientation-dependent magnetic dipolar interaction [9]. However, it is desirable to measure distances at physiological temperatures, where potential complications caused by the

freezing process and/or shifts in conformational equilibria due to the presence of cryoprotectant are eliminated. Attempts are being made to adapt DEER and DQC for room temperature distance measurements in proteins and DNA using the slowly-relaxing TAM radical as a spin label and immobilizing the macromolecule on a solid support [9–11]. For proteins, the maximum distance so far measured with this strategy is ≈ 30 Å. Room temperature DEER measurements have been made using a spirocyclohexyl nitroxyl spin label on a protein trapped in trehalose glass [10], but this approach does not eliminate a potential structural perturbation in the glassy environment.

Dipolar broadening of continuous wave (CW) lineshapes of interacting nitroxides can be employed to measure distances up to 20 Å at room temperature [12,13]. Hyde and coworkers recently extended this upper limit to 30 Å using non-adiabatic rapid sweep (NARS) EPR at S band [14]. The major barrier preventing the measurement of longer distances by CW methods is the difficulty in detecting very marginal spectral broadening.

An alternative approach to measuring distances at room temperature is based on the T_1 relaxation enhancement (RE) of nitroxyl radicals by Cu^{2+} . Using this strategy, Jun *et al.* reported that RE of an R1 nitroxide side chain in a rapidly tumbling peptide ($T_1 \approx 2$ μs) could measure Cu^{2+} -nitroxide distances up to ~ 25 Å [15]. Yang *et al.* recently verified the theoretical prediction that the maximum range of distance measured by Cu^{2+} -induced RE increases with increasing T_1 of the nitroxide (Fig. 1) [16]. In that study, increased T_1 was achieved using nitroxides R1p and RX ($T_1 \approx 6$ –8 μs).

Here we show that a new class of spin labels based on triarylmethyl (TAM) radicals that have $T_1 \approx 12$ μs when attached to a protein can in principle measure distances up to ≈ 50 Å at room temperature using RE (Fig. 1). For this purpose, a novel TAM-methanethiosulfonate (MTS) reagent that undergoes a selective and high yield reaction with cysteine was prepared. The reaction generates a TAM-labeled side chain designated TAM1 (Scheme 1); details of the TAM-MTS synthesis and reaction conditions with the protein are included in Supporting Information. In an earlier study, a TAM radical for use in DQC was introduced at cysteine via reaction of an activated disulfide TAM reagent [9]. That reagent was limited for general use by a low reactivity, a problem now solved by the highly reactive MTS function.

The general application of TAM relaxation enhancement by Cu^{2+} requires site-selective introduction of both TAM1 and a high affinity Cu^{2+} binding site in the protein of interest. Recent advances make this possible with the facile introduction of high-affinity Cu^{2+} binding motifs consisting either of a genetically engineered pentapeptide in loops [16] or histidine pairs introduced into regular secondary structure [17]. A problem encountered upon the introduction of TAM1 is a reduction of the protein solubility [9] at least for some small soluble proteins, a problem that is solved by tethering the protein to a solid support prior to labeling. In this communication, we demonstrate the combined application of the new TAM1 side chain, a high affinity Cu^{2+} motif and protein tethering to measure the longest range distances yet obtained by EPR methods at ambient temperatures.

In the present study, T4 Lysozyme (T4L) mutants containing the high affinity Cu^{2+} binding pentapeptide (GGGHG) inserted in loop sequences and a single cysteine residue for

introduction of TAM1 are used to illustrate the overall strategy; the site of peptide insertion is indicated with the residue number and the suffix L. Tethering of the mutants to a solid support is most simply carried out by direct covalent attachment to commercially available CNBr-activated Sepharose via protein lysine residues [18]. This procedure has the potential disadvantage for some studies that the protein is attached in random orientation with respect to the matrix. Site-selective attachment can be achieved with the use of genetically incorporated unnatural amino acids as recently described [19]. Here we employ para-acetyl phenylalanine (p-AcPhe) as the unnatural amino acid to which biotin is conjugated via a hydroxylamine derivative [20]. The biotinylated protein is quantitatively adsorbed in high yield to commercially available streptavidin Sepharose, yielding effective concentrations as high as 700 μM without aggregation [20]. Tethering by either approach results in complete immobilization of the protein on the CW EPR time scale [18,20]. Details of protein labeling and tethering are provided in the Supporting Information.

The three Cu^{2+} /TAM1 pairs investigated in this study are shown in Figure 2. Each was tethered using CNBr-activated sepharose, and one (23L/131TAM1) was additionally coupled site-selectively using p-AcPhe at site 65 for comparison (Fig. 2A). The CW spectra of the TAM-MTS reagent alone and attached to T4L at site 76 under the indicated conditions are shown in Figure 3. Spectral line-broadening due to the increased correlation time of TAM1 on the protein is evident, particularly in the tethered sample, where the overall linewidth is similar to those observed for TAM radicals in frozen solution [21], which suggests strong immobilization of the TAM radical with respect to the protein. The major contribution to the linewidth is unresolved anisotropic hyperfine interactions within TAM1 [22]. Similar spectra are obtained for TAM1 at site 131.

The spin lattice relaxation times for TAM1 in the tethered T4L mutants in the absence and presence of a stoichiometric amount of Cu^{2+} were measured using long-pulse Saturation Recovery (SR) EPR following established protocols [23,24]. Figure 4 shows representative SR relaxation curves for T4L 23L/131TAM1 attached site-selectively at residue 65.

For each mutant, copper-free relaxation curves were well-fit with single exponentials, whereas mono-exponential fits of copper-bound samples were poor. For immobilized proteins containing Cu^{2+} and TAM1, the apparent T_1 of TAM1 depends on the orientation of the interspin vector with respect to the external field. If the interacting spins can be assumed to have isotropic g values, the orientation-dependent T_1 is given by Hirsch *et al.* [25]:

$$\frac{1}{T_{1s}} - \frac{1}{T_{1s}^0} = \frac{3\pi^2 g_s^2 g_f^2 \beta_e^4}{h^2 r^6} \left[\left(\frac{1}{6} \right) \frac{T_{2f}}{1 + (\omega_f - \omega_s)^2 T_{2f}^2} (1 - 3 \cos^2 \theta)^2 + \frac{3T_{1f}}{1 + \omega_s^2 T_{1f}^2} \sin^2 \theta \cos^2 \theta + \left(\frac{3}{2} \right) \frac{T_{2f}}{1 + (\omega_f + \omega_s)^2 T_{2f}^2} \sin^4 \theta \right]$$

(Eq. 1)

where T_{1s} and T_{1s}^0 are the spin-lattice relaxation times for TAM in the presence and absence of Cu^{2+} , respectively, ω_s and ω_f are the resonant frequencies of TAM and Cu^{2+} , T_{1f} and T_{2f} are the spin relaxation times of Cu^{2+} , and β_e and h are the Bohr magneton and the Planck

constant, respectively. The TAM radical has an essentially isotropic g-tensor, but Cu²⁺ complexes do not. Nevertheless, it is found that use of $g_f = 2.1166$, the average g for a 3N1O-bound copper state [26] and $g_s = 2.0058$ for TAM (determined by characterization of TAM MTS; see SI) in Eq. 1 can account well for the experimental data and reasonably corresponds to TAM1 modeled in the protein (Figure 2 and SI), when taking $T_{1f} = T_{2f} = 3$ ns. The resonant frequency of Cu²⁺, ω_f , was calculated using the isotropic value of g_f . Averaging of the copper g-tensor may arise from localized fast motion of the Cu²⁺ complex in the protein, as suggested by MD simulations as well as DEER distance distributions between nitroxides and Cu²⁺ in binding loops [16]. The values for Cu²⁺ relaxation times have been reported to be in the range of 1 to 5 ns [27], and a value of 3 ns has been found to account for copper-induced relaxation of nitroxides in a number of earlier studies [15,16,28]. Calculation of r from relaxation enhancement requires fits of the experimental relaxation curves to a weighted sum of exponentials as a function of the angle θ as described in detail in the SI. The experimentally-determined copper-free intrinsic T₁ values and average interspin distances for each mutant studied are given in Table 1.

The method reported herein is only valid if the occupancy of the Cu²⁺ binding site is close to 100%. Low affinity sites (K_d in the μM range) can in principle be saturated with high Cu²⁺ concentrations, but in this case Heisenberg exchange with free Cu²⁺ will shorten T₁; the *i* and *i+4* bi-histidine helical motif employed by Voss *et al.* [28] has an affinity too low for the present method. Various high affinity Cu²⁺ peptide binding sites have been reported in the literature; for example, studies described in References [16], [17], and [29] all involve Cu²⁺ binding sites with nanomolar K_d values.

As mentioned above, the broad linewidth of the TAM1 in solid support-tethered T4L suggests strong immobilization of the spin label. The degree to which this reflects spatial localization of TAM1 was investigated by comparing an R1-R1 distance distribution with the corresponding R1-TAM1 distribution, both being obtained using DEER at cryogenic temperatures. If this comparison is done at sites where the R1-R1 distribution is narrow, a difference in the distribution widths can be tentatively assigned to disorder in the TAM1 side chain. The sample selected for this purpose was the mutant T4L 68C/109C which has a mono-modal distribution of narrow width ($<5 \text{ \AA}$ at half-height) when labeled with R1, as seen in Figure 5 [19].

To prepare the T4L R1-TAM hybrid, 25% of cysteines were labeled with R1, followed by reaction with excess TAM-MTS. Statistically, this yields 1/16 R1-R1, 9/16 TAM-TAM spin pair, and 3/8 R1-TAM spin pair. Consequentially, the R1-R1 spin pairs only contribute minimally in the R1-TAM distance distribution. The 4-pulse DEER data are shown in Figure 5. As shown in the figure inset, the pump (inversion) pulse was adjusted to excite primarily TAM spins, and the observe pulse was selected so that primarily R1 spins were observed. The resultant distance distribution of the R1-TAM pair shows essentially the same mean distance and a slightly broader distance distribution compared to that of the R1-R1 pair. The data suggest that the inherent flexibility of the TAM1 side chain is comparable to the R1 side chain commonly employed in DEER distance mapping in proteins.

In summary, the TAM1 side chain has useful advantages over a nitroxide side chain for distance mapping using T_1 relaxation enhancement. First, the long T_1 for TAM1 attached to a protein enables significantly longer distances to be measured at ambient temperatures, up to ≈ 50 Å (Fig. 1). Although the size of the T4L molecule did not permit sites to be selected with interspin distance measurement beyond ≈ 40 Å, the high signal to noise of the SR data should make the longer distance measurement feasible. Second, the narrow single central resonance line has a ~ 30 fold higher intensity compared to the same concentration of an immobile nitroxide, providing a comparably higher signal-to-noise in a saturation recovery experiment. Disadvantages of TAM1 compared to a nitroxide include the much larger steric bulk and the instability of the ester linked TAM due to hydrolysis. The half-life of hydrolysis for the ester linkage is ~ 50 hours and does not severely limit its use [30]. In future studies, other more stable linkages will be explored, and the degree to which the TAM1 side chain destabilizes the protein will be investigated. Regarding the latter point, we note that the measured TAM1-Cu²⁺ distances agree with those modeled in the native structure, so apparently the presence of TAM1 at surface sites does not lead to significant unfolding. The requirement for tethering the protein to a solid support may be viewed as a disadvantage, but in reality we advocate the general use of tethering in protein spin labeling due to the ability to provide high concentrations of spin with no perturbation to the protein structure and the absence of protein-protein interactions [20].

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institute of Health (USA) grants NIH/NEI EY005216, NIH/NEI EY00331 and Jules Stein Professorship Endowment (W.L.H.), and Russian Science Foundation (No. 14-14-00922), Russian Foundation for Basic Research (grant 14-03-93180) and the National Institute of Biomedical Imaging and Bioengineering (NIH No. 5P41EB002034) (V.M.T.).

References

1. Hubbell WL, López CJ, Altenbach C, Yang Z. Technological advances in site-directed spin labeling of proteins. *Curr Opin Struct Biol.* 2013; 23:725–733. [PubMed: 23850140]
2. Hubbell WL, Cafiso D, Altenbach C. Identifying conformational changes with site-directed spin labeling. *Nat Struct Biol.* 2000; 7:735–739. [PubMed: 10966640]
3. Todd AP, Cong J, Levinthal F, Levinthal C, Hubbell WL. Site-directed mutagenesis of colicin E1 provides specific attachment sites for spin labels whose spectra are sensitive to local conformation. *Proteins.* 1989; 6:294–305. [PubMed: 2560193]
4. Fleissner MR, Cascio D, Hubbell WL. Structural origin of weakly ordered nitroxide motion in spin-labeled proteins. *Protein Sci.* 2009; 18:893–908. [PubMed: 19384990]
5. Fawzi N, Fleissner MR, Anthis N, Kálai T, Hideg K, Hubbell WL, Clore GM. A rigid disulfide-linked nitroxide side chain simplifies the quantitative analysis of PRE data. *J Biomol NMR.* 2011; 51:105–114. [PubMed: 21947919]
6. Fleissner MR, Bridges MD, Brooks EK, Cascio D, Kálai T, Hideg K, Hubbell WL. Structure and dynamics of a conformationally constrained nitroxide side chain and applications in EPR spectroscopy. *Proc Nat Acad Sci USA.* 2011; 108:16241–16246. [PubMed: 21911399]
7. Jeschke G. DEER distance measurements on proteins. *Annu Rev Phys Chem.* 2012; 63:419–446. [PubMed: 22404592]

8. Borbat, P.; Freed, JH. Pulse Dipolar ESR: Distance Measurements. In: Harmer, J.; Timmel, C., editors. *Structural Information from Spin-Labels and Intrinsic Paramagnetic Centers in the Biosciences. Structure and Bonding*. Vol. 152. Springer, Heidelberg, Germany; New York, USA: 2014. p. 1
9. Yang Z, Liu Y, Borbat P, Zweier JL, Freed JH, Hubbell WL. Pulsed ESR dipolar spectroscopy for distance measurements in immobilized spin labeled proteins in liquid solution. *J Am Chem Soc*. 2012; 134:9950–9952. [PubMed: 22676043]
10. Meyer V, Swanson MA, Clouston LJ, Boraty ski PJ, Stein RA, Mchaourab HS, Rajca A, Eaton SS, Eaton GR. Room-temperature distance measurements of immobilized spin-labeled protein by DEER/PELDOR. *Biophys J*. 2015; 108:1213–1219. [PubMed: 25762332]
11. Shevelev GY, Krumkacheva OA, Lomzov AA, Kuzhelev AA, Rogozhnikova OY, Trukhin DV, Troitskaya TI, Tormyshev VM, Fedin MV, Pyshnyi DV, Bagryanskaya EG. Physiological-temperature distance measurement in nucleic acid using triarylmethyl-based spin labels and pulsed dipolar EPR spectroscopy. *J Am Chem Soc*. 2014; 136:9874–9877. [PubMed: 24963806]
12. Altenbach C, Oh KJ, Trabanino RJ, Hideg K, Hubbell WL. Estimation of inter-residue distances in spin labeled proteins at physiological temperatures: Experimental strategies and practical limitations. *Biochemistry*. 2001; 40:15471–15482. [PubMed: 11747422]
13. Rabenstein MD, Shin YK. Determination of the distance between two spin labels attached to a macromolecule. *Proc Nat Acad Sci USA*. 1995; 92:8239–8243. [PubMed: 7667275]
14. Kittell AW, Hustedt EJ, Hyde JS. Inter-spin distance determination using L-band (1–2 GHz) non-adiabatic rapid sweep electron paramagnetic resonance (NARS EPR). *J Magn Reson*. 2012; 221:51–56. [PubMed: 22750251]
15. Jun S, Becker JS, Yonkunas M, Coalson R, Saxena S. Unfolding of alanine-based peptides using electron spin resonance distance measurements. *Biochemistry*. 2006; 45:11666–11673. [PubMed: 16981726]
16. Yang Z, Jiménez-Osés G, López CJ, Bridges MD, Houk KN, Hubbell WL. Long-range distance measurements in proteins at physiological temperatures using saturation recovery EPR spectroscopy. *J Am Chem Soc*. 2014; 136:15356–15365. [PubMed: 25290172]
17. Cunningham TF, Putterman MR, Desai A, Horne WS, Saxena S. The double-histidine Cu²⁺-binding motif: a highly rigid, site-specific spin probe for electron spin resonance distance measurements. *Angew Chem Int Ed*. 2015; 54:6330–6334.
18. López CJ, Fleissner MR, Guo Z, Kusnetzow AK, Hubbell WL. Osmolyte perturbation reveals conformational equilibria in spin-labeled proteins. *Protein Sci*. 2009; 18:1637–1652. [PubMed: 19585559]
19. Fleissner MR, Brustad EM, Kálai T, Altenbach C, Cascio D, Peters FB, Hideg K, Peuker S, Schultz PG, Hubbell WL. Site-directed spin labeling of a genetically encoded unnatural amino acid. *Proc Nat Acad Sci USA*. 2009; 106:21637–21642. [PubMed: 19995976]
20. López CJ, Fleissner MR, Brooks EK, Hubbell WL. Stationary-phase EPR for exploring protein structure, conformation, and dynamics in spin-labeled proteins. *Biochemistry*. 2014; 53:7067–7075. [PubMed: 25333901]
21. Fielding AJ, Carl PJ, Eaton GR, Eaton SS. Multifrequency EPR of four triarylmethyl radicals. *Appl Magn Reson*. 2005; 28:231.
22. Owenius R, Eaton GR, Eaton SS. Frequency (250 MHz to 9.2 GHz) and viscosity dependence of electron spin relaxation of triarylmethyl radicals at room temperature. *J Magn Reson*. 2005; 172:168–175. [PubMed: 15589420]
23. Hyde, JS. Saturation recovery methodology. In: Kevan, L.; Schwartz, RN., editors. *Time domain electron spin resonance*. John Wiley & Sons; New York: 1979. p. 1-30.
24. Yang, Z.; Bridges, MD.; Lerch, MT.; Altenbach, C.; Hubbell, WL. Saturation Recovery EPR and Nitroxide Spin Labeling for Exploring Structure and Dynamics in Proteins. In *Electron Paramagnetic Resonance Investigations of Biological Systems by Using Spin Labels, Spin Probes, and Intrinsic Metal Ions, Part B*. In: Qin, PZ.; Warncke, K., editors. *Method Enzymol*. Vol. 564. Academic Press; UK: 2015. p. 1-27.
25. Hirsh DJ, Beck WF, Innes JB, Brudvig GW. Using saturation-recovery EPR to measure distances in proteins: Applications to photosystem II. *Biochemistry*. 1992; 31:532–541. [PubMed: 1310040]

26. Aronoff-Spencer E, Burns CS, Avdievich NI, Gerfen GJ, Peisach J, Antholine WE, Ball HL, Cohen FE, Prusiner SB, Millhauser GL. Identification of the Cu²⁺ Binding Sites in the N-Terminal Domain of the Prion Protein by EPR and CD Spectroscopy. *Biochemistry*. 2000; 39:13760–13771. [PubMed: 11076515]
27. Bertini, I.; Luchinat, C.; Parigi, G. *Solution NMR of Paramagnetic Molecules: Applications to metalloproteins and models*. Vol. 2. Elsevier Science; New York: 2001.
28. Voss J, Salwinski L, Kaback HR, Hubbell WL. A method for distance determination in proteins using a designed metal ion binding site and site-directed spin labeling: Evaluation with T4 Lysozyme. *Proc Nat Acad Sci USA*. 1995; 92:12295–12299. [PubMed: 8618888]
29. Cunningham TF, Shannon MD, Putterman MR, Arachchige RJ, Sengupta I, Gao M, Jaroniec CP, Saxena S. Cysteine-Specific Cu²⁺ Chelating Tags Used as Paramagnetic Probes in Double Electron Electron Resonance. *J Phys Chem B*. 2015; 119:2839–2843. [PubMed: 25608028]
30. Kuzhelev AA, Trukhin DV, Krumkacheva OA, Strizhakov RK, Rogozhnikova OY, Troitskaya TI, Fedin MV, Tormyshev VM, Bagryanskaya EG. Room-temperature electron spin relaxation of triarylmethyl radicals at the X- and Q-bands. *J Phys Chem B*. 2015; 119:13630–13640. [PubMed: 26001103]

Highlights

- A new cysteine-reactive triarylmethyl spin label for proteins is reported.
- Relaxation enhancement of the spin label (TAM1) by Cu^{2+} is distance dependent.
- Interspin distances up to ~ 50 Å are measurable at physiological temperature.




Figure 1. Maximum measurable spin label-to- Cu^{2+} distance measured by relaxometry as a function of spin label T_1 . The maximum distances were estimated by taking the minimum measurable drop in T_1 due to the presence of Cu^{2+} to be $0.5 \mu\text{s}$ with a signal-to-noise ratio $>50:1$, typical of the data acquired (cf. ref. 16). The arrows indicate the T_1 values for the indicated nitroxide side chains (cf. ref. 16) and for TAM1 attached to an immobile protein, reported here to be $\approx 12 \mu\text{s}$.




Figure 2.

Models of the TAM1-Cu²⁺ pairs investigated. The 23L/131TAM1 mutant was tethered to Sepharose randomly at native Lys residues by reaction with CNBr-activated Sepharose, or site-selectively using the unnatural amino acid p-AcPhe introduced at site 65, indicated by the magenta sphere in (A) (see text). The TAM1 spin label and Cu²⁺ in the binding loop were modeled as described in the Supporting Information.




Figure 3. Normalized CW EPR spectra for TAM-MTS in solution and for the T4 Lysozyme mutant 76TAM1 in various conditions. Spectral line broadening is apparent in the labeled protein samples due to the reduced rotational correlation time of the TAM1 side chain. The spectrum for the Sepharose-immobilized protein via the CNBr method is also shown with the vertical axis scaled up by 50 times (gray). The ¹³C-satellite lines are clearly evident in the TAM-MTS spectrum (cf. ref. 26).




Figure 4.

Representative relaxation enhancement data. The saturation recovery traces obtained for the 23L/65p-AcPhe/131TAM1 sample in the absence (black) and presence (red) of Cu²⁺ at a microwave observe power of 50 μW. A clear “enhancement” in the relaxation rate can be seen upon the addition of Cu²⁺. SR curves, best fits, and residual traces for the four mutants studied in all conditions are given the SI (Figures S1–S4).





Figure 5.

DEER data reveal the spatial localization of the TAM1 spin label. (A) The time domain signal of the 68C/109C mutant labeled with R1 (black trace) and with a R1-to-TAM1 ratio of 1:3 (red trace). (B) The distance distributions obtained from analysis of the DEER data in (A) for the R1-R1 (black) and R1-TAM1 (red) labeled mutants. The inset in (B) shows the field-swept electron spin echo spectra (200 G wide) at Q-band for the R1-R1 (black) and R1-TAM1 (red) labeled 68C/109C mutant. The arrows marked “p” and “o” indicate field positions of pump and observe pulses, respectively, for each experiment (see text).



Scheme 1.
Preparation of TAM-MTS reagent and TAM-labeling of protein

Table 1

Intrinsic copper-free T_1 values and interspin distances measured for four TAM1-spin labeled mutants of T4 Lysozyme, and modeled interspin distances (See Figure 2).

T4 Lysozyme Mutant ^a	Intrinsic T_1 , T_{1s}^0 (μ s) ^b	Interspin Distance, r (\AA)	Modeled Distance (\AA)
23L/131TAM1	12.0	37.9	39.5
23L/131TAM1 (p-AcPhe at 65)	12.5	37.7	39.5
37L/76TAM1	12.1	32.6	35.1
135L/76TAM1	12.1	32.8	36.1

^aAll proteins immobilized via random attachment to CNBr-activated Sepharose solid support, except for the sample labeled 'p-AcPhe at 65', which was site-specifically attached via biotin linkage.

^bReported intrinsic T_1 values obtained by measuring copper-free T_1 values at various observe powers with extrapolation to zero power; complete data in Supporting Information.