

## CMEARTICLE

## ST-segment changes with exercise stress


Yoke Ching Lim<sup>1</sup>, MBChB, MRCP, Swee-Guan Teo<sup>2</sup>, MRCP, FACC, Kian-Keong Poh<sup>1,3</sup>, FRCP, FACC

Fig. 1 ECG at peak stress during treadmill ECG stress test.

## CASE 1

## CLINICAL PRESENTATION

A 62-year-old woman presented with atypical angina for the past three months. Her past medical history includes hyperlipidaemia. Electrocardiography (ECG) performed at rest shows normal sinus rhythm. The patient underwent treadmill ECG stress test, and her ECG at highest heart rate during the treadmill stress test is shown in Fig. 1. What are the ECG abnormalities? What is the diagnosis?

## ECG STRESS TEST INTERPRETATION

The test was terminated at Stage 2 of the Bruce protocol with seven Metabolic Equivalents of Tasks (METs) of work due to reports of chest pain and breathlessness. The patient could not achieve the target heart rate of 135 beats per minute (bpm). ECG performed at her highest heart rate (Fig. 1) shows sinus tachycardia at a rate


of 112 bpm, with a 1-mm horizontal ST-segment depression at 80 ms after the J-point, in the inferolateral leads (II, III, aVF and V4–6), which is suggestive of ischaemia due to obstructive coronary artery disease (CAD). In addition, the patient also had a hypertensive response to exercise with a blood pressure of up to 206/96 mmHg.

## CLINICAL COURSE


In view of her symptoms and abnormal treadmill ECG stress test, the patient underwent coronary angiography, which showed severe left main and triple vessel disease. She had 90% ostial left main stenosis, severe diffuse distal disease in the left anterior descending (LAD) and circumflex arteries, as well as 90% ostial right coronary artery (RCA) stenosis with 70% stenosis in the posterior descending artery (Fig. 2). The patient was referred to a cardiothoracic surgeon, and coronary artery bypass graft surgery was performed.

<sup>1</sup>Department of Cardiology, National University Heart Centre, National University Health System, <sup>2</sup>Raffles Heart Centre, Raffles Hospital, <sup>3</sup>Yong Loo Lin School of Medicine, National University of Singapore, Singapore

**Correspondence:** A/Prof Poh Kian Keong, Department of Cardiology, National University Heart Centre, 1E Kent Ridge Road, NUHS Tower Block Level 9, Singapore 119228. kian\_keong\_poh@nuhs.edu.sg



**Fig. 2** Coronary angiograms show (a) 90% stenosis in the ostium of the right coronary artery (white arrow) in the left anterior oblique view; and (b) 90% stenosis in the ostium of the left main coronary artery (black arrow) with diffuse disease in the left anterior descending and left circumflex artery in the right anterior oblique cranial view.



**Fig. 3** ECG at peak stress during treadmill ECG stress test.

## CASE 2 CLINICAL PRESENTATION

A 51-year-old man was referred for treadmill ECG stress test due to breathlessness on exertion. He had a previous anterior ST-elevation myocardial infarction in the year 2005 and primary percutaneous intervention to his proximal LAD coronary artery. The patient's medical history also included diabetes mellitus and dyslipidaemia. ECG done at rest shows Q waves in the anterior lead V1–2, consistent with a previous anterior myocardial infarction. The patient underwent treadmill ECG stress test. The ECG in Fig. 3 was taken at his maximal heart rate during stress. What are the ECG abnormalities?

## ECG STRESS TEST INTERPRETATION

The patient achieved Stage 4 of the Bruce protocol, exercising for a

total of nine minutes 46 seconds with 11.2 METs of work. ECG at peak stress showed sinus tachycardia at a rate of 146 bpm. There were new horizontal and up-sloping ST-segment depressions in leads II, III, aVF and V4–6, which were suggestive of myocardial ischaemia. In addition, there was an isolated 1-mm ST-segment elevation in lead aVL. The ST segments returned to baseline at five minutes of recovery.

## CLINICAL COURSE

The patient subsequently underwent coronary angiography, which showed total occlusion in the ostial LAD artery, with further 90% stenosis in the left posterior descending artery (LPDA) (Fig. 4). He underwent successful coronary angioplasty with stenting of the ostial to proximal LAD arteries and balloon angioplasty of the LPDA. The patient's effort tolerance improved after angioplasty.



**CASE 3****CLINICAL PRESENTATION**

A 61-year-old man presented with a one-week history of angina. He had a history of hypertension and dyslipidaemia. The patient underwent myocardial perfusion treadmill ECG stress test. ECG at rest showed normal sinus rhythm. Fig. 5 shows the patient's ECG immediately after achieving target heart rate during stress. What are the ECG abnormalities?

**ECG STRESS TEST INTERPRETATION**


ECG performed immediately after achieving target heart rate (Fig. 5) showed sinus tachycardia with a heart rate of 100 bpm. The test was stopped, as the patient developed breathlessness at Stage 3 with a peak heart rate of 111 bpm, at 10.2 METs of work. ST-segment elevations of up to 2 mm were seen in the inferior leads II, III and aVF. In addition, there were also horizontal and down-sloping ST-segment depressions in leads I, aVL and V4–6.

**CLINICAL COURSE**

The patient's myocardial perfusion images showed a reversible perfusion defect in the mid-to-basal inferior wall, which is suspicious of obstructive CAD in the RCA territory. His coronary angiogram showed total occlusion of the proximal RCA, with Rentrop Grade 2 collaterals from the left coronary arteries to the distal RCA (Fig. 6). The patient underwent angioplasty with stenting to the RCA.

**DISCUSSION****Treadmill ECG stress test**

In 2014, ischaemic heart disease (IHD) was the third leading cause of death in Singapore, accounting for 16% of total deaths.<sup>(1)</sup> There is a need for early identification and improved treatment of IHD. The treadmill ECG stress test is commonly used as a screening test to identify myocardial ischaemia. During exercise, limitation




**Fig. 4** Coronary angiogram shows total occlusion of the ostial left anterior descending artery in the right anterior oblique caudal view. The occluded left anterior descending stent is also seen (arrow).

of coronary blood flow due to obstructive CAD may lead to myocardial ischaemia and its subsequent ECG changes. This is the basis of detecting significant CAD on the treadmill ECG stress test. In addition, the treadmill ECG stress test is also frequently used to assess for chronotropic response to exercise, exercise-induced arrhythmia and evaluation of symptoms during exercise, as well as the individual's exercise capacity. The sensitivity of the treadmill ECG stress test in diagnosing obstructive CAD was approximately 50%, with a specificity of approximately 90%.<sup>(2)</sup>


**Indications for the treadmill ECG stress test**

The American Heart Association previously published a guideline recommending stress tests for adult patients with intermediate pretest probability of CAD.<sup>(3)</sup> Table I provides a guide on the pretest probability of CAD for individuals based on age, gender and presenting symptoms.

The Ministry of Health, Singapore, published its own guidelines on screening for cardiovascular disease.<sup>(4)</sup> In addition



**Fig. 5** ECG done immediately after peak stress.



**Fig. 6** Coronary angiogram shows (a) 100% occlusion of the proximal right coronary artery (white arrow) in the left anterior oblique cranial view; and (b) collateral from the left coronary arteries (black arrow) in the left anterior oblique cranial view.

**Table I. Pretest probability of coronary artery disease based on age, gender and presenting symptoms.<sup>(3)</sup>**

Age (yr)	Gender	Typical angina pectoris	Atypical/probable angina pectoris	Non-angina chest pain	Asymptomatic
30–39	Male	Intermediate	Intermediate	Low	Very low
	Female	Intermediate	Very low	Very low	Very low
40–49	Male	High	Intermediate	Intermediate	Low
	Female	Intermediate	Low	Very low	Very low
50–59	Male	High	Intermediate	Intermediate	Low
	Female	Intermediate	Intermediate	Low	Very low
60–69	Male	High	Intermediate	Intermediate	Low
	Female	High	Intermediate	Intermediate	Low

Categories are very low: < 5%; low: ≥ 5% to < 10%; intermediate: 10%–90%; and high: > 90%. Typical angina is defined as retrosternal chest pain provoked by exertion or emotional stress and relieved by rest or nitroglycerin. Atypical angina lacks one of these three characteristics.

to evaluating symptomatic patients with intermediate pretest probability, the guidelines also recommended the treadmill ECG stress test for the detection of CAD in the following groups: (a) asymptomatic individuals with multiple risk factors; (b) asymptomatic men above 45 years of age and women above 55 years of age who are planning to start vigorous exercise; (c) individuals in occupations in which impairment may affect public safety (e.g. pilots); (d) individuals at high risk of coronary artery disease due to comorbidities; and (e) patients with diabetes mellitus who are planning to start vigorous exercise.

Patients whose baseline ECG shows more than 1-mm ST-depression or less than 1-mm ST-depression, and who are taking digoxin or fulfil the ECG criteria for left ventricular hypertrophy, electronically paced ventricular rhythm or complete left bundle branch block were considered not suitable for evaluation using the treadmill ECG stress test alone, as it would be difficult to interpret any changes in the ST segment during the test.

### Performing the treadmill ECG stress test

The Bruce protocol is the most commonly used exercise treadmill stress protocol. It consists of a total of seven stages, with a gradual increment in the speed and gradient of the treadmill. Each stage lasts three minutes to allow the patient to acclimatise to the specific speed and gradient before advancing to the next stage.

Exercise intensity during the treadmill ECG stress test is calculated in METs, which reflect the amount of oxygen consumed per minute. 1 MET is equal to 3.5 mL/min/kg of oxygen used. Watching television, for example, requires 1 MET, while brisk walking requires 4 METs.

Prior to starting the Bruce protocol, ECG may be performed with the patient hyperventilating in the standing position to assess for any ST-segment changes caused by hyperventilation. The MET value increases as the patient advances to higher stages of the Bruce protocol. Table II shows a standard Bruce protocol indicating the speed, gradient and MET values of each stage. Continuous ECG monitoring is carried out throughout the test and into the recovery period to assess for any changes in the ST segments, as well as for arrhythmia. The test is often stopped before completing Stage 7, when the patient achieves the target heart rate or is symptomatic, or when the test is positive. Other indications for termination are listed in Table III.

### Contraindications for exercise stress test

As with any investigation, it is important to ensure the safety of the patient undergoing the test, since the treadmill ECG stress test could be harmful to the patient in certain situations. Each laboratory has its own protocol regarding the safety and suitability of the patients undergoing such tests. Table IV shows the main contraindications for a treadmill ECG stress test.

**Heart rate response**

During exercise, the physiological increase in heart rate is due to a decrease in vagal tone followed by an increase in sympathetic tone. This, in turn, leads to increased oxygen supply to the vital organs in the body. However, the maximal heart rate achievable during exercise is heavily influenced by age. Most laboratories worldwide use a simple equation to predict the maximum heart rate: *Maximum predicted heart rate = 220 – age in years.*

Traditionally, sufficient effort during treadmill ECG stress testing is defined as achieving 85% of the age-predicted maximal heart rate. However, some studies have reported a significant variability in the maximum predicted heart rate among people of identical ages;<sup>(5,6)</sup> hence, this may not be used in isolation as a termination criterion. In addition to the evaluation of CAD, the treadmill ECG stress test is also often used to evaluate for chronotropic incompetency, in which the patient is unable to mount an adequate heart rate response to exercise, leading to symptoms such as effort-related dyspnoea and lethargy.

**Table II. Characteristics of each stage of the Bruce protocol.**

Stage	Time (min)	Speed (km/hr)	Gradient (%)	Metabolic equivalents of tasks
1	3	2.7	10	5
2	6	4.0	12	7
3	9	5.4	14	10
4	12	6.7	16	13
5	15	8.0	18	15
6	18	8.8	20	18
7	21	9.6	22	20

**Blood pressure response**

The increase in sympathetic tone and vasodilation during exercise generally causes a rise in blood pressure. A drop in systolic blood pressure > 10 mmHg is an indication to terminate the treadmill ECG stress tests, as it could be due to myocardial ischaemia leading to left ventricular dysfunction. This often signifies severe multi-vessel CAD, especially if there are accompanying symptoms and/or ST changes on the ECG. An exaggerated blood pressure response, defined as a peak systolic blood pressure of > 210 mmHg in men and > 190 mmHg in women according to the Framingham criteria,<sup>(7)</sup> was shown to predict the risk of hypertension in normotensive individuals.<sup>(8)</sup>

**Indications for termination of the treadmill ECG stress test**

In addition to achieving 85% of the age-predicted maximal heart rate, other indications for the termination of a treadmill ECG stress test are shown in Table III.

**Interpretation of the treadmill ECG stress test**

*ST-segment changes indicative of obstructive CAD*

A treadmill ECG stress test is considered abnormal when there is a horizontal or down-sloping ST-segment depression ≥ 1 mm at 60–80 ms after the J point.<sup>(9)</sup> Exercise ECGs with up-sloping ST-segment depressions are typically reported as an ‘equivocal’ test. In general, the occurrence of horizontal or down-sloping ST-segment depression at a lower workload (calculated in METs) or heart rate indicates a worse prognosis and higher likelihood of multi-vessel disease. The duration of ST-segment depression is also important, as prolonged recovery after peak

**Table III. Indications for termination of a treadmill ECG stress test.<sup>(9)</sup>**

Absolute indications	Relative indications
<ul style="list-style-type: none"> <li>• ST-segment elevation of &gt; 1 mm in leads without pre-existing Q waves due to a prior myocardial infarction, other than aVR, aVL and V1</li> <li>• Development of symptoms: moderate to severe angina, dizziness, near-syncope</li> <li>• Sustained ventricular tachycardia or other arrhythmia, including second or third degree heart block that interferes with maintenance of cardiac output during exercise</li> <li>• Drop in systolic blood pressure &gt; 10 mmHg despite an increased workload, accompanied by other evidence of ischaemia</li> <li>• Signs of poor perfusion (cyanosis, pallor)</li> <li>• Neurological symptoms</li> <li>• Individual’s request to stop</li> <li>• Technical difficulty in monitoring the ECG or systolic blood pressure</li> </ul>	<ul style="list-style-type: none"> <li>• ST-segment depression (horizontal or down-sloping of &gt; 2 mm at 60–80 ms after the J point in patient with suspected ischaemia</li> <li>• Drop in systolic blood pressure &gt; 10 mmHg despite an increased workload without any other evidence of ischaemia</li> <li>• Increasing chest pain, fatigue, dyspnoea, wheezing, leg cramps or claudication</li> <li>• Other tachyarrhythmia or bradyarrhythmia that may potentially interfere with haemodynamic instability (e.g. supraventricular tachycardia, multifocal ventricular ectopics)</li> <li>• Exaggerated hypertensive response defined as systolic blood pressure &gt; 250 mmHg or diastolic blood pressure &gt; 115 mmHg</li> <li>• Development of bundle branch block that cannot be immediately be distinguished from ventricular tachycardia</li> </ul>

**Table IV. Main contraindications for exercise stress test.**

Absolute contraindication	Relative contraindication
<ul style="list-style-type: none"> <li>• Acute myocardial infarction or unstable angina</li> <li>• Symptomatic severe aortic stenosis</li> <li>• Cardiac arrhythmia with haemodynamic instability</li> <li>• Active endocarditis</li> <li>• Decompensated heart failure</li> <li>• Acute myocarditis or pericarditis</li> <li>• Acute pulmonary embolism, pulmonary infarction or deep vein thrombosis</li> <li>• Acute aortic dissection</li> </ul>	<ul style="list-style-type: none"> <li>• Known obstructive left main coronary artery stenosis</li> <li>• Moderate to severe aortic stenosis</li> <li>• Cardiac arrhythmia with rapid ventricular rates, advanced or complete heart blocks</li> <li>• Hypertrophic obstructive cardiomyopathy</li> <li>• Recent stroke or transient ischaemic attack</li> <li>• Poorly controlled hypertension (resting blood pressure &gt; 200/110 mmHg)</li> <li>• Untreated medical conditions, such as anaemia and thyrotoxicosis</li> </ul>



stress is consistent with a positive treadmill ECG stress test. Another finding that is highly indicative of significant CAD is the occurrence of ST-segment elevation > 1 mm (often suggesting transmural ischaemia); these patients are frequently referred urgently for coronary angiography.

In Case 1, the patient had an abnormal treadmill ECG stress test with mild horizontal and down-sloping ST depression in the inferior-lateral leads at Stage 2 (low workload) of the Bruce protocol. In addition to being electrically positive, the test was also subjectively positive, with the patient reporting chest pain with exercise stress. These are highly suggestive of obstructive CAD, as proven in the subsequent coronary angiogram. Case 2 illustrated the use of the treadmill ECG stress test in a patient with a previous history of anterior myocardial infarction; the ST-segment depressions were seen not only at maximal heart rate, but also during the recovery period, indicating significant ischaemia. While ST-segment depression in the inferolateral leads during the treadmill ECG stress test indicates ischaemia from obstructive CAD, we cannot identify the diseased coronary artery from the ST-segment depression alone. However, as shown in Case 3, the ST-segment elevation during the treadmill ECG stress test can be used to identify the culprit coronary artery (where the stenosis causing ischaemia was located), which in this case was the RCA. Other possible causes of ST-segment elevation

during the treadmill ECG stress test are coronary artery spasm and the presence of existing Q waves due to previous myocardial infarction.

In conclusion, the treadmill stress ECG test is a reliable screening test for obstructive CAD, especially in patients with an intermediate pretest probability. It is readily available in many centres in Singapore.

## REFERENCES

1. Ministry of Health, Singapore. Singapore Health Facts: Principle Causes of Death [online]. Available at: [https://www.moh.gov.sg/content/moh\\_web/home/statistics/Health\\_Facts\\_Singapore/Principal\\_Causes\\_of\\_Death.html](https://www.moh.gov.sg/content/moh_web/home/statistics/Health_Facts_Singapore/Principal_Causes_of_Death.html). Accessed May 1, 2016.
2. Gibbons RJ, Balady GJ, Beasley JW, et al. ACC/AHA guidelines for exercise testing: executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Exercise Testing). *Circulation* 1997; 96:345-54.
3. Gibbons RJ, Balady GJ, Bricker JT, et al; American College of Cardiology/American Heart Association Task Force on Practice Guidelines. Committee to Update the 1997 Exercise Testing Guidelines. ACC/AHA 2002 guideline update for exercise testing: summary article. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). *J Am Coll Cardiol* 2002; 40:1531-40.
4. Ministry of Health, Singapore. Screening for Cardiovascular Disease and Risk Factors. MOH Clinical Practice Guidelines 1/2011. Singapore: Ministry of Health, 2011.
5. Pinkstaff S, Peberdy MA, Kontos MC, Finucane S, Arena R. Quantifying exertion level during exercise stress testing using percentage of age-predicted maximal heart rate, rate pressure product, and perceived exertion. *Mayo Clin Proc* 2010; 85:1095-100.
6. Jain M, Nkonde C, Lin BA, Walker A, Wackers FJ. 85% of maximal age-predicted heart rate is not a valid endpoint for exercise treadmill testing. *J Nucl Cardiol* 2011; 18:1026-35.
7. Lauer MS, Levy D, Anderson KM, Plehn JF. Is there a relationship between exercise systolic blood pressure response and left ventricular mass? The Framingham Heart Study. *Ann Intern Med* 1992; 116:203-10.
8. Singh JP, Larson MG, Manolio TA, et al. Blood pressure response during treadmill testing as a risk factor for new-onset hypertension. The Framingham heart study. *Circulation* 1999; 99:1831-6.
9. Fletcher GF, Ades PA, Kligfield P, et al; American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee of the Council on Clinical Cardiology, Council on Nutrition, Physical Activity and Metabolism, Council on Cardiovascular and Stroke Nursing, and Council on Epidemiology and Prevention. Exercise standards for testing and training: a scientific statement from the American Heart Association. *Circulation* 2013; 128:873-934.

**ABSTRACT** The treadmill electrocardiogram (ECG) stress test is widely used to screen for obstructive coronary artery disease (CAD). The presence of ST-segment changes, either depression or elevation, on the ECG during the treadmill test often suggests presence of CAD and warrants further management. We herein present three cases, with evidence of ischaemia on the treadmill ECG stress test. In addition, we discuss the use of the treadmill ECG stress test, including its indications, contraindications, reasons for termination and interpretation of the ST-segment changes, heart rate, as well as blood pressure responses to exercise.