

NANO EXPRESS

Open Access



Inhibition of HIV Virus by Neutralizing Vhh Attached to Dual Functional Liposomes Encapsulating Dapivirine

Scarlet Xiaoyan Wang^{1*}, Johan Michiels², Kevin K. Ariën², Roger New³, Guido Vanham² and Ivan Roitt¹

Abstract

Although highly active antiretroviral therapy (HAART) has greatly improved the life expectancy of HIV/AIDS patients, the treatment is not curative. It is a global challenge which fosters an urgent need to develop an effective drug or neutralizing antibody delivery approach for the prevention and treatment of this disease. Due to the low density of envelope spikes with restricted mobility present on the surface of HIV virus, which limit the antibody potency and allow virus mutation and escape from the immune system, it is important for a neutralizing antibody to form bivalent or multivalent bonds with the virus. Liposome constructs could fulfil this need due to the flexible mobility of the membrane with its attached antibodies and the capacity for drug encapsulation. In this study, we evaluated the neutralization activity of a range of liposome formulations in different sizes coated with anti-gp120 llama antibody fragments (Vhhs) conjugated via either non-covalent metal chelation or a covalent linkage. The non-covalent construct demonstrated identical binding affinity to HIV-1 envelope glycoprotein gp120 and neutralizing ability for HIV virus as free Vhh. Although covalently linked Vhh showed significant binding affinity to gp120, it unexpectedly had a lower neutralization potency. This may be due to the comparability in size of the viral and liposome particles restricting the number which can be bound to the liposome surface so involving only a fraction of the antibodies, whereas non-covalently attached antibodies dissociate from the surface after acting with gp120 and free the remainder to bind further viruses. Covalently conjugated Vhh might also trigger the cellular uptake of a liposome-virion complex. To explore the possible ability of the antibody-coated liposomes to have a further function, we encapsulated the hydrophobic antiviral drug dapivirine into both of the non-covalently and covalently conjugated liposome formulations, both of which revealed high efficacy in reducing viral replication *in vitro*. Thus, dual function liposomes may lead to a novel strategy for the prophylaxis of HIV/AIDS by combining the neutralizing activity of Vhh with antiviral effects of high drug concentrations.

Keywords: HIV/AIDS, Liposome, Neutralizing antibody, Drug delivery, Prophylaxis

Background

Human immunodeficiency virus (HIV), responsible for the acquired immunodeficiency syndrome (AIDS), is a lentivirus belonging to the retrovirus family. Infection causes extensive destruction of T-helper cells, macrophages, dendritic cells and other cellular components associated with cell-mediated immunity, eventually leading to AIDS [1, 2]. As a consequence, HIV-infected patients are more susceptible to opportunistic infections [3]. At present, HIV-1 and HIV-2 are the two known types of HIV [4].

HIV-1 is more pathogenic and transmissible and is mainly responsible for the global AIDS pandemic [4]. HIV infection has continued spreading by the transfer of body fluids due to exposure to blood or blood products, by homo- or heterosexual contact, or prenatally from mother-to-child. Since the start of the epidemic, around 78 million people have become infected with HIV and 35 million people have died of AIDS-related illnesses according to the global statistical report from UNAIDS in 2016. In addition, 2.1 million people became newly infected with HIV and 36.7 million people were living with HIV globally in 2015.

Currently, the major means of efficient management of HIV and prevention of its progression towards AIDS

* Correspondence: x.wang@mdx.ac.uk

¹Department of Natural Sciences, Middlesex University, London, UK
Full list of author information is available at the end of the article

is highly active antiretroviral therapy (HAART), which combines a minimum of three antiretroviral drugs from at least two classes. These include nucleoside reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors (PIs), fusion inhibitor, integrase inhibitors and/or entry inhibitors [5]. Although the therapy has improved long-term survival rate, there are associated toxic side effects and emergence of multi-drug resistance. Furthermore, there is a limited access to anti-HIV drugs in the developing countries due to their high costs [6]. Clearly, there remains an urgent demand for an effective and low-cost system for prophylaxis and treatment of HIV infection.

The challenge of developing an effective vaccine or entry-inhibitor drug for HIV infection lies in the rapid mutation of the glycoproteins comprising gp120 and gp41 and the structural features that facilitate antibody evasion [7–13]. Another recent explanation is the spatial arrangement of the envelope spikes on HIV virus, which contributes to its ability to escape the neutralization by antibodies [14]. As is known, most neutralization activities can be attributed to antibody-antigen interactions and the avidity resulting from bivalent binding between two Fabs and two physically linked antigens greatly increase the neutralization potency of the antibody [15–17]. The published studies indicated that there are 4 to 35 spikes per HIV virus particle [18–22] with very restricted mobility [23, 24]. This makes the antibody cross-linking very unlikely because the hinge (10–15 nm) between the two Fabs is not long enough to reach the two spikes randomly distributed on the surface of the virus. When the anti-HIV antibody b12 was tested as an IgA, IgM or IgG, equivalent neutralization potency was observed [14]. Actually, a number of studies have demonstrated that neutralizing antibodies against HIV cannot take the advantage of avidity effects resulting from the IgG cross-linking epitopes on the surface of virus [15–17], while antibodies have been shown to bind bivalently to other viruses comprising closely spaced epitopes [25, 26].

Liposomes are artificially prepared vesicles composed of a lipid bilayer and have been used extensively for drug delivery due to their unique properties [27]. Liposomes are also platforms for the development of multivalent antibody constructs by coupling antibody molecules onto liposome membranes. Chiu et al. have demonstrated that the antibody potency was increased 25-fold by employing the multivalent liposome formulation [28], which even induced cascade effects by cross-linking the antibody complex [29].

Recently, llama heavy-chain antibody fragments (Vhh), which compete with surface CD4 for binding to the HIV-1 envelope glycoprotein, have been described as potent cross-clade HIV-1 candidate microbicides [30, 31]. Such Vhh would be excellent candidates for multivalent

liposome platform due to their specificity, high stability and low production cost [31, 32]. Moreover, these antibody fragments showed very efficient ability to penetrate human tissue by diffusing into the mucosal layer when placed in a vaginal ring [31].

In this study, multivalent liposomal antibody constructs were developed by grafting Vhh (J3), which achieves broad and potent neutralization of HIV-1 via interaction with the CD4-binding site of HIV-1 env [33], onto the liposome membrane via either non-covalent metal chelation or a covalent linkage. The neutralization potency of multivalent presentation of these fragments by liposomes was analyzed by antiviral assays with TZM-bl cells. The results indicated that the avidity of the Vhh was significantly increased while the neutralization potency was compromised when the Vhh was covalently conjugated to the lipid. The additional ability of the liposome construct to effectively deliver dapivirine, a potent non-nucleoside reverse transcriptase inhibitor presently being developed as a vaginal HIV microbicide, was further investigated.

Methods

Materials

Soy phosphatidylcholine (PC), 1,2-dioleoyl-*sn*-glycero-3-[(*N*-(5-amino-1-carboxypentyl)iminodiacetic acid)succinyl] (18:1 DGS-NTA(Ni)) and 1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[maleimide(polyethylene glycol)-2000] (malDSPE-PEG₂₀₀₀) were purchased from Avanti Polar Lipids. Methoxypolyethyleneglycol-di-stearoyl-phosphatidylethanolamine (DSPE-PEG2000, with mPEG MW2000Da) was obtained from Genzyme. Cholesterol (Chol), PBS and Sephadex G25 and G50 were purchased from Sigma. Acetone, methanol and dichloromethane were from Fisher Scientific. RPMI1640, L-glutamine, penicillin-streptomycin and foetal bovine serum (FBS) were from Invitrogen Life Technologies. Dulbecco's minimum essential medium (DMEM) was purchased from Lonza. Vhhs, the variable region fragment of single heavy-chain llama antibodies, including J3 and lab5, were purchased from Quality in Antibodies (Netherlands). Dapivirine, a non-nucleoside reverse transcriptase inhibitor (NNRTI), was kindly provided by the authors Prof. Guido Vanham and Dr. Kevin Arien. CN54 gp140 was kindly provided by Prof. Robin Weiss from University College London.

Cells, Virus and Reagents

TZM-bl cells (obtained through the NIH AIDS Research and Reference Reagent Program, Germantown, MD) were used for the evaluation of the anti-HIV activity of free and liposome conjugated Vhh with and without dapivirine. TZM-bl cells express high densities of CD4, CCR5 and CXCR4. They contain a luciferase reporter gene under control of HIV LTR, which will be transcribed and translated into luciferase protein, if the cells

get infected with HIV and start producing Tat. TZM-bl cells were cultured in DMEM supplemented with 10 % heat-inactivated FBS and 50 µg gentamicin/ml medium, at 37 °C in a humidified 7 % CO₂ environment. Cells were split twice a week and plated at 10⁶ cells in tissue culture flasks and at 10⁴ cells/well in 96-well plates.

The subtype B CCR-5 using reference HIV-1 strain Ba-L was obtained from the NIH AIDS Research and Reference Reagent Program, Germantown, MD. It was grown and titered in human activated peripheral blood mononuclear cells (PBMC). PBMC were isolated from buffy coats from HIV-seronegative blood donors (provided by the Antwerp Blood Transfusion Centre) using Ficoll density gradient centrifugation. PBMC were stimulated for 48 h with 2 µg/ml of phytohemagglutinin (PHA) (Remel, Kent, UK) in RPMI 1640 medium containing 10 % FBS, 50 µg/ml gentamicin and 2 µg/ml polybrene (Sigma-Aldrich, Bornem, Belgium). Subsequently, PBMC were activated for 24 h with 1 ng/ml of interleukin-2 (IL-2) (Gentaur, Brussels, Belgium) in RPMI 1640 medium supplemented with 10 % FBS, 50 µg/ml gentamicin, 2 µg/ml polybrene and 5 µg/ml hydrocortisone (Calbiochem, Leuven, Belgium). PHA, IL-2 activated PBMC.

Liposome Preparation

Liposome compositions used in this study with diameters in brackets were as follows: (a) lipo1 (200 nm), PC/Chol/DSPE-PEG₂₀₀₀/18:1 DGS-NTA in proportions 65/33/1/1 mol%; (b) lipo2 (100 nm), PC/Chol/DSPE-PEG₂₀₀₀/18:1 DGS-NTA in proportions 65/33/1/1 mol% and (c) lipo3 (100 nm), PC/Chol/DSPE-PEG₂₀₀₀/mal-DSPE-PEG₂₀₀₀ in proportions 65/33.5/0/1.5 mol%. Briefly, the lipids with or without dapivirine were dissolved in methanol: dichloromethane 1:2 (v/v) at room temperature. The lipid mixtures were deposited on the side wall of a rotary glass vial by removing the solvent with nitrogen. The resulting dried lipid films were hydrated in 10-mM sodium phosphate buffer pH 7.4. This process led to the spontaneous formation of pegylated liposomes. The liposomes were then extruded through 0.2 or 0.1 µm Anotop 10 filters (Whatman). Excess dapivirine was removed on a Sephadex G25 column [34].

J3 is a neutralizing Vhh, while lab5 is an irrelevant Vhh (irr-Vhh). J3 and lab5 were polyhistidinylated (J3-His and lab5-His) for the non-covalent attachment to liposomes via nitrilotriacetic acid (NTA) linkage. Meanwhile another form of J3 (J3-cys) with extra cysteine in the C-terminus was generated for covalent conjugation to liposomes via interaction with a maleimide group. Vhhs were produced as described before [31]. Non-covalent immunoliposomes were produced by mixing Vhh-His with the liposome suspension in a 1:10 M ratio at room temperature for 1 h on a bench roller. Free thiol groups were generated by incubating J3-cys with

10 mM DTT for 30 min at room temperature. Covalent immunoliposomes were prepared via interaction between J3-cys and maleimide-derivatized lipid inserted in lipo3. The mixture of J3-cys and liposomes with the maleimide group was allowed to conjugate at room temperature overnight. Unreacted maleimide groups were subsequently saturated by adding cysteine. The immunoliposomes were purified by running through a Sephadex G50 column to remove excess cysteine or/and unbound protein.

Characterization of Liposomes

The phospholipid, mainly PC, in the liposome was quantified by the Stewart assay [34]. The number of liposomes per ml was estimated by assuming there were 80,000 phospholipid molecules per liposome of 100 nm or 320,000 phospholipid molecules per liposome of 200-nm diameter [35]. The mean particle size distribution and polydispersity index of liposome suspensions were determined by dynamic light scattering in a Malvern Zetasizer Nano-2S (Malvern Instruments, Malvern, UK). The buffer for zeta potential is MilliQ water pH 7. The stability of liposomes was evaluated by storing the liposomes at 4 °C for 1 month, and samples were taken for size measurement.

SDS-PAGE was employed to confirm the conjugation of Vhh to the liposomes. Fluorescamine was used to quantify the protein conjugated to liposomes [36]. Fluorescamine is a non-fluorescent compound, which reacts rapidly with primary amines in proteins, such as the terminal amino group of peptides and the ε-amino group of lysine, to form highly fluorescent moieties. Basically, the liposome suspension was diluted 1:20 in PBS pH 7. An aliquot of 25 µl of 2 mg/ml fluorescamine was added to a total assay volume of 100-µl diluted liposome suspension. The fluorescence was measured at ex 390 nm/em 475 nm with a cut-off at 455 nm, and the background was subtracted. The concentration of conjugated protein was calculated by comparison with standard samples of the same protein. The concentration of dapivirine in liposomes was determined by measuring the absorbance at 310 nm after incubating the liposomes in ethanol [37, 38].

Enzyme-Linked Immunosorbent Assay

An enzyme-linked immunosorbent assay (ELISA) was carried out to determine the interaction between liposome-linked Vhh and gp140 which is a derivative of gp160 that can be cleaved into gp120 and gp41. Briefly, 1 µg/ml CN54 gp140 was coated at 4 °C overnight on 96-well Maxisorp plates (NUNC, Denmark). After blocking with 200 µl/well Microwell Blocking Buffer with Stabilizer (Rockland), serial dilutions of immunoliposomes in PBS pH 7, free Vhh and negative control liposomes coated with irr-Vhh (lab5) were added to the plates for 1 h. After three washes with PBS,

anti-myc-horse radish peroxidase (HRP) (Roche), which targets the myc-tag in Vhh, was diluted 1:1000 in HRP Conjugate Stabilizer (Rockland) and added to detect the bound Vhh. 3,3',5,5'-Tetramethylbenzidine liquid (Sigma) substrate was added and left to develop for 30 min at room temperature; after which, the reaction was stopped with 2 M H₂SO₄. Absorbance of the product was measured at 450 nm.

In Vitro Cytotoxicity

Human cancer cells, Hela cells (Sigma-Aldrich, UK), were cultured in 96-well plates overnight and treated with different concentrations of liposomes for 48 h. Medium was then removed from the plates, and the cells washed twice with PBS. Fifty microlitres of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) was added per well. Cells were incubated for 30 min at 37 °C with 95 % O₂ and 5 % CO₂. MTT was removed carefully before 100- μ l propanol was added per well to dissolve crystals and incubated for at least 30 min. The absorbance of this coloured solution was quantified by measuring at a wavelength of 570 nm by FLUOstar Omega (BMG Labtech, Aylesbury, UK). Each treatment was conducted in triplicate.

Antiviral Assays for Antibody-Mediated Neutralization and Effectiveness of Liposomal Dapivirine Drug Delivery

The antiviral activity of the compounds was determined by pre-incubating 10⁴ TZMbl cells/well in a 96-well plate for 2 h at 37 °C and 7 % CO₂ to maintain the optimal pH with or without a serial dilution of compounds. Next, 200 TCID₅₀ of HIV-1 BaL virus was added to each well, and cultures were incubated for 48 h before luciferase activity was quantified. To this end, 120 μ L of supernatant was removed, 75 μ l of the luciferase substrate Steadylite (Perkin Elmer, Life Sciences, Zaventem, Belgium) was added to the wells, and the plates were incubated at room temperature on an orbital shaker for 10 min. Next, the luciferase activity was measured using a TriStar LB941 luminometer (Berthold Technologies GmbH & Co. KG., Bad Wildbad, Germany) and expressed in relative light units (RLU).

Each compound was tested in triplicate and each experiment was repeated in three independent runs. Antiviral activity was expressed as the percentage of viral growth compared to the control and plotted against the compound concentration. Next, non-linear regression analysis was used to calculate the EC₅₀, using GraphPad Prism 5.03 using non-linear regression (GraphPad Software, San Diego, CA, USA).

Results

Liposome Preparation and Characterization

Three groups of liposome suspensions: lipo1, lipo2 and lipo3, were prepared to investigate whether the size or

Vhh density of the liposomes will contribute to the neutralization efficiency of the liposomes for HIV virus. Lipo1 was of 200-nm diameter while lipo2 and lipo3 were of 100-nm diameter. His-tagged Vhh (J3-His and lab5-His) were conjugated to lipo1 or lipo2 by interaction with the nickel-chelating lipid 18:1 DGS-NTA-Ni which was incorporated into the liposome bilayer (Fig. 1a). The sulphhydryl-reactive lipid (malDSPE-PEG₂₀₀₀) in lipo3 allowed covalent attachment of the reduced J3-cys to the liposome (Fig. 1b). Immunoliposomes were then purified by size exclusion chromatography to remove the unbound Vhh and impurities. The conjugation and purification were successful in all cases, and the resulting products were analyzed by SDS-PAGE (Fig. 2). The conjugated Vhhs gave bands of higher molecular weight than the free Vhhs, and this was due to the contribution of the conjugated lipids including 18:1 DGS-NTA-Ni or malDSPE-PEG₂₀₀₀. More than 95 % of the Vhh was shown to be conjugated to the liposomes by the fluorescamine method described above [34, 36]. The density of Vhh on liposome surfaces was shown in Table 1.


Immunoliposomes with or without NNRTI dapivirine were produced to investigate whether conjugated Vhh could show higher avidity to antigen and improve the efficacy of Vhh in HIV viral inhibition, as compared to free Vhh. Dapivirine was dissolved in the solvent with lipid mixture when formulating the liposomes. Unincorporated dapivirine was removed by size exclusion chromatography, and the concentration of dapivirine was determined by UV spectrometry.

The final immunoliposomes were characterized, and the data was shown in Table 1. Phospholipid (PC) was detected in the liposomes by Stewart assay. It was estimated to result in approximately 160 Vhh molecules per 200-nm liposome or 95–125 Vhh molecules per 100-nm liposome on the basis that one 100-nm liposome contains 80,000 phospholipid molecules and one 200-nm liposome contains 320,000 phospholipid molecules [29, 35]. This resulted in similar Vhh density on the surface of the liposomes.

PEGylated liposome suspensions were prepared by the lipid film rehydration method [34] and extrusion technique, with a final mean diameter of 200–230 nm or 120–140 nm (polydispersity index of 0.1–0.2). The negative charge of the liposome formulations were attributed to the DSPE-PEG₂₀₀₀ or malDSPE-PEG₂₀₀₀ which accounts for 1–1.5 % mol of the lipid. The liposomes were stable for at least 1 month, with no significant change in size or charge (Fig. 3).

In Vitro Cytotoxicity of the Liposome

Potential cytotoxicity of the liposomes was estimated from cell viability relative to control Hela cells treated with medium only. MTT assay demonstrated that cell



viability was between 80 and 120 % in relation to the control samples (Fig. 4). One-way ANOVA statistical analysis showed there is no evidence of liposome toxicity at concentrations up to 1 mg/ml ($p > 0.05$).

Analysis of Binding of Liposomal Vhhs to Recombinant HIV gp140 by ELISA

As Vhh competed with soluble CD4 (sCD4) for binding to gp120 [32], an ELISA assay was carried out to determine whether the Vhh conjugated to liposomes showed compromised binding affinity or increased avidity to gp140 (CN54). Liposome-linked J3 was compared with

the free J3 at the same concentrations, together with the irrelevant Vhh (lab5-His) and liposome-coated the same irrelevant Vhh as negative controls. J3-His and J3-cys revealed similar binding affinities for gp140 as expected (Fig. 5). No improved or compromised affinity was observed for lipo1-J3 and lipo2-J3 compared to free J3-His, implying size differences are not influencing the avidity despite similar Vhh densities on the liposome surfaces. Meanwhile, one-way ANOVA statistical analysis showed that J3 conjugated to lipo3 demonstrated much higher avidity to gp140 than free J3-cys ($p < 0.001$), indicating covalent attachment greatly facilitated J3 interacting with




Table 1 Characterisation of the immunoliposomes

Liposome	PC conc. (mg/ml) ^a	Dapivirine (μM)	No. of Vhh molecules per liposome	Vhh (μg/ml)	Diameter (nm) ^a	Zeta potential (mV) ^a
Lipo1-lab5	11.937 ± 0.001	–	160	126.3	228.0 ± 1.0	–45.0 ± 0.9
Lipo1-J3	11.708 ± 0.008	–	160	118.4	223.4 ± 4.5	–37.4 ± 0.5
Lipo2-lab5	5.974 ± 0.005	–	95	150.0	130.9 ± 1.4	–38.7 ± 0.7
Lipo2-J3	6.149 ± 0.003	–	108	159.9	131.2 ± 1.9	–38.5 ± 0.2
Lipo2-dapi	6.478 ± 0.001	6.5	–	–	122.9 ± 1.8	–46.0 ± 0.9
Lipo2-dapi-J3	5.548 ± 0.002	5.5	125	169.6	123.4 ± 1.8	–38.5 ± 0.4
Lipo3 control	7.573 ± 0.003	–	–	–	130.9 ± 1.7	–42.5 ± 0.4
Lipo3-J3	6.456 ± 0.004	–	121	168.2	142.2 ± 3.2	–36.4 ± 0.4
Lipo3-dapi	6.675 ± 0.003	6.7	–	–	128.2 ± 2.6	–41.7 ± 0.5
Lipo3-dapi-J3	6.806 ± 0.001	6.8	108	157.0	139.1 ± 1.8	–37.8 ± 0.6

Dapi: dapivirine

^aData are means ± standard deviations of three replicate measurements in one representative experiment of three independent experiments

gp140. In contrast, irrelevant negative controls demonstrated no binding to gp140.


Characterization of Neutralizing Antibody by Antiviral Assay


The neutralization activity of liposomal Vhhs and free Vhhs for HIV virus was evaluated by subsequent incubation with TZM-bl cells. Lipo1-J3 and lipo2-J3 revealed similar potencies for neutralization compared to free J3. Surprisingly, lipo3-J3 only provided 60 % of viral inhibition at J3 concentration of 286 ng/ml, whereas lipo1-J3 and lipo2-J3 inhibited approximately 90 % of viral growth at the same concentration (Fig. 6a). No viral growth reduction was observed for control liposomes, liposomes coated with irrelevant Vhhs or free Vhhs alone. There was no significant difference in neutralizing activities between

lipo1-J3 and lipo2-J3 compared to free J3-His (Fig. 6b), revealing no contribution from the size differences. However, covalently conjugated J3 (lipo3-J3) had a significantly higher EC 50 (Fig. 6b), indicating less potency for neutralization than non-covalently attached J3 or free J3. The result is unexpected because the ELISA data revealed significantly increased binding affinity of lipo3-J3 to gp140 which should contribute to the neutralization activity of conjugated J3.


Comparison of Antiviral Effects of Liposomal Dapivirine with Free Dapivirine

We compared the antiviral effects of different formulations of liposomal dapivirine with free dapivirine. The dose-response curves indicated increased antiviral effect of liposomal dapivirine compared with free dapivirine,





although no significant difference was observed between lipo2 and lipo3 with or without neutralizing Vhhs (Fig. 7a). Furthermore, there was approximately 76 % of viral growth at a concentration of 1-nM free dapivirine, whereas only 50–56 % of viral growth was detected with liposomal dapivirine at this concentration. Calculated EC50 (Fig. 7b) confirmed that liposomal dapivirine is more effective than free dapivirine in inhibiting HIV virus. However, the antiviral effects were dominated by dapivirine since no difference was observed between the liposome preparations with or without neutralizing J3.




Discussion

The treatment of HIV remains a serious challenge due to the high genetic variation of the virus and associated toxic side effects of the antiviral drugs. HIV-1 neutralizing antibodies have been considered to be critical for vaccine development and prevention of HIV infection. Liposomes provided another approach shown to have potential for HIV prophylaxis [39]. In the present study, a combined platform with both liposome and antibody fragments was developed to investigate the antiviral efficiency of the multivalent liposomal formulation.

Previous studies have demonstrated that Vhhs produced by llamas (*Lama glama*) exhibited comparable affinity and specificity for gp120 to conventional immunoglobulins despite their much smaller size. This smaller size of Vhhs is an advantage over conventional immunoglobulins as it has been shown to reduce their immunogenicity [33]. In particular, J3 Vhh, which achieved broad and potent neutralization of HIV-1 via interaction with the CD4-binding site, neutralized 96 of 100 tested HIV-1 strains and chimeric simian-HIV strains with HIV subtypes B and C env [33]. Unlike most of the enveloped viruses which express large numbers of closely spaced spikes on the surface, HIV virus has a low spike surface density (4 to 35 spikes) and the arrangement of these spikes is likely to be static over the time periods relevant to neutralization [14]. Therefore, the location and arrangement of the neutralizing Vhhs on the liposome surfaces is critical for their interaction with the virus particles.


To optimize the size of the liposomal platform, J3 was non-covalently conjugated to the surface of two liposome formulations with diameters of 100 or 200 nm, each with approximately the same Vhh densities. Metal chelation via NTA has been employed to attach peptide and protein antigens to liposomes to compare with covalent linkage for their ability to elicit antibody responses [40]. Pegylated liposomes were employed to ensure the stability and minimal immunogenicity of the platform. To check that J3 was available and not masked by the PEG coating on the liposome surface, appropriate access of J3 to gp140 was verified by ELISA (Fig. 5). The identical binding curves demonstrated that lipo1-J3 and lipo2-J3 bound to gp140 with the same avidity as free J3, showing that the different geometrical location of J3 on the surface of spherical liposomes did not compromise its interaction with gp140. Surprisingly, no enhanced avidity was observed for these liposomal J3 constructs compared to free J3. This may be due to the charge on the nickel-histidine complex which would inhibit close association of the J3 antibody fragments to generate multivalent attachment to the gp140 on ELISA surface. This presumption is confirmed by the neutralization results indicating lipo1-J3 and lipo2-J3 inhibited HIV infection in a dose-response manner similar to free J3



in antiviral assay (Fig. 6). There was no significant difference in the EC50 between lipo1-J3, lipo2-J3 and free J3, suggesting no improved antiviral potency for the non-covalently anchored Vhh independent of liposome sizes.

J3-cys was then covalently attached to liposomes to produce lipo3-J3 (Fig. 1). One hundred-nanometre diameter liposomes were used for economic reasons and since liposomes of this size are broadly applied as drug delivery system [41, 42]. The binding curves for lipo2-J3 and lipo3-

J3 were distinct from each other, indicating that their interactions with gp140 were qualitatively different with superior avidity for lipo3-J3. This implied that a multivalent platform had been formed with cross-linking or multimerization of Vhhs. This is not surprising since the absence of the nickel complex allows the free movement and geometrical arrangement of J3-lipid complex within the lipid liposome membrane so permitting the formation of multivalent liposomal Vhh constructs through cross-linking the envelope



spikes which are sparsely distributed on the virion surface and remained static over the time periods. Unexpectedly, the antiviral potency of lipo3-J3 was remarkably reduced as evaluated in the antiviral assay compared to lipo1-J3, lipo2-J3 or free J3, the EC50 for lipo3-J3 being approximately four times higher than that for lipo2-J3 or free J3. We hypothesize that in the case of non-covalent linkage, J3 was dissociated from liposome after it bound to gp140 on the virion surface because the interaction between J3 and gp140 was so strong ($K \cong 10^{-9}$ M $\sim 10^{-12}$ M compared to $K \cong 10^{-5}$ M $\sim 10^{-6}$ M for attachment to the nickel-chelating lipid) [32, 40, 43], that J3 was pulled apart from the nickel complex inserted in the liposome membrane. The remaining surface antibody molecules were now free to react with further virus, the net effect being that lipo2-J3 neutralized the virus as efficiently as free J3. Although the covalent attachment of the J3 in lipo3 enhanced lipo3-J3-gp140 interaction, this fixed the viruses to the surface of the liposome and due to the comparability in size of the interacting particles, the number bound to the liposome surface will be restricted and the small number of viral spikes involved will leave a large fraction of the antibodies remaining uncombined which were otherwise available in the non-covalently linked constructs.

In addition to the neutralization capacity of antibody-coated liposomes, we investigated whether the liposomes could have the extra-function of delivering antiviral drugs intracellularly, in particular using high doses of otherwise poorly soluble hydrophobic drugs whose encapsulation in the liposome would limit its general toxicity. All of these dapivirine containing liposomes revealed higher efficacy in reducing viral replication than free dapivirine, although no significant differences were observed between them i.e. irrespective of the presence of irrelevant or J3 Vhh on the surface. This suggested that the antiviral activities of J3 were masked by the high dose of dapivirine and that the liposome platform per se is an efficient delivery system for dapivirine.

Conclusions

This study demonstrated that antibody-conjugated liposomes can provide anti-HIV viral defence in two ways. First, non-covalently conjugated liposomal J3 neutralized HIV virus as efficiently as free J3. It was also shown that incorporation of the lipophilic drug dapivirine in liposomes resulted in a higher level of virus inactivation in vitro than observed after exposure of cells to free drug dissolved in DMSO—an effective method of solubilizing the drug, but not one which can be used in vivo. Since we have demonstrated that immunoliposomes with covalently grafted anti-CD4 Vhhs facilitated cellular uptake of liposomes by lymphoma cells overexpressing CD4 on the surface (data not shown), it follows that a combined

liposome platform encapsulating dapivirine with non-covalently attached J3 and covalently linked anti-CD4 Vhh could be a powerful potential candidate for HIV prevention where J3 neutralizes free virus and dissociates from liposome while the anti-CD4 Vhh coated liposome efficiently delivers the antiviral drug to HIV target cells overexpressing CD4.

Acknowledgements

This work was supported by the CHAARM EU consortium to develop vaginal HIV microbicides. We would like to thank Dr. Lei Wang for his assistance in statistical analysis.

Authors' Contributions

SXW and IR are working in Middlesex University. SXW has designed the study, prepared the liposome formulations for all the experiments in this study and wrote the manuscript. IR provided consultation on the whole study including the design and use of the liposome and Vhh antibody fragments. JM, KKA and GV are working in Institute of Tropical Medicine, Belgium. JM has carried out the antiviral assay in this study. KKA has designed the antiviral assay. GV provided consultation on HIV infection and treatment. RN is the chief scientist in Proxima Concepts Ltd. He has provided consultation on the liposome preparation. All authors read and approved the final manuscript.

Competing Interests

The authors declare that they have no competing interests.

Author details

¹Department of Natural Sciences, Middlesex University, London, UK. ²Institute of Tropical Medicine, Antwerp, Belgium. ³Proxima Concepts. Ltd, London, UK.

Received: 3 February 2016 Accepted: 16 July 2016

Published online: 28 July 2016

References

- Wyatt R, Sodroski J (1998) The HIV-1 envelope glycoproteins: fusogens, antigens, and immunogens. *Science* 280:1884–8
- Chan DC, Kim PS (1998) HIV entry and its inhibition. *Cell* 93:681–4
- Wong HL, Chattopadhyay N, Wu XY, Bendayan R (2010) Nanotechnology applications for improved delivery of antiretroviral drugs to the brain. *Adv Drug Deliv Rev* 62:503–17
- Freed EO (2001) HIV-1 replication. *Somat Cell Mol Genet* 26:13–33
- Da Cunha J (2015) Impact of antiretroviral therapy on lipid metabolism of human immunodeficiency virus-infected patients: old and new drugs. *World J Virol* 4:56
- Rao KV1, Chitturi RT2, Kattappagari KK2, Kantheti LP2, Pooarla C2, Baddam V (2015) Impact of highly active antiretroviral therapy on oral manifestations of patients with human immunodeficiency virus_acquired immuno deficiency synd. *Indian J Sex Transm Dis* 36(1):35–39
- Wei X, Decker JM, Wang S et al (2003) Antibody neutralization and escape by HIV-1. *Nature* 422:307–12
- Kwong PD, Doyle ML, Casper DJ et al (2002) HIV-1 evades antibody-mediated neutralization through conformational masking of receptor-binding sites. *Nature* 420:678–82
- Kwong PD, Wyatt R, Robinson J et al (1998) Structure of an HIV gp120 envelope glycoprotein in complex with the CD4 receptor and a neutralizing human antibody. *Nature* 393:648–59
- Labrijn AF, Poignard P, Raja A et al (2003) Access of antibody molecules to the conserved coreceptor binding site on glycoprotein gp120 is sterically restricted on primary human immunodeficiency virus type 1. *J Virol* 77:10557–65
- Wyatt R, Desjardins E, Olshevsky U et al (1997) Analysis of the interaction of the human immunodeficiency virus type 1 gp120 envelope glycoprotein with the gp41 transmembrane glycoprotein. *J Virol* 71:9722–31
- Moore JP, Sodroski J (1996) Antibody cross-competition analysis of the human immunodeficiency virus type 1 gp120 exterior envelope glycoprotein. *J Virol* 70:1863–72

13. Starcich BR, Hahn BH, Shaw GM et al (1986) Identification and characterization of conserved and variable regions in the envelope gene of HTLV-III/LAV, the retrovirus of AIDS. *Cell* 45:637–48
14. Klein JS, Bjorkman PJ (2010) Few and far between: how HIV may be evading antibody avidity. *PLoS Pathog* 6:e1000908
15. Zhang M-Y, Xiao X, Sidorov IA et al (2004) Identification and characterization of a new cross-reactive human immunodeficiency virus type 1-neutralizing human monoclonal antibody. *J Virol* 78:9233–42
16. Klein JS, Gnanaprasadam PNP, Galimidi RP et al (2009) Examination of the contributions of size and avidity to the neutralization mechanisms of the anti-HIV antibodies b12 and 4E10. *Proc Natl Acad Sci U S A* 106:7385–90
17. Ofek G, Tang M, Sambor A et al (2004) Structure and mechanistic analysis of the anti-human immunodeficiency virus type 1 antibody 2F5 in complex with its gp41 epitope. *J Virol* 78:10724–37
18. Zhu P, Chertova E, Bess J et al (2003) Electron tomography analysis of envelope glycoprotein trimers on HIV and simian immunodeficiency virus virions. *Proc Natl Acad Sci U S A* 100:15812–7
19. Zhu P, Liu J, Bess J et al (2006) Distribution and three-dimensional structure of AIDS virus envelope spikes. *Nature* 441:847–52
20. Liu J, Bartschaghi A, Borgnia MJ et al (2008) Molecular architecture of native HIV-1 gp120 trimers. *Nature* 455:109–13
21. Chertova E, Bess JW, Crise BJ et al (2002) Envelope glycoprotein incorporation, not shedding of surface envelope glycoprotein (gp120/SU), is the primary determinant of SU content of purified human immunodeficiency virus type 1 and simian immunodeficiency virus. *J Virol* 76:5315–25
22. Layne SP, Merges MJ, Dembo M et al (1992) Factors underlying spontaneous inactivation and susceptibility to neutralization of human immunodeficiency virus. *Virology* 189:695–714
23. Yu X, Yuan X, Matsuda Z et al (1992) The matrix protein of human immunodeficiency virus type 1 is required for incorporation of viral envelope protein into mature virions. *J Virol* 66:4966–71
24. Bhatia AK, Kaushik R, Campbell NA et al (2009) Mutation of critical serine residues in HIV-1 matrix result in an envelope incorporation defect which can be rescued by truncation of the gp41 cytoplasmic tail. *Virology* 384:233–41
25. Smith TJ, Olson NH, Cheng RH et al (1993) Structure of a human rhinovirus-bivalently bound antibody complex: implications for viral neutralization and antibody flexibility. *Proc Natl Acad Sci U S A* 90:7015–8
26. Icenogle J, Shiwen H, Duke G et al (1983) Neutralization of poliovirus by a monoclonal antibody: kinetics and stoichiometry. *Virology* 127:412–25
27. Duncan R (2003) The dawning era of polymer therapeutics. *Nat Rev Drug Discov* 2:347–60
28. Chiu GNC, Edwards LA, Kapanen AI et al (2007) Modulation of cancer cell survival pathways using multivalent liposomal therapeutic antibody constructs. *Mol Cancer Ther* 6:844–55
29. Oliveira S, Schiffelers RM, van der Veeken J et al (2010) Downregulation of EGFR by a novel multivalent nanobody-liposome platform. *J Control Release* 145:165–75
30. Koh WWL, Steffensen S, Gonzalez-Pajuelo M et al (2010) Generation of a family-specific phage library of llama single chain antibody fragments that neutralize HIV-1. *J Biol Chem* 285:19116–24
31. Gorlani A, Brouwers J, McConville C, et al. (2011) Llama antibody fragments have good potential for application as HIV type 1 topical microbicides. *AIDS Res Hum Retroviruses* 28(2):198-205.
32. Forsman A, Beirnaert E, Aasa-Chapman MMI et al (2008) Llama antibody fragments with cross-subtype human immunodeficiency virus type 1 (HIV-1)-neutralizing properties and high affinity for HIV-1 gp120. *J Virol* 82:12069–81.
33. McCoy LE, Quigley AF, Strokappe NM et al (2012) Potent and broad neutralization of HIV-1 by a llama antibody elicited by immunization. *J Exp Med* 209:1091–103
34. New RR (1990) *Liposomes a practical approach*. 1st ed. Oxford: Oxford University Press.
35. Paolo GD, Wenk MR (2012) *Lipids*. 1st ed. San Diego: Academic press.
36. Shew RL, Deamer DW (1985) Hemoglobin and alkaline phosphatase were each encapsulated in phosphatidylcholine liposomes using a dehydration-rehydration cycle for liposome formation. In this method, liposomes prepared by sonication are mixed in aqueous solution with the solute desi. *Science* (80-) 816:1–8
37. das Neves J, Sarmento B, Amiji MM, Bahia MF (2010) Development and validation of a rapid reversed-phase HPLC method for the determination of the non-nucleoside reverse transcriptase inhibitor dapivirine from polymeric nanoparticles. *J Pharm Biomed Anal* 52:167–72
38. (2010) Linear order release polymer. <http://www.google.com/patents/EP2244782A2?cl=en>.
39. Malavia NK, Zurakowski D, Schroeder A et al (2011) Liposomes for HIV prophylaxis. *Biomaterials* 32:8663–8
40. Watson DS, Platt VM, Cao L et al (2011) Antibody response to polyhistidine-tagged peptide and protein antigens attached to liposomes via lipid-linked nitrilotriacetic acid in mice. *Clin Vaccine Immunol* 18:289–97
41. Clayton R, Ohagen A, Nicol F et al (2009) Sustained and specific in vitro inhibition of HIV-1 replication by a protease inhibitor encapsulated in gp120-targeted liposomes. *Antiviral Res* 84:142–9
42. Gaddy DF, Lee H, Zheng J et al (2015) Whole-body organ-level and kidney micro-dosimetric evaluations of (64)Cu-loaded HER2/ErbB2-targeted liposomal doxorubicin ((64)Cu-MM-302) in rodents and primates. *EJNMMI Res* 5:24
43. Nieba L, Nieba-Axmann SE, Persson A et al (1997) BIACORE analysis of histidine-tagged proteins using a chelating NTA sensor chip. *Anal Biochem* 252:217–28

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
