

PEARLS

Major Sensing Proteins in Pathogenic Fungi: The Hybrid Histidine Kinase Family

Anaïs Hérivaux¹, Yee-Seul So², Amandine Gastebois¹, Jean-Paul Latgé³, Jean-Philippe Bouchara^{1,4}, Yong-Sun Bahn^{2*}, Nicolas Papon^{1*}

1 Université d'Angers, Groupe d'Etude des Interactions Hôte-Pathogène, Angers, France, **2** Department of Biotechnology, College of Life Science and Biotechnology, Yonsei University, Seoul, Republic of Korea, **3** Unité des Aspergillus, Institut Pasteur, Paris, France, **4** Laboratoire de Parasitologie—Mycologie, Centre Hospitalier Universitaire d'Angers, Angers, France

* ysbahn@yonsei.ac.kr (YSB); nicolas.papon@univ-angers.fr (NP)

Milestones in the Discovery of Histidine Kinases

The pioneering discovery of histidine kinases (HKs) from *Escherichia coli* was made in the early 1980s with the identification of the *envZ* gene [1] (Fig 1A). Further biochemical characterization of the corresponding protein revealed a new type of protein kinase activity, namely HK, to add to the well-known serine/threonine and tyrosine kinases. For a decade, HKs were considered to be restricted to bacteria, but in the 1990s, HKs were identified in plants [2], fungi [3], archaea [4], cyanobacteria [5], and amoebae (Fig 1A) [6]. Soon after, evidence suggested that HKs regulate essential processes in pathogenic bacteria and fungi [7]. Although some HKs appear to be present in humans, typical bacterial or fungal HK-like sensor proteins have not been reported yet in mammals [8], promoting these proteins as ideal targets for future therapeutics [9].

It is now accepted that HKs are involved in cell signaling systems referred to as His-to-Asp phosphorelays and several canonical schemes depicting transduction pathways involving HKs in bacteria, amoebae, plants, and fungi have emerged (Fig 1B). To date, HKs act as primary sensors for various environmental stimuli, and, upon activation, initiate phosphate transfer events between various proteins, leading to an adaptive response. Although these mechanistic models are largely described in bacteria and plants, limited evidence is available for amoebae and fungi.

Structure and Classification of Fungal Histidine Kinases

The basic structure of fungal HKs is now well established. They are composed of three main regions (Fig 2A). The first region corresponds to a highly variable N-terminal sequence that determines which stimulus is perceived by the HK. This region is referred to as the “sensor domain.” The central region is the transmitter domain consisting of both histidine kinase A (HisKA) (dimerization/phosphoacceptor) (Fig 2A) and cognate histidine kinase-like ATPase catalytic (HATPase_c) (Fig 2A) subdomains. HisKA domains include an H-box, usually containing a phosphorylatable histidine (see Fig 2A), and an X-box. The HATPase_c subdomain displays four distinct boxes: N-, G1- (sometimes called D-box), F-, and G2-boxes. The third and final region, well conserved in fungal HKs, is the C-terminal receiver domain (RD) (Fig 2A) characterized by the presence of a three amino-acids signature (DDK) including a phosphorylatable aspartate residue (see Fig 2A) [10]. Another important aspect is that fungal HKs are generically defined as hybrid HKs since the transmitter domain is fused to the receiver domain (Fig 1B).



OPEN ACCESS

Citation: Hérivaux A, So Y-S, Gastebois A, Latgé J-P, Bouchara J-P, Bahn Y-S, et al. (2016) Major Sensing Proteins in Pathogenic Fungi: The Hybrid Histidine Kinase Family. *PLoS Pathog* 12(7): e1005683. doi:10.1371/journal.ppat.1005683

Editor: Deborah A. Hogan, Geisel School of Medicine at Dartmouth, UNITED STATES

Published: July 28, 2016

Copyright: © 2016 Hérivaux et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by National Research Foundation of Korea grants (2015R1A2A1A15055687) from MEST and the Strategic Initiative for Microbiomes in Agriculture and Food funded by Ministry of Agriculture, Food and Rural Affairs (916006-2) (to YSB). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.




Fig 1. Milestones in the discovery of histidine kinases (HKs) and currently accepted canonical signaling pathways involving HKs in prokaryotes, plants, amoebae, and fungi. (A) Historical timeline depicting the evolution of knowledge concerning HKs. In the order of appearance from left to right: the EnvZ osmosensor in *Escherichia coli* [1], the phytohormone ethylene receptor ETR1 in *Arabidopsis thaliana* [2], the Sln1 osmosensor in *Saccharomyces cerevisiae* [3], the RcaE cyanobacteriochrome [5], the discadenine receptor DhkA in *Dictyostelium discoideum* [6], the quorum sensing-associated Chk1 in *Candida albicans* [26], the virulence factor Fos-1 in *Aspergillus fumigatus* [27], the dimorphism-related Drk1 in *Blastomyces dermatitidis* [28], the Bos-1 osmosensor in *Botrytis cinerea* [29], the *Cryptococcus neoformans* Tco1 and Tco2 (a first functionally characterized dual HK) [30], and the *Metarhizium robertsii* Mhk1 [31]. (B) Canonical schemes depicting signaling pathways involving HKs in prokaryotes, amoebae, plants, and fungi. In prokaryotes, most signaling pathways involving HKs simply consist of two components. The perception of a stimulus by the sensor domain (grey box) induces the autophosphorylation of a conserved histidine (H, pink box) by the catalytic domain (N G₁ F G₂, yellow box) in the HK. The phosphate is then transferred to a conserved aspartate residue (D) located on a cytosolic response regulator (RR) and the activated RR governs the expression of response genes. In plant cells, most (but not all) HKs constitute the initial sensing proteins of a four-step phosphorelay signaling pathway involving phosphorylation events of two downstream elements, i.e., histidine phosphotransfer shuttle proteins (Hpt) and RRs. Note that a first phosphorylatable receiver domain (DDK) is fused to the catalytic domain (N G₁ F G₂) in the HK. As observed for the archetypal two-component system in prokaryotes, the activated RR governs the expression of response genes. In amoebae, similarly to plants, a four-step phosphorelay signaling pathway is observed, but this latter controls a downstream cyclic AMP pathway. Finally, in fungi, knowledge is very fragmented, but initial studies in *Saccharomyces cerevisiae* have demonstrated that HKs also constitute the initial sensing proteins of a four-step phosphorelay signaling pathway that governs a cascade of mitogen-activated protein (MAP) kinases.

doi:10.1371/journal.ppat.1005683.g001




Fig 2. Structure, classification, function, and distribution of fungal HKs at a glance. (A) Basic structure of fungal HKs. They are composed of three main regions: a highly variable N-terminal sequence that determines which stimulus is perceived by the HK ("sensor" domain), a central transmitter domain consisting of both histidine kinase A (HisKA) and cognate histidine kinase-like ATPase catalytic subdomains (HATPase_c), and a C-terminal receiver domain showing a three amino-acids signature (DDK). Fungal HKs are currently categorized in 16 groups according to the sequence analysis of two regions: the H-box signature (alignment of group III HKs from major pathogenic fungi are provided in the right panel) containing the phosphorylatable histidine (red background) and the combination of domains found in the N-terminus. Domains that compose the N-terminal sensor region of major HK groups, whose

functions have been at least partially characterized, are provided on the left panel. Abbreviations: HAMP, Histidine kinases-Adenylate cyclases-Methyl accepting proteins and Phosphatases; TH, Transmembrane Helix; PAS, Period circadian protein-Aryl hydrocarbon receptor nuclear translocator protein-Single-minded protein; GAF, cGMP-specific phosphodiesterases-Adenylyl cyclases-FhlA; PHY, Phytochrome; S/Tkrd, Serine/Threonine kinase related domain. (B) Some notable functions currently assigned to the prominent groups III, VI, VIII, X, and dual HKs (for more details see [11]). (C) Quantitative and qualitative distribution of HKs in fungal clades. The total number of HKs and the occurrence of major HK groups are provided for a panel of representative well-known pathogenic and non-pathogenic (n-p) fungi. A grey box signifies that the corresponding HK groups are not observed in the species. A colored box signifies that a unique member of the corresponding HK group is found in the species and the number of members is only indicated when many members are observed. Abbreviations: A, Ascomycota; B, Basidiomycota; Pe, Pezizomycotina; Sa, Saccharomycotina; Ta, Taphrinomycotina; Us, Ustilaginomycotina; Pu, Pucciniomycotina; Ag, Agaricomycotina; Mu, Mucoromycotina; Mi, Microsporidia.

doi:10.1371/journal.ppat.1005683.g002

Sequence analysis of the transmitter domain and the sensor domain have aided in categorizing fungal HKs into 16 groups [11]. More precisely, the H-box signature (pink region in Fig 2A) containing the phosphorylatable histidine (red residue in Fig 2A) is often sufficient to classify an HK into its respective group [12]. However, since several fungal HK groups are divergent [12], it is essential to verify that the N-terminus of a newly identified HK contains most of the domains commonly found in HKs already assigned to the corresponding group. Thus, this means that each fungal HK group is defined by a relatively well-conserved H-box signature and a specific combination of domains within the N-terminus. For instance, the extensively studied group III fungal HKs are characterized by both a highly conserved H-box signature (alignment of group III HKs from prominent pathogenic species is provided in the Fig 2A) and tandem repeats of Histidine kinases-Adenylate cyclases-Methyl accepting proteins and Phosphatases (HAMP) domains within the N-terminus. The combination of domains that compose sensor region of functionally well-characterized HK groups (see following section) is provided in Fig 2A.

Functions Currently Assigned to Fungal Histidine Kinases

HKs are key signaling proteins involved in the perception and the transduction of a wide range of environmental stimuli in prokaryotes, amoebae, and plants. HKs are also widespread in the fungal kingdom, but their precise roles in the regulation of physiological processes remain fragmentary. The function of several fungal HKs has been studied over the last 15 years using classical reverse genetic approaches, mainly by creating targeted mutants and comparing them with the wild type strain (for a review see [11]). This approach has given insight into the involvement of some fungal HK groups in the response and adaptation to environmental conditions. The most striking associations reported between fungal HKs and physiological processes are summarized below and in Fig 2B.

Some fungal HKs regulate osmotic adaptation by governing the high-osmolarity glycerol (HOG) mitogen-activated protein kinase (MAPK) pathway (Fig 1B). This includes members of HK groups VI and III (Fig 2A). Deletion of these groups of HKs may cause cell lethality not because of the absence of the osmosensing function, but because of hyperactivation of the HOG pathway causing overaccumulation of intracellular glycerol levels [13,14].

Some other fungal HKs display a key role in response to oxidative stress. This function is particularly attributed to the group X HKs (Fig 2A) but was also reported for a few HKs belonging to other groups. Interestingly, the group X HK members in pathogenic fungi that were initially found to be involved in chemical oxidant adaptation also protected the fungal cells against phagocytic cells of their host [15].

Fungal phytochromes are light-perceiving HK receptors, as is the case in bacteria and cyanobacteria but not in plants, where phytochromes curiously evolved for serine/threonine kinase activity [16]. These proteins commonly belong to group VIII (also called the Fph group) (Fig 2A) and aid red-light sensing in Ascomycota and Basidiomycota while other types of

photoreceptors mediate shorter wavelength light sensing (for a review see [17]). Notably, it has been recently reported that a phytochrome HK in *Aspergillus nidulans*, FphA, modulates sexual differentiation through the HOG MAPK pathway by sensing red-light [18].

Adding to the aforementioned chemical and physical stressors/stimuli that can be perceived by HKs, it is important to remember that a number of HKs were also associated with essential fungal developmental programs. Specifically, some HK deletion strains, notably from group III or X HKs (and to a lesser extent group VI), displayed strong physiological perturbations, such as altered hyphal growth and reduced conidiation in pathogenic fungi and morphogenetic switch inability in dimorphic fungal pathogens. For instance, while a *Candida albicans* group X HK mutant strain is unable to switch from yeast to hyphae, *Blastomyces dermatitidis*, *Histoplasma capsulatum*, or *Penicillium marneffeii* group III HK mutant strains are incapable of the hyphae to yeast transition. Furthermore, deletion of group III HK also diminishes filamentous growth and mating efficiency in *Cryptococcus neoformans*. It is important to point out that these morphogenetic perturbations seen in mutants are often associated with an alteration in cell wall composition or integrity resulting in a reduced virulence [19].

Finally, some other interesting roles were reported for a few series of fungal HKs, but they currently appear species-specific including melanin production, adaptation to hypoxia, regulation of secondary metabolism, and biofilm formation (for a review see [11]).

Distribution of Histidine Kinases in the Fungal Kingdom

HKs are widespread in all fungal clades with the exception of Pneumocystidiomycetes (e.g., *Pneumocystis jirovecii*, the causal agent of pneumocystosis, Fig 2C) and microsporidia (e.g., *Enterocytozoon bieneusi*, a causal agent of microsporidiasis, Fig 2C). In light of new genomic data, the initial conviction that Ascomycota harbor the largest number of HKs was recently revised. Surprisingly, some previously unexplored early diverging fungal lineages, such as Mucoromycotina (e.g., the fungal pathogen *Lichtheimia corymbifera*, an agent of human mucormycosis, Fig 2C) and its closely clade Entomophthoromycotina (e.g., *Basidiobolus sp.* and *Conidiobolus sp.*, agents of human entomophthoromycosis), includes species with about 15–20 predicted HKs [11]. Nevertheless, it is accepted that Ascomycete yeasts (e.g., *Candida albicans*, Fig 2C) contain fewer HKs than filamentous Ascomycete species (Fig 2C), in which the number of HKs appears particularly variable. Some filamentous Ascomycete families have species that display a large series of HKs, such as Dothideomycetes (e.g., the plant pathogen *Zymoseptoria tritici*, Fig 2C) or Orbiliomycetes (e.g., the nematode trapping fungus *Arthrobotrys oligospora*, Fig 2C). However, it is important to note that even within a family, the total number of HKs can drastically differ between species. For example, in Sordariomycetes, *Fusarium verticillioides* (a pathogen of maize producing deadly mycotoxins) encodes 16 HKs [12], whereas only 9 HK-encoding genes are present in the genome of *Beauveria bassiana* (a prominent insect pathogen, Fig 2C) [11]. Even more strikingly, in Leotiomycetes, *Botrytis cinerea* (the cause of the gray mold on many plants) encodes 20 HKs [12], whereas *Pseudogymnoascus destructans* (the causal agent of the white-nose syndrome in bats, Fig 2C) contains only 6 predicted HKs.

A recent compilation/classification based on genomic analysis of 50 species indicates that Basidiomycota species contains few (2 to 7) HK members and there is no marked trend between yeast and filamentous species from this clade [20]. For instance, the Agaricomycete yeast *C. neoformans* encodes 7 HKs, whereas 5 genes are predicted to encode HKs in *Serpula lacrymans*, a devastating wood degrading filamentous fungus from the same clade (Fig 2C).

Finally, major qualitative points concerning HK proteins found in Ascomycota, Basidiomycota, and other early diverging fungal lineages also require particular mention. Some HK groups are widely distributed among fungal clades (Fig 2C). This is the case of group III HKs,

which are present in most of species that contain HKs with the exception of Taphrinomycotina (e.g., *Taphrina deformans*, the causal agent of peach leaf curl, Fig 2C), the Schizosaccharomycetaceae (e.g., the model fission yeast *Schizosaccharomyces pombe*), and the Saccharomycetaceae Whole Genome Duplication (WGD) clade (e.g. *Candida glabrata*). Interestingly, this observation has led some research teams to use group III HK-expressing strains of *Saccharomyces cerevisiae* as cell-based reporters to develop high throughput screening for new antifungals (see [9] for a review). On the other hand, some HK groups seem to have emerged or been maintained in a clade-specific manner. This has been hypothesized for the dual HK group that only occurs in Basidiomycota [21], or group VI osmosensors, which appear to be restricted to Ascomycota. Finally, it remains important to note that the broad series of HKs (up to 15) observed in some clades/species originates mainly from gene duplication events (likely leading to functional redundancy) and, to a lesser extent, from acquisition of new structures (HK groups). This is particularly obvious for the HKs found in Mucoromycotina (e.g., *L. corymbifera*) (Fig 2C) and filamentous Ascomycete plant pathogens [12].

In conclusion, it is striking to observe such a high degree of diversity among HKs in different fungi, both in the number of genes and their structure and functional variations, but, unfortunately, the underlying mechanisms that drive such diversity remain unclear. Closely related pathogenic and non-pathogenic fungal species do not appear to have significantly varying number of HKs (Fig 2C). However, it is highly likely that this gene family may have undergone rapid gene expansion under certain selection pressures allowing fungal species to adapt to their specific niches.

Future Prospects in Fungal Histidine Kinase Research

Many functional and evolutionary aspects concerning HKs in the fungal kingdom are still largely uncharted. However, we can assume that the recent expansion of genomic resources along with improved genetic approaches for studying pathogenic fungi (see [22] for a review) will contribute to broadened knowledge about fungal HKs. It is likely that the availability of new genome sequences will aid in identifying unknown HK structures/groups, which will help complete the current classification of this protein family. In the same way, comparative genomics should also expedite comprehensive phylogenetic analyses, enabling us to decipher the evolutionary path that led to the appearance, transfer, duplication, and loss of HK genes in fungi. As recently observed for few species [23,24], the development of novel efficient gene disruption strategies now promotes systematic multiple gene deletions in molds that bear a large number of HKs. In the near future, these powerful genetic approaches will likely help in deciphering (1) the role of each HK in a species, (2) putative genetic interactions, and (3) the basis of their apparent functional redundancy. In addition, although basic HK-mediated signal transduction routes are well characterized, downstream interacting partners and the global architecture of the signaling pathway involving fungal HKs remain critical missing pieces in this field. For instance, fluorescence microscopy technics can provide insight into the subcellular localization of fungal HKs [25]. Notably, with the exception of group VI HKs harboring transmembrane regions, most HKs are not predicted to be localized on the cell surface. Therefore, it will be intriguing to study how an environmental signal is translated into intracellular input signal for intracellular HKs and even a possibility that HKs may collaborate with other sensor proteins. Additionally, it will be highly relevant to determine which of the known HK groups directly communicates with the only currently known downstream candidate, the phosphotransfer shuttle protein Ypd1 (see Fig 1B), and to identify unknown downstream interacting partners. It is puzzling how different signals coming from different HKs are converged into Ypd1 and a few (1 to 2) response regulators, implying that some adaptor or scaffold proteins,

hitherto uncharacterized, may play such roles. Overall, we can assume that all the aforementioned strategies and outlooks will accelerate acquisition of new basic knowledge concerning fungal HK properties and general knowledge about signaling pathways involving these proteins. This is of primary importance for the future development of innovative HK-targeted antifungal strategies.

Acknowledgments

We acknowledge the Broad Institute Fungal Genome Initiative and the MycoCosm/JGI Fungal program for making available the complete genome sequence of many fungal species. We thank Pr. Gérald Larcher and Christina Cuomo for providing the compilation of HK sequences from the bat pathogen *Pseudogymnoascus destructans* and *Cryptococcus amyloletus*, respectively. We apologize to those whose work was not referenced due to citation restrictions.

References

- Mizuno T, Wurtzel ET, Inouye M. Osmoregulation of gene expression. II. DNA sequence of the *envZ* gene of the *ompB* operon of *Escherichia coli* and characterization of its gene product. *J Biol Chem*. 1982 Nov 25; 257(22):13692–8. PMID: [6292200](#).
- Chang C, Kwok SF, Bleecker AB, Meyerowitz EM. *Arabidopsis* ethylene-response gene ETR1: similarity of product to two-component regulators. *Science*. 1993 Oct 22; 262(5133):539–44. PMID: [8211181](#).
- Ota IM, Varshavsky A. A yeast protein similar to bacterial two-component regulators. *Science*. 1993 Oct 22; 262(5133):566–9. PMID: [8211183](#).
- Rudolph J, Oesterhelt D. Chemotaxis and phototaxis require a CheA histidine kinase in the archaeon *Halobacterium salinarium*. *EMBO J*. 1995 Feb 15; 14(4):667–73. PMID: [7882970](#); PubMed Central PMCID: PMC398130.
- Kehoe DM, Grossman AR. Similarity of a chromatic adaptation sensor to phytochrome and ethylene receptors. *Science*. 1996 Sep 6; 273(5280):1409–12. PMID: [8703080](#).
- Schuster SC, Noegel AA, Oehme F, Gerisch G, Simon MI. The hybrid histidine kinase DokA is part of the osmotic response system of *Dictyostelium*. *EMBO J*. 1996 Aug 1; 15(15):3880–9. PMID: [8670893](#)
- Mascher T, Helmann JD, Unden G. Stimulus perception in bacterial signal-transducing histidine kinases. *Microbiol Mol Biol Rev*. 2006 Dec; 70(4):910–38. Review. PMID: [17158704](#); PubMed Central PMCID: PMC1698512.
- Attwood PV. Histidine kinases from bacteria to humans. *Biochem Soc Trans*. 2013 Aug; 41(4):1023–8. doi: [10.1042/BST20130019](#) Review. PMID: [23863173](#).
- Shor E, Chauhan N. A case for two-component signaling systems as antifungal drug targets. *PLoS Pathog*. 2015 Feb; 11(2):e1004632. doi: [10.1371/journal.ppat.1004632](#) eCollection 2015 Feb. PMID: [25723524](#); PubMed Central PMCID: PMC4344368.
- Grebe TW, Stock JB. The histidine protein kinase superfamily. *Adv Microb Physiol*. 1999; 41:139–227. Review. PMID: [10500846](#).
- Defosse TA, Sharma A, Mondal AK, Dugé de Bernonville T, Latgé J-P, Calderone R, et al. Hybrid histidine kinases in pathogenic fungi. *Mol Microbiol*. 2015 Mar; 95(6):914–24. doi: [10.1111/mmi.12911](#) Epub 2015 Jan 30. Review. PMID: [25560420](#).
- Catlett NL, Yoder OC, Turgeon BG. Whole-genome analysis of two-component signal transduction genes in fungal pathogens. *Eukaryot Cell*. 2003 Dec; 2(6):1151–61. PMID: [14665450](#); PubMed Central PMCID: PMC326637.
- Maeda T, Takekawa M, Saito H. Activation of yeast PBS2 MAPKK by MAPKKs or by binding of an SH3-containing osmosensor. *Science*. 1995 Jul 28; 269(5223):554–8. PMID: [7624781](#).
- Posas F, Wurgler-Murphy SM, Maeda T, Witten EA, Thai TC, Saito H. Yeast HOG1 MAP kinase cascade is regulated by a multistep phosphorelay mechanism in the SLN1-YPD1-SSK1 "two-component" osmosensor. *Cell*. 1996 Sep 20; 86(6):865–75. PMID: [8808622](#).
- Chauhan N, Latgé J-P, Calderone R. Signalling and oxidant adaptation in *Candida albicans* and *Aspergillus fumigatus*. *Nat Rev Microbiol*. 2006 Jun; 4(6):435–44. Review. PMID: [16710324](#).
- Yeh KC, Lagarias JC. Eukaryotic phytochromes: light-regulated serine/threonine protein kinases with histidine kinase ancestry. *Proc Natl Acad Sci U S A*. 1998 Nov 10; 95(23):13976–81. PMID: [9811911](#); PubMed Central PMCID: PMC24997.

17. Bayram O, Braus GH, Fischer R, Rodriguez-Romero J. Spotlight on *Aspergillus nidulans* photosensory systems. *Fungal Genet Biol*. 2010 Nov; 47(11):900–8. doi: [10.1016/j.fgb.2010.05.008](https://doi.org/10.1016/j.fgb.2010.05.008) Epub 2010 May 23. Review. PMID: [20573560](https://pubmed.ncbi.nlm.nih.gov/20573560/).
18. Yu Z, Armant O, Fischer R. Fungi use the SakA (HogA) pathway for phytochrome-dependent light signaling. *Nat Microbiol*. doi: [10.1038/nmicrobiol.2016.19](https://doi.org/10.1038/nmicrobiol.2016.19)
19. Boyce KJ, Andrianopoulos A. Fungal dimorphism: the switch from hyphae to yeast is a specialized morphogenetic adaptation allowing colonization of a host. *FEMS Microbiol Rev*. 2015 Nov; 39(6):797–811. doi: [10.1093/femsre/fuv035](https://doi.org/10.1093/femsre/fuv035) Epub 2015 Aug PMID: [26253139](https://pubmed.ncbi.nlm.nih.gov/26253139/)
20. Lavín JL, García-Yoldi A, Ramírez L, Pisabarro AG, Oguiza JA. Two-component signal transduction in *Agaricus bisporus*: a comparative genomic analysis with other basidiomycetes through the web-based tool BASID2CS. *Fungal Genet Biol*. 2013 Jun; 55:77–84. doi: [10.1016/j.fgb.2012.09.012](https://doi.org/10.1016/j.fgb.2012.09.012) Epub 2012 Oct 31. PMID: [23123423](https://pubmed.ncbi.nlm.nih.gov/23123423/).6. Review. 26253139.
21. Lavín JL, Sarasola-Puente V, Ramírez L, Pisabarro AG, Oguiza JA. Dual-histidine kinases in basidiomycete fungi. *C R Biol*. 2014 Feb; 337(2):111–6. doi: [10.1016/j.crv.2013.12.007](https://doi.org/10.1016/j.crv.2013.12.007) Epub 2014 Feb 10. PMID: [24581805](https://pubmed.ncbi.nlm.nih.gov/24581805/).
22. Cairns TC, Studholme DJ, Talbot NJ, Haynes K. New and improved techniques for the study of pathogenic fungi. *Trends Microbiol*. 2016 Jan; 24(1):35–50. doi: [10.1016/j.tim.2015.09.008](https://doi.org/10.1016/j.tim.2015.09.008) Epub 2015 Nov 5. Review. PMID: [26549580](https://pubmed.ncbi.nlm.nih.gov/26549580/).
23. Jacob S, Foster AJ, Yemelin A, Thines E. Histidine kinases mediate differentiation, stress response, and pathogenicity in *Magnaporthe oryzae*. *Microbiologyopen*. 2014 Oct; 3(5):668–87. doi: [10.1002/mbo3.197](https://doi.org/10.1002/mbo3.197) Epub 2014 Aug 8. PMID: [25103193](https://pubmed.ncbi.nlm.nih.gov/25103193/); PubMed Central PMCID: PMC4234259.
24. Chapeland-Leclerc F, Dilmaghani A, Ez-Zaki L, Boisnard S, Da Silva B, Gaslonde T, et al. Systematic gene deletion and functional characterization of histidine kinase phosphorelay receptors (HKRs) in the human pathogenic fungus *Aspergillus fumigatus*. *Fungal Genet Biol*. 2015 Nov; 84:1–11. doi: [10.1016/j.fgb.2015.09.005](https://doi.org/10.1016/j.fgb.2015.09.005) Epub 2015 Sep 10. PMID: [26365385](https://pubmed.ncbi.nlm.nih.gov/26365385/).
25. Foureau E, Clastre M, Montoya EJ, Besseau S, Oudin A, Glévarec G, et al. Subcellular localization of the histidine kinase receptors Sln1p, Nik1p and Chk1p in the yeast CTG clade species *Candida guilliermondii*. *Fungal Genet Biol*. 2014 Apr; 65:25–36. doi: [10.1016/j.fgb.2014.01.007](https://doi.org/10.1016/j.fgb.2014.01.007) Epub 2014 Feb 8. PMID: [24518307](https://pubmed.ncbi.nlm.nih.gov/24518307/).
26. Calera JA, Zhao XJ, De Bernardis F, Sheridan M, Calderone R. Avirulence of *Candida albicans* CaHK1 mutants in a murine model of hematogenously disseminated candidiasis. *Infect Immun*. 1999 Aug; 67(8):4280–4. PMID: [10417206](https://pubmed.ncbi.nlm.nih.gov/10417206/); PubMed Central PMCID: PMC96739.
27. Clemons KV, Miller TK, Selitrennikoff CP, Stevens DA. Fos-1, a putative histidine kinase as a virulence factor for systemic aspergillosis. *Med Mycol*. 2002 Jun; 40(3):259–62. PMID: [12146755](https://pubmed.ncbi.nlm.nih.gov/12146755/).
28. Nemecek JC, Wüthrich M, Klein BS. Global control of dimorphism and virulence in fungi. *Science*. 2006 Apr 28; 312(5773):583–8. PMID: [16645097](https://pubmed.ncbi.nlm.nih.gov/16645097/).
29. Viaud M, Fillinger S, Liu W, Polepalli JS, Le Pêcheur P, Kunduru AR, et al. A class III histidine kinase acts as a novel virulence factor in *Botrytis cinerea*. *Mol Plant Microbe Interact*. 2006 Sep; 19(9):1042–50. PMID: [16941908](https://pubmed.ncbi.nlm.nih.gov/16941908/).
30. Bahn Y-S, Kojima K, Cox GM, Heitman J. A unique fungal two-component system regulates stress responses, drug sensitivity, sexual development, and virulence of *Cryptococcus neoformans*. *Mol Biol Cell*. 2006 Jul; 17(7):3122–35. Epub 2006 May 3. PMID: [16672377](https://pubmed.ncbi.nlm.nih.gov/16672377/); PubMed Central PMCID: PMC1483045.
31. Zhou G, Wang J, Qiu L, Feng M-G. A Group III histidine kinase (mhk1) upstream of high-osmolarity glycerol pathway regulates sporulation, multi-stress tolerance and virulence of *Metarhizium robertsii*, a fungal entomopathogen. *Environ Microbiol*. 2012 Mar; 14(3):817–29. doi: [10.1111/j.1462-2920.2011.02643.x](https://doi.org/10.1111/j.1462-2920.2011.02643.x) Epub 2011 Nov 28. PMID: [22118192](https://pubmed.ncbi.nlm.nih.gov/22118192/).