

Published in final edited form as:

Eur J Endocrinol. 2015 August ; 173(2): R69–R83. doi:10.1530/EJE-14-0826.

Influence of maternal vitamin D status on obstetric outcomes and the foetal skeleton

Rebecca J Moon^{1,2}, Nicholas C Harvey^{1,3}, and Cyrus Cooper^{1,3,4}

¹MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton General Hospital, Tremona Road, Southampton, SO16 6YD, UK

²Paediatric Endocrinology, University Hospitals Southampton NHS Foundation Trust, Tremona Road, Southampton, SO16 6YD, UK

³NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Tremona Road, Southampton, SO16 6YD, UK

⁴NIHR Musculoskeletal Biomedical Research Unit, University of Oxford, Oxford, OX3 7LD UK

Abstract

Vitamin D status is increasingly associated with wide ranging clinical outcomes. There is now a wealth of observational studies reporting on its associations with obstetric complications, including preeclampsia, gestational diabetes and mode and timing of delivery. The findings are inconsistent and currently there is a lack of data from high quality intervention studies to confirm a causal role for vitamin D in these outcomes. This is similarly true with regards to fetal development, including measures of fetal size and skeletal mineralisation. The available data justify the instatement of high-quality randomised placebo controlled trials in a range of populations and health care settings to establish potential efficacy and safety of vitamin D supplementation to improve particular outcomes.

Keywords

Vitamin D; pregnancy; pre-eclampsia; gestational diabetes; osteoporosis; epidemiology

Introduction

The classical role of Vitamin D is in calcium and phosphate homeostasis: it is without doubt that severe vitamin D deficiency (VDD) can result in rickets, osteomalacia and hypocalcaemia. However, there is increasing suggestion that VDD is associated with wide ranging clinical outcomes, including pregnancy complications and adverse fetal development. As a result, a number of national guidelines recommend vitamin D

Corresponding Author and person to whom reprint requests should be addressed: Professor Cyrus Cooper, MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton General Hospital, Southampton. SO16 6YD, Tel: 023 8077 7624, Fax: 023 8070 4021, cc@mrc.soton.ac.uk.

Disclosures: NH has received consultancy, lecture fees and honoraria from Alliance for Better Bone Health, AMGEN, MSD, Eli Lilly, Servier, Shire, Consilient Healthcare and Internis Pharma; CC has received consultancy, lecture fees and honoraria from AMGEN, GSK, Alliance for Better Bone Health, MSD, Eli Lilly, Pfizer, Novartis, Servier, Medtronic and Roche.

supplementation during pregnancy^{1–3}, although this is not currently supported by the World Health Organisation (WHO)⁴. Here, we review the evidence basis for antenatal vitamin D supplementation to prevent obstetric complications, and the influence of vitamin D on fetal growth and skeletal development.

Vitamin D physiology and epidemiology in pregnancy

Vitamin D can be derived from the diet, as ergocalciferol (vitamin D₂) from plant sources, or cholecalciferol (vitamin D₃) from animal sources. However, the majority is formed endogenously within the skin from the action of ultraviolet B (290–315nm wavelength) to convert 7-dehydrocholesterol to pre-vitamin D₃. Hydroxylation within the liver produces 25-hydroxyvitamin D [25(OH)D]. This is the main circulating form of vitamin D, found either bound to vitamin D binding protein (VDBP), albumin or in the free form. 25(OH)D acts as a reservoir for conversion to 1,25-dihydroxyvitamin D [1,25(OH)₂D], primarily in the renal proximal tubular cells, but also within bone, the parathyroid gland and placenta. Whilst 1,25(OH)₂D is the active metabolite, its production is regulated in response to serum calcium and its half life is short at 4–6 hours. Conversely, hepatic 25-hydroxylation is not physiologically regulated and 25(OH)D has a half-life of approximately 2–3 weeks⁵. Therefore, serum 25(OH)D is currently considered the best marker of vitamin D status⁶.

The primary function of 1,25(OH)₂D is in calcium and phosphate homeostasis, which occurs in conjunction with parathyroid hormone (PTH). Thus, low serum ionised Ca²⁺ stimulates PTH release, which simultaneously increases renal calcium reabsorption in the distal tubule of the kidney, decreases proximal tubule phosphate reabsorption, and increases 1,25(OH)₂D synthesis. The main action of 1,25(OH)₂D is to increase uptake of dietary calcium through the intestinal enterocytes, but it also enables PTH induced mobilisation of calcium and phosphate from bone mineral⁷.

During pregnancy alterations to calcium and phosphate metabolism occur to allow the accretion of calcium within the fetal skeleton, particularly during the last trimester⁸. This occurs through increased maternal intestinal calcium absorption^{9, 10} and mobilization of calcium within the maternal skeletal¹¹, but without alteration to maternal serum ionized calcium concentration. Maternal calcitropic hormones, including 1,25(OH)₂D, likely have an important role in these adaptations, as total 1,25(OH)₂D increases during the second and third trimesters^{9, 12}, although this could also reflect the increase in VDBP from early through to late pregnancy^{10, 13}. The effect of pregnancy on 25(OH)D however is less well understood: Zhang et al. observed a reduction in 25(OH)D in late compared with early pregnancy, however as all subjects were recruited in summer months this might reflect seasonal variation¹³. In contrast, Ritchie et al. reported no significant differences in 25(OH)D measured in 14 women before pregnancy, in each trimester and during lactation¹⁰. Nonetheless, biochemically low levels of 25(OH)D are highly prevalent: In a cohort of predominantly Caucasian women in the United Kingdom (UK), 31% had a serum 25(OH)D less than 50nmol/l, which is widely considered to be insufficient, and 18% less than 25nmol/l, often considered deficient¹⁴. However in an ethnically more diverse UK population, 36% of women had a 25(OH)D <25nmol/l at pregnancy booking¹⁵. Indeed, dark skin pigmentation and extensive skin covering (eg for religious or cultural reasons) are the

strongest risk factors for vitamin D deficiency. Obesity is also associated with biochemically low 25(OH)D levels, whereas in pregnancy, use of vitamin D supplements may prevent deficiency¹⁴. Maternal 25(OH)D in pregnancy is an important consideration as the fetus is entirely dependent on the mother for 25(OH)D. 25(OH)D readily crosses the placenta, and maternal and umbilical cord venous blood 25(OH)D are moderately well correlated, with randomised controlled trials demonstrating that vitamin D supplementation in pregnancy can increase umbilical cord venous and neonatal serum 25(OH)D compared to placebo^{16–23}.

Obstetric Complications

Observational studies

There are numerous observational studies reporting associations between either vitamin D intake in pregnancy or serum measurement of 25(OH)D and pregnancy complications, including gestational hypertension (GHT) and preeclampsia (PET), gestational diabetes (GDM), timing and mode of delivery. The interpretation and comparison of these studies is limited by the timing of 25(OH)D measurements, ranging from first trimester to delivery, definition used for both VDD and the outcome, covariates adjusted for and study design (eg prospective cohort, case-control).

Gestational hypertension & preeclampsia

Although the aetiology of PET is poorly understood and likely multifactorial, there is some evidence that maternal calcium status might be important, and calcium supplementation can reduce PET risk particularly in women with low calcium intake²⁴. Thus, exploring a role for calcitropic hormones, including vitamin D, is a sensible approach. Several case-control and prospective cohort studies have demonstrated that women who developed PET had lower serum 25(OH)D compared to controls in early^{25–27}, mid^{28, 29} or late pregnancy^{25, 30, 31}, and that VDD increases the risk of PET^{25, 30, 32}. One case-control study suggested women with serum 25(OH)D < 37.5 nmol/l measured at less than 22 weeks gestation have a 5-fold higher risk of PET than women with a 25(OH)D > 37.5 nmol/l, independent of ethnicity, season, gestational age at sampling, pre-pregnancy body mass index (BMI), and educational achievement²⁵. Similarly, in a cohort of 23,425 pregnant women in Norway, lower vitamin D intake estimated from a food frequency questionnaire at 22 weeks gestation was associated with a significantly increased risk of PET³³. The lower vitamin D intake in women who developed PET was mostly due to a difference in vitamin D obtained from supplements, suggesting supplementation might prevent PET. However, these findings are not supported by all studies^{27, 34–41}, and indeed in a prospective cohort of 1591 women, for each additional 25 nmol/l increment in 25(OH)D in early pregnancy, the risk of GHT (without PET) increased by 30%, but no effect on PET risk was observed³⁸, highlighting possible detrimental effects of higher vitamin D status.

In recent years, there have been several published meta-analyses of the relationship between maternal vitamin D status and PET risk, as shown in table 142–48. Similarly to the observational studies, the conclusions of these are inconsistent. In our own meta-analysis, we found no significant reduction in the risk of PET with higher vitamin D status (Figure 1)⁴⁷. In contrast, Aghajafari et al. found that the increased risk of PET in VDD was only

observed in studies in which blood sampling was later than 16 weeks gestation and when VDD was defined as $25(\text{OH})\text{D} < 75 \text{ nmol/l}$ and not $< 50 \text{ nmol/l}$. However, Tabesh et al., including a larger number of studies defining VDD as less than 50 nmol/l , did demonstrate an increased risk of PET, which was not found when deficiency was defined as less than 38 nmol/l . Importantly, the total number of women included in these meta-analyses varied from 610-2485 (excluding those based on intake only and the most recent meta-analyses which included novel data⁴²). However, between January 2013 and July 2014 at least a further 14 case-control or prospective cohort studies with measurement of serum $25(\text{OH})\text{D}$ and assessing PET risk have been published^{27, 31, 32, 39–42, 48–55}. These newer studies include data for a further 21,000 women, considerably more than were included in the meta-analyses.

Gestational Diabetes

Similarly to PET, conflicting findings have been reported for $25(\text{OH})\text{D}$ status in case-control and prospective cohort studies of GDM risk: both lower^{48, 56–61} and similar serum $25(\text{OH})\text{D}$ ^{62, 63} during pregnancy in women with and without GDM have been reported. One study of women referred for GDM screening did not find a difference in the prevalence of GDM in women with $25(\text{OH})\text{D}$ above and below 50 nmol/l , but the women with $25(\text{OH})\text{D} < 50 \text{ nmol/l}$ did have higher fasting blood glucose, $\text{HBA}_{1\text{C}}$ and insulin resistance. However these women also had higher BMI, lower physical activity and were less likely to be Caucasian, which might have confounded the findings⁶⁴. Three separate meta-analyses of published studies all concluded that women with GDM had significantly lower mean $25(\text{OH})\text{D}$ than normoglycaemic women^{44, 46, 65} with the mean difference in $25(\text{OH})\text{D}$ ranging from 3.9 to 7.4 nmol/l . Furthermore, these meta-analyses suggested that the risk of GDM was increased by 40-60% in women with VDD^{44, 46, 65}, as shown in Figure 144. However, similarly to studies assessing PET risk, there is now substantially more data available than was used for these meta-analyses^{39, 48, 58–61, 63, 66} and whilst many of the smaller studies would support the previous conclusions, a large prospective cohort of women in Australia, including 5109 women, of whom 7.4% developed GDM, first trimester VDD (defined either as $< 25 \text{ nmol/l}$ or $< 37.5 \text{ nmol/l}$) was not associated with increased risk of GDM compared to $25(\text{OH})\text{D}$ $50\text{--}75 \text{ nmol/l}$ after adjustment for age, parity, smoking during pregnancy, maternal weight, previously diagnosed hypertension, diabetes, season at sampling, country of birth, or socioeconomic disadvantage⁴⁸. Furthermore in 1953 women in Southern China vitamin D sufficiency ($25(\text{OH})\text{D} > 75 \text{ nmol/l}$) at 16-20 weeks gestation was associated with a small, but statistically significant, increased risk of GDM (OR 1.02, 95% CI 1.00, 1.04)³⁹.

Caesarean Delivery

Unsurprisingly, in recent years, there has also been an increase in studies reporting maternal vitamin D status in relation to mode and timing of delivery. Again, these are inconsistent. After adjustment for potential confounding factors three studies which assessed $25(\text{OH})\text{D}$ in early pregnancy, when attending for GDM screening, and at delivery, reported an increased risk of Caesarean delivery^{64, 67, 68}. Conversely, two studies, which measured $25(\text{OH})\text{D}$ in the first trimester demonstrated no increased risk^{37, 39}. Assessment of the influence of VDD on mode of delivery is further complicated by the underlying cause for intervention;

however Savvidou et al. additionally categorised women requiring emergency caesarean delivery due to failure to progress and for fetal distress. Neither group had significantly different serum 25(OH)D in early pregnancy compared to women who delivered vaginally⁶⁹.

Preterm Delivery

More studies have concluded that maternal 25(OH)D status is not related to preterm birth^{34, 37, 48, 70–74}, than have shown VDD increases this risk^{64, 75, 76}. Furthermore, Zhou et al reported women with higher vitamin D status at 16–20 weeks gestation had a higher odds of preterm delivery³⁹, and similarly Hossain et al. found that cord blood 25(OH)D was higher in preterm (<37 weeks gestation) deliveries (mean 55nmol/l) compared to term pregnancies (mean 40nmol/l, $p=0.009$) in women in Pakistan⁷⁷. Interestingly, two of the studies which suggest VDD increased the risk of preterm delivery used a definition of less than 35 weeks gestation for preterm^{75, 76}, whereas all, but one⁷⁴, of the studies reporting either no relationship or VDD reduced the risk considered preterm delivery to be at less than 37 weeks gestation. Whilst this might suggest that VDD is particularly associated with an increased risk of very preterm birth, Schneuer et al, who prospectively studied first trimester 25(OH)D status in over 5000 women, found VDD did not increase the risk of either, all, or spontaneous, preterm birth <34 weeks gestation, before or after adjustment for potential confounding factors⁴⁸. However, differences in timing of 25(OH)D assessment, and one study showing increased risk including only twin pregnancies⁷⁵, could account for these different findings. Furthermore, Bodnar et al. observed that only non-white mothers had an increased risk of preterm birth with low 25(OH)D at 26 weeks gestation⁷⁶, suggesting stratification of women by ethnicity in future intervention studies might be necessary.

Intervention studies of vitamin D supplementation to reduce obstetric complications

Observational data cannot confirm a causal effect of vitamin D or justification for population wide supplementation, particularly as some studies have suggested possible detrimental effects of higher 25(OH)D^{38, 39, 77}. As 25(OH)D status is primarily determined by environmental factors, confounding and reverse causality need to be considered, and differences in covariates included in multivariate models might explain the inconsistent findings. For example, obese individuals have lower 25(OH)D status, and a higher incidence of GDM, GHT, PET, caesarean section and preterm delivery^{78, 79}. Similarly African-American women are more likely to require delivery by Caesarean section and to experience pre-eclampsia and preterm labour⁸⁰. Whether these outcomes can truly be attributed to lower 25(OH)D compared to Caucasian women and therefore prevented by vitamin D supplementation must be established through intervention studies.

Despite the expanse of observational data, there are currently few trials of antenatal vitamin D supplementation reporting on maternal outcomes other than maternal/neonatal vitamin D and calcium status⁸¹. In three of the five studies, the interventional product contained only vitamin D^{21, 82, 83}, whereas a further two assessed the effects of combined vitamin D and calcium supplementation^{84, 85} (Table 2). The interpretation of these two studies with regards to GHT and PET is limited as calcium supplementation is known to reduce the risk of PET²⁴. Nonetheless, high dose vitamin D supplementation, with or without calcium

supplementation, did not improve the incidence of GHT, PET, GDM, or preterm delivery compared to either usual care or low dose supplementation^{21, 82–85}. However these studies were most likely underpowered to detect a difference in these outcomes. GDM complicates approximately 4.5% of pregnancies in the UK⁸⁶. Thus, to detect a 50% reduction in this incidence with 80% power at the 5% significance level, 1010 women would be needed in each study arm. As PET occurs in 2–3% of pregnancies, even larger study numbers are needed.

Although trials of vitamin D supplementation have not yet demonstrated a reduction in the incidence of PET or GDM, there is some evidence to support effects on blood pressure and glucose metabolism when considered as continuous outcomes. For example, Marya et al. demonstrated a reduction in both systolic and diastolic BP in women randomised to vitamin D and calcium supplementation compared to those who received usual care⁸⁵. Confirmation of this finding using vitamin D alone is now needed. Three studies have assessed the effects of vitamin D supplementation on insulin resistance. In an unblinded study of 113 Iranian women randomised to one of three treatment groups (200 IU/day, 50,000 IU/month, 50,000 IU/fortnight) from 12 weeks gestation until delivery, insulin resistance, assessed by HOMA-IR, increased significantly from baseline to delivery in all three groups, but the rise was significantly less in women randomised to 50,000 IU/fortnight than in women who received 200 IU/day⁸⁷. In contrast, Yap et al found no difference in either fasting blood glucose or that measured two hours post glucose load in women randomised to either 400 IU/day or 5000 IU/day cholecalciferol, with similar results for HOMA-IR⁸³. Finally, in a small study of 54 women with a diagnosis of GDM, two doses of 50,000IU cholecalciferol 3 weeks apart did improve fasting blood glucose and insulin resistance compared to placebo. However the women randomised to vitamin D supplementation had significantly higher insulin resistance at baseline making these results difficult to interpret⁸⁸. Nonetheless, these findings support the need for further high quality large randomised controlled trials, and to concurrently determine if any effects on maternal physiology might also have beneficial effects on maternal and/or fetal morbidity, for example macrosomia or neonatal hypoglycaemia.

Fetal Development

Early rickets and symptomatic neonatal hypocalcaemia have been reported in infants born to mothers with VDD^{89–91}. However, these outcomes are rarely reported in infants of white mothers, and most commonly occur in those born to mothers with dark skin pigmentation, extensive skin covering and profound VDD. The fetus is dependent on the mother for accretion of approximately 30g of calcium to enable skeletal development. As such, a subclinical role for vitamin D and/or calcium in fetal growth and bone development has been considered, yet maternal supplementation with calcium alone does not appear to have beneficial effects on fetal bone mineral accrual⁸¹.

Size at birth

There are now a number of intervention studies assessing the effect of vitamin D supplementation on birth anthropometry, although the dose and timing of introduction of

vitamin D varied widely (Table 3). Most studies trialled supplementation with vitamin D alone and did not find a significant effect on birth weight, length or head circumference (Table 1) However, interestingly, vitamin D in combination with calcium did increase birth weight in three studies despite women in the control group also receiving calcium supplementation in two of these studies^{84, 92, 93}. Indeed the prevalence of VDD at baseline and mean 25(OH)D achieved was similar in a study of women in Bangladesh, who received 35,000 IU/day cholecalciferol from 26-30 weeks gestation¹⁹, to women participating in a study of 50,000 IU cholecalciferol per week in addition to 200mg elemental calcium supplementation in Iran⁹³. Both studies included a similar number of women. However, in the former study birth weight was similar in both intervention and control groups, whereas in the latter study mean birth weight in the intervention group was 170g greater than that in the control group. These differing findings might suggest that the effect of vitamin D is dependent on the availability of calcium, or could result from genetic/racial variation in response to vitamin D supplementation, but nonetheless highlight the importance of using data obtained from an appropriate population in the development of antenatal supplementation policies.

Skeletal Development

Currently, the data relating maternal 25(OH)D status to offspring bone development is largely observational in nature, but does span antenatal measurements to peak bone mass. Indeed, using gestational ultrasound, smaller femoral volumes⁹⁴ and widening of the distal femoral metaphysis relative to femur length has been demonstrated in fetuses of mothers with low levels of serum 25(OH)D⁹⁵.

A number of studies have demonstrated associations between maternal 25(OH)D status in pregnancy and offspring bone mineralisation in the neonatal period. In 71 Korean neonates, those born in summer (July-September) had 6% higher whole body bone mineral content (BMC) than infants born in winter (January-March), and neonatal 25(OH)D at delivery was correlated with whole body BMC in all children ($r=0.24$, $p=0.05$)⁹⁶. However, in three similar studies by the same author in North America a reversed pattern was observed with whole body BMC 8-12% lower in infants born in summer⁹⁷. The authors suggest that this difference reflects low uptake of vitamin D supplementation throughout pregnancy in Korea, but only during the first trimester in North America, thereby suggesting early pregnancy during winter might impact on skeletal development⁹⁷. However, Weiler et al. studied 50 Canadian infants born between August and April, with the majority of mothers taking vitamin D supplementation in pregnancy. Infants with a cord blood 25(OH)D < 37.5 nmol/l ($n=18$) were heavier and longer than those with a cord blood 25(OH)D above this cut-point, but skeletal size was not relatively increased, such that whole body and femur BMC relative to body weight were significantly lower⁹⁸. In a Finnish study, peripheral quantitative computed tomography (pQCT) was used to assess both BMC and bone geometry of the tibia in 98 neonates. In this analysis, the mean of two maternal 25(OH)D measurements in early pregnancy and 2 days postpartum was used to define maternal vitamin D status, and the median for the cohort used to establish two groups. BMC and bone cross-sectional area (CSA) were 13.9% and 16.3% higher, respectively, in infants of mothers with higher 25(OH)D⁹⁹. When these children were reassessed at 14 months of age, the difference in

tibial BMC was no longer present, but the greater CSA persisted¹⁰⁰. Conversely, in 125 Gambian mother-offspring pairs, no significant relationships were observed between maternal 25(OH)D at either 20 or 36 weeks gestation and offspring whole body BMC or bone area at 2, 13 or 52 weeks of age¹⁰¹. However, in contrast to the other studies, no mother had a 25(OH)D less than 50nmol/l, consistent with the notion that poorer skeletal mineralisation might only occur in fetuses of mothers with the lowest vitamin D levels.

There is evidence to support the persistence of these relationships outside of the neonatal period, although the data are less consistent. In the first study to report on the relationship between maternal 25(OH)D status and offspring bone mineralisation in childhood, Javaid et al demonstrated positive associations between late pregnancy 25(OH)D and offspring whole body and lumbar spine BMC, bone area and areal bone mineral density (aBMD) measured at 9 years (Figure 2)¹⁴. Positive relationships with umbilical venous calcium concentration were also observed, suggesting that the effect of vitamin D on skeletal development might be mediated through placental calcium transport¹⁴. This was initially supported by data from the Avon Longitudinal Study of Parents and Children (ALSPAC), in which maternal estimated ultraviolet B exposure in late pregnancy, used as a proxy measure of vitamin D status, was positively associated with offspring whole body less head (WBLH) BMC and bone area at 9-10 years of age in 6955 children¹⁰². However, subsequent re-analysis in a more limited subset of the ALSPAC cohort using serum 25(OH)D measured in pregnancy demonstrated no association with WBLH BMC or bone area¹⁰³. Interestingly there was strong collinearity between maternal gestational UVB exposure and offspring age at bone assessment, which limits the interpretation of these studies¹⁰⁴. Finally, data from the Raine cohort in Western Australia provide support for a positive relationship between maternal gestational vitamin D status and offspring bone development to peak bone mass: In this study, whole body BMC and aBMD were 2.7% and 1.7% lower, respectively, at 20 years of age in offspring of mothers with 25(OH)D<50nmol/l (compared with offspring of mothers >50nmol/l) at 18 weeks gestation after adjustment for sex, age, height and body composition at 20 years, maternal height and prepregnancy weight, age at delivery, parity, education, ethnicity, smoking during pregnancy, and season of maternal blood sampling¹⁰⁵.

Currently there is only one intervention study of the effect of vitamin D supplementation in pregnancy on offspring bone mineralisation. Congdon et al. assessed forearm BMC using single photon absorptiometry in 64 infants of Asian mothers living in the UK who participated in a non-randomised study of vitamin D and calcium supplementation in pregnancy¹⁰⁶. 19 women received 1000IU vitamin D and a calcium supplement (of unknown strength) during the last trimester, and were compared to 45 women who did not receive any supplement. No significant differences were identified between these two groups, but interpretation of the study findings is limited by the small study size, lack of randomisation and technique used to assess BMC. The ongoing Maternal Vitamin D Osteoporosis Study (MAVIDOS) in which over 1000 women were randomised to 1000 IU cholecalciferol or placebo daily from 14 weeks gestation till delivery, with assessment of offspring bone mineralisation at birth and 4 years of age by dual energy X-ray absorptiometry (DXA)¹⁰⁷, will provide much needed high quality evidence on the role of vitamin D supplementation in pregnancy in fetal skeletal development¹⁰⁸.

Conclusions

There is now a wealth of observational data relating vitamin D status in pregnancy to obstetric complications, fetal growth and offspring bone development. The findings of these studies are inconsistent and whilst justifying the need for assessment of vitamin D supplementation in high quality randomised controlled trials, observational data alone should not be used as a basis for population wide vitamin D supplementation in pregnancy. Indeed it is possible that the variability in findings of both observational and the few intervention studies reflects the wide heterogeneity in the populations studied (including prevalence of VDD and calcium status, ethnic diversity), dose of vitamin D and timing of initiation or assessment of 25(OH)D status and definition used for the outcomes considered. Thus any public health recommendations need to be based on an appropriate population. Furthermore, whilst currently available data does not suggest any short term detrimental effects for the mother or fetus, the long term safety of vitamin D supplementation, particularly at supra-physiological doses remains to be established.

References

1. National Institute for Health and Clinical Excellence. Antenatal care (NICE Clinical Guideline 62). 2010. www.guidance.nice.org.uk/cg62
2. Paxton GA, Teale GR, Nowson CA, Mason RS, McGrath JJ, Thompson MJ, Siafarikas A, Rodda CP, Munns CF. Vitamin D and health in pregnancy, infants, children and adolescents in Australia and New Zealand: a position statement. *Med J Aust.* 2013; 198:142–143. [PubMed: 23418693]
3. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2011; 96:1911–1930. [PubMed: 21646368]
4. World Health Organisation. Guideline: Vitamin D supplementation in pregnant women. Geneva: 2012.
5. Jones KS, Assar S, Harnpanich D, Bouillon R, Lambrechts D, Prentice A, Schoenmakers I. 25(OH)D₂ Half-Life Is Shorter Than 25(OH)D₃ Half-Life and Is Influenced by DBP Concentration and Genotype. *J Clin Endocrinol Metab.* 2014; 99:3373–3381. [PubMed: 24885631]
6. Hollis BW, Wagner CL. Clinical review: The role of the parent compound vitamin D with respect to metabolism and function: Why clinical dose intervals can affect clinical outcomes. *J Clin Endocrinol Metab.* 2013; 98:4619–4628. [PubMed: 24106283]
7. Bikle, DS.; Adams, J.; Christakos, S. Vitamin D: Production, Metabolism, Mechanism of Action, and Clinical Requirements. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism.* John Wiley & Sons, Inc; 2013. p. 235-248.
8. Kovacs CS. Calcium and Bone Metabolism in Pregnancy and Lactation*. *The Journal of Clinical Endocrinology & Metabolism.* 2001; 86:2344–2348. [PubMed: 11397820]
9. Cross NA, Hillman LS, Allen SH, Krause GF, Vieira NE. Calcium homeostasis and bone metabolism during pregnancy, lactation, and postweaning: a longitudinal study. *Am J Clin Nutr.* 1995; 61:514–523. [PubMed: 7872214]
10. Ritchie LD, Fung EB, Halloran BP, Turnlund JR, Van Loan MD, Cann CE, King JC. A longitudinal study of calcium homeostasis during human pregnancy and lactation and after resumption of menses. *Am J Clin Nutr.* 1998; 67:693–701. [PubMed: 9537616]
11. More C, Bhattoa HP, Bettembuk P, Balogh A. The effects of pregnancy and lactation on hormonal status and biochemical markers of bone turnover. *Eur J Obstet Gynecol Reprod Biol.* 2003; 106:209–213. [PubMed: 12551794]
12. Ardawi MS, Nasrat HA, HS BAA. Calcium-regulating hormones and parathyroid hormone-related peptide in normal human pregnancy and postpartum: a longitudinal study. *Eur J Endocrinol.* 1997; 137:402–409. [PubMed: 9368509]

13. Zhang JY, Lucey AJ, Horgan R, Kenny LC, Kiely M. Impact of pregnancy on vitamin D status: a longitudinal study. *Br J Nutr.* 2014;1–7.
14. Javaid MK, Crozier SR, Harvey NC, Gale CR, Dennison EM, Boucher BJ, Arden NK, Godfrey KM, Cooper C. Maternal vitamin D status during pregnancy and childhood bone mass at age 9 years: a longitudinal study. *The Lancet.* 2006; 367:36–43.
15. McAree T, Jacobs B, Manickavasagar T, Sivalokanathan S, Brennan L, Bassett P, Rainbow S, Blair M. Vitamin D deficiency in pregnancy - still a public health issue. *Matern Child Nutr.* 2013; 9:23–30. [PubMed: 23230904]
16. Brooke OG, Brown IR, Bone CD, Carter ND, Cleeve HJ, Maxwell JD, Robinson VP, Winder SM. Vitamin D supplements in pregnant Asian women: effects on calcium status and fetal growth. *Br Med J.* 1980; 280:751–754. [PubMed: 6989438]
17. Grant CC, Stewart AW, Scragg R, Milne T, Rowden J, Ekeroma A, Wall C, Mitchell EA, Crengle S, Trenholme A, Crane J, et al. Vitamin D During Pregnancy and Infancy and Infant Serum 25-Hydroxyvitamin D Concentration. *Pediatrics.* 2013
18. Mallet E, Gugi B, Brunelle P, Henocq A, Basuyau JP, Lemeur H. Vitamin D supplementation in pregnancy: a controlled trial of two methods. *Obstet Gynecol.* 1986; 68:300–304. [PubMed: 3755517]
19. Roth DE, Al Mahmud A, Raqib R, Akhtar E, Perumal N, Pezzack B, Baqui AH. Randomized placebo-controlled trial of high-dose prenatal third-trimester vitamin D3 supplementation in Bangladesh: the AViDD trial. *Nutr J.* 2013; 12:47. [PubMed: 23587190]
20. Yu CK, Sykes L, Sethi M, Teoh TG, Robinson S. Vitamin D deficiency and supplementation during pregnancy. *Clin Endocrinol (Oxf).* 2009; 70:685–690. [PubMed: 18771564]
21. Wagner CL, McNeil RB, Johnson DD, Hulsey TC, Ebeling M, Robinson C, Hamilton SA, Hollis BW. Health characteristics and outcomes of two randomized vitamin D supplementation trials during pregnancy: a combined analysis. *J Steroid Biochem Mol Biol.* 2013; 136:313–320. [PubMed: 23314242]
22. Hollis BW, Johnson D, Hulsey TC, Ebeling M, Wagner CL. Vitamin D supplementation during pregnancy: double-blind, randomized clinical trial of safety and effectiveness. *J Bone Miner Res.* 2011; 26:2341–2357. [PubMed: 21706518]
23. Dawodu A, Saadi HF, Bekdache G, Javed Y, Altaye M, Hollis BW. Randomized controlled trial (RCT) of vitamin D supplementation in pregnancy in a population with endemic vitamin D deficiency. *J Clin Endocrinol Metab.* 2013; 98:2337–2346. [PubMed: 23559082]
24. Hofmeyr GJ, Lawrie TA, Atallah AN, Duley L, Torloni MR. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *Cochrane Database Syst Rev.* 2014; 6:Cd001059. [PubMed: 24960615]
25. Bodnar LM, Catov JM, Simhan HN, Holick MF, Powers RW, Roberts JM. Maternal vitamin D deficiency increases the risk of preeclampsia. *J Clin Endocrinol Metab.* 2007; 92:3517–3522. [PubMed: 17535985]
26. Baker AM, Haeri S, Camargo CA Jr, Espinola JA, Stuebe AM. A nested case-control study of midgestation vitamin D deficiency and risk of severe preeclampsia. *J Clin Endocrinol Metab.* 2010; 95:5105–5109. [PubMed: 20719829]
27. Bodnar LM, Simhan HN, Catov JM, Roberts JM, Platt RW, Diesel JC, Klebanoff MA. Maternal vitamin D status and the risk of mild and severe preeclampsia. *Epidemiology.* 2014; 25:207–214. [PubMed: 24457526]
28. Wei SQ, Audibert F, Hidiroglou N, Sarafin K, Julien P, Wu Y, Luo ZC, Fraser WD. Longitudinal vitamin D status in pregnancy and the risk of pre-eclampsia. *Bjog.* 2012; 119:832–839. [PubMed: 22462640]
29. Robinson CJ, Alanis MC, Wagner CL, Hollis BW, Johnson DD. Plasma 25-hydroxyvitamin D levels in early-onset severe preeclampsia. *Am J Obstet Gynecol.* 2010; 203:366.e361–366. [PubMed: 20692641]
30. Xu L, Lee M, Jeyabalan A, Roberts JM. The relationship of hypovitaminosis D and IL-6 in preeclampsia. *Am J Obstet Gynecol.* 2014; 210:149.e141–147. [PubMed: 24080305]
31. Abedi P, Mohaghegh Z, Afshary P, Latifi M. The relationship of serum vitamin D with pre-eclampsia in the Iranian women. *Matern Child Nutr.* 2014; 10:206–212. [PubMed: 23782626]

32. Scholl TO, Chen X, Stein TP. Vitamin D, secondary hyperparathyroidism, and preeclampsia. *Am J Clin Nutr.* 2013; 98:787–793. [PubMed: 23885046]
33. Haugen M, Brantsaeter AL, Trogstad L, Alexander J, Roth C, Magnus P, Meltzer HM. Vitamin D supplementation and reduced risk of preeclampsia in nulliparous women. *Epidemiology.* 2009; 20:720–726. [PubMed: 19451820]
34. Shand AW, Nassar N, Von Dadelszen P, Innis SM, Green TJ. Maternal vitamin D status in pregnancy and adverse pregnancy outcomes in a group at high risk for pre-eclampsia. *Bjog.* 2010; 117:1593–1598. [PubMed: 21040394]
35. Powe CE, Seely EW, Rana S, Bhan I, Ecker J, Karumanchi SA, Thadhani R. First trimester vitamin D, vitamin D binding protein, and subsequent preeclampsia. *Hypertension.* 2010; 56:758–763. [PubMed: 20733087]
36. Seely EW, Wood RJ, Brown EM, Graves SW. Lower serum ionized calcium and abnormal calcitropic hormone levels in preeclampsia. *J Clin Endocrinol Metab.* 1992; 74:1436–1440. [PubMed: 1592891]
37. Fernandez-Alonso AM, Dionis-Sanchez EC, Chedraui P, Gonzalez-Salmeron MD, Perez-Lopez FR. First-trimester maternal serum 25-hydroxyvitamin D(3) status and pregnancy outcome. *Int J Gynaecol Obstet.* 2012; 116:6–9. [PubMed: 21959069]
38. Burris HH, Rifas-Shiman SL, Huh SY, Kleinman K, Litonjua AA, Oken E, Rich-Edwards JW, Camargo CA Jr, Gillman MW. Vitamin D status and hypertensive disorders in pregnancy. *Ann Epidemiol.* 2014; 24:399–403. e391. [PubMed: 24631523]
39. Zhou J, Su L, Liu M, Liu Y, Cao X, Wang Z, Xiao H. Associations between 25-hydroxyvitamin D levels and pregnancy outcomes: a prospective observational study in southern China. *Eur J Clin Nutr.* 2014
40. Dalmar A, Raff H, Chauhan SP, Singh M, Siddiqui DS. Serum 25-hydroxyvitamin D, calcium, and calcium-regulating hormones in preeclampsia and controls during first day postpartum. *Endocrine.* 2014
41. Yu CK, Ertl R, Skyfta E, Akolekar R, Nicolaides KH. Maternal serum vitamin D levels at 11-13 weeks of gestation in preeclampsia. *J Hum Hypertens.* 2013; 27:115–118. [PubMed: 22336906]
42. Hypponen E, Cavadino A, Williams D, Fraser A, Vereczkey A, Fraser WD, Banhidy F, Lawlor D, Czeizel AE. Vitamin D and pre-eclampsia: original data, systematic review and meta-analysis. *Ann Nutr Metab.* 2013; 63:331–340. [PubMed: 24603503]
43. Thorne-Lyman A, Fawzi WW. Vitamin D during pregnancy and maternal, neonatal and infant health outcomes: a systematic review and meta-analysis. *Paediatr Perinat Epidemiol.* 2012; 26 Suppl 1:75–90. [PubMed: 22742603]
44. Aghajafari F, Nagulesapillai T, Ronksley PE, Tough SC, O'Beirne M, Rabi DM. Association between maternal serum 25-hydroxyvitamin D level and pregnancy and neonatal outcomes: systematic review and meta-analysis of observational studies. *BMJ.* 2013; 346:f1169. [PubMed: 23533188]
45. Tabesh M, Salehi-Abargouei A, Tabesh M, Esmailzadeh A. Maternal vitamin D status and risk of pre-eclampsia: a systematic review and meta-analysis. *J Clin Endocrinol Metab.* 2013; 98:3165–3173. [PubMed: 23783101]
46. Wei SQ, Qi HP, Luo ZC, Fraser WD. Maternal vitamin D status and adverse pregnancy outcomes: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med.* 2013; 26:889–899. [PubMed: 23311886]
47. Harvey N, Holroyd C, Ntani G, Javaid M, Cooper P, Moon R, Cole Z, Tinati T, Godfrey K, Dennison E, Bishop N, et al. Vitamin D supplementation in pregnancy: a systematic review. *Health Technol Assess.* 2014; 18
48. Schneuer FJ, Roberts CL, Guilbert C, Simpson JM, Algert CS, Khambalia AZ, Tasevski V, Ashton AW, Morris JM, Nassar N. Effects of maternal serum 25-hydroxyvitamin D concentrations in the first trimester on subsequent pregnancy outcomes in an Australian population. *Am J Clin Nutr.* 2014; 99:287–295. [PubMed: 24257720]
49. Reeves IV, Bamji ZD, Rosario GB, Lewis KM, Young MA, Washington KN. Vitamin D deficiency in pregnant women of ethnic minority: a potential contributor to preeclampsia. *J Perinatol.* 2014

50. Xu L, Nicholson P, Wang Q, Alen M, Cheng S. Bone and muscle development during puberty in girls: a seven-year longitudinal study. *J Bone Miner Res.* 2009; 24:1693–1698. [PubMed: 19419294]
51. Bener A, Al-Hamaq AO, Saleh NM. Association between vitamin D insufficiency and adverse pregnancy outcome: global comparisons. *Int J Womens Health.* 2013; 5:523–531. [PubMed: 24043954]
52. Wetta LA, Biggio JR, Cliver S, Abramovici A, Barnes S, Tita AT. Is midtrimester vitamin D status associated with spontaneous preterm birth and preeclampsia? *Am J Perinatol.* 2014; 31:541–546. [PubMed: 24022379]
53. Ullah MI, Koch CA, Tamanna S, Rouf S, Shamsuddin L. Vitamin D deficiency and the risk of preeclampsia and eclampsia in Bangladesh. *Horm Metab Res.* 2013; 45:682–687. [PubMed: 23733167]
54. Robinson CJ, Wagner CL, Hollis BW, Baatz JE, Johnson DD. Association of maternal vitamin D and placenta growth factor with the diagnosis of early onset severe preeclampsia. *Am J Perinatol.* 2013; 30:167–172. [PubMed: 22875657]
55. Anderson CM, Ralph J, Johnson L, Scheett A, Wright ML, Taylor JY, Ohm JE, Uthus E. First trimester vitamin D status and placental epigenomics in preeclampsia among Northern Plains primiparas. *Life Sci.* 2014
56. Maghbooli Z, Hossein-Nezhad A, Karimi F, Shafaei AR, Larjani B. Correlation between vitamin D3 deficiency and insulin resistance in pregnancy. *Diabetes Metab Res Rev.* 2008; 24:27–32. [PubMed: 17607661]
57. Clifton-Bligh RJ, McElduff P, McElduff A. Maternal vitamin D deficiency, ethnicity and gestational diabetes. *Diabet Med.* 2008; 25:678–684. [PubMed: 18544105]
58. Lacroix M, Battista MC, Doyon M, Houde G, Menard J, Ardilouze JL, Hivert MF, Perron P. Lower vitamin D levels at first trimester are associated with higher risk of developing gestational diabetes mellitus. *Acta Diabetol.* 2014
59. McManus R, Summers K, de Vrijer B, Cohen N, Thompson A, Giroux I. Maternal, umbilical arterial and umbilical venous 25-hydroxyvitamin D and adipocytokine concentrations in pregnancies with and without gestational diabetes. *Clin Endocrinol (Oxf).* 2014; 80:635–641. [PubMed: 24102192]
60. Cho GJ, Hong SC, Oh MJ, Kim HJ. Vitamin D deficiency in gestational diabetes mellitus and the role of the placenta. *Am J Obstet Gynecol.* 2013; 209:560.e561–568. [PubMed: 23954530]
61. Wang O, Nie M, Hu YY, Zhang K, Li W, Ping F, Liu JT, Chen LM, Xing XP. Association between vitamin D insufficiency and the risk for gestational diabetes mellitus in pregnant Chinese women. *Biomed Environ Sci.* 2012; 25:399–406. [PubMed: 23026519]
62. Farrant HJ, Krishnaveni GV, Hill JC, Boucher BJ, Fisher DJ, Noonan K, Osmond C, Veena SR, Fall CH. Vitamin D insufficiency is common in Indian mothers but is not associated with gestational diabetes or variation in newborn size. *Eur J Clin Nutr.* 2009; 63:646–652. [PubMed: 18285809]
63. Whitelaw DC, Scally AJ, Tuffnell DJ, Davies TJ, Fraser WD, Bhopal RS, Wright J, Lawlor DA. Associations of circulating calcium and 25-hydroxyvitamin D with glucose metabolism in pregnancy: a cross-sectional study in European and South Asian women. *J Clin Endocrinol Metab.* 2014; 99:938–946. [PubMed: 24423329]
64. Perez-Ferre N, Torrejon MJ, Fuentes M, Fernandez MD, Ramos A, Bordiu E, del Valle L, Rubio MA, Bedia AR, Montanez C, Calle-Pascual AL. Association of low serum 25-hydroxyvitamin D levels in pregnancy with glucose homeostasis and obstetric and newborn outcomes. *Endocr Pract.* 2012; 18:676–684. [PubMed: 22548949]
65. Poel YH, Hummel P, Lips P, Stam F, van der Ploeg T, Simsek S. Vitamin D and gestational diabetes: a systematic review and meta-analysis. *Eur J Intern Med.* 2012; 23:465–469. [PubMed: 22726378]
66. Parildar H, Dogruk Unal A, Aksan Desteli G, Cigerli O, Guvener Demirag N. Frequency of Vitamin D deficiency in pregnant diabetics at Baskent University Hospital, Istanbul. *Pak J Med Sci.* 2013; 29:15–20. [PubMed: 24353500]

67. Merewood A, Mehta SD, Chen TC, Bauchner H, Holick MF. Association between vitamin D deficiency and primary cesarean section. *J Clin Endocrinol Metab.* 2009; 94:940–945. [PubMed: 19106272]
68. Scholl TO, Chen X, Stein P. Maternal vitamin D status and delivery by cesarean. *Nutrients.* 2012; 4:319–330. [PubMed: 22606373]
69. Savvidou MD, Makgoba M, Castro PT, Akolekar R, Nicolaides KH. First-trimester maternal serum vitamin D and mode of delivery. *Br J Nutr.* 2012; 108:1972–1975. [PubMed: 22360844]
70. Dunlop AL, Taylor RN, Tangpricha V, Fortunato S, Menon R. Maternal micronutrient status and preterm versus term birth for black and white US women. *Reprod Sci.* 2012; 19:939–948. [PubMed: 22527984]
71. Mehta S, Hunter DJ, Mugusi FM, Spiegelman D, Manji KP, Giovannucci EL, Hertzmark E, Msamanga GI, Fawzi WW. Perinatal outcomes, including mother-to-child transmission of HIV, and child mortality and their association with maternal vitamin D status in Tanzania. *J Infect Dis.* 2009; 200:1022–1030. [PubMed: 19673647]
72. Delmas PD, Glorieux FH, Delvin EE, Salle BL, Melki I. Perinatal serum bone Gla-protein and vitamin D metabolites in preterm and fullterm neonates. *J Clin Endocrinol Metab.* 1987; 65:588–591. [PubMed: 3497940]
73. Thorp JM, Camargo CA, McGee PL, Harper M, Klebanoff MA, Sorokin Y, Varner MW, Wapner RJ, Caritis SN, Iams JD, Carpenter MW, et al. Vitamin D status and recurrent preterm birth: a nested case-control study in high-risk women. *Bjog.* 2012; 119:1617–1623. [PubMed: 23078336]
74. Baker AM, Haeri S, Camargo CA Jr, Stuebe AM, Boggess KA. A nested case-control study of first-trimester maternal vitamin D status and risk for spontaneous preterm birth. *Am J Perinatol.* 2011; 28:667–672. [PubMed: 21500145]
75. Bodnar LM, Rouse DJ, Momirova V, Peaceman AM, Sciscione A, Spong CY, Varner MW, Malone FD, Iams JD, Mercer BM, Thorp JM Jr, et al. Maternal 25-hydroxyvitamin d and preterm birth in twin gestations. *Obstet Gynecol.* 2013; 122:91–98. [PubMed: 23743453]
76. Bodnar LM, Klebanoff MA, Gernand AD, Platt RW, Parks WT, Catov JM, Simhan HN. Maternal vitamin D status and spontaneous preterm birth by placental histology in the US Collaborative Perinatal Project. *Am J Epidemiol.* 2014; 179:168–176. [PubMed: 24124195]
77. Hossain N, Khanani R, Hussain-Kanani F, Shah T, Arif S, Pal L. High prevalence of vitamin D deficiency in Pakistani mothers and their newborns. *Int J Gynaecol Obstet.* 2011; 112:229–233. [PubMed: 21247568]
78. Li N, Liu E, Guo J, Pan L, Li B, Wang P, Liu J, Wang Y, Liu G, Baccarelli AA, Hou L, et al. Maternal prepregnancy body mass index and gestational weight gain on pregnancy outcomes. *PLoS One.* 2013; 8:e82310. [PubMed: 24376527]
79. Bautista-Castano I, Henriquez-Sanchez P, Aleman-Perez N, Garcia-Salvador JJ, Gonzalez-Quesada A, Garcia-Hernandez JA, Serra-Majem L. Maternal obesity in early pregnancy and risk of adverse outcomes. *PLoS One.* 2013; 8:e80410. [PubMed: 24278281]
80. Zhang S, Cardarelli K, Shim R, Ye J, Booker KL, Rust G. Racial disparities in economic and clinical outcomes of pregnancy among Medicaid recipients. *Matern Child Health J.* 2013; 17:1518–1525. [PubMed: 23065298]
81. Curtis EM, Moon RJ, Dennison EM, Harvey NC. Prenatal calcium and vitamin d intake, and bone mass in later life. *Curr Osteoporos Rep.* 2014; 12:194–204. [PubMed: 24740166]
82. Hossain N, Kanani FH, Ramzan S, Kausar R, Ayaz S, Khanani R, Pal L. Obstetric and neonatal outcomes of maternal vitamin D supplementation: Results of an open label randomized controlled trial of antenatal vitamin D supplementation in Pakistani women. *J Clin Endocrinol Metab.* 2014; jc20133491.
83. Yap C, Cheung NW, Gunton JE, Athayde N, Munns CF, Duke A, McLean M. Vitamin d supplementation and the effects on glucose metabolism during pregnancy: a randomized controlled trial. *Diabetes Care.* 2014; 37:1837–1844. [PubMed: 24760259]
84. Kalra P, Das V, Agarwal A, Kumar M, Ramesh V, Bhatia E, Gupta S, Singh S, Saxena P, Bhatia V. Effect of vitamin D supplementation during pregnancy on neonatal mineral homeostasis and anthropometry of the newborn and infant. *Br J Nutr.* 2012; 108:1052–1058. [PubMed: 22212646]

85. Marya RK, Rathee S, Manrow M. Effect of Calcium and Vitamin D Supplementation on Toxaemia of Pregnancy. *Gynecol Obstet Invest.* 1987; 24:38–42. [PubMed: 3623260]
86. NICE Clinical Guideline 63. Diabetes in Pregnancy. 2008
87. Soheilykhah S, Mojibian M, Moghadam MJ, Shojaoddiny-Ardekani A. The effect of different doses of vitamin D supplementation on insulin resistance during pregnancy. *Gynecol Endocrinol.* 2013; 29:396–399. [PubMed: 23350644]
88. Asemi Z, Hashemi T, Karamali M, Samimi M, Esmailzadeh A. Effects of vitamin D supplementation on glucose metabolism, lipid concentrations, inflammation, and oxidative stress in gestational diabetes: a double-blind randomized controlled clinical trial. *Am J Clin Nutr.* 2013; 98:1425–1432. [PubMed: 24132976]
89. Innes AM, Seshia MM, Prasad C, Al SS, Friesen FR, Chudley AE, Reed M, Dilling LA, Haworth JC, Greenberg CR. Congenital rickets caused by maternal vitamin D deficiency. *Paediatr Child Health.* 2002; 7:455–458. [PubMed: 20046322]
90. Anatoliotaki M, Tsilimigaki A, Tsekoura T, Schinaki A, Stefanaki S, Nicolaidou P. Congenital rickets due to maternal vitamin D deficiency in a sunny island of Greece. *Acta Paediatr.* 2003; 92:389–391. [PubMed: 12725557]
91. Orbak Z, Karacan M, Doneray H, Karakelleoglu C. Congenital rickets presenting with hypocalcaemic seizures. *West Indian Med J.* 2007; 56:364–367. [PubMed: 18198743]
92. Marya RK, Rathee S, Lata V, Mudgil S. Effects of vitamin D supplementation in pregnancy. *Gynecol Obstet Invest.* 1981; 12:155–161. [PubMed: 7239350]
93. Hashemipour S, Ziaee A, Javadi A, Movahed F, Elmizadeh K, Javadi EH, Lalooha F. Effect of treatment of vitamin D deficiency and insufficiency during pregnancy on fetal growth indices and maternal weight gain: a randomized clinical trial. *Eur J Obstet Gynecol Reprod Biol.* 2014; 172:15–19. [PubMed: 24210789]
94. Ioannou C, Javaid MK, Mahon P, Yaqub MK, Harvey NC, Godfrey KM, Noble JA, Cooper C, Papageorgiou AT. The effect of maternal vitamin D concentration on fetal bone. *J Clin Endocrinol Metab.* 2012; 97:E2070–2077. [PubMed: 22990090]
95. Mahon P, Harvey N, Crozier S, Inskip H, Robinson S, Arden N, Swaminathan R, Cooper C, Godfrey K, Group SWSS. Low maternal vitamin D status and fetal bone development: cohort study. *J Bone Miner Res.* 2010; 25:14–19. [PubMed: 19580464]
96. Namgung R, Tsang RC, Lee C, Han DG, Ho ML, Sierra RI. Low total body bone mineral content and high bone resorption in Korean winter-born versus summer-born newborn infants. *J Pediatr.* 1998; 132:421–425. [PubMed: 9544894]
97. Namgung R, Tsang RC. Factors affecting newborn bone mineral content: in utero effects on newborn bone mineralization. *Proc Nutr Soc.* 2000; 59:55–63. [PubMed: 10828174]
98. Weiler H, Fitzpatrick-Wong S, Veitch R, Kovacs H, Schellenberg J, McCloy U, Yuen CK. Vitamin D deficiency and whole-body and femur bone mass relative to weight in healthy newborns. *CMAJ.* 2005; 172:757–761. [PubMed: 15767609]
99. Viljakainen HT, Saarnio E, Hytinanti T, Miettinen M, Surcel H, Makitie O, Andersson S, Laitinen K, Lamberg-Allardt C. Maternal vitamin D status determines bone variables in the newborn. *J Clin Endocrinol Metab.* 2010; 95:1749–1757. [PubMed: 20139235]
100. Viljakainen HT, Korhonen T, Hytinanti T, Laitinen EK, Andersson S, Makitie O, Lamberg-Allardt C. Maternal vitamin D status affects bone growth in early childhood--a prospective cohort study. *Osteoporos Int.* 2011; 22:883–891. [PubMed: 21153404]
101. Prentice A, Jarjou LM, Goldberg GR, Bennett J, Cole TJ, Schoenmakers I. Maternal plasma 25-hydroxyvitamin D concentration and birthweight, growth and bone mineral accretion of Gambian infants. *Acta Paediatr.* 2009; 98:1360–1362. [PubMed: 19594476]
102. Sayers A, Tobias JH. Estimated maternal ultraviolet B exposure levels in pregnancy influence skeletal development of the child. *J Clin Endocrinol Metab.* 2009; 94:765–771. [PubMed: 19116232]
103. Lawlor DA, Wills AK, Fraser A, Sayers A, Fraser WD, Tobias JH. Association of maternal vitamin D status during pregnancy with bone-mineral content in offspring: a prospective cohort study. *The Lancet.* 2013; 381:2176–2183.

104. Harvey NC, Javaid MK, Inskip HM, Godfrey KM, Cooper C. Maternal vitamin D status during pregnancy and bone-mineral content in offspring. *Lancet*. 2013; 382:766. [PubMed: 23993185]
105. Zhu K, Whitehouse AJ, Hart P, Kusel M, Mountain J, Lye S, Pennell C, Walsh JP. Maternal Vitamin D Status During Pregnancy and Bone Mass in Offspring at 20 Years of Age: A Prospective Cohort Study. *J Bone Miner Res*. 2013
106. Congdon P, Horsman A, Kirby PA, Dibble J, Bashir T. Mineral content of the forearms of babies born to Asian and white mothers. *Br Med J (Clin Res Ed)*. 1983; 286:1233–1235.
107. Harvey NC, Javaid K, Bishop N, Kennedy S, Papageorghiou AT, Fraser R, Gandhi SV, Schoenmakers I, Prentice A, Cooper C. MAVIDOS Maternal Vitamin D Osteoporosis Study: study protocol for a randomized controlled trial. The MAVIDOS Study Group. *Trials*. 2012; 13:13. [PubMed: 22314083]
108. Harvey NC, Cooper C. Vitamin D: some perspective please. *BMJ*. 2012; 345:e4695. [PubMed: 22815431]
109. Wagner CL, McNeil R, Hamilton SA, Winkler J, Rodriguez Cook C, Warner G, Bivens B, Davis DJ, Smith PG, Murphy M, Shary JR, et al. A randomized trial of vitamin D supplementation in 2 community health center networks in South Carolina. *Am J Obstet Gynecol*. 2013; 208:137.e131–113. [PubMed: 23131462]
110. Marya RK, Rathee S, Dua V, Sangwan K. Effect of vitamin D supplementation during pregnancy on foetal growth. *Indian J Med Res*. 1988; 88:488–492. [PubMed: 3243609]




Figure 1.

Forest plot of the association between maternal vitamin D status and risk of preeclampsia (observational studies)

Reproduced from Harvey N, Holroyd C, Ntani G, Javaid M, Cooper P, Moon R, Cole Z, Tinati T, Godfrey K, Dennison E, Bishop N, Baird J & Cooper C. Vitamin D supplementation in pregnancy: a systematic review. *Health Technol Assess* 2014 **18**.




Figure 2. Meta-analysis of maternal serum 25(OH)D in pregnancy and gestational diabetes. Reproduced from Association between maternal serum 25-hydroxyvitamin D level and pregnancy and neonatal outcomes: systematic review and meta-analysis of observational studies, Aghajafari F et al, BMJ 2013;346:f1169 doi: [10.1136/bmj.f1169](https://doi.org/10.1136/bmj.f1169) with permission from BMJ Publishing Group Ltd.




Figure 3. Maternal 25(OH)D concentration in late pregnancy and childhood bone mass at age 9 years
Reprinted from The Lancet, Vol 367, Javaid MK et al, Maternal vitamin D status during pregnancy and childhood bone mass at age 9 years: a longitudinal study, Pages 36–43, Copyright (2014), with permission from Elsevier

Table 1
Meta-analyses of maternal vitamin D status (intake and serum 25-hydroxyvitamin D level) and risk of pre-eclampsia.

Author	Publication cut-off	Number of studies included	Number of women included	Comparison	Direction of effect	Risk of preeclampsia with low vitamin D status	Reported odds ratio (95% CI)
Vitamin D intake							
Thorne-Lyman, 2012 43	June 2011	2	25141	Highest vs lowest category of vitamin D intake	↔		0.95 (0.86, 1.06)
Hypponen, 2013 42	March 2013 + inclusion of novel data	2	77165	Self-supplementation vs unsupplemented	↑		1.23 (1.15, 1.33)
Serum 25(OH)D							
Aghajafari, 2013 44	August 2012	2	697	Serum 25(OH)D 50nmol/l vs <50nmol/l	↔		1.27 (0.67, 2.42)
		5	1165	Serum 25(OH)D 75nmol/l vs <75nmol/l	↑		2.11 (1.36, 3.27)
		7	1862	Higher serum 25(OH)D as defined by each study vs lower serum 25(OH)D	↑		1.79 (1.25, 2.58)
		7	1862	Higher serum 25(OH)D as defined by each study vs lower serum 25(OH)D, adjusted for "critical confounders"	↔		1.51 (0.89, 2.57)
Hypponen, 2013 42	March 2013 + inclusion of novel data	6	6864	Higher serum 25(OH)D as defined by each study vs lower serum 25(OH)D	↑		1.92 (1.12, 3.33)
Tabesh, 2013 45							
	December 2012	4	931	Serum 25(OH)D 38nmol/l vs <38nmol/l	↔		Actual odds ratios not reported
		5	1775	Serum 25(OH)D 50nmol/l vs <50nmol/l	↑		
		8	2485	Higher serum 25(OH)D as defined by each study vs lower serum 25(OH)D	↑		
Wei, 2013 46							
	October 2012	6	610	Serum 25(OH)D 50nmol/l vs <50nmol/l	↑		2.09 (1.50, 2.90)
		5	802	Serum 25(OH)D 75nmol/l vs <75nmol/l	↑		1.78 (1.23, 2.56)
Harvey, 2014 47							
	June 2012	4	628	Each 25nmol/l increase in serum 25(OH)D	↔		0.78 (0.59-1.05)

Table 2
Intervention studies of vitamin D supplementation (alone, and in combination with calcium supplementation) in pregnancy to reduce obstetric complications.

Study	Population	Gestation at randomisation	Interventional medicinal product (IMP)	Control	Effect of IMP vs control on incidence of obstetric events					
					Hypertensive disorders GHT	GDM	Preterm delivery	Caesarean section	Intrauterine death/stillbirth	
Vitamin D supplementation										
Hossain, 2014 82 (Karachi, Pakistan)	N=178	20 weeks	4000 IU/day oral cholecalciferol	Usual care	↔	↔	↔	↔	↔	↔
Wagner, 2013 21 (South Carolina, USA) ¹	N=504	12-16 weeks	2000 IU/day oral cholecalciferol (n=201) 4000 IU/day oral cholecalciferol (n=193)	400 IU/day oral cholecalciferol (n=111)	↔	↔	↔	↔	↔	↔
Yap, 2014 83 (Sydney, Australia)	N=179 25(OH)D<80nmol/l at baseline	< 20 weeks	5000 IU/day oral cholecalciferol	400 IU/day oral cholecalciferol	↔	↔	↔	↔	↔	↔
Vitamin D + Calcium supplementation										
Kalra, 2011 84 (Lucknow, India)	N=140	12-24 weeks	Group 1: 60,000 IU single dose oral cholecalciferol at recruitment + 1g elemental Ca/day until delivery (n=48) Group 2: 120,000 IU oral cholecalciferol at recruitment and 28 weeks gestation + 1g elemental Ca/day until delivery (n=49)	Usual care (n=43)	↔	↔	↔	↔	↔	↔
Marya, 1987 85 (Rothak, India)	N=400	20-24 weeks	1200 IU/day vitamin D + 375mg calcium	Usual care	↔	↔	↔	↔	↔	↔

↔ no effect shown, ↓ vitamin D supplementation reduced the incidence of the outcome; GHT – gestational hypertension; PET – preeclampsia; GDM – gestational diabetes mellitus. (1) This reported a combined analysis of data collected in two previous studies.22,109

Table 3
Intervention studies of the effect of vitamin D supplementation in pregnancy on offspring anthropometry at birth

Study	Population	Gestation at Allocation/Randomisation	Interventional medicinal product (IMP)	Control	Effect of vitamin D supplementation
					Birth Weight Birth Length Head Circumference
Vitamin D only					
Brooke 1980 16 (London, UK)	126 Asian women	28-32 weeks	1000 IU/day oral vitamin D	Placebo	↔ ↔ ↔
Mallet 1986 18 (France)	68 women	Last trimester	Group A: 1000 IU/day oral vitamin D Group B: 200,000 IU single dose in 7 th month of pregnancy	Usual care	↔
Marya 1988 110 (Rohtak, India)	200 Indian women	7 months	Single dose of 600000IU cholecalciferol in months 7 and 8 of pregnancy	Usual care	↑ ↑ ↑
Yu 2009 20 (London, UK)	180 women	27 weeks	Group A: 800 IU/day oral cholecalciferol Group B: 200000IU oral cholecalciferol single dose at 27 weeks gestation	Usual care	↔
Dawodu 2013 23 (Al Ain, UAE)	192 Arab women	12-16 weeks	Group A: 4000 IU/day oral cholecalciferol Group B: 2000 IU/day oral cholecalciferol	400 IU/day oral cholecalciferol	↔ ↔ ↔
Grant 2013 17 (Auckland, New Zealand)	260 women	26-30 weeks	Group A: 10000IU/day oral cholecalciferol Group B: 20000IU/day	Placebo	↔

Study	Population	Gestation at Allocation/Randomisation	Interventional medicinal product (IMP)	Control	Effect of vitamin D supplementation		
					Birth Weight	Birth Length	Head Circumference
Wagner 2013 21 (USA)	Combined analysis of two trials including a total of 513 women	12-16 weeks	oral cholecalciferol Group A: 2000IU/day oral cholecalciferol Group B: 4000IU/day oral cholecalciferol	400 IU/day oral cholecalciferol	↔	↔	↔
Roth, 2013 19 (Dhaka, Bangladesh)	148	26-30 weeks	35000 IU/week oral cholecalciferol	Placebo	↔	↔	↔
Vitamin D + calcium Marya 1981 92 (Rohtak, India)	120 Hindu women	Last trimester	Group A: 12000IU/day vitamin D + 375mg calcium during third trimester Group B: 600000IU vitamin D orally in the 7 th and 8 th months of pregnancy (n=20)	Usual care	↑		
Kalra 2011 84 (Lucknow, India)	140 women	12-24 weeks	Group A: 60,000IU oral cholecalciferol single dose at randomisation + 1g/day calcium carbonate Group B: 120,000IU oral cholecalciferol at randomisation and at 28 weeks gestation + 1g/day calcium carbonate	1g calcium carbonate/day	↑	↑	↑

Study	Population	Gestation at Allocation/Randomisation	Interventional medicinal product (IMP)	Control	Effect of vitamin D supplementation		
					Birth Weight	Birth Length	Head Circumference
Hashemipour 2014 93 (Qazvin, Iran)	109 women, 25(OH)D<75nmol/l	24-26 weeks	50,000 IU/week cholecalciferol for 8 weeks in addition to the supplement received by control group	400IU/day oral cholecalciferol; 200mg elemental calcium	↑	↑	↑
Hossain 2014 82 (Karachi, Pakistan)	198	20 weeks	4000IU/day oral cholecalciferol, 600mg calcium lactate & 200mg ferrous sulphate	600mg calcium lactate & 200mg ferrous sulphate	↔	↔	↔

↔ no effect shown,

↑ vitamin D supplementation increased the outcome,

↓ vitamin D supplementation the outcome