

Factors Associated With Emergency Department Visits: A Multistate Analysis of Adult Fee-for-Service Medicaid Beneficiaries

Parul Agarwal¹, Thomas K. Bias², Suresh Madhavan¹,
Nethra Sambamoorthi³, Stephanie Frisbee², and Usha Sambamoorthi¹

Abstract

Objective: The objective of this study was to examine the association of patient- and county-level factors with the emergency department (ED) visits among adult fee-for-service (FFS) Medicaid beneficiaries residing in Maryland, Ohio, and West Virginia.

Methods: A cross-sectional design using retrospective observational data was implemented. Patient-level data were obtained from 2010 Medicaid Analytic eXtract files. Information on county-level health-care resources was obtained from the Area Health Resource file and County Health Rankings file.

Results: In adjusted analyses, the following patient-level factors were associated with higher number of ED visits: African Americans (incidence rate ratios [IRR] = 1.47), Hispanics (IRR = 1.63), polypharmacy (IRR = 1.89), and tobacco use (IRR = 2.23). Patients with complex chronic illness had a higher number of ED visits (IRR = 3.33). The county-level factors associated with ED visits were unemployment rate (IRR = 0.94) and number of urgent care clinics (IRR = 0.96).

Conclusion: Patients with complex healthcare needs had a higher number of ED visits as compared to those without complex healthcare needs. The study results provide important baseline context for future policy analysis studies around Medicaid expansion options.

Keywords

access to care, emergency visits, primary care, community health, medical cost

Introduction

The Center for Disease Control and Prevention estimated 91.1 million visits to the emergency department (ED) in 2010 among adults aged 18 to 64 years.¹ Emergency Department visits for health care are a major concern because a majority of these visits are for the care of ambulatory care sensitive conditions (ACSCs) and can be prevented with timely primary care.² Emergency Department visits for ACSCs impose significant economic burden compared to other outpatient settings.^{3,4} Emergency department visits due to ACSCs account for US\$38 billion of total health-care spending in the United States.⁵

There is a common misperception that almost all ED users who have preventable conditions or are frequent ED users are uninsured.⁶ Emergency department users often have health insurance including Medicaid or Medicare.^{6,7} Emergency department visits by Medicaid beneficiaries accounted for

about 12% of the total health-care spending on ED services in 2012.⁸

Emergency department use among Medicaid beneficiaries has received considerable attention due to expanding Medicaid eligibility in 31 states. As there is some evidence of an inverse

¹ Department of Pharmaceutical Systems and Policy, School of Pharmacy, West Virginia University, Morgantown, WV, USA

² Department of Health Policy, Management and Leadership, School of Public Health, West Virginia University, Morgantown, WV, USA

³ School of Professional Studies, Northwestern University, Chicago, IL, USA

Submitted April 14, 2016. Accepted April 14, 2016.

Corresponding Author:

Thomas K. Bias, Department of Health Policy, Management, and Leadership, West Virginia University, PO Box 9136, Morgantown, WV 26501, USA.
Email: tbias@hsc.wvu.edu



relationship between ED visits and community-level supply of primary care providers and health centers,⁹ it could be hypothesized that providing health insurance coverage to the uninsured without corresponding increases in primary care availability may lead to increased ED visits for nonemergent care. However, published literature reveals mixed findings. Some investigations have found little or no change in the ED utilization after provision of insurance coverage to the uninsured,^{10,11} while others found that the Medicaid expansion significantly increased both preventable and nonpreventable ED utilization.¹² These studies have limited ability to provide conclusive evidence because they did not include a systematic adjustment for a comprehensive set of patient- and county-level factors.

The primary objective of the current study is to use patient-level administrative claims data to examine the association between the patient- and county-level factors and the ED visits. For the purposes of the study, data on ED visits by adult, fee-for-service (FFS) Medicaid beneficiaries residing in Maryland, Ohio, and West Virginia were selected. These states were chosen as part of a long-term plan to investigate differences between the impact of the Medicaid Expansion (Ohio did not immediately expand, while the other states did) and the Health Insurance Marketplace (Maryland created a state-based marketplace, Ohio a federally facilitated marketplace, and West Virginia a partnership model marketplace). All states have many subtle differences within their Medicaid programs and benefits, as well, which can be compared easily through the Kaiser Family Foundation's online database.¹³ Andersen's behavioral model (ABM) for health-care services usage was adopted to provide a conceptual framework for the study.¹⁴ The ABM model posits that the health-care services utilization (ED visits in the current study) can be influenced by *predisposing, enabling, need, external environment factors, and personal health practices* as explained in the measures section.

Methods

Study Design

This study used a retrospective cross-sectional design with observational data.

Data Sources

Medicaid Analytic Extract (MAX) Files-2010. MAX files are prepared and released by The Centers of Medicare and Medicaid in assistance with Research Data Assistance Center. These files include (1) personal summary, (2) inpatient claims, (3) other therapy claims, and (4) prescription drugs claims. The current study used data on Medicaid beneficiaries residing in Maryland, Ohio, and West Virginia.

Area Health Resources Files (AHRF). The Area Health Resources Files contain national-, state-, and county-level data on approximately 6000 variables.¹⁵

County Health Rankings Data. The County Health Rankings data provides information on health behaviors, clinical care, social

and economic factors, and physical environment for all counties in all states.¹⁶ This information is compiled from 50 different data sources.

Study Population

The inclusion criteria were: FFS Medicaid beneficiaries, aged 22 to 64 years, with continuous Medicaid enrollment, not eligible for Medicare, and alive during the entire observation period. Pregnant women were excluded from the analysis because they may have unique prenatal needs (N = 68 882).

Dependent Variable

Number of ED visits. Emergency department visits were identified from inpatient and outpatient claims. To identify ED visits from the outpatient claims, CPT codes (99281-85) were used. Emergency department visits from the inpatient claims were identified using revenue codes (450-52, 456, 459, and 981).

Independent variables. Predisposing factors were age, gender, and race/ethnicity. Enabling factors were patient-level Medicaid eligibility due to cash assistance/poverty (cash eligibility, no cash eligibility), county-level college education rate, primary care use (none, fragmented, continuous), and county-level unemployment rate. Primary care use was measured using the modified, modified continuity index (MMCI) developed by Magill and Senf,¹⁷ which ranged from 0 to 1. The MMCI was developed to solve limitations of the Usual Provider of Care and modified continuity index.

Need factors were patient-level complex chronic illness (physical health conditions, mental health conditions, physical and mental health conditions, none), Medicaid eligibility due to medical need/waiver, and polypharmacy (Yes, No). Polypharmacy was defined as concomitant use of multiple prescription drugs within a 90-day period and was based on number of prescription drugs 1 standard deviation above the mean.¹⁸ In this study, complex chronic illness was defined as those having both physical and mental health conditions. The selection of physical and mental health conditions was based on the framework provided by the Health and Human Services Office of the Assistant Secretary of Health.¹⁹ Both physical and mental health conditions were identified if patients had an inpatient or outpatient claim. Personal health practices were patient-level tobacco use (clinically coded into the Medicaid data) and county-level obesity rates.

External environment factors were measured at the county-level and included metro status, health professional shortage area (HPSA—no, partial, and complete shortage areas), number of hospitals with EDs, number of hospitals with psychiatric emergency services, number of rural health clinics, number of federally qualified health centers (FQHCs), number of community mental health centers, and number of urgent care clinics per 100 000 population.

Statistical Analysis

Frequencies were used to describe the characteristics of the study population. Mean, interquartile range (IQR), and 90th percentile were used to describe the frequency of ED visits. The frequency of ED visits can be analyzed using a variety of count data regression models. They include Poisson regression, negative binomial regression (NBR), zero inflated Poisson regression (ZIP), and zero inflated NBR (ZINB). After comparing the predicted and actual probabilities, and log likelihood from all 4 statistical models, NBR and ZINB models were deemed appropriate. The ZINB model is complex and difficult to interpret due to its 2-part structure, and many economists and statisticians discourage using ZINB models when NBR models fit well with the data.²⁰ Therefore, this study used both unadjusted and adjusted NBR models to examine the patient- and county-level factors associated with the number of ED visits.

The adjusted NBR models included predisposing, enabling, need, external environment factors, and personal health practices. The parameter estimates from the NBR models were converted to incidence rate ratios (IRRs) by exponentiating the regression coefficients, and 95% confidence intervals were estimated. Incidence rate ratio can be interpreted as the percent change in ED visits. Incidence rate ratio above 1.0 implies higher number of ED visits and IRR below 1.0 implies lower number of ED visits. The data consisted of all counties in Maryland, Ohio, and West Virginia (167) and patients nested within these counties. Therefore, the NBR models were adjusted for clustering due to counties using STATA version 14.

Results

During 2010, 46% of the study population had zero ED visits. Around, 36% beneficiaries had ≤3 ED visits, 14% had 4 to 9 ED visits, and remaining 4% had 10 or more ED visits. Table 1 summarizes the patient-level characteristics of the study population. The majority were females (56.2%), older adults aged 45 to 64 (58.1%), whites (85.3%), were eligible for Medicaid through cash assistance/poverty (81.4%), lived in counties designated as either whole/part county HPSA (79.5%), and had fragmented/no primary care use (89.4%). In all, 18% had polypharmacy and 38% had both physical and mental health conditions. About 6.6% of the study population were tobacco users.

The following were the range of county-level factors: college education rate 4% to 42%, unemployment rate 2.7% to 10.5%, obesity rate 18.8% to 35.7%, number of hospitals with psychiatric emergency services/100 000 population 0 to 5.9, number of EDs/100 000 population 0 to 13.1, number of rural health clinics/100 000 population 0 to 32.9, number of urgent care centers/100 000 population 0 to 13.1, number of FQHCs/100 000 population 0 to 64.1, and number of community mental health centers/100 000 population 0 to 3.6.

Overall, the mean number of ED visits were 2 with an IQR of 0-2. Nearly, 10% of the study population had 6 or more ED

Table 1. Description of the Study Population by Patient-Level Factors Adult Fee-for-Service Medicaid Beneficiaries Multistate Medicaid Analytical eXtract Files-2010.^a

	N	%
All	68 882	100.0
Predisposing factors		
Age		
22-34 years	16 124	23.4
35-44 years	12 708	18.4
45-54 years	20 182	29.3
55-64 years	19 868	28.8
Gender		
Female	38 694	56.2
Male	30 188	43.8
Race		
White	58 760	85.3
African Americans	9023	13.1
Hispanics	532	0.77
Other races	567	0.82
Enabling factors		
Medicaid cash eligibility		
Cash eligibility	56 074	81.4
No cash eligibility	12 808	18.6
Primary care use		
None	13 969	20.3
Fragmented	47 582	69.1
Continuous	7331	10.6
Need factors		
Complex chronic illness		
Physical health conditions	25 128	36.5
Mental health conditions	6994	10.2
Physical and mental health conditions	25 867	37.5
None	10 893	15.8
Polypharmacy		
Yes	12 469	18.1
No	56 413	81.9
Medicaid medical eligibility		
Medical eligibility	7064	10.3
No medical eligibility	61 818	89.7
Personal health practices		
Tobacco Use		
Yes tobacco use	4558	6.6
No tobacco use	64 324	93.4
County-level external environment factors		
Metro		
Metro	41 964	60.9
Non-metro	26 918	39.1

^aBased on 68 882 adult Medicaid fee-for-service beneficiaries aged 22 to 64 years and who are continuously enrolled for the year 2010, who are not Medicare and Medicaid eligible, who are alive and non-pregnant. County-level variables were extracted from the Area Health Resource Files and county health ranking data.

visits. The frequencies of ED visits for each subgroup are presented as well. The unadjusted NBRs revealed that many subgroups of the population had higher number of ED visits: adults in the age-group 35 to 44 (IRR = 1.25) and 45 to 54 (IRR = 1.22) years; African Americans (IRR = 1.46); Medicaid eligibility due to poverty/cash assistance (IRR = 1.35); college education rate (IRR = 1.02); fragmented primary care use

Table 2. Incidence Rate Ratios and 95% Confidence Intervals (CIs) From Multivariable Negative Binomial Regression of Adult Fee-for-Service Medicaid Beneficiaries Multistate Medicaid Analytical eXtract Files-2010.^a

	IRR	95% CI	Sig
Predisposing factors			
Age			
22-34 years		Ref	
35-44 years	0.96	0.92-1.01	
45-54 years	0.78	0.72-0.84	^b
55-64 years	0.66	0.61-0.71	^b
Gender			
Female	0.95	0.91-0.99	^c
Male		Ref	
Race			
Whites		Ref	
African American	1.47	1.23-1.76	^b
Hispanics	1.63	1.16-2.31	^d
Others	1.36	0.97-1.92	
Enabling factors			
Medicaid cash eligibility			
Cash eligibility	1.07	0.99-1.16	
No Cash eligibility		Ref	
County-level education			
Percent with college education	1.00	0.99-1.02	
County-level unemployment			
Percent unemployed	0.94	0.89-0.98	^d
Primary care use			
None	0.92	0.86-0.97	^d
Fragmented	1.26	1.18-1.35	^b
Continuous		Ref	
Need factors			
Complex chronic illness			
Physical health conditions	2.12	1.88-2.56	^b
Mental health conditions	1.53	1.39-1.69	^b
Both physical and mental health conditions	3.33	2.96-3.75	^b
None		Ref	
Polypharmacy			
Yes	1.89	1.80-1.99	^b
No		Ref	
Medicaid medical eligibility			
Medical eligibility	1.29	1.14-1.46	^b
No medical eligibility		Ref	
Personal health practices			
Tobacco use			
Yes tobacco use	2.23	2.09-2.40	^b
No tobacco use		Ref	
County-level obesity			
Obesity rate	0.96	0.93-1.00	
External environment factors			
Health professional shortage area			
No shortage	1.04	0.88-1.23	
Part county shortage	1.04	0.88-1.23	
Whole county shortage		Ref	
Metro			
Metro		Ref	
Nonmetro	0.92	0.82-1.03	
Emergency departments			
Number of ED/100 000 population	1.03	1.00-1.06	^c
Psychiatric emergency services			
Number of psychiatric ED/100 000 population	1.09	0.97-1.21	

(continued)

Table 2. (continued)

	IRR	95% CI	Sig
Rural health centers			
Number of rural health centers/100 000 population	1.00	0.99-1.02	
Urgent care centers			
Number of urgent care centers/100 000 population	0.96	0.92-0.99	^d
Federally qualified health centers			
Number of FQHCs/100 000 population	1.00	0.99-1.01	
Community Mental Health Centers			
Number of community mental health centers/100 000 population	0.97	0.84-1.12	

Abbreviations: ED: Emergency Department; FQHC: Federally Qualified Health Centers; Ref: Reference Group; Sig: Significance

^aBased on 68 882 adult Medicaid fee-for-service beneficiaries aged 22 to 64 years and who are continuously enrolled for the year 2010, who are not Medicare and Medicaid eligible, who are alive and non-pregnant. County-level variables were extracted from the Area Health Resource Files and county health ranking data. Significant subgroup differences in number of ED visits were tested with negative binomial regression, which adjusted for clustering of individuals within counties.

^b $P < .001$.

^c $.01 < P < .05$.

^d $.001 < P < .01$.

(IRR = 1.55); polypharmacy (IRR = 2.10); presence of physical health conditions (IRR = 2.46); presence of mental health conditions (IRR = 1.69); presence of both physical and mental health conditions (IRR = 4.23); Medicaid eligibility due to medical needs (IRR = 1.35); tobacco use (IRR = 2.54); and number of hospitals with psychiatric emergency services (IRR = 1.21). Some subgroups of the population had lower number of ED visits: female (IRR = 0.08); obesity rate (IRR = 0.95); nonmetro status (IRR = 0.81); number of rural health centers (IRR = 0.99); and number of FQHCs (IRR = 0.99). No associations were observed between ED visits and the following factors: those aged 55 to 64 years old; unemployment rate; no primary care use; no/partial county HPSA; number of hospitals with EDs; number of urgent care centers; and number of community mental health centers.

The IRRs and 95% confidence intervals from the adjusted NBR are summarized in Table 2. The relationship between ED visits and the following factors remained the same as observed in the unadjusted NBR models: females, African Americans, no medical eligibility, fragmented primary care use, polypharmacy, and presence of complex chronic illness. For example, those with polypharmacy had higher number of ED visits (IRR = 1.89) when compared to those individuals without polypharmacy. However, the association between ED visits and the following factors changed: age-group 45 to 54 (IRR = 0.78), and 55 to 64 (IRR = 0.66); unemployment rate (IRR = 0.94); no primary care use (IRR = 0.92); and number of urgent care centers (IRR = 0.96). No associations were observed between ED visits and the following factors: 35 to 44 years old, Medicaid eligibility due to poverty, college education rate, obesity rate, no/partial HPSA, nonmetro status of the county, number of hospitals with psychiatric emergency services, number of

rural health centers, number of FQHCs, and number of community mental health centers.

Discussion

The study findings revealed that a number of patient-level and few county-level factors were associated with the ED visits among adult FFS Medicaid beneficiaries. Among predisposing factors, the current study observed that African Americans and Hispanics had a higher number of ED visits when compared to whites. The findings on racial/ethnic disparities are consistent with the published literature on ED usage.^{9,21}

Among the enabling factors, county-level unemployment rate and primary care use were associated with the number of ED visits. A counterintuitive finding of this study is the inverse relationship between county-level unemployment rate and number of ED visits. The reasons behind lower number of ED visits by individuals living in counties with high unemployment rate are not known.

Findings from the current study support the published literature in which primary care use is inversely related to ED visits.²² Medicaid beneficiaries with fragmented primary care use had higher number of ED visits when compared to those with primary care continuity. These findings suggest that timely and continuous primary care in outpatient health-care settings can reduce the frequency of ED visits among adult FFS Medicaid beneficiaries. When primary care use is fragmented, the patient may choose to use ED for nonemergency services. It has been documented that individuals without adequate primary care may delay receiving appropriate care,²³ which may in turn lead to increased use of ED.

All the need factors were associated with higher number of ED visits. Adult FFS Medicaid beneficiaries with complex needs had a higher number of ED visits. This finding is consistent with prior studies in which ED users were found to have high medical needs,²⁴⁻²⁹ have chronic conditions,³⁰ suffer from mental illnesses,³¹⁻³⁸ and have greater number of psychotropic medications.³⁹ A plausible explanation for higher ED visits by those with chronic complex illness may be due to complications of chronic conditions, side effects, and adverse events due to multiple medications use,⁴⁰ fragmented care because of visits to multiple providers, and/or direct contraindications to therapy for one condition by other conditions themselves.⁴¹

Among personal health practices, individuals with tobacco use had a higher number of ED visits when compared to those without tobacco use. Future research may examine whether EDs can serve as appropriate health-care settings for the provision of smoking cessation interventions. Indeed, the results from a recently published randomized controlled trial revealed that a smoking cessation intervention offered in an ED declined smoking rates among low-income smokers significantly.⁴² It is plausible that ED users have limited access to other health-care providers who can encourage them to quit tobacco use.

Some county-level external environment factors were associated with ED visits among adult FFS Medicaid beneficiaries. For example, this study noted that with higher number of urgent

care centers significantly lower number of ED visits were observed. This finding is consistent with another study by Weinick et al that observed about 13.1% to 27.1% of ED visits can be prevented with increased use of urgent care centers and other health-care settings.⁴³ Studies that have compared urgent care and ED care have found that costs of care in urgent care centers are lower when compared to EDs.^{3,4}

Limitations

Considering the geographic population, policy, and resource differences typically seen across states, the results of this study represent only Maryland, Ohio, and West Virginia and not generalizable to the entire Medicaid population. As Medicaid beneficiaries enrolled under managed care plans were excluded from the analytical cohort, the study suffers selection bias. The study was conducted using observational data; therefore, it is difficult to account for selection bias, as ED users may have different attributes in unobserved variables compared to the non-ED users. The cross-sectional design of the study does not allow causal inferences. As all the independent variables were measured in the same as year as the outcome variable, temporal relationships cannot be established. Substance abuse leads to increased ED use; however, the current study could not account for alcohol use and drug abuse because of limited sample size. Additionally, administrative claims data were used, which are created for billing purposes rather than research. This may result in misclassification of diagnosis.

Conclusion

To conclude, the study is timely because many states have surveillance and other research projects to monitor the use of ED by Medicaid beneficiaries and are exploring policies and programs that can reduce preventable ED visits.⁴⁴ The current study highlighted that only very few county-level factors and many patient-level factors were associated with ED visits. These findings suggest that health-care delivery models that provide comprehensive care to complex patients may reduce the likelihood of ED visits. Implementing value-based insurance designs that provide financial incentives to promote primary care continuity may go a long way in reducing ED visits among adult FFS Medicaid beneficiaries. Our study also indicates urgent care centers may be a viable substitute for EDs in providing care for acute conditions and exacerbations of chronic conditions. Future studies need to examine whether the emerging healthcare delivery models such as medical homes, and accountable care organizations (ACO), which are specifically designed to take care of complex patients, can reduce the number of ED visits by the complex patients. Medicaid ACO demonstration projects, for example, may provide evidence about whether the new delivery models can reduce the use of ED as well as the relationship between individual-level unemployment, poverty, and ED visits.

This study provides important baseline information on Medicaid ED usage and the county- and individual-level factors

driving the number of visits. As data become available, it is expected that many studies will attempt to compare and contrast ED utilization among different states who chose various policy options related to Medicaid and the Health Insurance Marketplace. Our model presents a starting point and baseline data from which to begin that exploration.

Authors' Note

Data acquisition was made possible by funding from Claude Worthington Benedum Foundation. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or policy or position of West Virginia University (WVU) or any other affiliated organizations.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: SM and US were partially supported by the National Institute of General Medical Sciences, U54GM104942.

References

1. CDC. *Emergency Department Visits*. Atlanta, GA: CDC; 2014. Web site. <http://www.cdc.gov/nchs/fastats/emergency-department.htm>. Accessed May 2, 2016.
2. Uscher-Pines L, Pines J, Kellermann A, Gillen E, Mehrotra A. Emergency department visits for nonurgent conditions: systematic literature review. *Am J Manag Care*. 2013;19(1):47-59.
3. Thygeson M, Van Vorst KA, Maciosek MV, Solberg L. Use and costs of care in retail clinics versus traditional care sites. *Health Aff (Millwood)*. 2008;27(5):1283-1292.
4. Mehrotra A, Liu H, Adams JL, et al. Comparing costs and quality of care at retail clinics with that of other medical settings for 3 common illnesses. *Ann Intern Med*. 2009;151(5):321-328.
5. Network for Excellence in Health Innovation. A Matter of Urgency: Reducing Emergency Department Overuse. 2010. Web site. http://www.nehi.net/writable/publication_files/file/nehi_ed_overuse_issue_brief_032610final edits.pdf. Accessed May 4, 2016.
6. Pitts SR, Carrier ER, Rich EC, Kellermann AL. Where Americans get acute care: increasingly, it's not at their doctor's office. *Health Aff (Millwood)*. 2010;29(9):1620-1629.
7. Mortensen K, Song PH. Minding the gap: a decomposition of emergency department use by Medicaid enrollees and the uninsured. *Medical Care*. 2008;46(10):1099-1107.
8. AHRQ. *Emergency Room Services-Mean and Median Expenses per Person With Expense and Distribution of Expenses by Source of Payment: United States, 2012*. Medical Expenditure Panel Survey Household Component Data. 2012. Web site. http://meps.ahrq.gov/mepsweb/data_stats/tables_compendia_hh_interactive.jsp?_SERVICE=MEPSSocket0&_PROGRAM=MEPSPGM.TC.SAS&File=HCFY2012&Table=HCFY2012_PLEXP_E&VAR1=AGE&VAR2=SEX&VAR3=RACETH5C&VAR4=INSURCOV&VAR5=POVCAT12&VAR6=MSA&VAR7=REGION&VAR8=HEALTH&VARO1=4+17+44+64&VARO2=1&VARO3=1&VARO4=1&VARO5=1&VARO6=1&VARO7=1&VARO8=1&_Debug=. Accessed May 4, 2016.
9. Cunningham PJ. What accounts for differences in the use of hospital emergency departments across U.S. communities? *Health Aff (Millwood)*. 2006;25(5):w324-w336.
10. Miller S. The effect of the Massachusetts reform on health care utilization. *Inquiry*. 2012-2013;49(4):317-326.
11. Smulowitz PB, O'Malley J, Yang X, Landon BE. Increased use of the emergency department after health care reform in Massachusetts. *Ann Emerg Med*. 2014;64(2):107-115, 115 e101-e103.
12. Taubman SL, Allen HL, Wright BJ, Baicker K, Finkelstein AN. Medicaid increases emergency-department use: evidence from Oregon's Health Insurance Experiment. *Science*. 2014;343(6168):263-268.
13. Kaiser Family Foundation. Medicaid Benefits Data Collection. 2016. Web site. <http://kff.org/data-collection/medicaid-benefits/>. Accessed May 4, 2016.
14. Andersen RM. Revisiting the behavioral model and access to medical care: does it matter? *J Health Soc Behav*. 1995;36(1):1-10.
15. HRSA. *Area Health Resources Files (AHRF) National, State and County Health Resources Information Database*. Website. <http://ahrh.hrsa.gov/download.htm>. Accessed May 4, 2016.
16. Gortmaker SL, Swinburn BA, Levy D, et al. Changing the future of obesity: science, policy, and action. *Lancet*. 2011;378(9793):838-847.
17. Magill MK, Senf J. A new method for measuring continuity of care in family practice residencies. *J Fam Pract*. 1987;24(2):165-168.
18. Goldberg JF, Brooks JO III, Kurita K, et al. Depressive illness burden associated with complex polypharmacy in patients with bipolar disorder: findings from the STEP-BD. *J Clin Psychiatry*. 2009;70(2):155-162.
19. Goodman RA, Posner SF, Huang ES, Parekh AK, Koh HK. Defining and measuring chronic conditions: imperatives for research, policy, program, and practice. *Prev Chronic Dis*. 2013;10:e66.
20. Allison P. *Do We Really Need Zero-Inflated Models?* Web site. <http://statisticalhorizons.com/zero-inflated-models>. Accessed May 4, 2016.
21. Ruger JP, Richter CJ, Spitznagel EL, Lewis LM. Analysis of costs, length of stay, and utilization of emergency department services by frequent users: implications for health policy. *Acad Emerg Med*. 2004;11(12):1311-1317.
22. Gill JM, Mainous AG III, Nsereko M. The effect of continuity of care on emergency department use. *Arch Fam Med*. 2000;9(4):333-338.
23. Lukens G. State variation in health care spending and the politics of state Medicaid policy. *J Health Polit Policy Law*. 2014;39(6):1213-1251.
24. O'Brien GM, Stein MD, Zierler S, Shapiro M, O'Sullivan P, Woolard R. Use of the ED as a regular source of care: associated factors beyond lack of health insurance. *Ann Emerg Med*. 1997;30(3):286-291.
25. Sandoval E, Smith S, Walter J, et al. A comparison of frequent and infrequent visitors to an urban emergency department. *J Emerg Med*. 2010;38(2):115-121.

26. Blank FS, Li H, Henneman PL, et al. A descriptive study of heavy emergency department users at an academic emergency department reveals heavy ED users have better access to care than average users. *J Emerg Nurs*. 2005;31(2):139-144.
27. Zuckerman S, Shen YC. Characteristics of occasional and frequent emergency department users: do insurance coverage and access to care matter? *Medical Care*. 2004;42(2):176-182.
28. Friedman BW, Serrano D, Reed M, Diamond M, Lipton RB. Use of the emergency department for severe headache. A population-based study. *Headache*. 2009;49(1):21-30.
29. Cook LJ, Knight S, Junkins EP Jr, Mann NC, Dean JM, Olson LM. Repeat patients to the emergency department in a statewide database. *Acad Emerg Med*. 2004;11(3):256-263.
30. Lowe RA, Fu R, Ong ET, et al. Community characteristics affecting emergency department use by Medicaid enrollees. *Medical Care*. 2009;47(1):15-22.
31. Mandelberg JH, Kuhn RE, Kohn MA. Epidemiologic analysis of an urban, public emergency department's frequent users. *Acad Emerg Med*. 2000;7(6):637-646.
32. Sun BC, Burstin HR, Brennan TA. Predictors and outcomes of frequent emergency department users. *Acad Emerg Med*. 2003;10(4):320-328.
33. D'Onofrio G. Treatment for alcohol and other drug problems: Closing the gap. *Ann Emerg Med*. 2003;41(6):814-817.
34. Hansen GR. The drug-seeking patient in the emergency room. *Emerg Med Clin North Am*. 2005;23(2):349-365.
35. Wooden MD, Air TM, Schrader GD, Wieland B, Goldney RD. Frequent attenders with mental disorders at a general hospital emergency department. *Emerg Med Australas*. 2009;21(3):191-195.
36. Rockett IR, Putnam SL, Jia H, Smith GS. Assessing substance abuse treatment need: a statewide hospital emergency department study. *Ann Emerg Med*. 2003;41(6):802-813.
37. Coley KC, Saul MI, Seybert AL. Economic burden of not recognizing panic disorder in the emergency department. *J Emerg Med*. 2009;36(1):3-7.
38. Curran GM, Sullivan G, Williams K, Han X, Allee E, Kotrla KJ. The association of psychiatric comorbidity and use of the emergency department among persons with substance use disorders: an observational cohort study. *BMC Emerg Med*. 2008;8:17.
39. Doran KM, Raven MC, Rosenheck RA. What drives frequent emergency department use in an integrated health system? National data from the veterans health administration. *Ann Emerg Med*. 2013;62(2):151-159.
40. Hafner JW Jr., Belknap SM, Squillante MD, Bucheit KA. Adverse drug events in emergency department patients. *Ann Emerg Med*. 2002;39(3):258-267.
41. National Quality Forum. *Establishing a Measurement Framework for Multiple Chronic Conditions*. Washington, DC: National Quality Forum; 2011.
42. Bernstein SL, D'Onofrio G, Rosner J, et al. Successful Tobacco Dependence Treatment in Low-Income Emergency Department Patients: A Randomized Trial. *Ann Emerg Med*. 2015;66(2):140-147.
43. Weinick RM, Burns RM, Mehrotra A. Many emergency department visits could be managed at urgent care centers and retail clinics. *Health Aff (Millwood)*. 2010;29(9):1630-1636.
44. Mann C. *Targeting Medicaid super-utilizers to decrease costs and improve quality*. Washington, DC: Center for Medicare and Medicaid Services; 2013.

Author Biographies

Parul Agarwal, a PhD candidate at the West Virginia University (WVU) School of Pharmacy is completing her dissertation on factors and outcomes associated with emergency department utilization among Medicaid beneficiaries. She works as Graduate Research Assistant in the Health Research Center.

Thomas K. Bias, an assistant professor in the WVU School of Public Health, Department of Health Policy, Management, and Leadership, co-directs the WVU Health Research Center in a variety of projects related to health policy including insurance policy and community development and health.

Suresh Madhavan, MBA, PhD, FAPhA, is a professor and chairperson of the Department of Pharmaceutical Systems and Policy at West Virginia University (WVU). Dr Madhavan did his PhD from Purdue University and prior to that completed his BPharm and MBA degrees from India. Dr Madhavan's primary areas of research interests include health services research and health policy, with particular emphasis on improving access to and quality of health and preventive care services, and the cost-effective use of drugs in healthcare systems. In recent years, he and his graduate students have studied chronic conditions such as breast, colorectal, lung, and prostate cancers, cardiovascular disease, insomnia, diabetes, and autism looking at issues related to access, cost, affordability, and quality of health care.

Nethra Sambamoorthi, is a lead faculty in the graduate program, Master of Science in Predictive Analytics, in the School of Professional Studies at Northwestern University, Chicago. His research works cover statistical methodologies, computational methods, and innovative ways of analyzing new types of data for predictive and prescriptive purposes. He applies his talents in diverse fields, such as, health care, pharmaceutical marketing, financial services, credit cards, investment services, insurance, and education. He contributes to predictive analytics education around the world with his leadership work through Institute of Analytics (USA)(TM), a division of CRMportals Inc.

Stephanie Frisbee is an assistant professor in the Department of Health Policy, Management, and Leadership in the West Virginia University School of Public Health. Her research focuses on the factors that affect cardiovascular health in entire communities or populations. Dr. Frisbee evaluates the impact of factors such as community resources, health care resources, how communities are structured, environmental factors and exposures, and policies that affect the communities in which people live.

Usha Sambamoorthi, is a health services and outcomes researcher with extensive expertise in healthcare disparities, access, quality, outcomes, utilization, cost, and policy. Her research uses available secondary sources including healthcare surveys and Medicaid, Medicare, private insurance claims and Veterans Health Administration administrative data for. Currently, Dr Sambamoorthi's research focuses on the management of individuals with chronic physical and mental health conditions.