



# The roles of zinc and copper sensing in fungal pathogenesis

Elizabeth R Ballou and Duncan Wilson

All organisms must secure essential trace nutrients, including iron, zinc, manganese and copper for survival and proliferation. However, these very nutrients are also highly toxic if present at elevated levels. Mammalian immunity has harnessed both the essentiality and toxicity of micronutrients to defend against microbial invasion — processes known collectively as ‘nutritional immunity’. Therefore, pathogenic microbes must possess highly effective micronutrient assimilation and detoxification mechanisms to survive and proliferate within the infected host. In this review we compare and contrast the micronutrient homeostatic mechanisms of *Cryptococcus* and *Candida* — yeasts which, despite ancient evolutionary divergence, account for over a million life-threatening infections per year. We focus on two emerging arenas within the host–pathogen battle for essential trace metals: adaptive responses to zinc limitation and copper availability.

## Address

Aberdeen Fungal Group, MRC Centre for Medical Mycology, School of Medicine, Medical Sciences and Nutrition, University of Aberdeen, Institute of Medical Sciences, Aberdeen, UK

Corresponding author: Wilson, Duncan ([Duncan.Wilson@abdn.ac.uk](mailto:Duncan.Wilson@abdn.ac.uk))

Current Opinion in Microbiology 2016, 32:128–134

This review comes from a themed issue on **Host-microbe interactions: fungi**

Edited by **Elaine Bignell** and **Bart PHJ Thomma**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 18th June 2016

<http://dx.doi.org/10.1016/j.mib.2016.05.013>

1369-5274/© 2016 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Introduction

The concept of nutritional immunity, in terms of host-driven iron sequestration, has been appreciated for decades: our bodies maintain extremely low iron cation levels via intracellular sequestration (ferritin, haemoglobin, the hepcidin axis) and expression of extracellular high-affinity iron-binding proteins (transferrin, lactoferrin). Successful pathogens have in turn evolved effective assimilation mechanisms (high-affinity transporters, siderophores and specialised binding proteins) [1]. However, additional layers of nutritional immunity have been revealed in recent years: micronutrient restriction is not limited to

iron, with host zinc and manganese sequestration playing key roles in controlling infection [2]. Moreover, in certain host niches, pathogens can be exposed to potentially toxic levels of iron, zinc and copper. Therefore, beyond high-affinity iron uptake systems, microbial pathogens must also possess highly effective homeostatic mechanisms (both assimilation and detoxification) for other trace metals within their hosts [2].

Host nutritional immunity during fungal infection and the pathogen assimilation pathways employed in counterattack have been extensively discussed in a number of recent reviews [3–6]. In this mini-review we focus on two emerging areas: the adaptive responses of *Candida* and *Cryptococcus* species to changes in environmental zinc and copper.

## Divergent virulence factor intercalation of the zinc regulon

Zinc is the second most abundant transition metal in the human body but, like iron, its availability to microbial pathogens is strictly limited. Zinc restriction is further compounded during periods of inflammation due to the action of calprotectin. Calprotectin constitutes approximately half the cytoplasmic protein content of neutrophils, is a dominant component of NETs (*n*eutrophil *e*xtracellular *t*raps) and elicits antifungal activity via zinc chelation [7]. Calprotectin is a heterodimer of S100A8 and S100A9, members of the S100 family of low molecular weight proteins. Antifungal activity of another S100 family member, psoriasin (S100A7) has recently been reported to elicit antifungal activity via zinc-sequestration [8<sup>\*</sup>], suggesting that multiple members of this protein family may have similar functions. Furthermore, host cell internalisation and subcellular compartmentalisation further limit accessibility to extracellular and intracellular pathogens, respectively. For example, activated, *Histoplasma*-infected macrophages shuttle zinc to their Golgi, restricting fungal proliferation and promoting clearance [9].

Interestingly, contemporary pathogenic fungal species appear to simultaneously exhibit remarkably similar and divergent responses to changes in zinc availability. First described in the model yeast *Saccharomyces cerevisiae*, Zap1 is a zinc finger transcription factor that regulates the expression of zinc transporter-encoding genes and is essential for maintaining zinc homeostasis. Under zinc limitation, Zap1 exhibits positive auto-regulation, rapidly amplifying its own transcription and subsequent protein levels and positively regulating the expression of zinc

transporter-encoding genes, including *ZRT1*, *ZRT2* and *ZRT3*. For a detailed discussion of *S. cerevisiae* Zap1 function, readers are directed to an excellent recent review by Wilson and Bird [10].


It would appear that Zap1 orthologues function as conserved master regulators of zinc homeostasis in fungi: in the pathogenic fungal species *Candida albicans*, *Candida dubliniensis*, *Aspergillus fumigatus*, and *Cryptococcus deuterogattii*, Zap1 orthologues (also known as Csr1 in *Candida*, ZafA in *Aspergillus*) have been directly demonstrated to regulate the expression of zinc transporter-encoding genes. Given the evolutionary divergence of these species, it is plausible that orthologues of Zap1 regulate zinc homeostasis in many (if not all) fungi [11–14]. Underscoring the importance of fungal zinc homeostasis during infection, each fungal pathogen tested to date exhibits virulence attenuation or decreased infectivity upon deletion of their respective *ZAP1* orthologue in relevant animal models of infection (Figure 1).

Despite this apparent conservation of function as a regulator of zinc homeostasis and virulence, there may be differences in the Zap1-regulons of different contemporary fungal species. *Cryptococcus neoformans* and *C. deuterogattii* are closely related species that express three major virulence factors for infectivity: capsule formation, melanin production and growth capacity at 37 °C. Interestingly, while *C. deuterogattii* *ZAP1* is dispensable for capsule and melanin production [14], *C. neoformans* cells lacking the *ZAP1* orthologue (*ZAP104*) are deficient in capsule, melanin, and  $\alpha/\alpha$  cell fusion, suggesting rewiring [15,16\*]. While we cannot rule out that *C. deuterogattii*

Zap1 and zinc homeostasis may influence melanin and capsule formation under conditions not examined by Schneider *et al.* [14], Liu *et al.* and Jung *et al.* independently demonstrate *ZAP104* involvement in virulence factor expression (melanin and capsule synthesis) in *C. neoformans* [14,15,16\*]. These three studies together suggest that Zap1-mediated zinc homeostasis in *C. neoformans*, but not *C. deuterogattii*, may impinge on virulence factor (capsule/melanin) expression, and raise questions about the reasons for phenotypic differences. An examination of Zap1 and Zap104 protein sequences immediately reveals structural differences: where the Zap1 N-terminus encodes a single zinc-binding domain, Zap104 encodes two [14]. Whether these structural differences lead to differing sensitivities to zinc availability or differences in promoter binding remain to be explored.

In the *Candida* genus, Zap1 function has been investigated in three relatively closely related species [17,18]: *C. albicans*, *C. dubliniensis* and *C. tropicalis*. Despite diverging from basidiomycetes approximately 500 mya [19], *C. albicans*, like *C. neoformans*, appears to regulate major virulence attributes (hyphal morphogenesis, adhesion and biofilm maturation) via Zap1 (note that Zap1 in *Candida* is also known by the common name Csr1). Kim *et al.* [20] first reported that *ZAP1* deletion in *C. albicans* precluded hyphal morphogenesis under a range of environmental conditions. Moreover, overexpression of *ZRT1* or *ZRT2* (encoding the two predicted plasma membrane zinc transporters of *C. albicans*) promoted filamentous growth of the *zap1Δ* mutant on certain media, suggesting that the morphogenic defect may be due to perturbed

Figure 1



Phylogenetic and functional relationships of Zap1 orthologues in fungi. The predicted amino acid sequences of Zap1 orthologues were downloaded from FungiDB, *Aspergillus* Genome Database, *Saccharomyces* Genome Database and *Candida* Genome Database and aligned using Phylogeny.fr. Note that in all reported cases, Zap1 regulates the expression of zinc transporters and is required for virulence or infectivity in animal models of infection, but that confirmed pathogenicity factor expression is species-specific.

cellular zinc homeostasis rather than direct Zap1-regulation of hypha-formation.

A recent study of *C. dubliniensis* found that, in this close relative of *C. albicans*, Zap1 also positively regulates *ZRT1*, *ZRT2* and *PRA1* [12<sup>\*</sup>], reinforcing the concept that Zap1 is the universal regulator of zinc assimilation in the fungal kingdom (Figure 1). However, unlike *C. albicans*, the *C. dubliniensis zap1Δ* mutant was not defective for hypha formation. On the other hand, Zap1 does regulate filamentation in *C. tropicalis* [21].

Using a powerful combination of large-scale mutant library screening, transcriptional profiling, chromatin immunoprecipitation, regulatory network analysis and infection modelling, the group of Aaron Mitchell found that Zap1 regulates extracellular matrix production by *C. albicans* biofilms and governs the expression of adhesion molecules. The *zap1Δ* mutant efficiently formed biofilms; however, these structures produced highly elevated levels of β-glucan, indicating that Zap1 negatively regulates extracellular matrix production — a key attribute in biofilm maturation [11]. Zap1 occupies *ZRT1*, *ZRT2* and *PRA1* promoters and positively regulates their transcription, in agreement with Zap1 acting as the master regulator of zinc homeostasis in *C. albicans*. However, overexpression of these zinc assimilation factors in the *zap1Δ* mutant had no effect on β-glucan production, suggesting that cellular zinc homeostasis *per se* may not directly regulate extracellular matrix synthesis, and that Zap1 possesses additional regulatory function [11]. Indeed, Zap1 was also found to be required for the regulation of genes involved in adhesion — another key element of biofilm formation [22]. Despite this regulatory role, the *zap1Δ* mutant itself exhibited wild type levels of adhesion. Therefore, to test whether Zap1 functions redundantly in adherence regulation, these authors tested the effect of *ZAP1* overexpression in strains lacking other transcription factors important for adhesion. This approach demonstrated that *ZAP1* overexpression restored the adhesive capacity of multiple adhesion-defective transcription factor mutants. This was most likely due to Zap1-dependent transcriptional induction of CSTAR (cell surface targets of adherence regulators) genes, as *ZAP1* overexpression was found to increase the expression of these target genes in mutants lacking three direct regulators of adhesion (*zcf28Δ*, *try2Δ* and *try3Δ*). Therefore, in addition to zinc assimilation, *C. albicans* appears to have intercalated key virulence attributes — adhesion and biofilm maturation — into its Zap1 regulon. It should be noted that, in these studies, zinc limited media was not used, and Zap1 may regulate different subsets of genes, depending on the environmental zinc status.

In summary, it would appear that beyond zinc homeostasis, the Zap1 transcription factor plays different roles in different fungal species and, at least in *C. neoformans* and


*C. albicans*, these are directly associated with the expression of ‘classical’ pathogenicity factors. So why might fungal pathogens place control of key virulence attributes within their Zap1 regulon? One likely possibility is the relative zinc levels found in the natural reservoirs of these yeasts, compared to the infected host. Both arboreal environments and the mammalian digestive tract are relatively high zinc environments, whilst nutritional immunity creates extreme zinc depletion within infected tissues. Hardwiring virulence factor expression into their zinc starvation-responses may contribute to the high pathogenic potential of these species, similar to the situation for iron [23].

### Dynamic regulation of response to nutritional copper immunity

In contrast to zinc nutritional immunity, which functions primarily via micronutrient sequestration by the host, copper nutritional immunity appears to represent a highly dynamic system [3,23]. During *C. albicans* bloodstream infection, serum copper levels rise early in infection [24]. Upon kidney colonisation, an early spike in copper levels is followed by rapid copper sequestration. For *C. neoformans*, while the lungs are a high copper environment, alveolar macrophages induce host *CTR1* but repress the *ATP7A* transporter for phagosomal copper compartmentalization, resulting in copper starvation upon engulfment [25]. Consistent with this, *C. neoformans* requires the copper transporter *CTR4* for survival within alveolar macrophages but it is dispensable for growth in the lungs [26<sup>\*\*</sup>,27]. Although cerebral spinal fluid (CSF) copper levels are estimated at 100 μM under certain conditions, the CSF is generally copper poor and induces copper transporter expression [27,28]. Likewise, upon crossing the blood–brain-barrier, the low copper availability of the brain necessitates *CTR1* or *CTR4* for *C. neoformans* virulence [27]. Responding to this dynamic environment requires a coordinated transcriptional response on the part of the fungus.

Structurally, fungal copper responsive transcription factors fall into three classes typified by *S. cerevisiae* Ace1 and Mac1 and *S. pombe* Cuf1 (Figure 2). All maintain an N-terminal copper responsive (R/K)GRP motif and conserved N-terminus [29<sup>\*</sup>]. In ScAce1 and ScMac1 residues 1–40 encode a zinc-binding domain. ScAce1 residues 40–110 encode CXC and CX<sub>2</sub>C motifs necessary to coordinate a tetra-copper cluster for binding MRE promoter motifs. Mac1 is characterised instead by dual cysteine rich C-terminal motifs, REP-I (CXCX<sub>4</sub>CXCX<sub>2</sub>CX<sub>2</sub>H) and REP-II (CXCX<sub>4</sub>CXCX<sub>2</sub>CX<sub>2</sub>H), which sense copper and stabilise DNA binding, respectively[30,31]. Specificity is achieved through division of labour for high and low copper responses: ScMac1 regulates the expression of the *CTR1* and *CTR2* copper importers in response to copper limitation, while copper toxicity triggers Mac1 degradation and assembly of the ScAce1 tetra-copper cluster, activating

Figure 2



Phylogenetic and structural relationships of copper responsive transcription factors in fungi. The predicted amino acid sequences of the indicated orthologs were downloaded from FungiDB, Broad, NCBI, *Saccharomyces* Genome Database, Pombase, and *Candida* Genome Database and aligned using Phylogeny.fr. Structural schematics were constructed based on sequence analysis and functional analysis is based on the reported literature, discussed in the text.

DNA-binding and induction of *CUP1* and *CRS5* metallothioneins and *SOD1*, which in turn further negatively regulates Mac1 [32–34]. Likewise, the *C. glabrata* Ace1 orthologue Amt1 is required for high copper and metallothionein expression and *C. glabrata* encodes a putative Mac1 homolog with overall structural similarity to ScMac1 [35,36] (Figure 2).

SpCuf1 is representative of the third class, with a single, structurally distinct REP motif: CXCX<sub>3</sub>CXCX<sub>2</sub>CX<sub>2</sub>H. SpCuf1 is required for growth on limited copper through binding CuSE motifs in the *CTR4* promoter; in the presence of high copper, Cu-REP interaction induces a conformational change masking the N-terminal nuclear localisation signal and inhibiting function [37]. *S. pombe* lacks metallothioneins, but detoxifies copper through a dual function SOD chaperone, PCCS [38].

*C. albicans* responds to both low and high copper levels via CaMac1, which induces either Cu-*SOD1* in high copper or Mn-*SOD3* in low copper, along with the *CTR1* copper importer; both CaMAC1 and *CTR1* are required for hyphal growth, and *SOD1* is required for virulence [24,39–41].

*CaACE1/CUP2* is additionally required for growth on high copper and regulates *CUP1* and *CRD2* metallothioneins [42,43]. *C. dubliniensis* does encode *MAC1* and *AMT1* homologues that presumably act in a similar fashion, however their targets remain uncharacterised (Figure 2).

*Cryptococcus* Cuf1 is required for growth on both low and high copper and is structurally distinct from SpCuf1 and ScMac1/Ace1: in addition to the Cuf1 REP motif, a second cysteine-rich repeat (CCX3CX4CXCX3CCXCCXC) is conserved in both *C. neoformans* and *C. deneoformans*, estimated to have diverged 18–24 mya [44,45]. For the outbreak strain *C. deuterogattii*, Cuf1 sequences in multiple isolates (R265, LA55, RAM5) have undergone truncations that specifically excise the copper responsive REP motifs (Figure 2), although these motifs are maintained in *C. decagattii* (IND107), *C. gattii* (EJB2, Ru294, WM276) and *C. bacillisporus* (CA1873, Ca1280). The relationship between copper and Cuf1 activity has not yet been reported in these organisms.

For both *C. neoformans* and *C. deneoformans*, growth in low copper requires *CTR1/CTR2* (*CNAG\_07701*) and *CTR4*

[25,26<sup>\*\*</sup>,46,47]. While *CTR1* is constitutively expressed, *CTR4* is specifically induced by low copper in a *CUF1*-dependent manner. During growth on high copper, *CUF1* mediates *CMT1* and *CMT2* metallothionein expression [46]. Similarly, Lin *et al.* report that *C. deneoformans* Mac1/Cuf1 is required for growth on both high and low copper [48]. This dual activity represents a rewiring in comparison to Cuf1 activity in *S. pombe*, where high copper leads to the sequestration of Cuf1 in the cytoplasm [37]. In addition to growth, copper levels directly influence *Cryptococcus* virulence through capsule and melanin via Cuf1-independent and Cuf1-dependent mechanisms. In both species, Cuf1 is required for melanin production in a low copper environment, likely because melanin synthesis requires the *LAC1* copper-dependent oxidase and a *cuf1Δ* mutant is unable to acquire sufficient copper from the environment [46,48–50]. Lac1 copper loading is mediated by the *CCC2* P-type copper transporting ATPase, *ATX1* copper chaperone, and the *CIC-A* chloride channel. However, in copper-replete conditions, *Cncuf1Δ* melanin defects persist in the absence of growth defects [16<sup>\*</sup>,47], and the *cuf1Δ* null is more readily phagocytosed [51]. Although a complete transcriptional analysis of Cuf1 and copper homeostasis during virulence factor induction has not been performed, *CTR1* and *CTR4*, but not *CUF1*, are induced by the temperature shift sufficient to induce capsule during growth in DMEM [51,52]. In *C. deneoformans*, excess copper induces filamentation and is dependent on Ccc2 activity [48]. Interestingly, a *C. albicans* *gpa2Δ/Δ* mutant is deficient for *CTR1* and *FRE7* copper importer expression and inappropriately expresses *CRD2*, suggesting a role for cAMP/PKA in copper homeostasis and hyphal growth. In *C. neoformans*, cAMP/PKA regulates melanin and capsule via integration with the pH-responsive transcription factor Rim101 and ESCRT [53,54]. *Cnr101* and ESCRT mutants *vps25* and *rim20* are sensitive to low copper, although capsule and immune evasion defects in these mutants are likely due to changes in the cell wall rather than defects in expression [51,55]. *CnCCC2* and *ATX1* are also required for growth on low iron, likely due to the interaction between copper and iron uptake [49].

In summary, *Candida* and *Cryptococcus* species experience significant shifts in zinc and copper availability upon transitions between commensal or environmental and infective stages. It would appear that certain successful fungal pathogens have hardwired virulence factor expression into their metal ion sensing machinery and may use the metal ion environment of the host as a key signal. A major challenge for the future will be to understand how these signals are integrated, and whether we can therapeutically target these pathways to treat fungal infections.

## Acknowledgements

ERB is funded by the BBSRC (BB/M014525/1). DW is supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the

Royal Society (Grant Number 102549/Z/13/Z). We additionally acknowledge the MRC and University of Aberdeen for funding (MR/N006364/1) and the Wellcome Trust Strategic Award for Medical Mycology and Fungal Immunology (097377/Z/11/Z). Finally, we acknowledge FungiDB and the *Candida* Genome Database [56,57].

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Skaar EP: **The battle for iron between bacterial pathogens and their vertebrate hosts.** *PLoS Pathog* 2010, **6**:e1000949.
  2. Hood MI, Skaar EP: **Nutritional immunity: transition metals at the pathogen–host interface.** *Nat Rev Microbiol* 2012, **10**:525–537.
  3. Potrykus J, Ballou ER, Childers DS, Brown AJ: **Conflicting interests in the pathogen–host tug of war: fungal micronutrient scavenging versus mammalian nutritional immunity.** *PLoS Pathog* 2014, **10**:e1003910.
  4. Subramanian Vignesh K, Landero Figueroa JA, Porollo A, Caruso JA, Deepe GS Jr: **Zinc sequestration: arming phagocyte defense against fungal attack.** *PLoS Pathog* 2013, **9**:e1003815.
  5. Wilson D, Citiulo F, Hube B: **Zinc exploitation by pathogenic fungi.** *PLoS Pathog* 2012, **8**:e1003034.
  6. Crawford A, Wilson D: **Essential metals at the host–pathogen interface: nutritional immunity and micronutrient assimilation by human fungal pathogens.** *FEMS Yeast Res* 2015:15.
  7. Urban CF, Ermert D, Schmid M, Abu-Abad U, Goosmann C *et al.*: **Neutrophil extracellular traps contain calprotectin, a cytosolic protein complex involved in host defense against *Candida albicans*.** *PLoS Pathog* 2009, **5**:e1000639.
  8. Hein KZ, Takahashi H, Tsumori T, Yasui Y, Nanjoh Y *et al.*: **Disulphide-reduced psoriasis is a human apoptosis-inducing broad-spectrum fungicide.** *Proc Natl Acad Sci U S A* 2015, **112**:13039–13044.
  - Hein *et al.* found that another S100A7 protein family member, psoriasis, also elicits antifungal activity via zin sequestration, suggesting that multiple members of this large protein family may contribute to nutritional immunity.
  9. Subramanian Vignesh K, Landero Figueroa JA, Porollo A, Caruso JA, Deepe GS Jr: **Granulocyte macrophage-colony stimulating factor induced Zn sequestration enhances macrophage superoxide and limits intracellular pathogen survival.** *Immunity* 2013, **39**:697–710.
  10. Wilson S, Bird AJ: **Zinc sensing and regulation in yeast model systems.** *Arch Biochem Biophys* 2016.
  11. Nobile CJ, Nett JE, Hernday AD, Homann OR, Deneault JS *et al.*: **Biofilm matrix regulation by *Candida albicans* Zap1.** *PLoS Biol* 2009, **7**:e1000133.
  12. Bottcher B, Palige K, Jacobsen ID, Hube B, Brunke S: **Csr1/Zap1 maintains zinc homeostasis and influences virulence in *Candida dubliniensis* but is not coupled to morphogenesis.** *Eukaryot Cell* 2015, **14**:661–670.
  - Bottcher *et al.* reported that, in stark contrast to its close relative, *C. albicans*, the Zap1 transcription factor of *C. dubliniensis* does not regulate morphogenesis. This indicates that Zap1 governs different pathogenic properties in very closely related species.
  13. Moreno MA, Ibrahim-Granet O, Vicentefranqueira R, Amich J, Ave P *et al.*: **The regulation of zinc homeostasis by the ZafA transcriptional activator is essential for *Aspergillus fumigatus* virulence.** *Mol Microbiol* 2007, **64**:1182–1197.
  14. Schneider Rde O, Fogaca Nde S, Kmetzsch L, Schrank A, Vainstein MH *et al.*: **Zap1 regulates zinc homeostasis and modulates virulence in *Cryptococcus gattii*.** *PLoS One* 2012, **7**:e43773.
  15. Liu OW, Chun CD, Chow ED, Chen C, Madhani HD *et al.*: **Systematic genetic analysis of virulence in the human fungal pathogen *Cryptococcus neoformans*.** *Cell* 2008, **135**:174–188.

16. Jung KW, Yang DH, Maeng S, Lee KT, So YS *et al.*: **Systematic functional profiling of transcription factor networks in *Cryptococcus neoformans***. *Nat Commun* 2015, **6**:6757.  
 Jung *et al.* performed a comprehensive mutant analysis for 155 predicted *C. neoformans* zinc-finger transcription factors mutants, reporting the first phenotypic analysis for 85% of these ORFs across 32 conditions, including cell wall, temperature, and chemical stresses, virulence factor regulation, and impact on virulence in the *Galleria mellonella* and murine models of infection.
17. Butler G, Rasmussen MD, Lin MF, Santos MA, Sakthikumar S *et al.*: **Evolution of pathogenicity and sexual reproduction in eight *Candida* genomes**. *Nature* 2009, **459**:657-662.
18. Jackson AP, Gamble JA, Yeomans T, Moran GP, Saunders D *et al.*: **Comparative genomics of the fungal pathogens *Candida dubliniensis* and *Candida albicans***. *Genome Res* 2009, **19**:2231-2244.
19. Taylor JW, Berbee ML: **Dating divergences in the fungal tree of life: review and new analyses**. *Mycologia* 2006, **98**:838-849.
20. Kim MJ, Kil M, Jung JH, Kim J: **Roles of zinc-responsive transcription factor Csr1 in filamentous growth of the pathogenic yeast *Candida albicans***. *J Microbiol Biotechnol* 2008, **18**:242-247.
21. Zhang Q, Tao L, Guan G, Yue H, Liang W *et al.*: **Regulation of filamentation in the human fungal pathogen *Candida tropicalis***. *Mol Microbiol* 2016, **99**:528-545.
22. Finkel JS, Xu W, Huang D, Hill EM, Desai JV *et al.*: **Portrait of *Candida albicans* adherence regulators**. *PLoS Pathog* 2012, **8**:e1002525.
23. Kronstad JW, Hu G, Jung WH: **An encapsulation of iron homeostasis and virulence in *Cryptococcus neoformans***. *Trends Microbiol* 2013, **21**:457-465.
24. Samanovic MI, Ding C, Thiele DJ, Darwin KH: **Copper in microbial pathogenesis: meddling with the metal**. *Cell Host Microbe* 2012, **11**:106-115.
25. Li CX, Gleason JE, Zhang SX, Bruno VM, Cormack BP *et al.*: ***Candida albicans* adapts to host copper during infection by swapping metal cofactors for superoxide dismutase**. *Proc Natl Acad Sci U S A* 2015, **112**:E5336-E5342.
26. Ding C, Festa RA, Chen Y-L, Espart A, Palacios O *et al.*: ***Cryptococcus neoformans* copper detoxification machinery is critical for fungal virulence**. *Cell Host Microbe* 2013, **13**:265-276.  
*C. albicans* can defend itself against the oxidative burst of innate immunity in the absence of zinc by utilising a 'copper-only' superoxide dismutase, Sod5.
27. Waterman SR, Park YD, Raja M, Qiu J, Hammoud DA *et al.*: **Role of *CTR4* in the virulence of *Cryptococcus neoformans***. *mBio* 2012:3.
28. Sun TS, Ju X, Gao HL, Wang T, Thiele DJ *et al.*: **Reciprocal functions of *Cryptococcus neoformans* copper homeostasis machinery during pulmonary infection and meningoencephalitis**. *Nat Commun* 2014, **5**:5550.
29. Waggoner DJ, Drisaldi B, Bartnikas TB, Casareno RL, Prohaska JR *et al.*: **Brain copper content and cuproenzyme activity do not vary with prion protein expression level**. *J Biol Chem* 2000, **275**:7455-7458.  
 Sun and colleagues demonstrate *in vivo* the requirement for a dynamic response to changing copper availability during *C. neoformans* infection and carefully dissect conflicting data regarding Ctr1/4 function. They show that resistance to copper toxicity through deletion of *CTR4* leads to increased proliferation in the lung, independent of Cmt1/Cmt2 activity, and that Ctr activity is required for full virulence in the brain.
30. Beaudoin J, Labbe S: **The fission yeast copper-sensing transcription factor Cuf1 regulates the copper transporter gene expression through an Ace1/Amt1-like recognition sequence**. *J Biol Chem* 2001, **276**:15472-15480.
31. Keller G, Gross C, Kelleher M, Winge DR: **Functional independence of the two cysteine-rich activation domains in the yeast Mac1 transcription factor**. *J Biol Chem* 2000, **275**:29193-29199.
32. Voutsina A, Fragiadakis GS, Boutla A, Alexandraki D: **The second cysteine-rich domain of Mac1 is a potent transactivator that modulates DNA binding efficiency and functionality of the protein**. *FEBS Lett* 2001, **494**:38-43.
33. Zhu Z, Labbe S, Pena MM, Thiele DJ: **Copper differentially regulates the activity and degradation of yeast Mac1 transcription factor**. *J Biol Chem* 1998, **273**:1277-1280.
34. Winge D: **Copper-regulatory domain involved in gene expression**. *Prog Nucleic Acid Res Mol Biol* 1997, **58**:165-195.
35. Wood LK, Thiele DJ: **Transcriptional activation in yeast in response to copper deficiency involves copper-zinc superoxide dismutase**. *J Biol Chem* 2009, **284**:404-413.
36. Koch KA, Thiele DJ: **Autoactivation by a *Candida glabrata* copper metalloregulatory transcription factor requires critical minor groove interactions**. *Mol Cell Biol* 1996, **16**:724-734.
37. Thorvaldsen JL, Sewell AK, McCowen CL, Winge DR: **Regulation of metallothionein genes by the ACE1 and AMT1 transcription factors**. *J Biol Chem* 1993, **268**:12512-12518.
38. Beaudoin J, Labbe S: **Copper induces cytoplasmic retention of fission yeast transcription factor cuf1**. *Eukaryot Cell* 2006, **5**:277-292.
39. Laliberte J, Whitson LJ, Beaudoin J, Holloway SP, Hart PJ *et al.*: **The *Schizosaccharomyces pombe* Pccs protein functions in both copper trafficking and metal detoxification pathways**. *J Biol Chem* 2004, **279**:28744-28755.
40. Hwang CS, Rhie GE, Oh JH, Huh WK, Yim HS *et al.*: **Copper- and zinc-containing superoxide dismutase (Cu/ZnSOD) is required for the protection of *Candida albicans* against oxidative stresses and the expression of its full virulence**. *Microbiology* 2002, **148**:3705-3713.
41. Marvin ME, Mason RP, Cashmore AM: **The CaCTR1 gene is required for high-affinity iron uptake and is transcriptionally controlled by a copper-sensing transactivator encoded by CaMAC1**. *Microbiology* 2004, **150**:2197-2208.
42. Huang GH, Nie XY, Chen JY: **CaMac1, a *Candida albicans* copper ion-sensing transcription factor, promotes filamentous and invasive growth in *Saccharomyces cerevisiae***. *Acta Biochim Biophys Sin (Shanghai)* 2006, **38**:213-217.
43. Homann OR, Dea J, Noble SM, Johnson AD: **A phenotypic profile of the *Candida albicans* regulatory network**. *PLoS Genet* 2009, **5**:e1000783.
44. Schwartz JA, Olarte KT, Michalek JL, Jandu GS, Michel SL *et al.*: **Regulation of copper toxicity by *Candida albicans* GPA2**. *Eukaryot Cell* 2013, **12**:954-961.
45. Jiang N, Liu X, Yang J, Li Z, Pan J *et al.*: **Regulation of copper homeostasis by Cuf1 associates with its subcellular localization in the pathogenic yeast *Cryptococcus neoformans* H99**. *FEMS Yeast Res* 2011, **11**:440-448.
46. Hagen F, Khayhan K, Theelen B, Kolecka A, Polacheck I *et al.*: **Recognition of seven species in the *Cryptococcus gattii*/*Cryptococcus neoformans* species complex**. *Fungal Genet Biol* 2015, **78**:16-48.
47. Ding C, Yin J, Tovar EM, Fitzpatrick DA, Higgins DG *et al.*: **The copper regulon of the human fungal pathogen *Cryptococcus neoformans* H99**. *Mol Microbiol* 2011, **81**:1560-1576.
48. Waterman SR, Hacham M, Hu G, Zhu X, Park YD *et al.*: **Role of a CUF1/CTR4 copper regulatory axis in the virulence of *Cryptococcus neoformans***. *J Clin Invest* 2007, **117**:794-802.
49. Lin X, Huang JC, Mitchell TG, Heitman J: **Virulence attributes and hyphal growth of *C. neoformans* are quantitative traits and the MATalpha allele enhances filamentation**. *PLoS Genet* 2006, **2**:e187.
50. Walton FJ, Idnurm A, Heitman J: **Novel gene functions required for melanization of the human pathogen *Cryptococcus neoformans***. *Mol Microbiol* 2005, **57**:1381-1396.
51. Zhu X, Gibbons J, Zhang S, Williamson PR: **Copper-mediated reversal of defective laccase in a *Deltavph1* avirulent mutant**

- of *Cryptococcus neoformans*. *Mol Microbiol* 2003, **47**:1007-1014.
52. Chun CD, Madhani HD: **Ctr2 links copper homeostasis to polysaccharide capsule formation and phagocytosis inhibition in the human fungal pathogen *Cryptococcus neoformans***. *PLoS ONE* 2010:5.
53. Haynes BC, Skowrya ML, Spencer SJ, Gish SR, Williams M *et al.*: **Toward an integrated model of capsule regulation in *Cryptococcus neoformans***. *PLoS Pathog* 2011, **7**:e1002411.
54. O'Meara TR, Xu W, Selvig KM, O'Meara MJ, Mitchell AP *et al.*: **The *Cryptococcus neoformans* Rim101 transcription factor directly regulates genes required for adaptation to the host**. *Mol Cell Biol* 2014, **34**:673-684.
55. O'Meara TR, Holmer SM, Selvig K, Dietrich F, Alspaugh JA: ***Cryptococcus neoformans* Rim101 is associated with cell wall remodeling and evasion of the host immune responses**. *mBio* 2013:4.
56. Stajich JE, Harris T, Brunk BP, Brestelli J, Fischer S *et al.*: **FungiDB: an integrated functional genomics database for fungi**. *Nucleic Acids Res* 2012, **40**:D675-D681.
57. Inglis DO, Arnaud MB, Binkley J, Shah P, Skrzypek MS *et al.*: **The *Candida* Genome Database incorporates multiple *Candida* species: multispecies search and analysis tools with curated gene and protein information for *Candida albicans* and *Candida glabrata***. *Nucleic Acids Res* 2012, **40**:D667-D674.