

RESEARCH ARTICLE

How Can the Health System Retain Women in HIV Treatment for a Lifetime? A Discrete Choice Experiment in Ethiopia and Mozambique

Margaret E. Kruk^{1*}, Patricia L. Riley², Anton M. Palma³, Sweta Adhikari³, Laurence Ahoua⁴, Carlos Arnaldo⁵, Dercio F. Belo⁶, Serena Brusamento⁴, Luisa I. G. Cumba⁶, Eric J. Dziuban², Wafaa M. El-Sadr⁴, Yoseph Gutema⁴, Zelalem Habtamu⁷, Thomas Heller⁸, Aklilu Kidanu⁹, Judite Langa¹⁰, Epifanio Mahagaja⁶, Carey F. McCarthy², Zenebe Melaku⁴, Daniel Shodell¹⁰, Fatima Tsiouris⁴, Paul R. Young², Miriam Rabkin⁴

1 Department of Global Health and Population, Harvard TH Chan School of Public Health, Boston, Massachusetts, United States of America, **2** Division of Global HIV/AIDS, US Centers for Disease Control and Prevention, Atlanta, Georgia, United States of America, **3** Department of Health Policy and Management, Mailman School of Public Health, Columbia University, New York, New York, United States of America, **4** ICAP at Columbia University, Mailman School of Public Health, Columbia University, New York, New York, United States of America, **5** Center for Population and Health Research, Maputo, Mozambique, **6** Mozambique Ministry of Health, Maputo, Mozambique, **7** Oromio Regional Health Bureau, Ethiopia Ministry of Health, Addis Ababa, Ethiopia, **8** Centers for Disease Control and Prevention, Addis Ababa, Ethiopia, **9** Miz-Hasab Research Center, Addis Ababa, Ethiopia, **10** Centers for Disease Control and Prevention, Maputo, Mozambique

* mkruk@hsph.harvard.edu



OPEN ACCESS

Citation: Kruk ME, Riley PL, Palma AM, Adhikari S, Ahoua L, Arnaldo C, et al. (2016) How Can the Health System Retain Women in HIV Treatment for a Lifetime? A Discrete Choice Experiment in Ethiopia and Mozambique. PLoS ONE 11(8): e0160764. doi:10.1371/journal.pone.0160764

Editor: Jagat Kumar Roy, Banaras Hindu University, INDIA

Received: January 29, 2016

Accepted: July 25, 2016

Published: August 23, 2016

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: Data are available in a supporting information file.

Funding: Funding for this research was provided by the U.S. President's Emergency Plan for AIDS Relief (PEPFAR) through cooperative agreement 5U2GPS001537 from the U.S. Centers for Disease Control and Prevention (CDC), Division of Global HIV/AIDS (DGHA) and by the NIH's National Institute for Allergy & Infectious Diseases under award number T32AI114398. The findings and conclusions in this paper are those of the author(s) and do not necessarily represent the official position of the U.S.

Abstract

Introduction

Option B+, an approach that involves provision of antiretroviral therapy (ART) to all HIV-infected pregnant women for life, is the preferred strategy for prevention of mother to child transmission of HIV. Lifelong retention in care is essential to its success. We conducted a discrete choice experiment in Ethiopia and Mozambique to identify health system characteristics preferred by HIV-infected women to promote continuity of care.

Methods

Women living with HIV and receiving care at hospitals in Oromia Region, Ethiopia and Zambezia Province, Mozambique were shown nine choice cards and asked to select one of two hypothetical health facilities, each with six varying characteristics related to the delivery of HIV services for long term treatment. Mixed logit models were used to estimate the influence of six health service attributes on choice of clinics.

Results

2,033 women participated in the study (response rate 97.8% in Ethiopia and 94.7% in Mozambique). Among the various attributes of structure and content of lifelong ART

Centers for Disease Control and Prevention, the NIH or the Governments of Ethiopia and Mozambique.

Competing Interests: The authors have declared that no competing interests exist.

services, the most important attributes identified in both countries were respectful provider attitude and ability to obtain non-HIV health services during HIV-related visits. Availability of counseling support services was also a driver of choice. Facility type, i.e., hospital versus health center, was substantially less important.

Conclusions

Efforts to enhance retention in HIV care and treatment for pregnant women should focus on promoting respectful care by providers and integrating access to non-HIV health services in the same visit, as well as continuing to strengthen counseling.

Introduction

The international community committed to eliminating the transmission of HIV from mother to child by 2015, with a target of 90% reduction of HIV infections among children, and to maintaining the health of mothers, with a target of 50% reduction in AIDS-related maternal deaths [1]. This ambitious plan was informed by the dramatic decrease in pediatric HIV and HIV-related maternal deaths in wealthy countries through universal HIV testing of pregnant women and use of antiretroviral treatment (ART) for eligible women. In the 21 African countries comprising nearly 90% of pregnant women living with HIV in the world, only 65% of HIV-infected pregnant women received antiretrovirals for prevention of mother-to-child-transmission (PMTCT) in 2012 [2].

Option B+, a new treatment approach that provides ART to all pregnant and breastfeeding HIV-infected women irrespective of disease stage or CD4+ cell count and continues such treatment for life, has been widely adopted as a means to protect women's health and rapidly reduce vertical transmission and sexual transmission of HIV [3–5]. Because Option B+ calls for life-long ART for all pregnant and breastfeeding HIV-infected women, it markedly increases the number of women eligible for lifelong ART, posing a potential health systems challenge.

Ethiopia and Mozambique have an estimated 760,000 and 1.6 million persons living with HIV, respectively [6, 7]. Rates of HIV transmission from mother to child are high; in 2012, 9,500 children were newly infected in Ethiopia and 14,000 were infected in Mozambique (Table 1) [8, 9]. At the same time, both countries suffer from severe health worker shortages, with health worker-population ratios of 0.28 per 1,000 population in Ethiopia and 0.37 per 1,000 in Mozambique, compared to 12.2 per 1,000 in the United States [6]. In implementing Option B+, Ethiopia and Mozambique have embarked on decentralizing ART services from hospitals to lower level clinics that provide antenatal and post-natal care with the goal of scaling up the necessary services [10, 11]. However, while Option B+ has increased access to ART for HIV-infected pregnant women, findings from Malawi indicated challenges in retention of the women in care, with 17% of pregnant women lost to follow-up 6 months after initiating ART. Compared to women starting ART for their own health (i.e., women with more advanced illness) Option B+ clients were five times more likely to never return after a first visit [12]. The authors suggest a number of reasons for lower retention, including that women at earlier stages of HIV disease perceive themselves as healthy and thus not in need of treatment, reluctance to disclose HIV status to family, and lack of support and counseling in busy clinics [12, 13]. A smaller qualitative study of Option B+ clients in South Africa identified work conflicts that interfered with keeping appointments, stigma, and negative treatment by staff as barriers to retention in care [14].

Table 1. Demographic and health profiles of study countries and regions.

| National | Ethiopia | Year | Mozambique | Year |
|--|-----------------------------------|-------------------|-------------------|-------------------|
| Demographics | | | | |
| Population (thousands) | 91,729 | 2012 ¹ | 25,203 | 2012 ¹ |
| Percent urban | 17 | 2012 ¹ | 38 | 2012 ¹ |
| Literacy rate among adults aged ≥ 15 years (%) | 47 | 2011 ² | 44 | 2011 ³ |
| Health profile | | | | |
| Life expectancy (years) | 65 | 2012 ¹ | 53 | 2012 ¹ |
| HIV prevalence among adults aged 15 to 49 (%) | 1.5 | 2012 ¹ | 11.5 | 2009 ³ |
| Under 5 mortality (per 1,000 live births) | 68 | 2011 ² | 97 | 2011 ³ |
| Maternal mortality ratio (per 100,000 live births) | 420 | 2013 ¹ | 480 | 2013 ¹ |
| Health system | | | | |
| Total health expenditure per capita (international \$) | 43.7 | 2011 ¹ | 65.8 | 2012 ¹ |
| Midwives, nurses and doctors (per 1,000 population) | 0.28 | 2012 ¹ | 0.37 | 2008 ¹ |
| Antiretroviral therapy (ART)* coverage (%) | 61 | 2012 ¹ | 45 | 2010 ¹ |
| Antenatal care (ANC) [§] coverage (≥1 visit; ≥4 visits) (%) | 28; 12 | 2012 ⁴ | 89; 53 | 2012 ⁴ |
| Births attended by a skilled health worker (%) | 10 | 2011 ² | 54 | 2011 ¹ |
| Contraceptive prevalence (%) | 29 | 2011 ¹ | 12 | 2011 ¹ |
| Regional | Oromia region (excl. Addis Ababa) | | Zambézia province | |
| Demographics | | | | |
| Population (thousands) | 26,994 | 2007 ⁵ | 3,891 | 2007 ³ |
| Percent urban | 12 | 2007 ⁵ | 17 | 2007 ³ |
| Literacy rate among adults aged ≥ 15 years (%) | 39 | 2011 ² | 28 | 2011 ³ |
| Health profile | | | | |
| HIV prevalence among adults aged 15 to 49 (%) | 1.9 | 2011 ² | 12.6 | 2011 ³ |
| Under 5 mortality (per 1,000 live births) | 112 | 2011 ² | 142 | 2011 ³ |

[§] Antenatal care (ANC) is defined as attended healthcare visits during pregnancy.

* Antiretroviral therapy (ART) consists of antiretroviral drugs to suppress HIV virus and stop progression of HIV disease. Percent ART coverage is calculated as: No. of individuals enrolled in ART / No. of individuals eligible for ART * 100

Sources:

¹ World Health Organization. Global Health Observatory Data Repository: Ethiopia and Mozambique statistics summary. Geneva, Switzerland. 2013.

UNAIDS. AIDSInfo Country Profiles. 2013.

² Central Statistical Agency [Ethiopia], ICF International. Ethiopia Demographic and Health Survey 2011. Calverton, Maryland, USA: 2012.

³ Instituto Nacional de Estatística. Panorama Sócio-Demográfico de Moçambique. Maputo, Mozambique: 2013.

⁴ UNFPA. State of the World's Midwifery, 2011: Country Profiles. 2012.

⁵ Central Statistical Agency [Ethiopia]. Ethiopia Population Census: Oromiya region profile: 2007.

doi:10.1371/journal.pone.0160764.t001

These emerging concerns about retaining patients in lifelong treatment underscore the importance of identifying what HIV-infected women want from the health system and how services can best be organized to promote their retention in care and to optimize health outcomes. This is particularly urgent in resource-constrained settings where trade-offs in health system investments will inevitably be required. Discrete choice experiments (DCE) are a research tool for examining consumers' stated preferences for health services. DCEs can be used to present hypothetical health care scenarios, each with different attributes (e.g., level of health facility, cost, availability of counseling services) to respondents who are then asked to select their preferred scenario. From these data the relative importance of each attribute can be estimated. One advantage of DCEs is the ability to test services and design changes ahead of their implementation [15, 16].

We conducted two discrete choice experiments (DCEs) in Ethiopia and Mozambique to identify the preferences of HIV-infected women of childbearing age for attributes of outpatient visits for ART in the context of lifelong care. The study was conducted in Ethiopia and Mozambique given the governments' commitment to rapidly implement Option B+ and to learn whether preferences could be generalized across settings with different HIV prevalence. The findings from our study will inform policymakers in tailoring the provision of Option B+ to meet the needs of women and, in turn, increase uptake and retention in care.

Methods

Study setting and sample

Data for this study were collected using a cross-sectional survey that assessed health service utilization and preferences for HIV service delivery among HIV-infected women in the Oromia region of Ethiopia and the Zambézia province of Mozambique, two rural regions in which prior research has demonstrated that numerous well-known barriers to retention in HIV care and ANC persist [17–21]. We recruited study participants from health centers and hospitals supported by ICAP at Columbia University, an implementing partner for the President's Emergency Plan for AIDS Relief (PEPFAR) supporting HIV-related programming in both countries. Facilities selected for the study were the four ICAP-supported facilities with the greatest volume of patients on ART in their respective regions, with between 1,500 to 3,000 women actively enrolled in ART at each site in 2013 [22]. Both Mozambique and Ethiopia introduced Option B+ in study clinics in June 2013, approximately one year prior to this study.

For this study, we recruited HIV-infected women aged 15 to 49 who were receiving care in selected clinics and who were either pregnant or indicated their intention to have children in the future. Participants were recruited from ART clinics, antenatal care (ANC) clinics, and in Mozambique, from clinics providing services for children born to HIV-infected women and their mothers as well as other high-risk groups. Given the marked difference in HIV prevalence between the two countries, we expected that 50% of our Mozambican sample and 10% of our Ethiopian sample would be pregnant women, consistent with prior trends we observed from clinic census data.

Researchers selected a systematic random sample by inviting women who were exiting their clinic appointments to participate in the study at regular time intervals during clinic operating hours. All eligible women were informed of the purpose of the study and their right to refuse participation. Interviews were performed after receipt of written consent from the participant or, in the case of minors, upon receipt of assent from the participant and consent from the guardian.

The study was approved by the ethics review boards at Columbia University Medical Center, the US Centers for Disease Control and Prevention, the National Research Ethics Review Committee at the Ethiopia Ministry of Science and Technology and the National Bioethics Committee for Health at the Mozambique Ministry of Health.

DCE design

We conducted a literature review and held discussions with policymakers and clinical managers in both countries to determine an initial list of attributes for Option B+ services that were thought to be important to women and amenable to policy interventions. Where possible, similar attributes were selected to permit comparison between study findings in Mozambique and Ethiopia. Subsequently, we conducted four focus groups in each country with HIV-infected women in the reproductive age group in a subset of study clinics to narrow down the attribute list. Women were asked to choose HIV service attributes relevant to Option B+ most important

to them and to suggest others not on the list. Based on these rankings, we identified two sets of attributes and levels for the DCE, one each for Mozambique and Ethiopia. The final attributes were: type of facility (hospital, clinic, and in Mozambique, mobile clinic), provider attitude, availability of integrated non-HIV health services (e.g. blood pressure testing, newborn care and family planning), availability of mother support groups (Ethiopia only), availability of counselor support, involvement of husband or family in care (Mozambique only), and total cost of the visit (Table 2). Cost of visit was included to permit estimation of willingness to pay. Though care and treatment for HIV infection is provided free of charge in both countries, women may bear other costs, such as transportation, food, and, if purchased privately, medications.

The attributes and levels generated 160 possible alternatives in Ethiopia and 240 alternatives in Mozambique, which is too many for women to evaluate. Thus, we selected a fractional set of 16 alternatives or 8 choice sets using experimental design that maximized D-efficiency in Sawtooth Software, a software program for designing discrete choice experiments [23, 24]. To further enhance statistical efficiency and maximize study power we used five different versions of the DCE in each country. Each participant was presented with 9 choice cards, each showing two health facilities. One of the 9 cards was a “fixed choice” task that was used to assess the predictive validity of the model and not included in estimation of preference. After being read the

Table 2. Attributes and levels of health services used in discrete choice experiment.

| <i>Ethiopia</i> | | <i>Mozambique</i> | |
|-------------------------|---|----------------------------|---|
| Attribute | Level | Attribute | Level |
| Type of health facility | Hospital | Type of health facility | Hospital |
| | Health center | | Health center |
| | | | Mobile clinic |
| Provider attitude | Health providers are respectful and welcoming | Provider attitude | Health providers are respectful and pleasant |
| | Health providers are not respectful and welcoming | | Health providers are not respectful and pleasant |
| Non-HIV services | Non-HIV services available at the same consultation | Non-HIV services | Non-HIV services available at the same consultation |
| | Non-HIV services not available at the same consultation | | Non-HIV services not available at the same consultation |
| Mother support groups | Mother support groups available | Husband/family involvement | Clinic involves husband or family in care |
| | Mother support groups not available | | Clinic does not involve husband or family in care |
| Counselor support | Counselor to help you stay on treatment | Counselor support | Counselor to help you stay on treatment |
| | No counselor to help you stay on treatment | | No counselor to help you stay on treatment |
| Cost* | 0 Birr for each visit | Cost | 0 MTn for each visit |
| | 50 Birr for each visit | | 50 MTn for each visit |
| | 100 Birr for each visit | | 100 MTn for each visit |
| | 200 Birr for each visit | | 200 MTn for each visit |
| | 300 Birr for each visit | | 400 MTn for each visit |

Note. Introductory text: “In this section I will show you 9 cards. Each card describes two imaginary health facilities. These cards are not describing this facility or other facilities that you have used. Imagine that you are now pregnant and taking antiretroviral medicines (ART) to treat your HIV infection and to prevent your baby from getting HIV. You will need to take these medicines every day, throughout pregnancy and for the rest of your life. You will also need to visit a health facility ever 3 to 6 months for checkups and laboratory tests and to obtain medicines. You may need to come more frequently at the beginning. Please tell us which of the two facilities you would prefer to go to for your HIV care. Please keep in mind that you would need to return to this facility frequently so consider what services would be most important in helping you stay in treatment over many years. There are no right or wrong answers to these questions and remember we will not share your information with anyone. We are only interested in learning about what is important to you about health facilities.”

* 100 Ethiopian Birr = 5.12 USD and 100 Mozambican MTn = 3.20 USD; cost includes all expenses related to the care visit, including transportation and food.

doi:10.1371/journal.pone.0160764.t002

introductory text and the attributes for each alternative, women were asked to select their preferred health facility to obtain Option B+ services and lifelong care for HIV infection. Each choice scenario was accompanied by a standardized script, which was read by the interviewer. Sample DCE cards are shown in [S1 Fig](#).

Non-DCE variables

The survey accompanying the DCE included questions about participants' demographic and household characteristics, general and reproductive health history, HIV history, knowledge and stigma, and health system use and satisfaction with health services. We constructed a relative index of household wealth status using principal component analysis of a set of 20 questions on household assets [25].

Survey fielding

The survey and consent forms were developed in English, translated to Portuguese in Mozambique, and Afan Oromo and Amharic in Ethiopia, then back-translated to English and pre-tested to ensure accuracy. The surveys were piloted, revised, and administered by trained interviewers. The interviews lasted 45 to 60 minutes and were conducted using hand-held electronic tablets with SurveyCTO software (Dobility, Inc., Boston, United States of America). Data were collected over a period of 8 weeks (16 April to 14 June 2014 in Ethiopia, and 8 April to 6 June 2014 in Mozambique).

Statistical analysis

Data were cleaned (typographical errors corrected, variables recoded as necessary) and transferred to Stata v.12 (StataCorp LP, College Station, TX, USA). We calculated descriptive statistics for survey variables. We then fit mixed logit regression models with DCE attributes and levels as explanatory variables using Stata's `mixlogit` command to estimate relative utility of each DCE attribute. Mixed logit models are commonly used to analyze discrete choice data as they account for taste heterogeneity by allowing attribute coefficients to vary across respondents; they also control for intra-individual correlations due to repeated responses [26]. Mathematically, mixed logit differs from the standard logit model in that the researcher does not know the value of β_n or ε_{in} . For this reason, the solution of the equation requires that L_{nit} be integrated over all possible values of β weighted by the density distribution—usually a standard normal. The unconditional probability of the observed sequence of choices for a given choice set t is given by $P_{nit}(\theta) = \int L_{nit}(\beta)f(\beta|\theta)d\beta$.

Probabilities were estimated with a simulated maximum likelihood estimator. The output of a mixed logit model includes mean attribute utilities and standard deviations of the random coefficients, the latter reflecting the degree of preference heterogeneity among respondents.

We assessed the predictive validity of these base mixed logit models by comparing the model-predicted choice of health facility in the fixed choice task with the respondent's actual choice. Lastly, we fit additional models with interaction terms between each attribute and pregnancy status to explore whether pregnant respondents, respondents not on ART, and women who were potentially eligible for Option B+, and younger women (under 25) had different preferences. We defined potential Option B+ clients as women who were pregnant or breastfeeding and not on ART or were on ART only during this or past pregnancies (i.e., pregnant or breastfeeding women not on ART for their own health). In all models, as per standard practice, costs of the visit were specified as fixed to ensure ease of interpretation of willingness to pay, and all other variables were specified as random [27].

Results

Sample characteristics

Of 1,036 women eligible to participate in Ethiopia, 1,013 participated in the survey, for a response rate of 97.8%. In Mozambique, 1,020 of 1,077 eligible women participated in the survey, for a response rate of 94.7%. Demographic and health characteristics of the respondents are summarized in Table 3. The mean age was 30.5 years in Ethiopia and 25.3 years in Mozambique. Most respondents were on ART at the time of the interview (94.8% in Ethiopia and 85.7% in Mozambique). Women known to be pregnant at the time of the interview

Table 3. Characteristics of study population by country, Ethiopia and Mozambique, 2014.

| | Ethiopia (n = 1,013) | Mozambique (n = 1,020) |
|--|----------------------|------------------------|
| | n (%) | n (%) |
| Demographic | | |
| Age, mean (SD) | 30.5 (4.9) | 25.3 (5.4) |
| Respondent is head of household | 222 (24) | 165 (16.2) |
| Currently married or living with partner | 725 (72) | 754 (73.9) |
| No. of children, mean (SD) | 1.6 (1.2) | 2.5 (1.7) |
| Educational attainment: | | |
| <i>No education</i> | 235 (23) | 120 (12) |
| <i>Any primary</i> | 379 (38) | 467 (46) |
| <i>Any secondary</i> | 395 (39) | 433 (43) |
| Literate ¹ | 723 (71) | 710 (70) |
| Primary occupation as farmer or homemaker ² | 494 (49) | 712 (70) |
| Resides in village (vs. town) | 117 (12) | 640 (63) |
| Household assets | | |
| Household has electricity | 931 (92) | 581 (57) |
| Household has radio | 824 (81) | 521 (51) |
| Household has television | 713 (70) | 519 (51) |
| Household has a mobile phone | 931 (92) | 747 (73) |
| Wealth quintiles | | |
| Highest | 202 (20.0) | 202 (20.0) |
| Fourth | 201 (20.0) | 202 (20.0) |
| Middle | 203 (20.1) | 202 (20.0) |
| Second | 202 (20.0) | 202 (20.0) |
| Lowest | 203 (20.1) | 203 (20.1) |
| Pregnancy and ART status | | |
| Currently pregnant | 117 (12) | 499 (49) |
| Not on ART now ³ | 53 (5) | 146 (14) |
| Potential Option B+ client ⁴ | 55 (5) | 370 (36) |

Note: Data are no. (%) of subjects, unless otherwise indicated. For some rows, denominators differ from country totals owing to missing data.

¹ Has completed primary school or was able to read all or part of a sentence in the national language (Amharic or Affan Oromo in Ethiopia; Portuguese in Mozambique)

² Includes homemakers, farmers and house cleaners.

³ Women not on ART now include women who initiated ART at the current visit. This includes pregnant and non-pregnant women, i.e., current and future potential Option B+ clients.

⁴ Potential Option B+ clients include women who are pregnant or breastfeeding and not on ART or are on ART but have only taken ART during pregnancies. These women are not on ART for their health.

doi:10.1371/journal.pone.0160764.t003

comprised 11.5% (n = 117) of the sample from Ethiopia and 48.9% (n = 499) of the sample from Mozambique.

Among our sample, 55 (5.4%) women in Ethiopia and 370 (36.2%) women in Mozambique could be characterized as currently eligible for Option B+—i.e., they were pregnant/breastfeeding and not currently on ART or had been on ART only during pregnancies. The length of diagnosed HIV infection (calculated as the difference between the self-reported date of first HIV diagnosis and date of interview) among these women was shorter than for women on treatment for their own health (mean number of months since HIV diagnosis in Ethiopia: 51.2 vs. 64.9; Mozambique: 6.6 vs. 34.6). The majority of Option B+ clients (98.2% in Ethiopia and 100.0% in Mozambique) were recruited from antenatal clinics. Their average age was 28.0 years in Ethiopia and 24.2 years in Mozambique.

DCE utility estimates

Utility estimates for DCE attributes are shown in Table 4. All of the attribute levels were significantly associated with health facility choice at $\alpha = 0.05$. In the main effects model, five of the attributes—provider type, provider attitude, availability of non-HIV services in the same consultation, availability of counselor, availability of mother support groups (Ethiopia only), involvement of husband or family in care (Mozambique only)—had positive mean coefficients representing positive preferences for these attributes. As expected, increasing cost had an aversive effect on preference, indicated by the negative coefficient. Our models correctly predicted a high proportion of alternatives selected in the fixed choice cards: 86.7% in Ethiopia and 91.1% in Mozambique; confirming that the models included important characteristics and were thus well-specified.

The two most important attributes for women in both countries were respectful provider attitude and availability of non-HIV services (such as blood pressure measurement, newborn care and family planning) in the same consultation. To illustrate the magnitude of utility

Table 4. Results of mixed logit regression models.

| Ethiopia | | | | | Mozambique | | | | |
|---|-------------------|-----------------|-------|--------|---|-------------------|-----------------|-------|--------|
| Attribute | Mean ¹ | SE ² | SD | SE | Attribute | Mean ¹ | SE ² | SD | SE |
| Non-HIV services available at the same consultation | 2.31 | 0.12** | 2.08 | 0.12** | Non-HIV services available at the same consultation | 1.06 | 0.07** | 1.27 | 0.08** |
| Providers are respectful and welcoming | 1.78 | 0.09** | 1.51 | 0.09** | Providers are respectful and pleasant | 1.61 | 0.08** | 1.38 | 0.08** |
| Mother support groups available | 1.06 | 0.06** | 0.66 | 0.10** | Providers involve husband/family in care | 0.67 | 0.05** | 0.91 | 0.08** |
| Counseling services available | 0.94 | 0.07** | -0.57 | 0.14** | Counseling services available | 0.57 | 0.05** | 0.76 | 0.06** |
| Hospital (vs. health center) | 0.37 | 0.06** | 0.88 | 0.09** | Health center (vs. mobile clinic) | 0.16 | 0.06** | -0.25 | 0.18 |
| | | | | | Hospital (vs. mobile clinic) | 0.15 | 0.05** | 0.38 | 0.14** |
| Cost (continuous in 100 Birr) ³ | -0.46 | 0.03** | | | Cost (continuous in 100 MTn) ³ | -0.19 | 0.02** | | |
| Model diagnostics | | | | | Model diagnostics | | | | |
| Number of respondents | 1,013 | | | | Number of respondents | 1,020 | | | |
| Number of observations | 16,192 | | | | Number of observations | 16,156 | | | |
| Log-likelihood | -3565.6 | | | | Log-likelihood | -4183.7 | | | |
| Likelihood ratio χ^2 | 939.92 | | | | Likelihood ratio χ^2 | 635.79 | | | |

¹ Mean β coefficients show estimated utility of each attribute, where positive coefficients indicate positive preference.

² ** $p < .01$

³ Currency equivalents in USD are 100 Ethiopian Birr = 5.12 USD and 100 Mozambican MTn = 3.20 USD, using period average exchange rates for the dates of data collection, extracted from OANDA.com (Ethiopia: 16 Apr 2014 to 12 Jun 2014; Mozambique: 8 Apr 2014 to 23 May 2014).

gain for women associated with these attributes, good provider attitude was twice as important to women compared to availability of counseling services when choosing a health facility for HIV care in Ethiopia and three times as important as counseling in Mozambique (Ethiopia: $\beta = 1.781$ vs. 0.936 ; Mozambique: $\beta = 1.606$ vs. 0.573). Availability of non-HIV services in the same consultation was the most valued attribute in Ethiopia and was also two to three times as important as counseling services, though in different order of importance between the two countries (Ethiopia: $\beta = 2.312$ vs. 0.936 ; Mozambique: $\beta = 1.058$ vs. 0.573). By contrast the facility level was consistently the least important attribute (Ethiopia: $\beta = 0.368$ for hospital vs. health center; Mozambique: $\beta = 0.163$ and $\beta = 0.150$ for health center and hospital vs. mobile clinic, respectively). This means that women place relatively less emphasis on type of health facility where they obtain HIV care, holding all other attributes constant.

Supplementary analyses containing results from models that included interaction terms between service attributes and pregnancy status, ART status, current potential B+ clients, and young age (under 25 years) are presented in [S1–S4](#) Tables. Pregnant women in Ethiopia valued mother support groups more than did non-pregnant women, and in Mozambique they valued respectful providers more than did non-pregnant women. In Mozambique the involvement of husband or family was less important to pregnant women than to their non-pregnant counterparts. Analyses with interaction terms for ART status indicate that in Ethiopia there were no differences in preferences between women currently on ART vs. not currently on ART. In Mozambique, women on ART preferred availability of non-HIV services more than did women not on ART, and they were also more averse to increased cost. Women identified as current potential B+ clients (pregnant or breastfeeding and not on ART or on ART during pregnancies only) were compared to other women. In Ethiopia, these women valued mother support groups and receiving care at hospitals vs. health centers more than did non-B+ clients. In Mozambique, B+ clients preferred availability of non-HIV services and respectful providers more than did non-B+ clients. Lastly, analysis that included interaction terms between service attributes and young age (under 25 years) show no differences between younger women and older women (see [S1–S4](#) Tables). Analysis of dominant attributes, i.e., when respondents always selected alternatives with a specific level of one attribute, indicated that up to one-fourth of respondents had dominant preferences for the top attributes of good provider attitude and availability of non-HIV services. We elected to retain these respondents in the analysis as a dominant or non-trading response pattern may be consistent with random utility theory and deleting these responses may thus remove valid information about preferences. Removing respondents with dominant preferences can also induce selection bias, reducing generalizability to the broader population, and reduce statistical efficiency of the model [28].

Discussion

This is the first DCE to assess women's preferences for the structure and content of lifelong antiretroviral treatment in the context of Option B+ scale up in sub-Saharan Africa. We found that HIV-infected pregnant women and women desiring a future pregnancy in Ethiopia and Mozambique placed greatest value on respectful providers and the ability to obtain non-HIV services in their clinic visits for HIV treatment. To illustrate the magnitude of preference, these two service attributes were approximately twice as important to Ethiopian women as the availability of mother support groups and counseling services. In Mozambique, women valued respectful providers more than twice as much as providers who involved husband/family in care and valued the availability of non-HIV services in the same visit twice as much as obtaining counseling services.

These findings suggest that women's choice of clinics for lifelong ART in the context of the Option B+ strategy is influenced by desire for respectful care and convenient access to non-HIV services alongside HIV services—characteristics that are not specific to HIV care but rather reflect an effectively functioning, patient-centered health system. Additional analysis showed that the utility of respectful providers was even greater for pregnant women in Mozambique than non-pregnant respondents.

The high value for provider respect is consistent with DCE studies of obstetric care: women in Tanzania and Ethiopia were highly influenced by having a respectful provider in deciding between hypothetical facilities for delivery [29, 30]. Consistent with this, researchers found that harsh treatment from nurses was identified as a barrier to retention in care by women enrolled in Option B+ in South Africa [14]. Researchers in Malawi observed that provision of Option B+ in large clinics was associated with greater loss to follow-up than in smaller clinics, hypothesizing that this may have been in part due to overworked staff who had less time for clients [12].

Women in both countries in our study highly valued the ability to obtain needed non-HIV related services in their HIV clinic visit. This suggests that women would like to improve the efficiency of their encounters with the health system, a particularly trenchant concern given the need to return to clinic many times for HIV care and treatment [31]. This highlights the importance of better organizing outpatient care to permit access to more than one service and is consistent with recent PEPFAR priorities that promote integration, sustainability and coordination between countries [32]. Situating Option B+ services in maternal and child health, which is the model in Mozambique and other countries, may facilitate service integration and potentially promote retention in care. While much has been written on the benefits and costs of integration of HIV services with other health care, the finding that clients with HIV might choose their facility based on convenient access to a range of services is novel and of substantial policy relevance [33–35].

Availability of counseling, mother support groups (Ethiopia), and counseling and providers who involved the husband or family in care (Mozambique) were positively associated with clinic preference. Pregnant women in Ethiopia had a stronger preference for mother support groups than non-pregnant respondents. The value placed on peer support is consistent with the work of Assefa et al. who found that community involvement in care was an important driver of high retention in care after service decentralization in Ethiopia [36].

Pregnant respondents in Mozambique indicated a lower preference for provider efforts to involve the husband and family compared to non-pregnant women. Reluctance to disclose HIV status is likely influenced by prevailing stigma [37, 38]. However, more work is required to understand how this might differ for pregnant versus non-pregnant women and how clinicians can support disclosure in settings where stigma remains a reality. Mozambique is currently considering the introduction of peer support groups to promote adherence and reduce stigma. Our findings from Ethiopia buttress this as mother support groups were valued by respondents there.

Study participants in both countries indicated that the type of facility—hospital versus health center (or mobile clinic in Mozambique) was much less important to them than other factors. This is reassuring for policymakers as decentralization of HIV care to health centers and clinics is proceeding rapidly in both countries. The important caveat is that the indifference to service setting is conditional on receiving other valued attributes on the list: respectful care, access to non-HIV services, counseling, etc. Thus facilities selected for Option B+ services—no matter what their level—need to be well-functioning and include a basic package of non-HIV services [39].

The study had several strengths. This is the first discrete choice experiment exploring preferences for lifelong receipt of HIV treatment. The study involved a large number of HIV-

infected women and included both current and potentially future clients for Option B+. Finally, the robustness tests that compared model-predicted with actual choices of DCE facilities suggested that the experiment was well specified, capturing important drivers of preference. This study had also several limitations. First, all stated preference studies indicate respondents' preferences for hypothetical choices; validating this work with revealed preference data from a policy experiment is important to confirm the findings. The non-DCE variables were self-reported and are subject to recall bias. Second, the study was done in a limited number of facilities in Ethiopia and Mozambique and the results cannot be extrapolated to other populations. However, the concordance of preferences across the two countries, particularly the strong preference for the top two attributes, suggests our findings may be applicable to demographically and geographically comparable populations facing similar health systems challenges. Third, interviews were only conducted with women at health facilities, and the vast majority of participants were already receiving ART. Therefore we could not capture HIV care preferences of infected women who are not yet diagnosed or have discontinued care. Finally, the sample of pregnant women who met criteria for Option B+ was relatively small in Ethiopia, limiting inference.

An important implication of our findings is that health systems need to make structural changes such as improved staffing and service availability in order to enhance lifetime retention in HIV care and improve the quality of interpersonal care and access to a range of health services [40]. The emphasis on service availability and convenience is consistent with the fact that women with HIV perceive their HIV infection as one of several health needs that might require medical care.

Another central findings was the high value placed by women on respectful treatment. The demand for better interpersonal quality of care may reflect that women experience brusque and disrespectful treatment by providers in resource-constrained health systems [40, 41]. The need for effective communication and more empathetic treatment has been amply demonstrated in HIV care although the evidence on what works to ensure respectful treatment by providers is sparser [42, 43]. Research on underlying causes of disrespectful treatment and on interventions that address this is urgently needed. Potential approaches that have been documented in the literature include increased staffing—which reduces provider perceptions of feeling overburdened—thereby improving their ability to deliver care, pre-service and in-service training on respectful care, supportive supervision, task-shifting of services to other cadres to reduce workloads on clinicians, building interpersonal skills and efforts to improve facilities' accountability to their communities, such as community-facility committees and community charters [44, 45]. Our findings show that it is essential for policymakers and providers in HIV care to embrace the respectful care agenda now being promoted in other areas of health [31].

Option B+ offers an important opportunity for enhancing health outcomes for HIV-infected pregnant women and their infants. However, none of these benefits can be achieved without high rates of retention of the women in care. Our study findings indicate that in order to create people-centered health systems and to ensure that the services provided are services that people will use, policymakers must know what users value and shape the health systems in a manner to address those preferences [46, 47].

Supporting Information

S1 Dataset. B+ discrete choice experiment dataset.

(DTA)

S1 Fig. Sample DCE cards.

(DOCX)

S1 File. B+ discrete choice experiment dataset codebook.
(XLSX)

S1 Table. Results of mixed logit regression models with interaction terms with pregnancy status (pregnant vs. non-pregnant).
(DOCX)

S2 Table. Results of mixed logit regression models with interaction terms with ART status (Not currently on ART vs. on ART).
(DOCX)

S3 Table. Results of mixed logit regression models with interaction terms with current potential B+ clients (women pregnant or breastfeeding and not on ART or on ART during pregnancies only (pMTCT) vs. others).
(DOCX)

S4 Table. Results of mixed logit regression models with interaction terms with age less than 25 (<25 vs. 25+ years old).
(DOCX)

Acknowledgments

The authors are indebted to the women in Ethiopia and Mozambique who participated in the study. We wish to thank Boaventura Cau, Hawi Fita, Estevao Manhice, and Semret Zenebe for their capable supervision of the fieldwork for the study as well as the interviewers who collected the data. We appreciate the input and support of health authorities and health workers at the participating health facilities in Oromia region and Zambézia province as well as the support of national ministries of health in Ethiopia and Mozambique.

Author Contributions

Conceived and designed the experiments: MEK PLR AMP SA LA CA SB EJD TH AK JL
CFM ZM DS FT PRY MR.

Performed the experiments: CA AK AP SA MEK.

Analyzed the data: MEK AP SA MR.

Wrote the paper: MEK PLR AP SA LA CA DFB SB LIGC EJD WME YG ZH TH AK JL EM
CFM ZM DS FT PRY MR.

References

1. UNAIDS. Countdown to zero: global plan towards the elimination of new HIV infections among children by 2015 and keeping their mothers alive. Geneva: 2011.
2. World Health Organization. Global update on HIV treatment 2013: results, impact and opportunities. Geneva, Switzerland: 2013.
3. World Health Organization. PMTCT Strategic Vision 2010–2015—Preventing mother-to-child transmissions of HIV to reach the UNGASS and Millennium Development Goals. Geneva, Switzerland: WHO, 2010.
4. World Health Organization. Consolidated Guidelines on the use of Antiretroviral Drugs for Treating and Prevention of HIV Infection—Recommendations for a public health approach. Geneva, Switzerland: 2013.
5. Business Leadership Council, UNICEF, Clinton Global Health Initiative. A business case for options B and B+. New York, NY: 2012.

6. World Health Organization. Global Health Observatory Data Repository: WHO African Region: Ethiopia and Mozambique statistics summary (2002—present). Geneva, Switzerland: 2013.
7. UNAIDS. Global report: UNAIDS report on the global AIDS epidemic 2013. Geneva, Switzerland: 2013.
8. UNAIDS. Together we will end AIDS. 2012.
9. UNAIDS. UNAIDS 2013 World AIDS Day Report. Geneva, Switzerland: 2013.
10. MEASURE DHS. Mozambique Demographic and Health Survey 2011, Final Report. Calverton, MD: 2011.
11. MEASURE DHS. Ethiopia Demographic and Health Survey 2011, Final Report. Calverton, MD: 2011.
12. Tenthani L, Haas A, Tweya H, Jahn A, van Oosterhout J, Chimbandira F, et al. Retention in care under universal antiretroviral therapy for HIV-infected pregnant and breastfeeding women ('Option B+') in Malawi. *AIDS*. 2014; 28(4):589–98. doi: [10.1097/QAD.000000000000143](https://doi.org/10.1097/QAD.000000000000143) PMID: [24468999](https://pubmed.ncbi.nlm.nih.gov/24468999/)
13. Curran K, Ngure K, Shell-Duncan B, Vusha S, Mugo N, Heffron R, et al. 'If I am given antiretrovirals I will think I am nearing the grave': Kenyan HIV serodiscordant couples' attitudes regarding early initiation of antiretroviral therapy. *AIDS*. 2014; 28(2):227–33. doi: [10.1097/QAD.000000000000025](https://doi.org/10.1097/QAD.000000000000025) PMID: [24413310](https://pubmed.ncbi.nlm.nih.gov/24413310/)
14. Clouse K, Schwartz S, Van Rie A, Bassett J, Yende N, Pettifor A. "What they wanted was to give birth; nothing else": Barriers to retention in Option B+ HIV care among postpartum women in South Africa. *J Acquir Immune Defic Syndr*. 2014; 67(1):e12–8. doi: [10.1097/QAI.0000000000000263](https://doi.org/10.1097/QAI.0000000000000263) PMID: [24977376](https://pubmed.ncbi.nlm.nih.gov/24977376/).
15. Ryan M, Scott D, Reeves C, Bate A, van Teijlingen E, Russell E, et al. Eliciting public preferences for healthcare: a systematic review of techniques. *Health technology assessment (Winchester, England)*. 2001; 5(5):1–186.
16. Ryan M, Farrar S. Using conjoint analysis to elicit preferences for health care. *BMJ*. 2000; 320(7248):1530–3. PMID: [10834905](https://pubmed.ncbi.nlm.nih.gov/10834905/)
17. Audet CM, Chire YM, Vaz LM, Bechtel R, Carlson-Bremer D, Wester CW, et al. Barriers to Male Involvement in Antenatal Care in Rural Mozambique. *Qualitative health research*. 2015. Epub 2015/04/10. doi: [10.1177/1049732315580302](https://doi.org/10.1177/1049732315580302) PMID: [25854615](https://pubmed.ncbi.nlm.nih.gov/25854615/); PubMed Central PMCID: [PMC4598282](https://pubmed.ncbi.nlm.nih.gov/PMC4598282/).
18. da Silva M, Blevins M, Wester CW, Manjolo J, Jose E, Gonzalez LC, et al. Patient loss to follow-up before antiretroviral therapy initiation in rural Mozambique. *AIDS Behav*. 2015; 19(4):666–78. Epub 2014/08/07. doi: [10.1007/s10461-014-0874-0](https://doi.org/10.1007/s10461-014-0874-0) PMID: [25096897](https://pubmed.ncbi.nlm.nih.gov/25096897/).
19. Feyissa TR, Melka AS. Demand for modern family planning among married women living with HIV in western Ethiopia. *PLoS One*. 2014; 9(11):e113008. Epub 2014/11/13. doi: [10.1371/journal.pone.0113008](https://doi.org/10.1371/journal.pone.0113008) PMID: [25390620](https://pubmed.ncbi.nlm.nih.gov/25390620/); PubMed Central PMCID: [PMC4229307](https://pubmed.ncbi.nlm.nih.gov/PMC4229307/).
20. Paulin HN, Blevins M, Koethe JR, Hinton N, Vaz LM, Vergara AE, et al. HIV testing service awareness and service uptake among female heads of household in rural Mozambique: results from a province-wide survey. *BMC Public Health*. 2015; 15:132. Epub 2015/04/17. doi: [10.1186/s12889-015-1388-z](https://doi.org/10.1186/s12889-015-1388-z) PMID: [25881182](https://pubmed.ncbi.nlm.nih.gov/25881182/); PubMed Central PMCID: [PMC4339241](https://pubmed.ncbi.nlm.nih.gov/PMC4339241/).
21. Melaku Z, Lamb MR, Wang C, Lulseged S, Gadisa T, Ahmed S, et al. Characteristics and outcomes of adult Ethiopian patients enrolled in HIV care and treatment: a multi-clinic observational study. *BMC Public Health*. 2015; 15:462. Epub 2015/05/03. doi: [10.1186/s12889-015-1776-4](https://doi.org/10.1186/s12889-015-1776-4) PMID: [25934178](https://pubmed.ncbi.nlm.nih.gov/25934178/); PubMed Central PMCID: [PMC4455051](https://pubmed.ncbi.nlm.nih.gov/PMC4455051/).
22. ICAP. ICAP supported clinic inventory data. New York, NY:2013.
23. Sawtooth Software Inc. SMRT. 4.18 ed. Sequim, WA:2008.
24. Hole A. Modelling heterogeneity in patients' preferences for the attributes of a general practitioner appointment. *J Health Econ*. 2008; 27(4):1078–94. doi: [10.1016/j.jhealeco.2007.11.006](https://doi.org/10.1016/j.jhealeco.2007.11.006) PMID: [18179837](https://pubmed.ncbi.nlm.nih.gov/18179837/)
25. Filmer D, Pritchett L. Estimating wealth effects without expenditure data—or tears: An application to educational enrollments in states of India. *Demography*. 2001; 38(1):115–32. PMID: [11227840](https://pubmed.ncbi.nlm.nih.gov/11227840/)
26. Kjaer T, Gyrd-Hansen D. Preference heterogeneity and choice of cardiac rehabilitation program: results from a discrete choice experiment. *Health Policy*. 2008; 85(1):124–32. PMID: [17728004](https://pubmed.ncbi.nlm.nih.gov/17728004/)
27. Revelt D, Train K. Mixed logit with repeated choices: households' choices of appliance efficiency levels. *Review of Economics and Statistics*. 1998; 80:647–57.
28. Lancsar E, Louviere J. Deleting 'irrational' responses from discrete choice experiments: a case of investigating or imposing preferences. *Health Econ*. 2006; 15:797–811. PMID: [16615039](https://pubmed.ncbi.nlm.nih.gov/16615039/)
29. Kruk M, Paczkowski M, Mbaruku G, de Pinho H, Galea S. Women's preferences for place of delivery in rural Tanzania: a population-based discrete choice experiment. *Am J Public Health*. 2009; 99(9):1666–72. PMID: [19608959](https://pubmed.ncbi.nlm.nih.gov/19608959/). doi: [10.2105/AJPH.2008.146209](https://doi.org/10.2105/AJPH.2008.146209)

30. Kruk ME, Paczkowski MM, Tegegn A, Tessema F, Hadley C, Asefa M, et al. Women's preferences for obstetric care in rural Ethiopia: a population-based discrete choice experiment in a region with low rates of facility delivery. *J Epidemiol Community Health*. 2010; 64(11):984–8. Epub 2009/10/14. jech.2009.087973 [pii] doi: [10.1136/jech.2009.087973](https://doi.org/10.1136/jech.2009.087973) PMID: [19822558](https://pubmed.ncbi.nlm.nih.gov/19822558/).
31. Freedman LP, Kruk ME. Disrespect and abuse of women in childbirth: challenging the global quality and accountability agendas. *Lancet*. 2014; 384(9948):e42–4. Epub 2014/06/27. doi: [10.1016/S0140-6736\(14\)60859-X](https://doi.org/10.1016/S0140-6736(14)60859-X) PMID: [24965825](https://pubmed.ncbi.nlm.nih.gov/24965825/).
32. Office of the US Global AIDS Coordinator. PEPFAR 3.0—Controlling the Epidemic: Delivering on the Promise of an AIDS-free Generation. Washington, DC: 2014.
33. Thomas J, Reynolds H, Bevc C, Tsegaye A. Integration opportunities for HIV and family planning services in Addis Ababa, Ethiopia: an organizational network analysis. *BMC health services research*. 2014; 14:22. D—NLM: PMC3923232 EDAT- 2014/01/21 06:00 MHDA- 2014/01/21 06:00 CRDT- 2014/01/21 06:00 PHST- 2013/07/06 [received] PHST- 2014/01/08 [accepted] PHST- 2014/01/18 [aheadof-print] AID—1472-6963-14-22 [pii] AID—10.1186/1472-6963-14-22 PST—epublish. doi: [10.1186/1472-6963-14-22](https://doi.org/10.1186/1472-6963-14-22) PMID: [24438522](https://pubmed.ncbi.nlm.nih.gov/24438522/)
34. Howard A, El-Sadr W. Integration of tuberculosis and HIV services in sub-Saharan Africa: lessons learned. *Clin Infect Dis*. 2015; 50 Suppl 3:S238–44.
35. Wiegert K, Dinh T, Mushavi A, Mugurungi O, Kilmarx P. Integration of Prevention of Mother-to-Child Transmission of HIV (PMTCT) Postpartum Services with Other HIV Care and Treatment Services within the Maternal and Child Health Setting in Zimbabwe, 2012. *PLoS One*. 2014; 9(6):e98236. D—NLM: PMC4051591 EDAT- 2014/06/11 06:00 MHDA- 2014/06/11 06:00 CRDT- 2014/06/11 06:00 PHST- 2014 [ecollection] PHST- 2013/12/18 [received] PHST- 2014/04/29 [accepted] PHST- 2014/06/10 [epublish] AID—10.1371/journal.pone.0098236 AID—PONE-D-13-51314 [pii] PST—epublish. doi: [10.1371/journal.pone.0098236](https://doi.org/10.1371/journal.pone.0098236) PMID: [24915422](https://pubmed.ncbi.nlm.nih.gov/24915422/)
36. Assefa Y, Alebachew A, Lera M, Lynen L, Wouters E, Van Damme W. Scaling up antiretroviral treatment and improving patient retention in care: lessons from Ethiopia, 2005–2013. *Global Health*. 2014; 10:43. 1744-8603-10-43 [pii] doi: [10.1186/1744-8603-10-43](https://doi.org/10.1186/1744-8603-10-43) PMID: [24886686](https://pubmed.ncbi.nlm.nih.gov/24886686/); PubMed Central PMCID: PMC4046386.
37. Anglewicz P, Chintsanya J. Disclosure of HIV status between spouses in rural Malawi. *AIDS Care*. 2011; 23(8):998–1005. doi: [10.1080/09540121.2010.542130](https://doi.org/10.1080/09540121.2010.542130) PMID: [21390889](https://pubmed.ncbi.nlm.nih.gov/21390889/); PubMed Central PMCID: PMC3371657.
38. Seid M, Wasie B, Admassu M. Disclosure of HIV positive result to a sexual partner among adult clinical service users in Kemissie district, northeast Ethiopia. *Afr J Reprod Health*. 2012; 16(1):97–104. Epub 2012/07/13. PMID: [22783673](https://pubmed.ncbi.nlm.nih.gov/22783673/).
39. Decroo T, Koole O, Remartinez D, dos Santos N, Dezembro S, Jofrisse M, et al. Four-year retention and risk factors for attrition among members of community ART groups in Tete, Mozambique. *Trop Med Int Health*. 2014; 19(5):514–21. Epub 2014/06/06. doi: [10.1111/tmi.12278](https://doi.org/10.1111/tmi.12278) PMID: [24898272](https://pubmed.ncbi.nlm.nih.gov/24898272/).
40. Haddad S, Fournier P, Machouf N, Yatara F. What does quality mean to lay people? Community perceptions of primary health care services in Guinea. *Soc Sci Med*. 1998; 47(3):381–94. PMID: [9681908](https://pubmed.ncbi.nlm.nih.gov/9681908/).
41. Avortri G, Beke A, Abekah-Nkrumah G. Predictors of satisfaction with child birth services in public hospitals in Ghana. *Int J Health Care Qual Assur*. 2011; 24(3):223–37. Epub 2011/09/24. PMID: [21938968](https://pubmed.ncbi.nlm.nih.gov/21938968/).
42. Lettenmaier C, Kraft J, Raisanen K, Serlemitsos E. HIV Communication Capacity Strengthening: A Critical Review. *J Acquir Immune Defic Syndr*. 2014; 66 Suppl 3:S300–5. doi: [10.1097/QAI.000000000000238](https://doi.org/10.1097/QAI.000000000000238) PMID: [25007200](https://pubmed.ncbi.nlm.nih.gov/25007200/)
43. Tomori C, Risher K, Limaye R, Van Lith L, Gibbs S, Smelyanskaya M, et al. A Role for Health Communication in the Continuum of HIV Care, Treatment, and Prevention. *J Acquir Immune Defic Syndr*. 2014; 66 Suppl 3:S306–10. doi: [10.1097/QAI.000000000000239](https://doi.org/10.1097/QAI.000000000000239) PMID: [25007201](https://pubmed.ncbi.nlm.nih.gov/25007201/)
44. Callaghan M, Ford N, Schneider H. A systematic review of task- shifting for HIV treatment and care in Africa. *Human resources for health*. 2010; 8:8. doi: [10.1186/1478-4491-8-8](https://doi.org/10.1186/1478-4491-8-8) PMID: [20356363](https://pubmed.ncbi.nlm.nih.gov/20356363/); PubMed Central PMCID: PMC2873343.
45. Schneider H, Blaauw D, Gilson L, Chabikuli N, Goudge J. Health Systems and Access to Antiretroviral Drugs for HIV in Southern Africa: Service Delivery and Human Resources Challenges. *Reprod Health Matters*. 2006; 14(27):12–23. doi: [10.1016/s0968-8080\(06\)27232-x](https://doi.org/10.1016/s0968-8080(06)27232-x) PMID: [16713875](https://pubmed.ncbi.nlm.nih.gov/16713875/)
46. Cape Town Statement from the Third Global Symposium on Health Systems Research. The science and practice of people-centered health systems. Cape Town2014.
47. McIntyre D, Thiede M, Birch S. Access as a policy-relevant concept in low- and middle-income countries. *Health Econ Policy Law*. 2009; 4:179–93. Epub 2009/02/04. S1744133109004836 [pii] doi: [10.1017/S1744133109004836](https://doi.org/10.1017/S1744133109004836) PMID: [19187569](https://pubmed.ncbi.nlm.nih.gov/19187569/).