

CORRECTION

Correction: Fluorinated Cannabidiol Derivatives: Enhancement of Activity in Mice Models Predictive of Anxiolytic, Antidepressant and Antipsychotic Effects

Aviva Breuer, Christeene G. Haj, Manoela V. Fogaça, Felipe V. Gomes, Nicole Rodrigues Silva, João Francisco Pedrazzi, Elaine A. Del Bel, Jaime C. Hallak, José A. Crippa, Antonio W. Zuardi, Raphael Mechoulam, Francisco S. Guimarães

The first step of Fig 2 appears erroneously in the publication as a repetition of Fig 1. The correct [Fig 2](#) should show the oxidation of cannabidiol diacetate to 9-hydroxy-cannabidiol diacetate. Please see the correct [Fig 2](#) here.




OPEN ACCESS

Citation: Breuer A, Haj CG, Fogaça MV, Gomes FV, Silva NR, Pedrazzi JF, et al. (2016) Correction: Fluorinated Cannabidiol Derivatives: Enhancement of Activity in Mice Models Predictive of Anxiolytic, Antidepressant and Antipsychotic Effects. PLoS ONE 11(8): e0162087. doi:10.1371/journal.pone.0162087

Published: August 25, 2016

Copyright: © 2016 Breuer et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Step A.



Step B




Fig 2. Synthesis of HUF-102(2). Reagents and conditions: (a) SeO₂, t-BuOOH, CH₂Cl₂, r.t; (b) DAST, CH₂Cl₂, 0°C.

doi:10.1371/journal.pone.0162087.g001

Reference

1. Breuer A, Haj CG, Fogaça MV, Gomes FV, Silva NR, Pedrazzi JF, et al. (2016) Fluorinated Cannabidiol Derivatives: Enhancement of Activity in Mice Models Predictive of Anxiolytic, Antidepressant and Antipsychotic Effects. PLoS ONE 11(7): e0158779. doi: [10.1371/journal.pone.0158779](https://doi.org/10.1371/journal.pone.0158779) PMID: [27416026](https://pubmed.ncbi.nlm.nih.gov/27416026/)