

SCIENTIFIC REPORTS



OPEN

MicroRNA393 is involved in nitrogen-promoted rice tillering through regulation of auxin signal transduction in axillary buds

Xiang Li¹, Kuaifei Xia¹, Zhen Liang², Kunling Chen², Caixia Gao² & Mingyong Zhang¹

Received: 31 May 2016
Accepted: 03 August 2016
Published: 30 August 2016

Rice tillering has an important influence on grain yield, and is promoted by nitrogen (N) fertilizer. Several genes controlling rice tillering, which are regulated by poor N supply, have been identified. However, the molecular mechanism associated with the regulation of tillering based on N supply is poorly understood. Here, we report that rice microRNA393 (*OsmiR393*) is involved in N-mediated tillering by decreasing auxin signal sensitivity in axillary buds. Expression analysis showed that N fertilizer causes up-regulation of *OsmiR393*, but down-regulation of two target genes (*OsAFB2* and *OsTB1*). *In situ* expression analysis showed that *OsmiR393* is highly expressed in the lateral axillary meristem. *OsmiR393* overexpression mimicked N-mediated tillering in wild type Zhonghua 11 (ZH11). Mutation of *OsMIR393* in ZH11 repressed N-promoted tillering, which simulated the effects of limited N, and this could not be restored by supplying N fertilizer. Western blot analysis showed that *OslAA6* was accumulated in both *OsmiR393*-overexpressing lines and N-treated wild type rice, but was reduced in the *OsMIR393* mutant. Therefore, we deduced that N-induced *OsmiR393* accumulation reduces the expression of *OsTIR1* and *OsAFB2*, which alleviates sensitivity to auxin in the axillary buds and stabilizes *OslAA6*, thereby promoting rice tillering.

Rice tillering (branching) is an important agronomic trait, as the number of tillers per plant determines the panicle number, which is a key factor of the rice grain yield^{1,2}. The rice tiller represents a form of shoot branching that only exists in monocotyledonous plants at the jointing stage. Unlike in *Arabidopsis*, the dominance of the apical meristem in rice is weak, so tillering occurs during the vegetative stage upon inhibition by the apical meristem or in previously activated buds, which is not sufficiently strong. However, when the panicles of the main stems begin to differentiate, the buds formed at the elongated upper internodes become dormant³. The dormant bud is activated and later transformed into an activated, growing tiller. This process is complicated and intricate. Rice tillering is mediated by the interplay of the environment and endogenous signals such as phytohormones. It is already known that auxin, cytokinin (CK), and strigolactones (SLs) are specific regulators involved in bud outgrowth regulation⁴. Auxin and SLs inhibit bud outgrowth, whereas CK activates and promotes this process^{5,6}. Nitrogen (N) can considerably increase CK levels in rice tiller buds and nodes, and can augment IAA (early auxin responsive genes) in rice nodes⁷. The tiller number per plant is strongly affected by N fertilizer availability. High N fertilizer allows rice to produce more tillers⁷, and the *Arabidopsis* root system architecture is mediated by N availability⁸.

More is known of the mechanism by which limited N reduces branching than how abundant N promotes branching. In *Arabidopsis*, nitrate limitation reduces shoot branching by both inhibiting bud initiation and weakening the basipetal sequence of bud initiation that is caused by flowering⁹. Auxin was the first phytohormone to be identified as an important regulator of plant branching. Bud outgrowth is strongly inhibited by the apex of a whole plant. If the apex is removed, previously inactivated axillary buds become active and plant branching is initiated¹⁰. However, auxin cannot enter axillary meristem cells and therefore, its inhibitory functions are deemed to be indirect¹¹. Auxin suppresses bud development through at least two different processes. First, auxin

¹Key Laboratory of South China Agricultural Plant Molecular Analysis and Genetic Improvement & Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China. ²State Key Laboratory of Plant Cell and Chromosome Engineering, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China. Correspondence and requests for materials should be addressed to M.Z. (email: zhangmy@scbg.ac.cn)

is synthesized in the plant apex and is transported basipetally towards the root, which is known as polar auxin transport (PAT), and this inhibits bud outgrowth¹². Because auxin cannot enter the axillary meristem cells, the second mechanism is regarded as the regulation of other signalling molecules (CK and SLs) by auxin; these proteins can enter axillary meristem cells and regulate axillary bud initiation and outgrowth^{13,14}. The PAT model suggests that polar auxin transport and high auxin concentrations inhibit axillary bud development^{14,15}. The second messenger model suggests that auxin regulates the distribution and activity of CKs and SLs in the axillary meristem to control plant branching. Both of these models are supported by recent studies¹⁶.

Several genes have been identified as regulators that control rice tillering and branching. *MOC1*¹⁷ controls initiation and outgrowth of axillary meristems at both the vegetative and reproductive stages. *LAX1* is a regulator that controls axillary meristem initiation and/or maintenance during rice reproductive development¹⁸. *OsMADS57* was reported to interact with *OsTBI1*, and targets *D14* (Dwarf 14) to control the outgrowth of axillary buds¹⁹. *OsmiR156* targets *OsSPL14* and mediates its degradation to control rice tillering²⁰. In *Arabidopsis*, auxin and SL signalling are required to coordinate shoot branching with respect to N supply⁹.

MicroRNAs (miRNAs) are a class of oligonucleotides, 20 to 24 nucleotides long, and are endogenous small RNAs that are involved in post-transcriptional gene regulation in multicellular organisms, through influencing both the stability and translation of mRNAs²¹. miRNAs have important regulatory roles in the uptake, assimilation, and translocation of nutrients in plants²². miR395 and miR399 can regulate the distribution and homeostasis of sulphur and phosphate, respectively, in *Arabidopsis* and rice^{23,24}. miR169 has specific roles in the plant's response to N deficiency²⁵. miR167 and miR393 were reported to regulate NO₃⁻ signalling during lateral root development^{8,26}. *MIR393* is conserved among different plant species²⁷, and recent studies revealed that it has multiple functions in plant growth and development, such as controlling root architecture⁸, regulation of leaf development²⁸, antibacterial resistance to pathogen attack²⁷, tolerance to stress²⁹, and maintenance of normal plant growth³⁰. In *Arabidopsis*, miR393 is encoded by two distinct loci: *MIR393a* and *MIR393b*. In aerial tissues such as leaves, miR393 is mainly transcribed from *MIR393b*, suggesting distinct roles for *MIR393a* and *MIR393b*²⁸. Moreover, miR393-guided target cleavage processes generate secondary small interfering RNAs (siRNAs) from miR393 target transcripts. Feedback regulates the targeting of neo-generated siRNA, guiding the cleavage of miR393 targets²⁸. Similar to that in *Arabidopsis*, the rice *OsmiR393* family is encoded by two loci, *OsMIR393a* and *OsMIR393b*^{31,32}. Rice *OsMIR393a* and *OsMIR393b* also showed different expression patterns, which suggests that conserved mechanisms were adopted in monocots and eudicots during plant development³². To date, the exact mechanism of rice tillering regulation by *OsmiR393* has not been reported.

miR393 was shown to target auxin (IAA) receptor genes *TIR1* and *AFB* in different plants, including *Arabidopsis* and rice^{31,33,34}. *TIR1* and *AFBs* encode F-box proteins, which combine with three other proteins, ASK1, CUL1, and RBX, for assembly into the ubiquitin degradative complex (SCF^{TIR1}) to degrade specific substrates during auxin signaling^{35–37}. AUX/IAAs represent a class of proteins that represses auxin signalling. AUX/IAA proteins can bind to and repress ARFs (auxin response factor) to activate downstream auxin-responsive genes. *TIR1* and *AFBs* recognize and bind IAA and degrade AUX/IAA proteins via the SCF^{TIR1} complex³⁷ to ensure correct auxin signalling.

Here, we present findings showing that N-induced rice tillering is caused by attenuating the sensitivity of tiller buds to auxin through microRNA *OsmiR393*-mediated auxin signal transduction.

Results

Nitrogen supply promotes *OsmiR393* accumulation in rice. In *Arabidopsis*, miR393, which targets *AFB3*, is a unique nitrogen (N) responsive module that mediates root system architecture in response to external and internal N availability⁸. To investigate whether *OsmiR393* expression also responds to exogenous N fertilizer in rice, we analysed the accumulation of *OsmiR393* with various levels of NH₄NO₃ fertilizer as the N source. The level of NH₄NO₃ (1.43 mM) in the IRRI nutrient solution³⁸ was set to the normal level (1 N) of N fertilizer. *OsmiR393* accumulation increased following exposure to elevating levels of N fertilizer, from 0 mM (0 N) to 5.72 mM (4 N) NH₄NO₃, as assessed by a quantitative reverse transcription polymerase chain reaction (qRT-PCR) (Fig. 1A) and by a small RNA gel blot (Supplemental Fig. S1). Under conditions of 1 N and 4 N NH₄NO₃, *OsmiR393* expression increased by 1.4- and 3.5-fold, respectively, compared to that of the 0 N condition (Fig. 1A). These results showed that *OsmiR393* accumulation in rice is induced by high N fertilizer levels, similar to that observed in *Arabidopsis*⁸.

Overexpression of *OsmiR393* mimics N-promoted rice tillering. High N fertilizer can promote tillering in rice, and we found that *OsmiR393*-overexpressing rice produced more tillers than wild type rice with normal levels of N fertilizer³⁹. Given that high N fertilizer promotes *OsmiR393* accumulation (Fig. 1A, Supplemental Fig. S1), we hypothesized that *OsmiR393* is involved in mediating N-mediated rice tillering. To test this hypothesis, we first measured the tiller number in wild type *japonica* rice cultivar Zhonghua 11 (ZH11) with different levels of N fertilizer (Fig. 1B,C). Tiller numbers in ZH11 were increased with elevating levels of NH₄NO₃. When grown under low N conditions, specifically, 0 and 0.18 mM NH₄NO₃ (0 N and 1/8 N, respectively), ZH11 produced 2.6 tillers on average. At normal N levels (1.43 mM NH₄NO₃; 1 N), it produced 5.5 tillers, which is 2-fold greater than that of the 0 N condition. The tiller number reached approximately 10 when plants were cultivated with high N (5.72 mM NH₄NO₃; 4 N), which is 4-fold greater than that of the 0 N condition. Furthermore, tillering was analysed in three *OsmiR393*-overexpressing rice lines³⁹, relative to N content. A small RNA gel blot assay confirmed that *OsmiR393* accumulated in the three lines (*OX393-6*, *OX393-10*, and *OX393-31*) compared to expression in wild type ZH11, under normal N conditions (1 N) (Fig. 1D). When grown under normal N (1 N), all three *OsmiR393*-overexpressing lines produced more tillers than ZH11 (Fig. 1E,F). At 1 N, tiller numbers in the three *OsmiR393*-overexpressing lines reached approximately 8.3, which was about 1.5-fold higher than that observed in ZH11. These results demonstrate that tillering with *OsmiR393* overexpression at normal N levels




Figure 1. Nitrogen fertilizer triggers *OsmiR393* accumulation and overexpression resulting in an increased number of tillers. (A and D), qRT-PCR (A) and small RNA gel blot (D) analysis of *OsmiR393* expression level in ZH11 with elevating NH_4NO_3 levels (A) and in *OsmiR393*-overexpressing rice lines (OX393) under normal nitrogen (N) levels (1 N) (D). The level of NH_4NO_3 (1.43 mM) in the IRR1 nutrient solution³⁸ was set as the normal (1 N) level of N fertilizer; other N levels represent multiples of the 1 N level. Vertical bars in (A) indicate standard error from three individual repeats. Rice *U6* was used as an internal control. The asterisk in (A) indicates a mean fold change of greater than 2 vs the 0 N condition. (B,C) and (E,F), Tiller numbers of ZH11 with elevating NH_4NO_3 levels (B,C) and in *OsmiR393*-overexpressing rice plants with a normal N concentration (1 N) (E,F) respectively, when plants were grown in a pot. Scale bar = 2 cm in (B) and (E). Tiller numbers in (C) and (F) were counted and analysed in excel. Vertical bars in (C) and (F) indicate standard error ($n \geq 15$ plants for every sample and each experiment was repeated three times). The asterisk indicates significant differences (* $P \leq 0.05$ and ** $P \leq 0.01$) compared to ZH11 at 0 N (C), and at 1 N (F) as determined by a *t*-test.

mimics that of ZH11 at higher N levels. Third, to investigate whether *OsmiR393* overexpression at different N levels increases tillers as in ZH11, we treated OX393-6 with a gradient of N levels (Supplemental Fig. S2A,B). Although the tiller number in OX393-6 increased with elevating N levels, it did not reach the maximum observed with ZH11 (Fig. 1B,C). With 4 N conditions, the tiller number reached 7.4 in ZH11, but averaged only 5 in OX393-6. Moreover, at each N level (except 4 N), OX393-6 produced more tillers than ZH11. However, OX393-6 and ZH11 reached approximately the same tiller numbers when they were grown at 4 N. At the 1/8 N level, OX393-6 produced on average 5.2 tillers, whereas ZH11 produced 3.8 tillers. At the 1 N level, OX393-6 produced on an average 6.8 tillers and ZH11 produced 4.4 tillers. At the 4 N level, the tiller number for both OX393-6 and ZH11 reached approximately 9.5. In summary, our data suggests that *OsmiR393* overexpression is sufficient to increase tillering at almost all N concentrations, indicating that it is involved in N-promoted tillering. However, *OsmiR393* overexpression cannot fully mimic N-induced tillering at 4 N, suggesting that other factors/signalling pathways are involved in this process.

N fertilizer cannot restore the reduced tiller phenotype of an *OsmiR393* mutant. To confirm that *OsmiR393* is involved in N-mediated rice tillering, *OsmiR393* mutants were generated using a clustered regularly interspaced short palindromic repeats (CRISPR) approach^{40,41}. An sgRNA sequence was designed to target the mature *OsmiR393* sequence and ultimately disrupt this gene. Next, the sgRNA was linked to the sgRNA-Cas9 vector⁴² and transformed into ZH11. Three knock-out *OsmiR393* mutant lines were selected and designated as *CRP-9*, *CRP-15*, and *CRP-18*. A small RNA gel blot assay showed that *OsmiR393* accumulation was significantly decreased (Fig. 2A). To inspect the tillering phenotype in these *OsmiR393* mutants, we grew them in a controlled paddy that was managed conventionally. Contrary to *OsmiR393* overexpressing lines, which produced more tillers (Fig. 1E,F), *OsmiR393* mutants produced fewer tillers than ZH11 (Fig. 2B,C). ZH11 produced on




Figure 2. *OsMIR393* mutants develop fewer tillers (A–C) under 1 N condition and external application (4N) of NH_4NO_3 does not restore tillers in *OsMIR393* mutants (D–F). (A and D), Small RNA gel blot analysis of *OsmiR393* expression in *OsMIR393* mutants under normal nitrogen (N) levels (1 N) (A) and with 4 N levels of N fertilizer (D). (B and C) Tiller numbers for the *OsMIR393* mutant grown in a controlled field and supplied with 1 N fertilizer (NH_4NO_3). (E and F) Tiller numbers of *OsMIR393* mutants grown in a controlled field and supplied with 4 N fertilizer (NH_4NO_3). Vertical bars in (C) and (F) indicate standard error ($n \geq 15$ plants for every sample and each experiment was repeated three times). The asterisk indicates significant differences (* $P \leq 0.05$ and ** $P \leq 0.01$) compared to the ZH11 control, as assessed by a *t*-test. Scale bar = 3 cm in (B) and (E).

average 12.2 tillers, whereas all *OsMIR393* mutants (*CRP-9*, *CRP-15*, and *CRP-18*) produced approximately 6–7 tillers, approximately half that of ZH11. Combining these results, we concluded that *OsmiR393* regulates rice tillering. To investigate whether *OsmiR393* is associated with N fertilizer in rice tillering, and if tillering could be restored in *OsMIR393* mutants by N fertilizer, we grew *OsMIR393* mutants at a 4 N fertilizer level (Fig. 2E,F). *OsMIR393* mutants were planted in a controlled paddy with the aforementioned fertilizer level. A high N level (4 N) did not restore *OsmiR393* accumulation in *OsMIR393* mutants (Fig. 2D), and the tiller number in ZH11 was still much higher than that in *OsMIR393* mutants (Fig. 2E,F). This indicated that N fertilizer and *OsmiR393* have overlapping functions in rice tillering. However, tiller numbers in *OsMIR393* mutants grown at 4 N were still much higher than in those grown at 1 N. *OsMIR393* mutants produced on average 11 tillers at the 4 N level (Fig. 2F), and produced approximately 6 tillers at the 1 N fertilizer level (Fig. 2C). This implies that *OsmiR393* is required, but not sufficient, for N-mediated rice tillering. Thus, there could be some other mechanisms, distinct from *OsmiR393*, involved in rice tillering in the presence of N.

Auxin signal transduction, mediated by *OsmiR393*, is involved in N-mediated rice tillering.

Two auxin receptors *OsAFB2* and *OsTIR1* have been verified as target genes of *OsmiR393*, and seven genes were predicted to be candidate *OsmiR393* targets in rice^{32,39}. To better understand the actual targets of N fertilizer involved in rice tillering, plant materials were collected comprising regions between the shoot and root containing the SAM (shoot apical meristem), an area of the plant where tillers are formed, for RNA extraction and qRT-PCR detection. qRT-PCR results showed that *OsAFB2* and *OsTIR1* were down-regulated in the tillering region of *OX393-6*, and dramatically up-regulated in *CRP-9*, when compared to levels in ZH11, at N levels of 1 N (Fig. 3A).

To determine whether *OsAFB2* and *OsTIR1* also decreased with elevating N levels, qRT-PCR assays were conducted in wild type ZH11 (Fig. 3B). In contrast to over-expression of *OsmiR393* with elevating N levels (Fig. 1A), *OsAFB2* and *OsTIR1* transcripts decreased (Fig. 3B). This demonstrated that the two target genes (*OsAFB2* and *OsTIR1*) and *OsmiR393* are reciprocally expressed with elevating N levels, indicating that *OsAFB2* and *OsTIR1* are involved in the response to N supply in rice. However, the expression of another putative target gene (*LOC_Os03g52320*) was up-regulated in response to high N fertilizer levels in ZH11, suggesting that it was not a target




Figure 3. Inverse expression of OsmiR393 and its target genes (*OsAFB2* and *OsTIR1*) with elevating nitrogen (N) levels. (A) qRT-PCR verification of OsmiR393 targets in OsmiR393 overexpressing line (*OX393-6*), and *OsMIR393* mutant (*CRP-9*) with 1 N fertilizer. (B) qRT-PCR expression analysis of putative OsmiR393 target genes in wild type rice ZH11 with elevating N levels. Results are presented as the means of three experiments. Vertical bars indicate standard error. The asterisk mark indicates a mean fold change greater than 2 or less than 0.5 between ZH11 and the mutants (A), or the elevating N treated ZH11 (B).

of OsmiR393. LOC_Os03g52320 was predicted to encode a GRF1-interacting factor involved in various aspects of tissue differentiation and organ development^{43,44}. We speculated that its expression was enhanced, either directly or indirectly, by N fertilizer to control rice seedling growth. One putative target, LOC_Os10g39790 was not amplified by PCR, indicating that it was not expressed in the region that we collected.

During auxin signal transduction, TIR1 and AFB2 perceive the auxin signal and degrade repressors of auxin signalling via the SCF (SKP1-CULLIN/cdc53-F-box) complex mediated by the 26S proteasome⁴⁵. OsmiR393 overexpression and high levels of N, with OsmiR393 accumulation, might desensitize rice plants to exogenous auxin. In contrast, *OsMIR393* mutants should become hypersensitive under these conditions. To test this hypothesis, we treated ZH11, *OX393-6*, and *CRP-9* with NAA and observed their responses. The application of 5 mg L⁻¹ NAA typically inhibited the outgrowth of rice tiller buds (Supplemental Fig. S3A, ZH11, and ZH11 mock). However, *OX393-6* and 4 N-treated ZH11 plants showed obvious resistance. Tiller buds in *CRP-9* were dramatically inhibited, exhibiting hypersensitivity (Supplemental Fig. S3A). This result further confirmed N-mediated rice tillering resulting from attenuated auxin signalling through OsmiR393-mediated cleavage of *OsTIR1/OsAFB2*. The callus induction rate is an important index reflecting auxin/cytokinin cross-interactions⁴⁶. To further substantiate our hypothesis, ZH11, *OX393-6*, and *CRP-9* plants were induced to form calluses, and their callus induction rates were compared (Supplemental Fig. S3B). Calluses were induced in ZH11 at a faster rate than in OsmiR393-overexpressing lines, but at a slower rate than in the *OsMIR393* mutants. Moreover, adventitious root development is a typical process involved in auxin signaling^{32,47}. The OsmiR393-overexpressing line, *OX393-6*, showed a significant reduction in adventitious roots, whereas that in *CRP-9* increased (Supplemental Fig. S3C,D). The above results implied that auxin signalling interferes with adventitious root formation in OsmiR393-overexpressing lines, but is attenuated in *OsMIR393* mutants.

N fertilizer supply and OsmiR393 overexpression promote axillary bud outgrowth. To further investigate the relationship between OsmiR393 overexpression and N-mediated rice tillering, we explored the axillary bud formation process in ZH11 in response to different N levels and in the OsmiR393-overexpressing line, *OX393-6*, by tissue section (Fig. 4A–E; Supplemental Table S2). After 12 d of cultivation with different levels of N fertilizer, the axillary meristem numbers in ZH11 were inspected. No axillary meristems were observed when ZH11 was grown without N fertilizer (0 N, Fig. 4A), one was observed when ZH11 was grown at 1 N (Fig. 5B), and two were found with 4 N fertilizer (Fig. 4C). With an excessive N level (8 N), only one axillary meristem was observed, while the other had already formed an axillary bud (Fig. 4D). In contrast, the axillary meristem number for *OX393-6* grown in 1 N fertilizer was 2 (Fig. 4E), which resembled that of ZH11 with 4 N fertilizer (Fig. 4C). This suggests that OsmiR393 overexpression has the same effect on axillary bud growth as growth with 4 N fertilizer, in ZH11 plants, and that both N fertilizer supply and OsmiR393 expression promote outgrowth of the axillary bud. OsmiR393 expression in rice organs was also analysed by qRT-PCR (Fig. 4G). The results showed that OsmiR393 was expressed in all rice organs, and at a higher level in the roots and booting panicle, indicating that OsmiR393 might play a role in booting panicle development and lateral root growth. To investigate whether OsmiR393 is expressed in the axillary meristem, *in situ* RNA hybridization experiments were performed (Fig. 4F). The results showed that OsmiR393 was strongly expressed in the lateral axillary meristem




Figure 4. OsmiR393 affects axillary bud outgrowth and is expressed at a high level in the lateral axillary meristem. (A–D) Tissue section of wild type ZH11 rice grown with different nitrogen (N) levels using an NH_4NO_3 fertilizer. (E) Tissue section of OsmiR393-overexpressing line (OX393-6) under normal N levels (1 N) using NH_4NO_3 fertilizer. Red arrows in (A–E) indicate the formed axillary meristem. Pictures are representative of sections from 20 plants. (F) *In situ* hybridization of OsmiR393 in the ZH11 stem base with 1 N NH_4NO_3 fertilizer. The red arrow in (F) indicates intense OsmiR393 expression. Bar = 100 μm in (A–F). (G) qRT-PCR detection of OsmiR393 levels in different organs of ZH11 under normal N levels (1 N) using NH_4NO_3 fertilizer. Vertical bars indicate standard error from three repeats. 1L, the first leaf; 2L, the second leaf; 3L, the third leaf; FL, flag leaf; R, root; C, culm; BP, booting panicle; SS, shoot sheath.

(Fig. 4F). These results demonstrate that N fertilizer and OsmiR393 overexpression promote outgrowth of axillary buds, and that OsmiR393 is involved in axillary bud formation.

N fertilizer facilitates rice tillering through preventing the degradation of OsIAA6. TIR1 and AFB2 are two members of the SCF complex, which degrades auxin signalling repressors, to ensure correct auxin signaling^{45,48}. To determine which proteins are degraded by the SCF^{TIR1} complex, a construct with cMYC fused to OsTIR1 was generated and transformed into ZH11. Subsequently, a co-immunoprecipitation assay was employed and the precipitated proteins were analysed by mass spectrometry. We found that levels of OsIAA6, a protein belonging to the AUX/IAA class of auxin signalling repressors, changed tremendously between 35S:cMYC-TIR1 and OX393-6 plants. OsIAA6 was reported to enhance rice drought tolerance and regulate tiller outgrowth⁴⁹. Furthermore, AUX/IAA proteins are a substrate of the SCF^{TIR1} complex⁵⁰. We suspected that OsmiR393 might block degradation of OsIAA6 by the OsTIR1 and OsAFB2-mediated SCF^{TIR1} complex to control rice tillering. To investigate whether OsIAA6 expression changed following altered OsmiR393 expression, or in N-treated ZH11, we generated a construct containing an OsIAA6 full-length protein fused to a MYC epitope tag. This construct was transformed into protoplasts of OsmiR393-overexpressing rice plants, *OsMIR393* mutants, and N-treated ZH11. Western blot analysis showed that OsIAA6 was remarkably increased in the three OsmiR393 overexpressing lines (Fig. 5A), and in ZH11 with elevating N levels (Fig. 5B), but was reduced in *OsMIR393* mutants (Fig. 5C). These results suggest that OsmiR393 might affect auxin signal transduction through regulation of OsIAA6 accumulation.

Discussion

Availability of N fertilizer strongly affects rice tillering⁷, which is a major determinant of grain yield. In the past decades, several key regulators of rice tillering have been cloned and characterized, including *MOC1*¹⁷, *TAD1*⁵¹, *LAX1* and *SPA*¹⁸, *OsTBI*⁵², *OsmiR156*²⁰, and genes in the strigolactone signalling pathway^{53–56}. However, the mechanisms through which N fertilizer contributes to rice tiller formation remains elusive. In this study, we revealed a novel role for OsmiR393 in affecting rice tillering in response to N fertilizer. Our results show that adding N fertilizer increases the tiller number in rice (Fig. 1B). This is achieved by enhancing the initiation and activation of tiller bud outgrowth (Fig. 4A–D). High N fertilizer causes OsmiR393 accumulation in tiller bud (Figs 1A and 4F; Supplemental Fig. S1). OsmiR393 accumulation then decreases the transduction of auxin signalling (Figs 3 and 5), resulting in decreased sensitivity to auxin in the tiller buds (Supplemental Fig. S3).




Figure 5. OsIAA6 protein level in ZH11 plants with different nitrogen (N) levels and in different transgenic rice with 1 N levels. (A–C) Western blot analysis of OsIAA6 in the protoplast of OsmiR393-overexpressing lines (OX393) (A) N-treated ZH11 (B) and *OsMIR393* mutants (CRP) (C). Protoplasts were prepared from 2-week old seedlings. HSC70 was used as a loading control.

Consequently, dampening the response to auxin signalling in the tillers with high N fertilizer might promote initiation and outgrowth of the tiller bud.

OsmiR393 responses to N fertilizer. *Arabidopsis* miR393 is induced by nitrate, and the nitrate-responsive miR393/AFB3 regulatory module mediates root system architecture⁸. In this study, the addition of N fertilizer also caused the accumulation of OsmiR393 in rice seedlings (Fig. 1A; Supplemental Fig. S1). This result suggests that in plants, a similar mechanism has been adopted in monocots and dicots, to respond to environmental changes in N levels. Over time, expression of its target genes (*OsTIR1* and *OsAFB2*) was repressed (Fig. 3B). This suggests that OsmiR393 is also involved in the response to exogenous N in rice. However, unlike *Arabidopsis* miR393, wherein expression induced by NO_3^- was specifically localized to the root⁸, the expression of OsmiR393 in rice was induced by NH_4NO_3 in the leaf and root tissue (Figs 1A and 4F). This discrepancy between rice and *Arabidopsis* might be due to different N treatments, or it might imply that there are functional differences for miR393 between dicots and monocots.

OsmiR393 affects rice tillering. miR393 is a plant-conserved miRNA that participates in many processes during plant development. Recent studies have demonstrated that miR393 plays a role in the establishment of root system architecture in response to nitrate⁸, the regulation of leaf development²⁸, and auxin signaling⁵⁷, and is involved in antibacterial resistance in response to pathogens²⁷. In this study, overexpression of OsmiR393 resulted in an increased number of tillers, compared to that in wild type plants (Fig. 1), which was similar to previous findings³⁹. In contrast, an *OsMIR393* mutant produced fewer tillers than wild type rice (Fig. 2). Furthermore, supplementation with N did not rescue this defect in *OsMIR393* mutants. These results suggest that *OsMIR393* is a regulator that participates in modulating rice tillering.

In addition, OsmiR393 altered tillering in response to N Fertilizer. OsmiR393 was induced by NH_4NO_3 treatment in rice (Fig. 1A), and was highly expressed in the bud meristem (Fig. 4F). When OsmiR393 was overexpressed, the increase in tiller numbers induced by N fertilizer (Supplemental Fig. 2) was not as high as that observed in wild type rice (Fig. 1B,C). However, with the same N level, OsmiR393-overexpressing rice produced more tillers than wild type rice (Fig. 1E,B compared to Supplemental Fig. S2). In contrast, the mutation of *OsMIR393* resulted in fewer tillers than that in wild type with high and normal N fertilizer levels (Fig. 2). These results suggest that *OsMIR393* is involved in the regulation of rice tillering in response to N fertilizer.

Rice varieties exhibit high diversity with respect to N response. According to the degree of responsiveness, rice varieties were classified into two groups, less-responsive varieties (such as, ZH11, QZL2, and Balila) and highly-responsive varieties (such as Minghui 63 and Nanjing 6)⁵⁸. Highly-responsive rice varieties tend to produce more tillers than less-responsive varieties under the same level of N fertilizer⁵⁸. *OsmiR393* expression was compared (Supplemental Fig. S4) in these two groups. However, the expression of *OsmiR393* was not obviously different. Different *DEP1* alleles confer variable N responses among less responsive and highly responsive rice cultivars, and *DEP1* has been the subject of artificial selection during *Oryza sativa* spp. *japonica* rice domestication⁵⁸. This result might imply that *OsmiR393* is a general regulator of rice tillering that responds to N fertilizer, but is not involved in the N-response variability observed among different rice varieties.

***OsmiR393* affects auxin signal transduction in response to N fertilizer treatment.** Plants synthesize auxin in the growing apex, wherein most of their energy is expended⁵⁹. Auxin flows from the stem towards the root, basipetally, a phenomenon known as polar auxin transport (PAT). This PAT inhibits axillary bud growth in terms of apical dominance⁶⁰. It is well known that plants establish their architecture and apical dominance through the effects of PAT.

In this study, we revealed that N-fertilizer promotes *OsmiR393* accumulation (Fig. 1A) and decrease expression of *OsTIR1* and *OsAFB2* (Fig. 3B), and increases tiller number (Fig. 1B) in wild type ZH11 rice. This N-promoted tillering is mimicked by *OsmiR393* overexpression (Fig. 1E). Moreover, high N-treatment in wild type ZH11 and overexpression of *OsmiR393* resulted in a decrease in *OsmiR393*-target gene (*OsTIR1* and *OsAFB2*) expression (Fig. 3). The reduced expression of *OsTIR1* and *OsAFB2* might attenuate auxin perception and signalling, and alleviate the effects of apical dominance. TIR1 and AFB2 are two important substrate recognition subunits of the well-known SKP1-CULLIN-F-box (SCF) ubiquitin ligase complex, which functions in auxin perception to promote the 26S proteasome-mediated degradation of Aux/IAA transcriptional repressors⁴⁷. In addition, the response of *MIR393* to environmental signals is conserved across plant species, through the auxin signalling pathway^{61,62}. It has been shown that auxin cannot enter axillary meristem cells¹¹; however, auxin from the top buds could inhibit export of auxin that is synthesized in the axillary buds^{11,12}. This would increase the auxin concentration in the axillary buds, and inhibit bud outgrowth. High *OsmiR393* accumulation after N treatment (Fig. 1A) might decrease sensitivity in the axillary buds to high levels of auxin.

We proved that *OsmiR393* overexpression and N treatment prevents degradation of *OsIAA6* (Fig. 5), which is an important regulator of axillary bud formation^{49,63}. This strongly supports the hypothesis that *OsmiR393* influences axillary bud formation following N fertilizer treatment through auxin signal transduction. Auxin signalling components have been conserved throughout land plant evolution, and have evolved to control specific developmental processes³². Plant genomes encode large numbers of F-box proteins (FBPs), and there are approximately 700 FBPs in *Arabidopsis*⁶⁴. The role of *miR393*/*TIR1*/*AFBs* in the plant's response to auxin represents a ubiquitous model for adaption and acclimation. Therefore, we concluded that N fertilizer promotes *OsmiR393* accumulation, which interferes with auxin signalling, finally triggering tiller production in rice.

N fertilizer can promote cytokinin biosynthesis/signalling and stimulates rice to produce more tillers^{65,66}. There is evidence to suggest that exogenous application of N fertilizer can promote cytokinin biosynthesis⁶⁷ and stimulate lateral bud outgrowth⁶⁸. In *Arabidopsis*, the expression of *AtIPT3* and *AtIPT5* is in response to N^{69,70}; these genes are key determinants of CK biosynthesis that respond to rapid changes in NO₃⁻ availability. Our present results showed that *OsmiR393* also responds to N fertilizer, interferes with auxin signalling, and attenuates apical dominance, ultimately leading to tiller production in rice. We did not analyse whether these two N-mediated tillering pathways overlap. Thus, the effect of N fertilizer on rice tillering might be through these two pathways, or in parallel, or through other undiscovered pathways. This also explains why exogenous N-treated *OsMIR393* mutants still produced more tillers than wild type plants in normal N conditions (Fig. 2B,C,E,F).

Materials and Methods

Plant materials and growth conditions. *Oryza sativa japonica* Zhonghua11 (ZH11) was used as the wild type and the source for transgenic plants. Plant materials used in this study were ZH11, three *OsmiR393* overexpressing lines (*OX393-6*, *OX393-10*, and *OX393-31*)³⁹, three *OsMIR393* knock-out lines (*CRP-9*, *CRP-15*, and *CRP-18*), and the distinct N responsive varieties (QZL2, Balila, Minghui 63, and Nanjing 6)⁵⁸. Seeds were dried at 37 °C for 24 h before germination at 25 °C, and were supplemented with water in the dark for 48 h. For phenotypic observations, germinated seeds were grown under controlled field conditions or in boxes filled with sand supplemented with hydroponic medium. For nutrient treatments, plants were grown in hydroponic cultures using the International Rice Research Institute (IRRI) liquid culture medium recipe (IRRI nutrient solution)³⁸. Except for modified NH₄NO₃, all nutrients were kept at the same concentration in all cultures. The NH₄NO₃ content was based on that of the IRRI nutrient solution, which was designated as the normal level; other N fertilizer levels were based on this concentration. For example, NH₄NO₃ content in 4-fold N fertilizer was quadruple that of the normal level. For paddy culture, seeds were germinated and sown on soil and then conventionally managed.

Vector construction for rice transformation. To generate the *cMYC-OsTIR1* construct, the coding region of *OsTIR1* was amplified by PCR, cloned into the pEASY-Blunt vector (Transgene, China), and subcloned into the pE3n vector⁷¹. The *cMYC-OsTIR1* fragment was then inserted into the pCAMBIA2300 vector downstream of the maize (*Zea mays*) *Ubiquitin* promoter after *KpnI* and *BamHI* digestion. To generate the *cMYC-OsIAA6* construct, the coding region of *OsIAA6* was amplified by PCR, cloned into the pEASY-Blunt vector (Transgene, China), and subcloned into the pE3n vector. The *cMYC-OsIAA6* fragment was then inserted into the pCAMBIA2300 vector downstream of the maize (*Zea mays*) *Ubiquitin* promoter after *BamHI* and *NotI* digestion. All of the primers used to generate the aforementioned constructs are listed in Supplemental Table S1, and all of the constructs were

confirmed by sequencing. The constructed vectors were transformed by *Agrobacterium tumefaciens* strain EHA105. Wild type ZH11 calli were used as the recipients for *Agrobacterium*-mediated transformation as described⁷².

RNA extraction and quantitative RT-PCR. Total RNA was extracted using an RNA extraction kit (Invitrogen, China) and digested with DNase I (Takara, China) according to the manufacturers' instructions. The RNA quality and integrity were analysed by agarose gel electrophoresis and the RNA concentration was determined using a biophotometer (METASH, B-500, China). cDNA was synthesized from total RNA using AMV Reverse Transcriptase (Promega, China). Small RNA was extracted using an RNAiso kit for small RNA (Takara, China) and digested with DNase I (Takara, China) according to the product manuals. Reverse transcription was performed with a cDNA Synthesis Kit (Promega, China) in combination with a stem-loop RT-PCR technique⁷³. Quantitative RT-PCR was performed on a 7500 RT-qPCR system (Applied Biosystems, USA) with SYBR Green Real-time PCR Master Mix (Toyobo, China) according to the manufacturer's instructions. Gene expression was normalized to that of rice *ACTIN1*. Primers used for qRT-PCR are presented in Supplemental Table S1.

Small RNA gel blot assay. Small RNA gel blot analysis was performed as described in Liu *et al.*⁷⁴. Briefly, total RNA was extracted from rice seedlings, and total RNA samples (approximately 20 µg) were separated on denaturing 15% polyacrylamide gels and transferred electrophoretically to Hybond-N⁺ membranes (<http://www.gelifsciences.com/>). The gel was stained with ethidium bromide before transfer to confirm equal loading. Hybridizations were performed at 42 °C in PerfectHyb Plus buffer with DNA oligonucleotide probes labelled by T4 polynucleotide kinase (New England Biolabs, <https://www.neb.com/>). Hybridization signals were detected with a phosphorimager (GE Healthcare Life Sciences, <http://www.gelifsciences.com/>). The sequences of the probes are provided in Supplemental Table S1.

Tissue sectioning and *in situ* hybridization. Seedlings of ZH11 plants were grown with different N fertilizer levels, and OX393-6 was grown with normal levels of N fertilizer for 15 d, after which SAM tissues and axillary buds were fixed and sectioned at a thickness of 7 mm. The sections were then stained with toluidine blue for light microscopic analysis (Zeiss, Germany). *In situ* hybridization was performed as described⁷⁵. The OsmiR393 probe was synthesized and labelled with digoxigenin. Shoots containing SAM and axillary meristems were used for hybridization assays. Slides were photographed under a microscope (Zeiss, Germany).

CRISPR-mediated mutation of *OsMIR393*. An sgRNA: AAGGATCAATGCGATCCCTTTGG was designed to target OsmiR393. The sgRNA was inserted into the *AarI* site of the p2300-rCas9-U3-gRNA vector, which contains a rice-codon optimized Cas9 driven by a 2 × 35S promoter⁴², and the sgRNA was activated by an *OsU3* promoter. The primer used for cloning is shown in Supplemental Table S1 (CRISPR-F and CRISPR-R). After introducing the CRISPR/Cas plasmid into rice varietal ZH11 through *Agrobacterium*-mediated transformation⁷², the T₀ generation mutants were screened with G418 (Invitrogen). All regenerated T₀ transgenic plants were genotyped using the primer (CRISPR-g-F and CRISPR-g-R) to select positive transgenic lines, and a small RNA gel blot analysis was performed to subsequently detect OsmiR393 expression and confirm the knock-out of OsmiR393 in mutated lines.

Co-IP and mass spectrum analysis. To validate the protein interaction profile of OsTIR1 *in vivo*, a Co-IP assay was employed using an immunoprecipitation kit (Sigma-Aldrich, China). The cDNA for *OsTIR1* was amplified using the primers cMYC-TIR1-F and cMYC-TIR1-R. Amplified cDNA was inserted into pRT107-6XMyc between the *BamHI* and *KpnI* sites to generate the expression vector 2 × 35S:6XMyc-OsTIR1. The co-immunoprecipitation procedure was performed in accordance with the manufacturer's instructions using an anti-c-Myc antibody (Sigma-Aldrich, China). Mass spectrum analysis was performed by the Boxin biotechnology company in Guangzhou. Primers used are listed in Supplemental Table S1.

Protoplast isolation and western blot analysis. To generate the cMYC-OsIAA6 sequence, the *OsIAA6* coding sequence was amplified with the gene specific primers, cMYC-OsIAA6-F and cMYC-OsIAA6-R containing the restriction enzyme sites *BamHI* and *NotI*, and ligated into a rice transformation vector for constitutive expression. Constructs were introduced into rice protoplasts of 2-week seedlings as previously reported⁷⁶. The generated constructs were then transfected into the isolated protoplasts using polyethyleneglycol-mediated transformation⁷⁶. To examine OsIAA6 protein expression, a western blot was performed as described in Niu *et al.*⁷⁷.

Tiller bud and callus induction with NAA treatments. Five-leaf seedlings were sprayed with 5 mg/L of NAA, and the length of tiller buds was measured as indicated. Callus induction with 2 mg/L of NAA was performed as previously described³⁹.

Tiller count and data analysis. Rice plants were planted in previously described conditions and grown for approximately 30 days before the headings and tillers were counted; all data were analysed in Excel using a *t*-test.

Sequence data from this study can be found in the GenBank/EMBL data libraries under accession numbers *OsAFB2* (LOC_Os04g32460), *OsTIR1* (LOC_Os05g05800), *OsIAA6* (LOC_Os01g53880), and *ACTIN1* (LOC_Os03g50885).

References

- Xing, Y. & Zhang, Q. Genetic and molecular bases of rice yield. *Annu. Rev. Plant Biol.* **61**, 421–442, doi: 10.1146/annurev-arplant-042809-112209 (2010).
- Liang, W. H., Shang, F., Lin, Q. T., Lou, C. & Zhang, J. Tillering and panicle branching genes in rice. *Gene* **537**, 1–5, doi: 10.1016/j.gene.2013.11.058 (2014).
- Wang, Y. & Li, J. The plant architecture of rice (*Oryza sativa*). *Plant Mol. Biol.* **59**, 75–84, doi: 10.1007/s11103-004-4038-x (2005).

4. Dun, E. A., de Saint Germain, A., Rameau, C. & Beveridge, C. A. Antagonistic action of strigolactone and cytokinin in bud outgrowth control. *Plant Physiol.* **158**, 487–498, doi: 10.1104/pp.111.186783 (2012).
5. Dun, E. A., Brewer, P. B. & Beveridge, C. A. Strigolactones: discovery of the elusive shoot branching hormone. *Trends Plant Sci.* **14**, 364–372, doi: 10.1016/j.tplants.2009.04.003 (2009).
6. Leyser, O. The control of shoot branching: an example of plant information processing. *Plant Cell Environ.* **32**, 694–703, doi: 10.1111/j.1365-3040.2009.01930.x (2009).
7. Xu, J. *et al.* The interaction between nitrogen availability and auxin, cytokinin, and strigolactone in the control of shoot branching in rice (*Oryza sativa* L.). *Plant Cell Environ.* **34**, 1647–1662, doi: 10.1007/s00299-015-1815-8 (2015).
8. Vidal, E. A. *et al.* Nitrate-responsive miR393/AFB3 regulatory module controls root system architecture in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **107**, 4477–4482, doi: 10.1073/pnas.09095711107 (2010).
9. de Jong, M. *et al.* Auxin and strigolactone signaling are required for modulation of *Arabidopsis* shoot branching by nitrogen supply. *Plant Physiol.* **166**, 384–395, doi: 10.1104/pp.114.242388 (2014).
10. Ongaro, V. & Leyser, O. Hormonal control of shoot branching. *J. Exp. Biol.* **59**, 67–74, doi: 10.1093/jxb/erm134 (2008).
11. Booker, J., Chatfield, S. & Leyser, O. Auxin acts in xylem-associated or medullary cells to mediate apical dominance. *Plant Cell* **15**, 495–507 (2003).
12. Li, C. & Bangerth, F. Auto-inhibition of indole-acetic acid transport in the shoot of two-branched pea (*Pisum sativum*) plants and its relationship to correlative dominance. *Physiol. Plant.* **106**, 415–420 (1999).
13. Bangerth, F. Response of cytokinin concentration in the xylem exudates of bean (*Phaseolus vulgaris* L.) plants to decapitation and auxin treatment, and relationship to apical dominance. *Planta* **194**, 439–442 (1994).
14. Brewer, P. B., Dun, E. A., Ferguson, B. J., Rameau, C. & Beveridge, C. A. Strigolactone acts downstream of auxin to regulate bud outgrowth in pea and *Arabidopsis*. *Plant Physiol.* **150**, 482–493, doi: 10.1104/pp.108.134783 (2009).
15. Balla, J., Kalousek, P., Reinohl, V., Friml, J. & Prochazka, S. Competitive canalization of PIN-dependent auxin flow from axillary buds controls pea bud outgrowth. *Plant J.* **65**, 571–577, doi: 10.1111/j.1365-313X.2010.04443.x (2011).
16. Domagalska, M. A. & Leyser, O. Signal integration in the control of shoot branching. *Nat. Rev. Mol. Cell Biol.* **12**, 211–221, doi: 10.1038/nrm3088 (2011).
17. Li, X. *et al.* Control of tillering in rice. *Nature* **422**, 618–621, doi: 10.1038/nature01518 (2003).
18. Komatsu, K. *et al.* LAX and SPA: major regulators of shoot branching in rice. *Proc. Natl. Acad. Sci. USA* **100**, 11765–11770, doi: 10.1073/pnas.1932414100 (2003).
19. Guo, S. *et al.* The interaction between OsMADS57 and OsTB1 modulates rice tillering via DWARF14. *Nat. Commun.* **4**, 1566, doi: 10.1038/ncomms2542 (2013).
20. Jiao, Y. *et al.* Regulation of OsSPL14 by OsmiR156 defines ideal plant architecture in rice. *Nat. Genet.* **42**, 541–544, doi: 10.1038/ng.591 (2010).
21. Rogers, K. & Chen, X. Biogenesis, turnover, and mode of action of plant microRNAs. *Plant Cell* **25**, 2383–2399, doi: 10.1105/tpc.113.113159 (2013).
22. Kuo, H. F. & Chiou, T. J. The role of microRNAs in phosphorus deficiency signaling. *Plant Physiol.* **156**, 1016–1024, doi: 10.1104/pp.111.175265 (2011).
23. Liang, G., Yang, F. & Yu, D. MicroRNA395 mediates regulation of sulfate accumulation and allocation in *Arabidopsis thaliana*. *Plant J.* **62**, 1046–1057, doi: 10.1111/j.1365-313X.2010.04216.x (2010).
24. Hu, B. *et al.* MicroRNA399 is involved in multiple nutrient starvation responses in rice. *Front. Plant Sci.* **6**, 188 (2015).
25. Zhao, M., Ding, H., Zhu, J. K., Zhang, F. & Li, W. X. Involvement of miR169 in the nitrogen-starvation responses in *Arabidopsis*. *New Phytol.* **190**, 906–915, doi: 10.1111/j.1469-8137.2011.03647.x (2011).
26. Kinoshita, N. *et al.* IAA-Ala Resistant3, an evolutionarily conserved target of miR167, mediates *Arabidopsis* root architecture changes during high osmotic stress. *Plant Cell* **24**, 3590–3602, doi: 10.1105/tpc.112.097006 (2012).
27. Navarro, L. *et al.* A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Science* **312**, 436–439, doi: 10.1126/science.1126088 (2006).
28. Si-Ammour, A. *et al.* miR393 and secondary siRNAs regulate expression of the TIR1/AFB2 auxin receptor clade and auxin-related development of *Arabidopsis* leaves. *Plant Physiol.* **157**, 683–691, doi: 10.1104/pp.111.180083 (2011).
29. Sunkar, R. & Zhu, J. K. Novel and stress-regulated microRNAs and other small RNAs from *Arabidopsis*. *Plant Cell* **16**, 2001–2019, doi: 10.1105/tpc.104.022830 (2004).
30. Chen, Z. H. *et al.* Regulation of auxin response by miR393-targeted transport inhibitor response protein 1 is involved in normal development in *Arabidopsis*. *Plant Mol. Biol.* **77**, 619–629, doi: 10.1007/s11103-011-9838-1 (2011).
31. Jones-Rhoades, M. W. & Bartel, D. P. Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Mol. Cell* **14**, 787–799, doi: 10.1016/j.molcel.2004.05.027 (2004).
32. Bian, H. *et al.* Distinctive expression patterns and roles of the miRNA393/TIR1 homolog module in regulating flag leaf inclination and primary and crown root growth in rice (*Oryza sativa*). *New Phytol.* **196**, 149–161, doi: 10.1111/j.1469-8137.2012.04248.x (2012).
33. Dharmasiri, N., Dharmasiri, S. & Estelle, M. The F-box protein TIR1 is an auxin receptor. *Nature* **435**, 441–445, doi: 10.1038/nature03543 (2005).
34. Kepinski, S. & Leyser, O. The *Arabidopsis* F-box protein TIR1 is an auxin receptor. *Nature* **435**, 446–451, doi: 10.1038/nature03542 (2005).
35. Gray, W. M., Kepinski, S., Rouse, D., Leyser, O. & Estelle, M. Auxin regulates SCF(TIR1)-dependent degradation of AUX/IAA proteins. *Nature* **414**, 271–276, doi: 10.1038/35104500 (2001).
36. Lechner, E., Achard, P., Vansiri, A., Potuschak, T. & Genschik, P. F-box proteins everywhere. *Curr. Opin. Plant Biol.* **9**, 631–638, doi: 10.1016/j.pbi.2006.09.003 (2006).
37. Calderon Villalobos, L. I. *et al.* A combinatorial TIR1/AFB-Aux/IAA co-receptor system for differential sensing of auxin. *Nat. Chem. Biol.* **8**, 477–485, doi: 10.1038/nchembio.926 (2012).
38. Yoshida, S., Forno, D. A., Cook, J. H. & Gomez, K. A. Routine procedures for growing rice plants in culture solution. In: Yoshida, S., Forno, D. A., Cook, J. H., Gomez, K. A. eds. *Laboratory Manual for Physiological Studies of Rice: International Rice Research Institute, Los Banos, Philippines*, 61–66 (1976).
39. Xia, K. *et al.* OsTIR1 and OsAFB2 downregulation via OsmiR393 overexpression leads to more tillers, early flowering and less tolerance to salt and drought in rice. *PLoS One* **7**, e30039, doi: 10.1371/journal.pone.0030039 (2012).
40. Jinek, M. *et al.* A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816–821, doi: 10.1126/science.1225829 (2012).
41. Kim, H. & Kim, J. S. A guide to genome engineering with programmable nucleases. *Nat. Rev. Genet.* **15**, 321–334, doi: 10.1038/nrg3686 (2014).
42. Shan, Q. *et al.* Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat. Biotechnol.* **31**, 686–688, doi: 10.1038/nbt.2650 (2013).
43. van der Knaap, E., Kim, J. H. & Kende, H. A novel gibberellin-induced gene from rice and its potential regulatory role in stem growth. *Plant Physiol.* **122**, 695–704 (2000).
44. Liu, J., Rice, J. H., Chen, N., Baum, T. J. & Hwezi, T. Synchronization of developmental processes and defense signaling by growth regulating transcription factors. *PLoS One* **9**, e98477, doi: 10.1371/journal.pone.0098477 (2014).

45. Wang, R. & Estelle, M. Diversity and specificity: auxin perception and signaling through the TIR1/AFB pathway. *Curr. Opin. Plant Biol.* **21**, 51–58, doi: 10.1016/j.pbi.2014.06.006 (2014).
46. Nyaboga, E. N., Njiru, J. M. & Tripathi, L. Factors influencing somatic embryogenesis, regeneration, and Agrobacterium-mediated transformation of cassava (*Manihot esculenta* Crantz) cultivar TME14. *Front. Plant Sci.* **6**, 411, doi: 10.3389/fpls.2015.00411 (2015).
47. Gutierrez, L. *et al.* Phenotypic plasticity of adventitious rooting in Arabidopsis is controlled by complex regulation of AUXIN RESPONSE FACTOR transcripts and microRNA abundance. *Plant Cell* **21**, 3119–3132, doi: 10.1105/tpc.108.064758 (2009).
48. Pan, X., Chen, J. & Yang, Z. Auxin regulation of cell polarity in plants. *Curr. Opin. Plant Biol.* **28**, 144–153, doi: 10.1016/j.pbi.2015.10.009 (2015).
49. Jung, H., Lee, D. K., Choi, Y. D. & Kim, J. K. OsIAA6, a member of the rice Aux/IAA gene family, is involved in drought tolerance and tiller outgrowth. *Plant Sci.* **236**, 304–312, doi: 10.1016/j.plantsci.2015.04.018 (2015).
50. Leyser, O. Auxin signalling: the beginning, the middle and the end. *Curr. Opin. Plant Biol.* **4**, 382–386 (2001).
51. Xu, C. *et al.* Degradation of MONOCULM 1 by APC/C (TAD1) regulates rice tillering. *Nat Commun.* **3**, 750, doi: 10.1038/ncomms1743 (2012).
52. Takeda, T. *et al.* The *OsTB1* gene negatively regulates lateral branching in rice. *Plant J.* **33**, 513–520 (2003).
53. Lin, H. *et al.* DWARF27, an iron-containing protein required for the biosynthesis of strigolactones, regulates rice tiller bud outgrowth. *Plant Cell* **21**, 1512–1525, doi: 10.1105/tpc.109.065987 (2009).
54. Lu, Z. *et al.* Genome-wide binding analysis of the transcription activator ideal plant architecture1 reveals a complex network regulating rice plant architecture. *Plant Cell* **25**, 3743–3759, doi: 10.1105/tpc.113.113639 (2013).
55. Cardoso, C. *et al.* Natural variation of rice strigolactone biosynthesis is associated with the deletion of two *MAX1* orthologs. *Proc. Natl. Acad. Sci. USA* **111**, 2379–2384, doi: 10.1073/pnas.1317360111 (2014).
56. Sang, D. *et al.* Strigolactones regulate rice tiller angle by attenuating shoot gravitropism through inhibiting auxin biosynthesis. *Proc. Natl. Acad. Sci. USA* **111**, 11199–11204, doi: 10.1073/pnas.1411859111 (2014).
57. Parry, G. *et al.* Complex regulation of the *TIR1/AFB* family of auxin receptors. *Proc. Natl. Acad. Sci. USA* **106**, 22540–22545, doi: 10.1073/pnas.0911967106 (2009).
58. Sun, H. *et al.* Heterotrimeric G proteins regulate nitrogen-use efficiency in rice. *Nat. Genet.* **46**, 652–656, doi: 10.1038/ng.2958 (2014).
59. Smit, M. E. & Weijers, D. The role of auxin signaling in early embryo pattern formation. *Curr. Opin. Plant Biol.* **28**, 99–105, doi: 10.1016/j.pbi.2015.10.001 (2015).
60. Dun, E. A., Ferguson, B. J. & Beveridge, C. A. Apical dominance and shoot branching. divergent opinions or divergent mechanisms? *Plant Physiol.* **142**, 812–819, doi: 10.1104/pp.106.086868 (2006).
61. Windels, D. & Vazquez, F. miR393: integrator of environmental cues in auxin signaling? *Plant Signal Behav.* **6**, 1672–1675, doi: 10.4161/psb.6.11.17900 (2011).
62. Windels, D. *et al.* miR393 is required for production of proper auxin signalling outputs. *PLoS One* **9**, e95972, doi: 10.1371/journal.pone.0095972 (2014).
63. Lu, G. *et al.* *OsPIN5b* modulates rice (*Oryza sativa*) plant architecture and yield by changing auxin homeostasis, transport and distribution. *Plant J.* **83**, 913–925, doi: 10.1111/tpj.12939 (2015).
64. Yu, H. *et al.* Untethering the TIR1 auxin receptor from the SCF complex increases its stability and inhibits auxin response. *Nature Plants* **1**, doi: 10.1038/nplants.2014.30 (2015).
65. Shimizu-Sato, S., Tanaka, M. & Mori, H. Auxin–cytokinin interactions in the control of shoot branching. *Plant Mol. Biol.* **69**, 429–435, doi: 10.1007/s11103-008-9416-3 (2009).
66. Ding, C., You, J., Chen, L., Wang, S. & Ding, Y. Nitrogen fertilizer increases spikelet number per panicle by enhancing cytokinin synthesis in rice. *Plant Cell Rep.* **33**, 363–371, doi: 10.1007/s00299-013-1536-9 (2014).
67. Hirose, N. *et al.* Regulation of cytokinin biosynthesis, compartmentalization and translocation. *J. Exp. Bot.* **59**, 75–83, doi: 10.1093/jxb/erm157 (2008).
68. Barazesh, S. & McSteen, P. Hormonal control of grass inflorescence development. *Trends Plant Sci.* **13**, 656–662, doi: 10.1016/j.tplants.2008.09.007 (2008).
69. Miyawaki, K., Matsumoto–Kitano, M. & Kakimoto, T. Expression of cytokinin biosynthetic isopentenyltransferase genes in Arabidopsis: tissue specificity and regulation by auxin, cytokinin, and nitrate. *Plant J.* **37**, 128–138 (2004).
70. Takei, K. *et al.* *ATIPT3* is a key determinant of nitrate-dependent cytokinin biosynthesis in Arabidopsis. *Plant Cell Physiol.* **45**, 1053–1062, doi: 10.1093/pcp/pch119 (2004).
71. Exner, V., Hirsch–Hoffmann, M., Gruißem, W. & Hennig, L. PlantDB – a versatile database for managing plant research. *Plant Methods* **4**, 1, doi: 10.1186/1746-4811-4-1 (2008).
72. Lin Y. J. *et al.* Establishment of high efficiency *Agrobacterium*-mediated genetic transformation system of Mudanjiang. *Acta Agron. Sin.* **28**, 294–300 (2002).
73. Chen C. *et al.* Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res.* **33**, e179, doi: 10.1093/nar/gni178 (2005).
74. Liu, B. *et al.* Loss of function of *OsDCL1* affects microRNA accumulation and causes developmental defects in rice. *Plant Physiol.* **139**, 296–305, doi: 10.1104/pp.105.063420 (2005).
75. Xu, Y. Y. *et al.* Activation of the *WUS* gene induces ectopic initiation of floral meristems on mature stem surface in Arabidopsis thaliana. *Plant Mol. Biol.* **57**, 773–784, doi: 10.1007/s11103-005-0952-9 (2005).
76. Zhang, Y. *et al.* A highly efficient rice green tissue protoplast system for transient gene expression and studying light/chloroplast-related processes. *Plant Methods* **7**, 30, doi: 10.1186/1746-4811-7-30 (2011).
77. Niu, L., Zhang, Y., Pei, Y., Liu, C. & Cao, X. Redundant requirement for a pair of *PROTEIN ARGININE METHYLTRANSFERASE4* homologs for the proper regulation of Arabidopsis flowering time. *Plant Physiol.* **148**, 490–503, doi: 10.1104/pp.108.124727 (2008).

Acknowledgements

This work was supported by Guangdong Science and Technology Department of China (grant nos. 2015B020231009 and 2014A020208081), Guangdong Agriculture Department of China (grant no. 2014492), and the National Natural Science Foundation of China (grant no. 31371604/31272240). We thank Xiexiu Zhang of the South China Botanical Garden (SCBG) for technical support. We also thank Dr. Hao Chen, Ze Yun and Dr. Xun Zeng of SCBG for comments on the manuscripts.

Author Contributions

X.L. and K.X. performed most of the experiments; Z.L., K.C. and C.G. provided CRISPR technical assistance to X.L. X.L. and M.Z. designed the experiments and analyzed the data; X.L. and M.Z. conceived the project and wrote the article with contributions of all the authors.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Li, X. *et al.* MicroRNA393 is involved in nitrogen-promoted rice tillering through regulation of auxin signal transduction in axillary buds. *Sci. Rep.* **6**, 32158; doi: 10.1038/srep32158 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016